

- 3 Francium-208 is radioactive and emits α -particles with a kinetic energy of $1.07 \times 10^{-12} \text{ J}$ to form nuclei of astatine, as illustrated in Fig. 3.1.

For
Examiner's
Use




Fig. 3.1

- (a) State the nature of an α -particle.

.....
 [1]

- (b) Show that the initial speed of an α -particle after the decay of a francium nucleus is approximately $1.8 \times 10^7 \text{ m s}^{-1}$.

[2]

- (c) (i) State the principle of conservation of linear momentum.

.....

 [2]

- (ii) The Francium-208 nucleus is stationary before the decay. Estimate the speed of the astatine nucleus immediately after the decay.

For
Examiner's
Use

speed = m s^{-1} [3]

- (d) Close examination of the decay of the francium nucleus indicates that the astatine nucleus and the α -particle are not ejected exactly in opposite directions.

Suggest an explanation for this observation.

.....
.....
..... [2]

- 5 (a) Distinguish between the structure of a metal and of a polymer.

metal:

.....

.....

polymer:

.....

..... [4]

- (b) Latex is a natural form of rubber. It is a polymeric material.

- (i) Describe the properties of a sample of latex.

.....

.....

..... [2]

- (ii) The process of heating latex with a small amount of sulphur creates cross-links between molecules. Natural latex has very few cross-links between its molecules.

Suggest how this process changes the properties of latex.

.....

.....

..... [2]

- 8 The radioactive decay of nuclei is both spontaneous and random.

Explain what is meant by

- (a) *radioactive decay* of a nucleus,

.....
.....
..... [2]

- (b) *spontaneous decay*,

.....
.....
..... [2]

- (c) *random decay*.

.....
.....
..... [2]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

- 2 The Brownian motion of smoke particles in air may be observed using the apparatus shown in Fig. 2.1.

For
Examiner's
Use




Fig. 2.1

- (a)** Describe what is seen when viewing a smoke particle through the microscope.

.....

[2]

- (b)** Suggest and explain what difference, if any, would be observed in the movement of smoke particles when larger smoke particles than those observed in **(a)** are viewed through the microscope.

.....

[2]

- 8 Fig. 8.1 shows the position of Neptunium-231 ($^{231}_{93}\text{Np}$) on a diagram in which nucleon number (mass number) A is plotted against proton number (atomic number) Z .

For
Examiner's
Use




Fig. 8.1

- (a) Neptunium-231 decays by the emission of an α -particle to form protactinium. On Fig. 8.1, mark with the symbol Pa the position of the isotope of protactinium produced in this decay. [1]
- (b) Plutonium-243 ($^{243}_{94}\text{Pu}$) decays by the emission of a β -particle (an electron). On Fig. 8.1, show this decay by labelling the position of Plutonium-243 as Pu and the position of the daughter product as D. [2]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

- 7 The α -particle scattering experiment provided evidence for the existence of a nuclear atom.

For
Examiner's
Use

(a) State what could be deduced from the fact that

- (i) most α -particles were deviated through angles of less than 10° ,

.....
.....
..... [2]

- (ii) a very small proportion of the α -particles was deviated through angles greater than 90° .

.....
.....
..... [2]