

Influence of School Closure on COVID-19 Contaminations in Ireland and Repercussions across Age Groups

Edgar Morgenroth^{a,*}, Damien Dupré^{a,**}

^a*Glasnevin, Dublin 9, Ireland*

Abstract

TBD

1. Introduction

Since the onset of Covid-19, contradicting messages are circulating about the spread of the virus in schools. Sometimes presented as a safe environment (Falk et al., 2021; Walger et al., 2020), at other times designed as an aggravating factor of the pandemic (Lopez et al., 2020; Meuris et al., 2021), child care facilities and schools have been closed multiple times in most countries to protect the public health. However, the efficacy of school closure is still unknown and little is known about how Covid-19 Cases spread across age cohort and especially from the younger cohort being at school to older cohorts including parents, grand parents and relatives. Some evidence show that school is an influential environment where the virus is likely to spread. Once contaminated at school, a common spread believe is that the children and teenagers will then contaminate their parents. Therefore, among the political actions to stop the spread of Covid-19, the closure of primary and secondary schools was widely adopted worldwide. This research aims to investigate 1) the effect of school closure on the evolution of Covid-19 cases and 2) the temporal relationship between the rise of cases in children and teenagers with the rise of cases of older age group.

1.1. Influence of School Closure

One of the strongest assumption of government actions is that by reducing social contact will reduce the spread of the virus. Therefore, it is legitimate to believe that by closing schools, a reduction of the contaminations would be observed in the younger age groups. However, the efficiency of school closure on

*Corresponding Author

**Equal contribution

Email addresses: edgar.morgenroth@dcu.ie (Edgar Morgenroth), damien.dupre@dcu.ie (Damien Dupré)

the reduction of Covid-19 cases is still questioned. While some research have observed that school closures contribute to limit or to reduce the growth rate of confirmed cases after implementation (Stage et al., 2021; Sugishita, 2020), other did not observe a change in the evolution of Covid-19 cases (Chang et al., 2020; Iwata et al., 2020). For instance, a controlled comparison between similar localities in Japan with schools closed and school open did not revealed any evidence that school closures reduced the spread of Covid-19 (Fukumoto et al., 2021). If the school closure had a real impact on the evolution of confirmed Covid-19 cases, it should be possible to observe a decrease or a least an inflection in the trend of its evolution among younger age groups.

1.2. Causal Relationship Between Age Groups

A second implicit believe regarding the effect of school closure on the spread of Covid-19 is that school not only has an effect on to the spread of the virus in children and teenagers but also has a knock-on effect on the spread of the virus in older age groups also called Secondary Attack Rate (SAR). The contaminated children and teenagers would bring the virus back at home and, then, they will contaminate their parents and relatives. For example, a research investigating the contamination in the household network not only revealed an exceptional high rate of secondary contamination but also that these contamination happened when the school were closed (Soriano-Arandes et al., 2021).

Despite being reported in several clinical and epidemiological studies (Siebach et al., 2021; Zhen-Dong et al., 2020), multiple research have shown that the SAR from children to household members was, in fact, lower than expected (Heavey et al., 2020; Hoek et al., 2020; Kim et al., 2021; Ludvigsson, 2020). However, the SAR of children and teenagers to the household member is likely to be age-dependent, with difference between infants, primary and secondary school children, and college students (Gras-Le Guen et al., 2021). If a secondary transmission from children and teenagers to household member has a significant influence, then a temporal causality relationship between their evolution should be observed.

2. Method

2.1. Observations

The data were collected from the official daily publication of Covid-19 cases by the Irish Department of Health. They include the amount of daily cases for 10 age groups from March 04, 2020 to January 01, 2022. For each day during this period, school closure either due to a government decision or a public holiday is determined.

2.2. Data Analysis

2.2.1. Generalised Additive Model

Four periods of school closure longer than 14 days have been identified. These four periods are used to fit a Generalised Additive Model (or GAM) using the

R package *mgcv* (Wood, 2017) in order to test the hypothesis of a significant change in the evolution of cases among age groups from 1 to 4, from 5 to 14, and from 15 to 24 (Eq 1).

By estimating the degree of smoothness of a Bayesian spline smoothing using restricted maximum likelihood estimation (Wood, 2011), GAM identifies dynamic patterns underlying the evolution of Covid-19 cases reported while taking into account the random effect of different age groups as follows:

$$Y_{is} = \alpha_i + f(X_s) + a_{is} + \epsilon_{is} \quad (1)$$

where i is an age group among the 10 age groups investigated and s is the date corresponding to the confirmed Covid-19 cases. Y_{is} represents the confirmed Covid-19 cases assuming a negative binomial distribution for the fitting (Loader, 2006). The response variable Y_{is} includes a specific intercept for each age group (α_i). A smooth effect over time $f(X_s)$ is applied to model (Eq 2) to predict the nonlinear evolution of Covid-19 cases. This smooth effect $f(X_s)$ is built up in basic components, called the basis functions $b_j(X_s)$, such that:

$$f(X_s) = \sum_{j=1}^k \beta_j \times b_j(X_s) \quad (2)$$

where the regression parameters β_j are estimated by penalized likelihood maximization.

The model also includes the random effects term $a_{is} = Zb_i$ where Z is a random effects matrix and b_i is a vector of random effects described by $b_i \sim N(0, D)$. In this, D represents a covariance matrix. The error term ϵ_{is} is assumed to be normally and independently distributed $\epsilon_{is} \sim N(0, \sigma^2)$.

2.2.2. Transfer Entropy

Transfer Entropy (TE) can be used to infer the temporal relationship between two time series I and J . This measure indicates whether J can be used to reduce the uncertainty on the future of I and, consequently, that J causes I (Schreiber, 2000). As such, Granger causality is a special case of TE applied to times series that are jointly Gaussian distributed (Barnett et al., 2009). Therefore, TE is a more robust analysis of times series especially when applied to the impact of age cohort in pandemic transmission (Kissler et al., 2020).

The influence of the evolution in Covid-19 cases across all age groups is evaluated using Shannon's transfer entropy is given by:

$$T_{J \rightarrow I}(k, l) = \sum_{i,j} p(i_{t+1}, i_t^{(k)}, j_t^{(l)}) \cdot \log \left(\frac{p(i_{t+1} | i_t^{(k)}, j_t^{(l)})}{p(i_{t+1} | i_t^{(k)})} \right), \quad (3)$$

where $T_{J \rightarrow I}$ consequently measures the influence of the change dynamic from an age group J to another age group I (Eq 3).

In order to observe how increase in daily numbers reported from a specific age group affects the other age groups, the day-by-day difference in Covid-19 confirmed cases (i.e., differences with a lag 1) is calculated. The use of day-by-day difference in Covid-19 confirmed cases satisfies the requirement of stationary for the calculation of Shannon’s Transfer Entropy (Shannon, 1948). The Transfer Entropy analysis of younger age groups daily changes in Covid-19 cases on older age groups is done using the R package *RTransferEntropy* (Behrendt et al., 2019).

3. Results

Overall, the trend of confirmed Covid-19 cases in Ireland is similar to those of other European countries (Figure 1). Indeed it is possible to observe a significant increase corresponding to each of the 4 waves of transmission (March-April 2020, December 2020-January 2021, September-October 2021 and November-December 2021).




Figure 1: Evolution of confirmed Covid-19 cases in Ireland alongside with school closures.

The evolution of Covid-19 cases reveals some similarities across all age groups. However, the influence of each waves on the each groups has also some particularities (Figure 2). For example, the first wave was more important among the oldest age groups whereas the third wave was more important among the youngest age groups.




Figure 2: Evolution of confirmed Covid-19 cases by age groups in Ireland: raw data (bars) and 7 day rolling average (lines).




Figure 3: Standardized effect of the smooth term in Generalized Additive Model by age group.

In order to evaluate the shape of the trend in the numbers of Covid-19 cases reported after the three school closure longer than 14 consecutive days, a GAM was fitted taking into account numbers for 1 to 4, 5 to 14, and 15 to 24 year old. The obtained results satisfy the requirements to fit this model which explains 99.6% of the deviance in Covid-19 cases. These results show no significant change in the evolution of cases for 1 to 4 year old ($\chi^2(1) = 2.15$, $p = 0.143$) as well as for 5 to 14 year old ($\chi^2(1) = 1.61$, $p = 0.204$). However, the smooth term of the GAM is significant for 15 to 24 year old ($\chi^2(5.02) = 201.27$, $p < 0.001$), which indicates that the evolution of cases is not linear and, therefore, the school closure had a significant impact on the evolution of cases.

Before the Transfer Entropy analysis, an Augmented Dickey-Fuller has been applied on each age group in order to insure that the daily changes in Covid-19 cases are stationary.

4. Discussion

Knowledge about the transmission of the virus significantly improved with the amount of studies performed, especially in the case of how the virus behaves with children. From the early analyses showing that the virus was instantness in children (Li et al., 2020), the results has changed to more nuanced position which states that the spread of the virus in children is moderate.

This indicate mix results, while the school closure was ineffective in child care settings and primary school, the closure of secondary schools and colleges had an impact on Covid-19 transmission.




Figure 4: Matrix of Transfer Entropy coefficient and their corresponding p-value according every age group combination. Age groups on the x-axis are influencing the age groups on the y-axis ($X \rightarrow Y$). * $p < .05$, ** $p < .01$, *** $p < .001$.

5. Conclusion

References

- Barnett, L., Barrett, A.B., Seth, A.K., 2009. Granger causality and transfer entropy are equivalent for gaussian variables. *Phys. Rev. Lett.* 103, 238701. doi:10.1103/PhysRevLett.103.238701
- Behrendt, S., Dimpfl, T., Peter, F.J., Zimmermann, D.J., 2019. RTransferEntropy — quantifying information flow between different time series using effective transfer entropy. *SoftwareX* 10, 100265. doi:10.1016/j.softx.2019.100265
- Chang, S.L., Harding, N., Zachreson, C., Cliff, O.M., Prokopenko, M., 2020. Modelling transmission and control of the COVID-19 pandemic in australia. *Nature communications* 11, 1–13. doi:10.1038/s41467-020-19393-6
- Falk, A., Benda, A., Falk, P., Steffen, S., Wallace, Z., Høeg, T.B., 2021. COVID-19 cases and transmission in 17 k–12 schools—wood county, wisconsin, august 31–november 29, 2020. *Morbidity and Mortality Weekly Report* 70, 136–140. doi:10.15585/mmwr.mm7004e3
- Fukumoto, K., McClean, C.T., Nakagawa, K., 2021. No causal effect of school closures in japan on the spread of COVID-19 in spring 2020. *Nature medicine* 27, 2111–2119. doi:10.1038/s41591-021-01571-8
- Gras-Le Guen, C., Cohen, R., Rozenberg, J., Launay, E., Levy-Bruhl, D., Delacourt, C., 2021. Reopening schools in the context of increasing COVID-19 community transmission: The french experience. *Archives de Pédiatrie* 28, 178–185. doi:10.1016/j.arcped.2021.02.001
- Heavey, L., Casey, G., Kelly, C., Kelly, D., McDarby, G., 2020. No evidence of secondary transmission of COVID-19 from children attending school in ireland, 2020. *Eurosurveillance* 25. doi:10.2807/1560-7917.ES.2020.25.21.2000903
- Hoek, W. van der, Backer, J.A., Bodewes, R., Friesema, I., Meijer, A., Pijnacker, R., Reukers, D.F.M., Reusken, C., Roof, I., Rots, N., Te Wierik, M.J.M., Gageldonk-Lafeber, A. van, Waegemaekers, C., Hof, S. van den, 2020. The role of children in the transmission of SARS-CoV-2. *Nederlands tijdschrift voor geneeskunde* 164.
- Iwata, K., Doi, A., Miyakoshi, C., 2020. Was school closure effective in mitigating coronavirus disease 2019 (COVID-19)? Time series analysis using bayesian inference. *International Journal of Infectious Diseases* 99, 57–61. doi:10.1016/j.ijid.2020.07.052
- Kim, J., Choe, Y.J., Lee, J., Park, Y.J., Park, O., Han, M.S., Kim, J.-H., Choi, E.H., 2021. Role of children in household transmission of COVID-19. *Archives of Disease in Childhood* 106, 709–711. doi:10.1136/archdischild-2020-319910
- Kissler, S.M., Viboud, C., Grenfell, B.T., Gog, J.R., 2020. Symbolic transfer entropy reveals the age structure of pandemic influenza transmission from high-volume influenza-like illness data. *Journal of the Royal Society Interface* 17, 20190628. doi:10.1098/rsif.2019.0628
- Li, X., Xu, W., Dozier, M., He, Y., Kirolos, A., Theodoratou, E., others, 2020. The role of children in transmission of SARS-CoV-2: A rapid review. *Journal of global health* 10. doi:10.7189/jogh.10.011101

- Loader, C., 2006. Local regression and likelihood. Springer, Boston, MA.
- Lopez, A.S., Hill, M., Antezano, J., Vilven, D., Rutner, T., Bogdanow, L., Claflin, C., Kracalik, I.T., Fields, V.L., Dunn, A., others, 2020. Transmission dynamics of COVID-19 outbreaks associated with child care facilities—salt lake city, utah, april–july 2020. *Morbidity and Mortality Weekly Report* 69, 1319. doi:10.15585/mmwr.mm6937e3
- Ludvigsson, J.F., 2020. Children are unlikely to be the main drivers of the COVID-19 pandemic – a systematic review. *Acta Paediatrica* 109, 1525–1530. doi:https://doi.org/10.1111/apa.15371
- Meuris, C., Kremer, C., Geerinck, A., Locquet, M., Bruyère, O., Defèche, J., Meex, C., Hayette, M.-P., Duchene, L., Dellot, P., Azarzar, S., Maréchal, N., Sauvage, A.-S., Fripiat, F., Giot, J.-B., Léonard, P., Fombellida, K., Moutschen, M., Durkin, K., Artesi, M., Bours, V., Faes, C., Hens, N., Darcis, G., 2021. Transmission of SARS-CoV-2 after COVID-19 screening and mitigation measures for primary school children attending school in liège, belgium. *JAMA Network Open* 4, e2128757–e2128757. doi:10.1001/jamanetworkopen.2021.28757
- Schreiber, T., 2000. Measuring information transfer. *Phys. Rev. Lett.* 85, 461–464. doi:10.1103/PhysRevLett.85.461
- Shannon, C.E., 1948. A mathematical theory of communication. *The Bell system technical journal* 27, 379–423.
- Siebach, M.K., Piedimonte, G., Ley, S.H., 2021. COVID-19 in childhood: Transmission, clinical presentation, complications and risk factors. *Pediatric Pulmonology* 56, 1342–1356. doi:https://doi.org/10.1002/ppul.25344
- Soriano-Arandes, A., Gatell, A., Serrano, P., Biosca, M., Campillo, F., Capdevila, R., Fàbrega, A., Lobato, Z., López, N., Moreno, A.M., Poblet, M., Riera-Bosch, M.T., Rius, N., Ruiz, M., Sánchez, A., Valdepérez, C., Vilà, M., Pineda, V., Lazcano, U., Díaz, Y., Reyes-Urueña, J., Soler-Palacín, P., Catalonia Research Group, C.P.D. in, 2021. Household severe acute respiratory syndrome coronavirus 2 transmission and children: A network prospective study. *Clinical Infectious Diseases* 73, e1261–e1269. doi:10.1093/cid/ciab228
- Stage, H.B., Shingleton, J., Ghosh, S., Scarabel, F., Pellis, L., Finnie, T., 2021. Shut and re-open: The role of schools in the spread of COVID-19 in europe. *Philosophical Transactions of the Royal Society B* 376, 20200277. doi:10.1098/rstb.2020.0277
- Sugishita, J.A.S., Yoshiyuki AND Kurita, 2020. Effects of voluntary event cancellation and school closure as countermeasures against COVID-19 outbreak in japan. *PLOS ONE* 15, 1–10. doi:10.1371/journal.pone.0239455
- Walger, P., Heininger, U., Knuf, M., Exner, M., Popp, W., Fischbach, T., Trapp, S., Hübner, J., Herr, C., Simon, A., others, 2020. Children and adolescents in the CoVid-19 pandemic: Schools and daycare centers are to be opened again without restrictions. The protection of teachers, educators, carers and parents and the general hygiene rules do not conflict with this. *GMS hygiene and infection control* 15. doi:10.3205/dgkh000346
- Wood, S.N., 2011. Fast stable restricted maximum likelihood and marginal likelihood estimation of semiparametric generalized linear models. *Journal of the Royal Statistical Society (B)* 73, 3–36.

- Wood, S.N., 2017. Generalized additive models: An introduction with r. CRC press.
- Zhen-Dong, Y., Gao-Jun, Z., Run-Ming, J., Zhi-Sheng, L., Zong-Qi, D., Xiong, X., Guo-Wei, S., 2020. Clinical and transmission dynamics characteristics of 406 children with coronavirus disease 2019 in china: A review. *Journal of Infection* 81, e11–e15. doi:10.1016/j.jinf.2020.04.030