




PROF. DR. NURSAN ÇINAR

THE LANCET

"If a new vaccine became available that could prevent one million or more child deaths a year, and that was moreover cheap, safe, administered orally, and required no cold chain, it would become an immediate public health imperative. Breastfeeding can do all this and more."

Eğer yılda bir milyondan fazla çocuk ölümünü engelleyecek, ayrıca diğerlerine göre daha ucuz, güvenilir, oral yol ile uygulanan ve soğuk zincir gerektirmeyen yeni bir aşı mevcut olsaydı, bu aşı acil bir halk sağlığı zorunluluğu haline gelecekti. EMZİRME, BÜTÜN BUNLARI VE DAHA FAZLASINI YAPABİLİR....

TNSA VERİLERİNE GÖRE İLK 6 AY SADECE ANNE SÜTÜ İLE BESLENME ORANLARI



WHO/NMH/NHD/14.7

Global Nutrition Targets 2025

Breastfeeding Policy Brief**TARGET:**

Increase the rate of exclusive breastfeeding in the first 6 months up to at least 50%



Gates/Jake Lyell

WHAT'S AT STAKE

In 2012, the World Health Assembly Resolution 65.6 endorsed a *Comprehensive implementation plan on maternal, infant and young child nutrition* (1), which specified six global nutrition targets for 2025 (2). This policy brief covers the fifth target: **Increase the rate of exclusive breastfeeding in the first 6 months up to at least 50%.** The purpose of this policy brief is to increase attention to, investment in, and action for a set of cost-effective interventions and policies that can help Member States and their partners in improving exclusive breastfeeding rates among infants less than six months.

DÜNYA SAĞLIK
ÖRGÜTÜ 2025
HEDEFİ:
İLK 6 AY
SADECE ANNE
SÜTÜ İLE
BESLENME
ORANI EN AZ

%50



Mucizevi
Besin
Anne
Sütü

Her bebek için en uygun besin
kendi annesinin sütüdür.

- Emzirme, hem anne hem bebek için en iyi sonuçları veren en ideal beslenme yöntemidir (Assarian et al. 2014).




Bebeğin doğumdan hemen sonra emzirmeye başlanması, ilk altı ay **tek başına** anne sütü ile beslenmesi, altıncı aydan sonra ek besinlerle birlikte anne sütü ile beslenmenin devam etmesi ve emzirmenin **iki yaşın sonuna** kadar sürdürülmesi; bebeğe sayısız yararlar sağlar.

Laktasyon


Prolaktin

Oksitosin



Bu döngüye müdahale süt üretimini azaltır

OKSİTOSİN REFLEKSİ



Son yıllarda yapılan çalışmalar oksitosin salınımının anne-bebek arasındaki bağlanmayı olumlu etkilediğini ortaya koymaktadır (Galbally ve ark. 2011; Zelkowitz ve ark. 2014).

Bunlar refleksel YARDIM eder.

Bunlar reflekse ENGEL olur

Psychosocial stress moderates the relationships between oxytocin, perinatal depression, and maternal behavior.

Zelkowitz P¹, Gold J², Feeley N³, Hayton B⁴, Carter CS⁵, Tulandi T⁶, Abenhaim HA⁷, Levin P⁸.

⊕ Author information

Abstract

The hormone oxytocin (OT) is of particular interest in the study of childbearing women, as it has a role in the onset and course of labor and breastfeeding. Recent research has linked OT to maternal caregiving behavior towards her infant, and to postpartum depressive symptomatology. There is also evidence that psychosocial adversity affects the oxytocin system. The present study investigated the relationship of endogenous OT in women during pregnancy and at 8weeks postpartum to psychosocial stress, maternal symptoms of depression, and maternal sensitive behavior. It was hypothesized that OT would mediate the effects of maternal depressive symptoms on maternal interactive behavior. We also tested the hypothesis that psychosocial stress would moderate the relationship between OT and maternal depressive symptoms and sensitive behavior. A community sample of 287 women was assessed at 12-14weeks of gestation, 32-34weeks of gestation, and 7-9weeks postpartum. We measured plasma OT, maternal symptoms of depression and psychosocial stress. At the postpartum home visit, maternal behavior in interaction with the infant was videotaped, and then coded to assess sensitivity. In the sample as a whole, OT was not related to maternal depressive symptoms or to sensitive maternal behavior. However, among women who reported high levels of psychosocial stress, higher levels of plasma OT were associated with fewer depressive symptoms and more sensitive maternal behavior. These results suggest that endogenous OT may act as a buffer against the deleterious effects of stress, thereby protecting high risk women from developing depressive symptoms and promoting more sensitive maternal interactive behavior.

Copyright © 2014 Elsevier Inc. All rights reserved.

Zelkowitz ve arkadaşlarının (2014) yaptıkları çalışma sonucunda stresin zararlı etkilerine karşı koruyan oksitosinin, doğum sonrasında salınımı artarak anne- bebek arasında duyarlı ilişkinin sürdürülmesine katkı sağladığı belirtilmiştir.

- * Bebeklerde normal emzirilmeye rağmen doğumu izleyen 8-12. Günlerde, 3-4 haftalarda ve 3. ayda sık sık ağlama görülebilir.



- * Bu durumun nedeni annenin sütünün yetersizliği değil, bebeğin söylenilen sürelerdeki hızlı büyümelerinden kaynaklanan iştah artmasıdır.



Daha sık ve daha uzun emme isteğine karşın bebek hep aç gibidir.

Mama verilmesi sorunu arttırabilir!

Birkaç gün içinde annenin sütünün artacağı ve duruma adapte olacağı bilinmelidir.

Beslenme sıklığı arttırılmalıdır.

KİLO KAYBI
ENDİŞESİ VE
GEREKSİZ
FORMÜL MAMA
BAŞLANMASI

Yenidoğan bebeğin midesi

İlk gün	Kiraz/Bilye kadar (~5-7 ml)
3. gün	Ceviz kadar (~ 22-27 ml)
1. hafta	Yumurta kadar (~45-60 ml)
1. ay	1-1,5 çay bardağı (~80-150 ml)

İlk haftada
%7-10 kilo
kaybı
normaldir

Anne Sütünün Yararları

- Sağlık

- Çocuğa ilişkin

Kısa Vadede

- Anneye ilişkin

Uzun Vadede

- Ekonomik- Sosyal

- Çevre



Anne st bebeęin hayata tutunduęu ilk canlı daldır.

- Her canlının stnn kendisine ve bebeęine zel olması anne stn yavrusu iin benzersiz bir besin maddesi yapan bir özelliktir...

ANNE SÜTÜ İLE BESLENEN BEBEKLER

- Doğumdan sonra emzirilen bebekler enfeksiyonlardan korunur ve mortalite azalır
- Anne sütü, bir çocuğun 6 -12 ay arasındaki enerji ihtiyacının yarısından daha fazlasını, 12-24 ay arasındaki enerjinin ise üçte birini sağlayabilir.
- Bebeklik döneminde emzirilen çocukların kilolu olma veya obez olma eğilimleri daha düşüktür.



465631||

The effect of breastfeeding on children's cognitive and noncognitive development ☆

Cristina Borra^{a,*}, Maria Iacovou^b, Almudena Sevilla^c

[Show more](#)

<https://doi.org/10.1016/j.labeco.2012.05.007>

Abstract

This paper uses propensity score matching methods to investigate the relationship between breastfeeding and children's cognitive and noncognitive development. We find that breastfeeding for four weeks is positively and statistically significantly associated with higher cognitive test scores, by around one tenth of a standard deviation. The association between breastfeeding and noncognitive development is weaker, and is restricted to children of less educated mothers. We conclude that interventions which increase breastfeeding rates would improve not only children's health, but also their cognitive skills, and possibly also their noncognitive development.

ACTA PÆDIATRICA
NURTURING THE CHILD

Acta Pædiatrica ISSN 0803-5253

REGULAR ARTICLE

Breastfeeding and early child development: a prospective cohort study

Wendy H. Oddy (wendyo@ichr.uwa.edu.au)¹, Monique Robinson¹, Garth E. Kendall^{1,2}, Jianghong Li³, Stephen R. Zubrick³, Fiona J. Stanley¹
Copyright © 2010 International Pediatric Research Foundation, Inc.

Impact of Breast Milk on Intelligence Quotient, Brain Size, and White Matter Development

ELIZABETH B. ISAACS, BRUCE R. FISCHL, BRIAN T. QUINN, WUI K. CHONG, DAVID G. GADIAN, AND ALAN LUCAS

- Yapılan çalışmalarda emzirmenin erken çocukluk döneminde beyin gelişimini desteklediği saptanmıştır.

RESEARCH ARTICLE

Does Breastfeeding Help to Reduce the Risk of Childhood Overweight and Obesity? A Propensity Score Analysis of Data from the KiGGS Study

Maïke Miriam Grube*, Elena von der Lippe, Martin Schlaud, Anna-Kristin Brettschneider

Department of Epidemiology and Health Monitoring, Robert Koch Institute, Berlin, Germany

* maike.grube@posteo.de

Abstract

Background

ACTA PÆDIATRICA
NURTURING THE CHILD

Acta Paediatrica ISSN 0803-5253

REVIEW ARTICLE

Long-term consequences of breastfeeding on cholesterol, obesity, systolic blood pressure and type 2 diabetes: a systematic review and meta-analysis

Bernardo L. Horta (blhorta@gmail.com), Christian Loret de Mola, Cesar G. Victora

Postgraduate Program in Epidemiology, Universidade Federal de Pelotas, Pelotas, Brazil

Keywords

Blood lipids, blood pressure, breastfeeding, meta-analysis, obesity, systematic review, type 2 diabetes

Correspondence

Bernardo L. Horta, Postgraduate Program in Epidemiology, Universidade Federal de Pelotas, Rua Marechal Deodoro, 1160 – 3o andar, 96020-220 – Pelotas, Brazil.
Tel: +55 (53) 3284 – 1300 |
Fax: +55 (53) 3284 – 1300 |
Email: blhorta@gmail.com

Received

13 May 2015; revised 25 June 2015;
accepted 14 July 2015

DOI:10.1111/apa.13133

ABSTRACT

Aim: To systematically review the evidence on the associations between breastfeeding and overweight/obesity, blood pressure, total cholesterol and type 2 diabetes.

Methods: Two independent literature searches were carried out using the MEDLINE, LILACS, SCIELO and Web of Science databases. Studies restricted to infants and those without an internal comparison group were excluded. Fixed- and random-effects models were used to pool the estimates.

Results: Breastfed subjects were less likely to be considered obese/overweight [pooled odds ratio: 0.74 (95% confidence interval (CI): 0.70; 0.78)] ($n = 113$). Among the 11 high-quality studies, the association was smaller [pooled odds ratio: 0.87 (95%CI: 0.76; 0.99)]. Total cholesterol ($n = 46$) was independent of breastfeeding [pooled mean difference: -0.01 mmol/L (95%CI: -0.05 ; 0.02)]. Systolic blood pressure ($n = 43$) was lower among breastfed subjects [mean difference: -0.80 (95%CI: -1.17 ; -0.43)], but no association was observed among larger studies, and for diastolic blood pressure ($n = 38$) [mean difference: -0.24 (95%CI: -0.50 ; 0.02)]. For type 2 diabetes ($n = 11$), the odds ratio was lower among those subjects who had been breastfed [pooled odds ratio: 0.65 (95%CI: 0.49; 0.86)].

Conclusion: Breastfeeding decreased the odds of type 2 diabetes and based on high-quality studies, decreased by 13% the odds of overweight/obesity. No associations were found for total cholesterol or blood pressure.

Yapılan
çalışmalarda
emzirmenin
özelikle
çocuklarda
obezite riskini
azalttığı
saptanmıştır.

Systematic review with meta-analysis: early infant feeding and coeliac disease – update 2015

H. Szajewska^{*}, R. Shamir[†], A. Chmielewska^{*}, M. Pieścik-Lech^{*}, R. Auricchio[‡], A. Ivarsson[§], S. Kolacek[¶], S. Koletzko^{**}, I. Korponay-Szabo^{††}, M. L. Mearin^{‡‡}, C. Ribes-Koninckx^{§§} & R. Troncone[‡] on behalf of the PREVENTCD Study Group^a

Szajewska ve ark.
(2015) yaptıkları
meta analiz de,
emzirmenin
çocuklukta çölyak
hastalığı
gelişiminde risk
oluşturmadığı
saptanmıştır.

SUMMARY

Background

New evidence emerged on early feeding practices and the risk of coeliac disease.

Aim

To systematically update evidence on these practices to find out whether there is a need to revise current recommendations.


Methods

MEDLINE, EMBASE and the Cochrane Library were searched from July 2012 (end of last search) to February 2015 for studies of any design that assessed the effect of gluten consumption and breastfeeding on the development of coeliac disease and/or coeliac disease-related autoimmunity.

Results

We identified 21 publications, including two, new, large, randomised controlled trials performed in high-risk infants. Exclusive or any breastfeeding, as well as breastfeeding at the time of gluten introduction, did not reduce the risk of developing coeliac disease during childhood. For infants at high risk of developing coeliac disease, gluten introduction at 4 months of age in very small amounts, or at 6 or 12 months of age, resulted in similar rates of coeliac disease diagnosis in early childhood. Later gluten introduction was associated with later development of coeliac specific autoimmunity and coeliac disease during childhood, but not total risk reduction. Observational studies indicate that consumption of a higher amount of gluten at weaning may increase the risk for coeliac disease development.

Conclusions



- Çalışmalarda, anne sütü içeriğinin bebeği kanserden koruyucu maddeler (Hamlet-Human Alpha-lactalbumin Made Lethal to Tumours) içerdiği saptanmıştır

Gustafsson, L., Hallgren, O., Mossberg, A. K., Pettersson, J., Fischer, W., Aronsson, A., & Svanborg, C. (2005). HAMLET kills tumor cells by apoptosis: structure, cellular mechanisms, and therapy. *The Journal of nutrition*, 135(5), 1299-1303

Rath, E. M., Duff, A. P., Håkansson, A. P., Vacher, C. S., Liu, G. J., Knott, R. B., & Church, W. B. (2015). Structure and Potential Cellular Targets of HAMLET-like Anti-Cancer Compounds made from Milk Components. *Journal of Pharmacy & Pharmaceutical Sciences*, 18(4), 773-824.

Mossberg, A. K., Hun Mok, K., Morozova-Roche, L. A., & Svanborg, C. (2010). Structure and function of human α -lactalbumin made lethal to tumor cells (HAMLET)-type complexes. *FEBS journal*, 277(22), 4614-4625.

ANNENİN SAĞLIĞINI KORUR

Hours Mothers Sleep



Emziren annenin uykü süresi daha uzundur

Emziren annenin günlük enerji ihtiyacı daha fazladır



Mothers' Daily Energy




Emziren annenin
doğum sonrası
depresyon riski
daha azdır



Mothers' Depression



Angry or Irritable



Emziren annede
doğum sonrası öfke
ve sinirlilik daha
azdır

Anne st hibir zaman birebir taklit edilemez.

- 
- Bu nedenle emzirmeyen anneler hangi mamanın bebekleri iin daha iyi olduėu konusunda oraya buraya kořuřtururlarken, emzirmeyi seen anneler bebeklerinin en iyi " besinle" beslendiklerinden emindirler.

Emziren annenin "**temiz su**" bulma endiřesi yoktur.



Emzirmek anne iin doęal bir sakinleřtiricidir.

- Emzirme anne bebek arasında bağlanmayı güçlendiren eşsiz bir etkileşimdir (Evcili ve ark. 2014; Himani ve Kumar 2011).



- Erken anne-bebek bağlanması, beyinde emzirme ile ilgili hormonların tetiklediği kimyasal nörotransmitterlerin salınmasını sağlayan bir dizi değişikliğe neden olur (Azza Abul-Fadl et al.2005)

ANNE VE BEBEK BAĞLANMAMASI

- Anne ve bebek bağlanmasını olumsuz etkileyen faktörler;
- Annenin sosyal veya psikolojik hastalıklar
- Yetersiz aile desteği
- Depresyon
- Düşük benlik saygısı
- Düşük sosyal sınıf



Emzirme ve
Bebek Bakımı etkilenir



İHMAL VE İSTİSMAR

The Relationship Between Maternal Attachment, Perceived Social Support and Breast-Feeding Sufficiency

Nursan Cinar, Dilek Köse and Sevin Altinkaynak

ABSTRACT

Objective: To determine the relationship between maternal attachment, perceived social support and breast-feeding sufficiency.

Study Design: Descriptive correlational design.

Place and Duration of Study: A state hospital and two family health centers in Sakarya, Turkey, between June to December 2011.

Methodology: The sample was 122 voluntary mothers who had healthy babies of 1 - 2 months old. The data were collected by a Personal Information Form, Maternal Attachment Inventory (MAI), Multidimensional Scale of Perceived Social Support (MSPSS) and Breast-feeding Self-Efficacy Scale-Short Form (BSES-SF). The data collected were analysed by percentage distribution, mean square, independent sample t-test, Mann-Whitney U, Kruskal-Wallis and Pearson correlation.

Results: The mean age of the mothers was 25.01 ± 2.2 years, and 48.4% of them were primary school graduates. BSES-SF was 61.02 ± 8.44 (16 - 70), MAI was 99.07 ± 7.19 (56 - 100) and MSPSS was 66.40 ± 13.58 (37 - 84). There was a positive, medium level, significant relationship between the total scores of BSES-SF and MAI ($r=0.370$, $p < 0.001$). There was a positive, medium level, significant relationship between the total score of BSES-SF and the score from family subdimension of MSPSS ($r=0.255$, $p < 0.01$). There was a positive, medium level, significant relationship between the total score of MAI and the total score of family subdimension of MSPSS ($r=0.339$, $p < 0.001$).

Conclusion: Successful maternal attachment and familial support positively affected the breast-feeding sufficiency of the mother.

Key Words: Breast-feeding. Self-sufficiency. Attachment. Maternal attachment. Social support.

Yapılan bir alıřmada, anne st ile bebeęini besleyen annelerde, annelik memnuniyetinin arttıęı saptanmıřtır.

Sevimli Gler, D. "Annelerin Algıladıkları Sosyal Destek ve Annelik Deneyimleri Arasındaki İliřki", Sakarya niversitesi Saęlık Bilimleri Enstits, Sakarya-2016 (Tez Danıřmanı: Nursan INAR)

Anne Sütü - Çocuk Ruh Sağlığı

Yapılan çalışmalarda bebeklik döneminde anne sütü almamış olan veya 2 haftadan kısa süreli alanlarda şizofreni, alkol bağımlılığı, dikkat eksikliği- hiperaktivite bozukluğu ve düşük benlik saygısı görülme olasığının daha yüksek olduğu tespit edilmiştir.

Sorensen H, Mortensen E, Reinisch J, Mednick S. Breastfeeding and Risk of schizophrenia in the Copenhagen perinatal cohort. *Acta Psychiatr Scand* 2005;112:26-9.

Sorensen H, Mortensen E, Reinisch J, Mednick S. Early weaning and hospitalization with alcohol-related diagnoses in adult life. *Am J Psychiatry* 2006;163:704-9.

Çakaloğlu B, Akay A. Dikkat eksikliği ve yıkıcı davranış bozukluklarında doğum öncesi, doğum ve doğum sonrasında yaşanan sorunların ve anne sütü alışı sürelerinin psikopatoloji gelişimine katkıları. *İzmir Çocuk ve Gençlik Ruh Sağlığı Dergisi* 2005;12:(1):3-10.

Kwok MK, Leung GM, Schooling CM. Breast feeding and early adolescent behaviour, self-esteem and depression: Hong Kong's 'Children of 1997' birth cohort. *Arch Dis Child* 2013; 98(11):887-94.

Çoğul bebekler de başarıyla
emzirilebilir.



ANNE SÜTÜ



*Düşük doğum
ağırlıklı
*Erken doğum



**Anne sütü ile
beslenme önemli**

Tek çocuğa göre çoğul bebeklerin emzirilmesi anne için:

- Daha yorucu, daha çok uykusuzluğa sebep olması ve annenin sütünün yetersiz olacağını düşünmesi, bakım eksikliği ve beslenme pozisyonu bilgisizliği nedeniyle göğüs uçlarındaki yaralar, kendisine beslenme süresince yardım edecek birinin olmaması gibi nedenlerle bebeklerini anne sütü verme oranları düşüktür.
- Oysa ki çoğul bebekler içinde yeterli miktar ve kalitede süt üretildiği çalışmalarla belgelenmiştir

(Rimon 2006, Gromada ve Spangle, 1998).

Süt üretiminin bir arz-talep ilişkisi



(Kielbratowska ve ark. 2010)
İkiz bebek sahibi annelerin doğumdan sonraki altı ay her gün **1-2 kg arasında süt** üretebildiği, üçüz bebek sahibi annelerde ise üretilen süt miktarının **3 kg'dan fazla** olduğunu bildirilmektedir.

- 
- ✓ bebeklerin sık emzirilmesi
 - ✓ annenin sağlıklı beslenmesi
 - ✓ yeterli sıvı alması ve
 - ✓ stresin azaltılması



Mothers' Attitudes Toward Feeding Twin Babies in the First Six Months of Life: A Sample From Sakarya, Turkey

Cinar N et al.

Table 1. Breastfeeding Attitudes by Baby and Month

	First Baby						Second Baby					
	1st mo	2nd mo	3rd mo	4th mo	5th mo	6th mo	1st mo	2nd mo	3rd mo	4th mo	5th mo	6th mo
Breast milk	3 (10.0)	2 (6.7)	1 (3.3)	1 (3.3)	1 (3.3)	-	2 (6.7)	3 (10.0)	3 (10.0)	1 (10.0)	1 (3.3)	-
Formula food	2 (6.7)	5 (16.7)	8 (20)	10 (33.3)	12 (40)	13 (43.3)	2 (6.7)	5 (16.7)	5 (16.7)	7 (23.3)	11 (36.7)	11 (40)
Breast milk and formula food	25 (83.3)	23 (76.7)	23 (76.7)	19 (63.3)	15 (50)	7 (23.3)	28 (86.7)	22 (73.4)	22 (73.4)	20 (66.7)	16 (53.3)	8 (26.6)
Breast milk, formula food, and weaning	-	-	-	-	1 (3.3)	7 (23.3)	-	-	-	-	1 (3.3)	7 (23.3)
Formula food and weaning	-	-	-	-	1 (3.3)	2 (6.7)	-	-	-	-	1 (3.3)	2 (6.7)
Breast milk and weaning	-	-	-	-	-	1 (3.3)	-	-	-	-	-	1 (3.3)

who stated that they had no previous experience of breastfeeding twin babies and had received training to breastfeed multiple babies was 17 (56.7%). Twenty-seven (90%) of the mothers had had caesarean sections, and half of the babies were preterm. Only a few of the babies were fed breast milk for five months. During the following months, the breastfeeding regimen was as follows: 5 babies were breastfed for a month, 5 babies for 2 months, 4 babies for 3 months, 4 babies for 4 months, 2 babies for 5 months, and no babies in the sixth month.

Conclusions: It was found that the number of twin babies who were only breastfed in the first six months of life was low. According to the literature, a method of breast milk is sufficient for some babies. Mothers are encouraged to breastfeed their babies for at least 6 months.

Sadece 5 bebek ilk ay boyunca, 5 bebek 2 ay boyunca, 4 bebek 4 ay boyunca, 2 bebek 5 ay boyunca anne sütü aldığı bulundu.
6. Ayda hiçbir bebek anne sütü almıyordu

Anne Sütü
Mucizedir!



- Emzirme annenin uterusunun eski haline dönmesini hızlandırır.
- Emziren annelerin doğum sonrası kanama riski önemli ölçüde azalır.

oEmziren anneler doğum sonrası daha kolay kilo verirler.

oEmzirme doğal bir gebelikten korunma yöntemidir.



- Emziren annelerde demir eksikliği anemisi ortaya çıkma riski azalır.



- ↓
- Emziren annelerde doğum sonrası kanama miktarı daha az olduğundan ve emzirmeye devam ettikleri sürece adet görme olasılıkları daha düşük olduğundan bu anneler, doğumda kaybettikleri demir depolarını daha kısa zamanda tekrar oluştururlar.

ANNENİN SAĞLIĞINI KORUR

- Emziren annelerde ileride yumurtalık, meme ve endometrium kanserine yakalanma riski daha azdır.



World Health Organization. (2009). Infant and young child feeding: model chapter for textbooks for medical students and allied health professionals. *Infant and young child feeding: model chapter for textbooks for medical students and allied health professionals*. Updated September 2016

ANNENİN SAĞLIĞINI KORUR

- Kalp hastalığı riski azalır.
- Emzirme sırasında,
total kolesterol, LDL↓kolesterol ve
trigliserid düzeyleri azalırken,
faydalı HDL düzeyi yüksek kalır.



Duration and exclusiveness of breastfeeding and risk of childhood atopic diseases.

Elbert NJ^{1,2}, van Meel ER^{1,3,4}, den Dekker HT^{1,3,4}, de Jong NW⁵, Nijsten TEC², Jaddoe VVW^{1,4,6}, de Jongste JC³, Pasmans SGMA², Duijts L^{3,7}.

⊕ Author information

Abstract

BACKGROUND: Breastfeeding may have immune modulatory effects that influence the development of childhood allergic sensitization and atopic diseases. We aimed to examine the associations of breastfeeding with childhood allergic sensitization, inhalant or food allergy and eczema, and whether any association was affected by disease-related modification of the exposure or modified by maternal history of maternal history of allergy, eczema or asthma.

METHODS: This study among 5,828 children was performed in a population-based prospective cohort from fetal life onwards. We collected information on duration (<2 months, 2-4 months, 4-6 months and ≥6 months) and exclusiveness (non-exclusive vs. exclusive for 4 months) of breastfeeding in infancy by postal questionnaires. At age 10 years, inhalant and food allergic sensitization were measured by skin prick tests, and physician-diagnosed inhalant and food allergy by a postal questionnaire. Data on parental-reported eczema were available from birth until age 10 years.

RESULTS: We observed no association of breastfeeding with any allergic sensitization, physician-diagnosed allergy, or combination of these outcomes. Shorter breastfeeding duration was associated with an overall increased risk of eczema (p-value for trend <0.05). Non-exclusively breastfed children had an overall increased risk of eczema (aOR (95% CI): 1.11 (1.01, 1.23)), compared with children exclusively breastfed for 4 months. Risk period-specific sensitivity analyses, additional adjustment for ointment use for eczema at age 2 months, and cross-lagged modeling showed no consistent results for disease-related modification of the exposure. Results were not modified by maternal history of allergy, eczema or asthma (lowest p-value for interaction =0.13).

CONCLUSIONS: Shorter duration or non-exclusiveness of breastfeeding is associated with a weak overall increased risk of eczema but not allergic sensitization or physician-diagnosed allergy at age 10 years This article is protected by copyright. All rights reserved.

Elbert ve arkadaşlarının (2017) yaptıkları çalışmada, kısa veya aralıksız emzirme zayıf düzeyde egzema riskine neden olsa da alerjik duyarlılık veya tıbbi tanı almış alleji vakalarına neden olmadığı saptanmıştır.



Maternal food restrictions during breastfeeding

Goun Jeong, MD, Sung Won Park, MD, Yeon Kyung Lee, MD, PhD, Sun Young Ko, MD, PhD
 Department of Pediatrics, Cheil General Hospital and Women's Healthcare Center, Dankook University College of Medicine, Seoul, Korea

Purpose: This study investigated self-food restriction during breastfeeding, reviewed the literature showing the effect of maternal diet on the health of breast-fed infants, and explored the validity of dietary restrictions.

Methods: Questionnaire data were collected from breastfeeding Korean mothers who visited the pediatric clinic of Cheil General Hospital & Women's Healthcare Center from July 2015 through August 2015. The survey included items assessing maternal age, number of children, maternal educational attainment, household income, degree of difficulty with self-food restriction, types of self-restricted foods, dietary customs during breastfeeding, and sources of information about breastfeeding.

Results: The questionnaire was completed by 145 mothers. More than a third ($n=56$, 39%) had discomfort from and usually avoided 4–5 types of food (mean, 4.92). Mothers younger than 40 years had more discomfort (odds ratio [OR], 12.762; $P=0.017$). Primiparas felt less discomfort than multiparas (OR, 0.436; $P=0.036$). Dietary practices were not influenced by maternal educational attainment or household income. The most common self-restricted foods were caffeine ($n=131$, 90.3%), spicy foods ($n=124$, 85.5%), raw foods ($n=109$, 75.2%), cold foods ($n=100$, 69%), and sikhye (traditional sweet Korean rice beverage) ($n=100$, 69%). Most mothers ($n=122$, 84.1%) avoided foods for vague reasons.

Conclusion: Most mothers restricted certain foods unnecessarily. Literature review identified no food that mothers should absolutely avoid during breastfeeding unless the infant reacts negatively to the food.

- Jeong ve ark. (2017) yaptıkları çalışmada, çoğu annenin bazı gıdaları gereksiz yere kısıtladığını, bebeğin gıdaya olumsuz tepki vermediği sürece emzirme sırasında annelerin kesinlikle kaçınması gereken yiyeceklerin literatürde de tanımlanmadığına dikkat çekmişlerdir.

RESEARCH

Open Access

Breast milk DHA levels may increase after informing women: a community-based cohort study from South Dakota USA



Brian A. Juber^{1†}, Kristina Harris Jackson^{2†}, Kristopher B. Johnson³, William S. Harris^{2,3} and Michelle L. Baack^{3,4*}

Abstract

Background: Docosahexaenoic acid (DHA), an omega-3 fatty acid found in breast milk, has many health benefits for both mother and baby. A 2007 meta-analysis found U.S. women had breast milk DHA levels (0.20% of total fatty acids) below the worldwide mean (0.32%). In 2008, international dietary recommendations were made for pregnant and lactating women to consume 200 mg of DHA per day. This community-based study aimed to define current milk DHA levels from upper Midwest USA lactating mothers and to determine if providing information about their own level along with dietary recommendations would incite changes to increase breast milk DHA content.

Methods: New mothers attending lactation classes or using hospital pumping rooms in Sioux Falls, South Dakota, USA participated by providing one drop of breast milk on a card for fatty acid analysis at baseline and 1 month after initial reporting. DHA levels were analyzed by gas chromatography. Mothers received a report of their own breast milk level along with dietary recommendations on DHA intake for lactating women. Median baseline and follow-up DHA levels were determined and differences were compared by Wilcoxon signed-rank test.

Results: At baseline, breast milk DHA content ($n = 84$) was highly variable (range 0.05 to 0.73%) with a median of 0.18% (IQR, 0.13, 0.28; mean $\pm$ SD, $0.22 \pm 0.13\%$), well below the worldwide average (0.32%). Women who reported taking DHA supplements ($n = 43$) had higher levels than those who did not (0.23% vs. 0.15%, $P < 0.0001$). In a subset of 60 mothers who submitted a second sample, median breast milk DHA content increased from 0.19 to 0.22% ($P < 0.01$).

Conclusions: Findings suggest that providing nursing mothers with their breast milk DHA level and education about DHA intake while breastfeeding motivates change to increase DHA levels.

Keywords: Breast milk fatty acids, Docosahexaenoic acid, Omega-3 fatty acids, Long chain polyunsaturated fatty acids, Breastfeeding

Juber ve ark. (2017) yaptıkları çalışmada, anne sütündeki DHA düzeyi (omega 3) ve DHA takviyesi konusunda eğitim verilen annelerin bebeklerini emzirmeye daha eğilimli oldukları saptanmıştır.

EMZİRMENİN AİLEYE FAYDALARI

- Bebeğin emzirildiđi ailelerde telaş azalır



Emzirme aileye zaman kazandırır.

Emzirilen bebekler daha az hasta olacađı için ailenin sađlık harcamaları azalır.



Hasta çocuklarıyla uğraşmayan ebeveynlerin iş gücü ve iş verimliliği artar.



EMZİRMENİN SOSYAL AÇIDAN FAYDALARI

- Kadına bebeği olsa dahi sosyal yaşama tam olarak katılıma devam etmesi için imkan tanır.
- Anne emzirme yoluyla ağlayan bebeği kolayca susturabileceği için gürültü açısından çevresini rahatsız edici durumlar azalır.
- Emziren anneler arasında bebeklerini besleme açısından sosyo-demografik farklar ortadan kalkar.


RESEARCH ARTICLE

Human breast milk is a rich source of multipotent mesenchymal stem cells

Satish PATKI,¹ Sachin KADAM,² Vikash CHANDRA² and Ramesh BHONDE²¹Patki Research Foundation and Hospital, Shivajinagar, Kolhapur, and ²National Center for Cell Science, Ganeshkhind, Pune, IndiaCytotechnology (2013) 65:38
DOI 10.1007/s10616-012-9492-6

ORIGINAL RESEARCH

Exploring the stem cell and non-stem cell constituents of human breast milk

S. Indumathi · M. Dhanasekaran ·
J. S. Rajkumar · D. Sudarsanam

- Yapılan çalışmalarda, anne sütünün birçok insan dokusunu oluşturmak için potansiyel olarak yeniden programlanabilen mezenkimal kök hücreleri (CD44, CD29, CD 90, CD 105, CD 73), barındırdığı saptanmıştır ve otolog kök hücre tedavisi için anne sütünün kök hücre kaynağı olabileceği düşünülmektedir (Patki et al., 2010; Indumath et al., 2013).

Emzirme ile daha sađlıklı nesiller yetiřtirilir.



Sevgi,
merhamet,
vefa...Bana
annemden
řifa!



Rome

June 22-25 2017



Rome (Italy)

22 - 25 June 2017

Preliminary Program

Daily programme

DAY 3: FRIDAY 23 JUNE 2017 - MORNING SESSIONS

11:50-13:30

Nutrition - Room A

Moderator: Ilaria Sanl (Italy)

- 11:50 **Determinants of iron deficiency anemia among hospitalized children aged 6-59 months in Armenia: a case-control study**
Anush Mnatsakanyan (Yerevan)
School of Public Health, American University of Armenia, N2 Department of Pediatrics, Yerevan State Medical University, Yerevan - Armenia
- 12:10 **The mother-newborn skin-to-skin contact questionnaire (MSSCQ) development and psychometric evaluation among Iranian midwives**
Sedighe Sadat Tavafian (Tehran)
Professor of Health Education & Health Promotion, Tarbiat Modares University - Iran
- 12:30 **Efficacy of the prescription of physical activity in obese child population**
Edgar Debray Hernandez (Bogota)
Corporal Human Movement, National University of Colombia, Bogota - Colombia
- 12:50 **Breast milk perception scale for adults: development, reliability, and validity study**
Nursan Cinar (Sakarya)
Department of Pediatric Nursing, Faculty of Health Sciences, Sakarya University - Turkey
- 13:10 **Determination the packaged ready food consumption patterns of parents and the factors that affect**
Sibel Kucuk, Ara .Gör. Dilek Uludasdemir (Ankara)
Ankara Yildirim Beyazit University - Turkey

Erişkinlerin
Anne Sütü ile
İlgili Algı
Ölçeği

EMZİRMENİN EKONOMİYE FAYDALARI

- Anne sütü bebekler için **en ucuz** ve **en iyi** gıdadır.
- Diğer mama ve gıdalar oldukça pahalıdır.
- Anne sütü ise bebek için değeri parayla ölçülemeyecek kadar yararlıdır.





- Emzirilen bebekler daha sağlıklı bebeklik ve çocukluk geçireceđi için ebeveynlerin işe devamsızlıđı azalacaktır.

Bu işverenin emzirmeyi desteklemesi için önemli bir sebeptir.

EMZİRMENİN ÇEVREYE FAYDALARI



- **Emzirme** doğaldır ve çevre dostudur ancak,
- **yapay mamalar** için bir çok fosil yakıt tüketilerek ve plastik ambalajlar kullanılarak çevreye zarar verilmektedir.

Emzirme atıkları azaltır.

- Emzirmede şişelere, kutulara , bunların paketlerine, etiketlere ihtiyaç yoktur.
- Piyasadaki biberonların çoğu plastik içerir.
- **Plastikse önemli bir çevre kirleticidir.**



- Yapay bebek s tleri ve i lenmi  bebek mamaları  retim, dađıtım ve kullanımının her a amasında ekolojik hasar olu turan yenilenebilir olmayan  r nlerdir.
- Biberonla besleme bebeklerden daha fazla orman kaybı, toprak erozyonu, hava kirliliđi (dioksin ve diđer toksinlerin dahil) iklim deđi iklikleri ve kaynakların israf demektir.

Emzirme atıkları azaltır.

Anne st daha iyi
sindirileceęi iin
bebeklerde dıřk az olur
dolays ile daha az bez
deęiřtirilir.



- Hammadde, paketleme malzemesi ve bunların sevkiyatının yapılması gerekli değildir.
- Birçok ülke dünyanın diğer ucundan bebek maması ve biberon gibi ürünler getirtir.
- Bu büyük yakıt harcanması anlamına gelir ve her yerde **hava kirliliğine** neden olur.

- Anne s  t   d  nya   apında sevk olmak zorunda deęildir; gittięi her yerde, her annenin hazır bir kaynaęı vardır ...

✓ Bu malzemelerin geri d  n    m   de enerji t  ketir.



✓ Besleme malzemelerini almak i  in maęazalara gidi   geli   gerekli deęildir.

○Emzirme için çoğu pahalı ve çok yer kaplayacak malzemelere gerek yoktur. (ölçüm cihazları, sterilizatör v.b)

- Anne sütü bebek için uygun sıcaklıkta ısıtmak gerekmez.
- Anne sütü dışındaki besinlerin hazırlanması daha çok atık üretir.



- Mamalar hazırlanırken suyun ısıtılması için çoğu ülke de odun kullanılır bu da yılda **73 kilo oduna** denk gelir.



COMMENTARY

Open Access

Breastfeeding in public: "You can do it?"

Lisa H Amir

Abstract

On a regular basis there is an outcry about a mother who has been told to cover up or move away from a public area while she is breastfeeding. Mothers should feel free to breastfeed whenever they need to. However, the increasing market for "nursing covers" to hide the breast while feeding is evidence of changing perceptions. Discomfort with the idea of breastfeeding in public has been cited as a reason for some women choosing not to initiate breastfeeding or planning a shorter duration of breastfeeding. Other women are choosing to express and bottle-feed their expressed milk when they are in public. In many cultures today there is a conflict between the concept of breast milk being pure (like tears), and contaminated or "dirty" (like genital secretions or vomit). In these settings the female breast may be considered primarily a sexual organ, and therefore a private part of the body, which needs to be invisible in the public arena. In order to increase breastfeeding initiation and duration and to reduce health inequities breastfeeding needs to be more visible. Let's strive together to make breastfeeding in public unremarkable.

Keywords: Breastfeeding in public, Public perception, Determinants of breastfeeding

- Anneler ihtiyaç duydukları her yerde emzirmekten çekinmemelidirler! Emzirmeyi halk arasında popüler kılmak için beraber çalışmalıyız!...



Breastfeeding in public:
"You can do it?"



You can do it in a train
You can do it on a plane
You can do it in a pool
You can do it at a school
You can do it on the grass
You can do it really fast
You can do it in a park
You can do it in the dark
You can do it at a fair
You can do it ANYWHERE!

*Apologies to Dr. Seuss

In Australian federal law breastfeeding is a right, not a privilege

www.breastfeeding.asn.au

Photos courtesy of Greg D. Buck

PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Breastfeeding: What Do We Know, and Where Do We Go From Here?

Lydia Furman

Pediatrics; originally published online March 27, 2017;

DOI: 10.1542/peds.2017-0150

The online version of this article, along with updated information and services, is located on the World Wide Web at:

[/content/early/2017/03/23/peds.2017-0150.full.html](http://content.early/2017/03/23/peds.2017-0150.full.html)

- Annelerin emzirmeyi seçme hakkına ve seçim yaptıkları taktirde başarılı olma şansını yakalama için yeterli kaynaklara sahip olmalarını nasıl sağlayabiliriz?



Interventions for promoting the initiation of breastfeeding (Review)

Balogun OO, O'Sullivan EJ, McFadden A, Ota E, Gavine A, Garner CD, Renfrew MJ, MacGillivray S


- Yedi ülkede 107.362 kadının yer aldığı ve 28 araştırmanın incelendiği çalışmada, ebe, hemşire ve doktorlar da dahil olmak üzere emzirme eğitimi alan sağlık uzmanları ve eğitilmiş gönüllüler, bebeklerini emzirmeye başlayan kadınların sayısını artırmak için eğitim oturumları düzenleyebilir, danışma ve mesleki destek sağlayabilir. Düşük ve orta gelirli ülkelerde hangi stratejilerin kadınları doğumdan hemen sonra emzirmeye teşvik edeceğini bulmak için daha fazla çalışma yapılması gerekmektedir.



«Breast milk makes the world healthier, smarter, and more equal: these are the conclusions of a new Lancet Series on breastfeeding. The deaths of 823 000 children and 20 000 mothers each year could be averted through universal breastfeeding, along with economic savings of US\$300 billion.»

Emzirme konusunda yeni Lancet Serisinin sonuçları: **Anne sütü, dünyayı daha sağlıklı, daha akıllı ve adil bir yer haline getirir!**
Her yıl 823 000 çocuğun ve 20.000 annenin ölümünün önlenmesi ile birlikte 300 milyar dolar tasarrufun sağlanması evrensel düzeyde emzirmeye gerçekleştirilebilir.

The lancet



Emzirme, çocuklar, kadınlar ve toplum için kısa vadeli ve uzun vadeli sağlık, ekonomik ve çevresel avantajlar sağlar.



Bu kazanımları gerçekleştirmek, emzirmeyi korumak, teşvik etmek ve desteklemek için politik destek ve finansal yatırıma ihtiyaç vardır.



Sevgi,bağlılık

Vitamin

Kemer

İlaç

Can suyu

Mucize

Anne sütü.....gibidir/benzer

Merhamet

Bağ

Büyük haz

Şifa

Oksijen

Anne sütü deyince aklınıza ne geliyor diye sorduk;



«Olmazsa ölmezsin ama olursa bir başka yaşarsın»

«Anneciğim emzir, sütünle karnımı sevginle ruhumu doyur»

«Ana kucağı gibi güvenli, saf, temiz bir aşı, bebeğini sütünle sağlıklı yarınlara taşı»

«Tek katkı maddesi sevgi olan gıda»

ANNE SÜTÜ....



«Bebek için en iyisi
Var mı anne sütü gibisi
İçinde anne sevgisi
Her bebeğe gerekli

Bebeğin ilk aşısı
Kalmaz ağrı sızısı
Şifa her damlası
Hem doğal, hem sağlıklı
Sevgi dolu anne sütü. . . .»

Berna Müge ÇNAR

Emzirme

- Bebek,
- Anne,
- Aile,
- Toplum,
- Ekonomí ve Çevre için en iyi seçenektir.

