

DUALITY IN MONOIDAL CATEGORIES

A tale of how details are important sometimes

Based on `arXiv:2301.03545` with Sebastian Halbig.




2023-05-23

Tony Zorman

`tony.zorman@tu-dresden.de`

The Goal: establishing connections



Notions of duality

Closed monoidal categories

Definition

A (strict) monoidal category $(\mathcal{C}, \otimes, 1)$ is called *(left) closed*, if for every $x \in \mathcal{C}$ there exists an adjunction

$$- \otimes x : \mathcal{C} \rightleftarrows \mathcal{C} : [x, -]$$

with unit $\eta^x : - \Rightarrow [x, - \otimes x]$ and counit $\varepsilon^x : [x, -] \otimes x \Rightarrow -$.

Closed monoidal categories

Definition

A (strict) monoidal category $(\mathcal{C}, \otimes, 1)$ is called *(left) closed*, if for every $x \in \mathcal{C}$ there exists an adjunction

$$- \otimes x : \mathcal{C} \rightleftarrows \mathcal{C} : [x, -]$$

with unit $\eta^x : - \Rightarrow [x, - \otimes x]$ and counit $\varepsilon^x : [x, -] \otimes x \Rightarrow -$.

Example

(Cat, \times) ,

Closed monoidal categories

Definition

A (strict) monoidal category $(\mathcal{C}, \otimes, 1)$ is called *(left) closed*, if for every $x \in \mathcal{C}$ there exists an adjunction

$$- \otimes x : \mathcal{C} \rightleftarrows \mathcal{C} : [x, -]$$

with unit $\eta^x : - \Rightarrow [x, - \otimes x]$ and counit $\varepsilon^x : [x, -] \otimes x \Rightarrow -$.

Example

(\mathbf{Cat}, \times) , $(\mathbf{k}\text{-Mod}, \otimes_{\mathbf{k}})$,

Closed monoidal categories

Definition

A (strict) monoidal category $(\mathcal{C}, \otimes, 1)$ is called *(left) closed*, if for every $x \in \mathcal{C}$ there exists an adjunction

$$- \otimes x : \mathcal{C} \rightleftarrows \mathcal{C} : [x, -]$$

with unit $\eta^x : - \Longrightarrow [x, - \otimes x]$ and counit $\varepsilon^x : [x, -] \otimes x \Longrightarrow -$.

Example

(\mathbf{Cat}, \times) , $(\mathbf{k}\text{-Mod}, \otimes_{\mathbf{k}})$, $([\mathcal{C}^{\text{op}}, \mathbf{Set}], \star)$,

Closed monoidal categories

Definition

A (strict) monoidal category $(\mathcal{C}, \otimes, 1)$ is called *(left) closed*, if for every $x \in \mathcal{C}$ there exists an adjunction

$$- \otimes x : \mathcal{C} \rightleftarrows \mathcal{C} : [x, -]$$

with unit $\eta^x : - \Longrightarrow [x, - \otimes x]$ and counit $\varepsilon^x : [x, -] \otimes x \Longrightarrow -$.

Example

(\mathbf{Cat}, \times) , $(\mathbf{k}\text{-Mod}, \otimes_{\mathbf{k}})$, $([\mathcal{C}^{\text{op}}, \mathbf{Set}], \star)$, $(\mathbb{S}\text{-Mod}, \circ)$.

(Co)evaluation morphisms in closed categories

Every object x in a closed category \mathcal{C} is equipped with natural *(co)evaluation* morphisms

$$\mathrm{coev}_x := \eta_1^x: 1 \longrightarrow [x, x],$$

$$\mathrm{ev}_x := \varepsilon_1^x: [x, 1] \otimes x \longrightarrow 1$$

(Co)evaluation morphisms in closed categories

Every object x in a closed category \mathcal{C} is equipped with natural *(co)evaluation* morphisms

$$\mathrm{coev}_x := \eta_1^x: 1 \longrightarrow [x, x],$$

$$\mathrm{ev}_x := \varepsilon_1^x: [x, 1] \otimes x \longrightarrow 1$$

satisfying e.g.,

$$\begin{array}{ccc} x & \xrightarrow{\eta_1^x \otimes x} & [x, x] \otimes x \\ & \searrow \mathrm{id}_x & \downarrow \varepsilon_x^x \\ & & x \end{array}$$

(Co)evaluation morphisms in closed categories

Every object x in a closed category \mathcal{C} is equipped with natural *(co)evaluation* morphisms

$$\mathrm{coev}_x := \eta_1^x: 1 \longrightarrow [x, x], \qquad \mathrm{ev}_x := \varepsilon_1^x: [x, 1] \otimes x \longrightarrow 1$$

satisfying e.g.,

$$\begin{array}{ccc} x & \xrightarrow{\eta_1^x \otimes x} & [x, x] \otimes x \\ & \searrow \mathrm{id}_x & \downarrow \varepsilon_x^x \\ & & x \end{array}$$

We need a coherent isomorphism $\phi_x: [x, x] \longrightarrow x \otimes [x, 1]$.

Rigid monoidal categories

Definition

A closed monoidal category \mathcal{C} is *(left) rigid monoidal*

Rigid monoidal categories

Definition

A closed monoidal category \mathcal{C} is *(left) rigid monoidal* if for every $x \in \mathcal{C}$ there exists a natural isomorphism $\phi_y^x: [x, y] \xrightarrow{\sim} y \otimes [x, 1]$,

Rigid monoidal categories

Definition

A closed monoidal category \mathcal{C} is *(left) rigid monoidal* if for every $x \in \mathcal{C}$ there exists a natural isomorphism $\phi_y^x: [x, y] \xrightarrow{\sim} y \otimes [x, 1]$, compatible with the evaluation and coevaluation; e.g.,

$$\begin{array}{ccc} [x, x] \otimes x & & \\ \phi_x^x \otimes x \downarrow & \searrow \varepsilon_x^x & \\ x \otimes [x, 1] \otimes x & \xrightarrow{x \otimes \varepsilon_1^x} & x \end{array}$$

Rigid monoidal categories

Definition

A closed monoidal category \mathcal{C} is *(left) rigid monoidal* if for every $x \in \mathcal{C}$ there exists a natural isomorphism $\phi_y^x: [x, y] \xrightarrow{\sim} y \otimes \overset{\vee}{x}$, compatible with the evaluation and coevaluation; e.g.,

$$\begin{array}{ccc} [x, x] \otimes x & & \\ \phi_x^x \otimes x \downarrow & \searrow \varepsilon_x^x & \\ x \otimes \overset{\vee}{x} \otimes x & \xrightarrow{x \otimes \varepsilon_1^x} & x \end{array}$$

Rigid monoidal categories

Definition

A closed monoidal category \mathcal{C} is *(left) rigid monoidal* if for every $x \in \mathcal{C}$ there exists a natural isomorphism $\phi_y^x: [x, y] \xrightarrow{\sim} y \otimes {}^{\vee}x$, compatible with the evaluation and coevaluation; e.g.,

$$\begin{array}{ccc} [x, x] \otimes x & & \\ \phi_x^x \otimes x \downarrow & \searrow \varepsilon_x^x & \\ x \otimes {}^{\vee}x \otimes x & \xrightarrow{x \otimes \varepsilon_1^x} & x \end{array} \quad \rightsquigarrow \quad \begin{array}{ccc} x & \xrightarrow{\text{coev}_x \otimes x} & x \otimes {}^{\vee}x \otimes x \\ & \searrow \text{id}_x & \downarrow x \otimes \text{ev}_x \\ & & x \end{array}$$

Rigid monoidal categories

Definition

A closed monoidal category \mathcal{C} is *(left) rigid monoidal* if for every $x \in \mathcal{C}$ there exists a natural isomorphism $\phi_y^x: [x, y] \xrightarrow{\sim} y \otimes {}^{\vee}x$, compatible with the evaluation and coevaluation; e.g.,

$$\begin{array}{ccc} [x, x] \otimes x & & \\ \phi_x^x \otimes x \downarrow & \searrow \varepsilon_x^x & \\ x \otimes {}^{\vee}x \otimes x & \xrightarrow{x \otimes \varepsilon_1^x} & x \end{array} \quad \rightsquigarrow \quad \begin{array}{ccc} x & \xrightarrow{\text{coev}_x \otimes x} & x \otimes {}^{\vee}x \otimes x \\ & \searrow \text{id}_x & \downarrow x \otimes \text{ev}_x \\ & & x \end{array}$$

Example

$(\text{vect}_k, \otimes_k),$

Rigid monoidal categories

Definition

A closed monoidal category \mathcal{C} is *(left) rigid monoidal* if for every $x \in \mathcal{C}$ there exists a natural isomorphism $\phi_y^x: [x, y] \xrightarrow{\sim} y \otimes {}^{\vee}x$, compatible with the evaluation and coevaluation; e.g.,

$$\begin{array}{ccc}
 [x, x] \otimes x & & \\
 \phi_x^x \otimes x \downarrow & \searrow \varepsilon_x^x & \\
 x \otimes {}^{\vee}x \otimes x & \xrightarrow{x \otimes \varepsilon_1^x} & x
 \end{array}
 \quad \rightsquigarrow \quad
 \begin{array}{ccc}
 x & \xrightarrow{\text{coev}_x \otimes x} & x \otimes {}^{\vee}x \otimes x \\
 \searrow \text{id}_x & & \downarrow x \otimes \text{ev}_x \\
 & & x
 \end{array}$$

Example

$$(\text{vect}_k, \otimes_k), \text{Ad}_C^\infty \leq ([C, C], \circ).$$

Grothendieck–Verdier categories

Definition

A *Grothendieck–Verdier* (also called **-autonomous*) category comprises a monoidal category \mathcal{C} and an object $d \in \mathcal{C}$,

Grothendieck–Verdier categories

Definition

A *Grothendieck–Verdier* (also called **-autonomous*) category comprises a monoidal category \mathcal{C} and an object $d \in \mathcal{C}$, such that there exists an equivalence $D: \mathcal{C}^{\text{op}} \rightarrow \mathcal{C}$, and a natural isomorphism

$$\mathcal{C}(x \otimes y, d) \cong \mathcal{C}(x, Dy).$$

Grothendieck–Verdier categories

Definition

A *Grothendieck–Verdier* (also called **-autonomous*) category comprises a monoidal category \mathcal{C} and an object $d \in \mathcal{C}$, such that there exists an equivalence $D: \mathcal{C}^{\text{op}} \rightarrow \mathcal{C}$, and a natural isomorphism

$$\mathcal{C}(Dy \otimes y, d) \cong \mathcal{C}(Dy, Dy).$$

Grothendieck–Verdier categories

Definition

A *Grothendieck–Verdier* (also called **-autonomous*) category comprises a monoidal category \mathcal{C} and an object $d \in \mathcal{C}$, such that there exists an equivalence $D: \mathcal{C}^{\text{op}} \rightarrow \mathcal{C}$, and a natural isomorphism

$$\mathcal{C}(Dy \otimes y, d) \cong \mathcal{C}(Dy, Dy) \quad \frac{\text{id}: Dy \rightarrow Dy}{\quad}$$

Grothendieck–Verdier categories

Definition

A *Grothendieck–Verdier* (also called **-autonomous*) category comprises a monoidal category \mathcal{C} and an object $d \in \mathcal{C}$, such that there exists an equivalence $D: \mathcal{C}^{\text{op}} \rightarrow \mathcal{C}$, and a natural isomorphism

$$\mathcal{C}(Dy \otimes y, d) \cong \mathcal{C}(Dy, Dy) \qquad \frac{\text{id}: Dy \rightarrow Dy}{\text{ev}_y: Dy \otimes y \rightarrow d}$$

Grothendieck–Verdier categories

Definition

A *Grothendieck–Verdier* (also called **-autonomous*) category comprises a monoidal category \mathcal{C} and an object $d \in \mathcal{C}$, such that there exists an equivalence $D: \mathcal{C}^{\text{op}} \rightarrow \mathcal{C}$, and a natural isomorphism

$$\mathcal{C}(x \otimes y, d) \cong \mathcal{C}(x, Dy)$$

Example

$(\Lambda M, \cap, 0),$

Grothendieck–Verdier categories

Definition

A *Grothendieck–Verdier* (also called **-autonomous*) category comprises a monoidal category \mathcal{C} and an object $d \in \mathcal{C}$, such that there exists an equivalence $D: \mathcal{C}^{\text{op}} \rightarrow \mathcal{C}$, and a natural isomorphism

$$\mathcal{C}(x \otimes y, d) \cong \mathcal{C}(x, Dy)$$

Example

$(\Lambda M, \cap, 0)$, $(\text{Chu}(\mathcal{C}, d), \otimes_{\mathcal{C}}, d)$,

Grothendieck–Verdier categories

Definition

A *Grothendieck–Verdier* (also called **-autonomous*) category comprises a monoidal category \mathcal{C} and an object $d \in \mathcal{C}$, such that there exists an equivalence $D: \mathcal{C}^{\text{op}} \rightarrow \mathcal{C}$, and a natural isomorphism

$$\mathcal{C}(x \otimes y, d) \cong \mathcal{C}(x, Dy)$$

Example

$(\Lambda M, \cap, 0)$, $(\text{Chu}(\mathcal{C}, d), \otimes_{\mathcal{C}}, d)$, $(eCe, \otimes_{\mathcal{C}}, De)$,

Grothendieck–Verdier categories

Definition

A *Grothendieck–Verdier* (also called **-autonomous*) category comprises a monoidal category \mathcal{C} and an object $d \in \mathcal{C}$, such that there exists an equivalence $D: \mathcal{C}^{\text{op}} \rightarrow \mathcal{C}$, and a natural isomorphism

$$\mathcal{C}(x \otimes y, d) \cong \mathcal{C}(x, Dy)$$

Example

$(\Lambda M, \cap, 0)$, $(\text{Chu}(\mathcal{C}, d), \otimes_{\mathcal{C}}, d)$, $(eCe, \otimes_{\mathcal{C}}, De)$, $(\text{ban}_{\mathbf{R}}, \otimes_{\mathbf{R}}, \mathbf{R})$.

Tensor representability

Every rigid category yields adjunctions

$$- \otimes x : \mathcal{C} \rightleftarrows \mathcal{C} : - \otimes {}^{\vee}x.$$

Tensor representability

Starting with an adjunction

$$- \otimes x : \mathcal{C} \rightleftarrows \mathcal{C} : - \otimes Rx, \quad (1)$$

what can we say about the category \mathcal{C} ?

Tensor representability

Starting with an adjunction

$$- \otimes x : \mathcal{C} \rightleftarrows \mathcal{C} : - \otimes Rx, \quad (1)$$


what can we say about the category \mathcal{C} ?

Definition


A monoidal category is *tensor representable*, if for every $x \in \mathcal{C}$ there is an adjunction as in Equation (1).

Tensor representability in context


Relationships among different types of duality




Relationships among different types of duality




Relationships among different types of duality




Relationships among different types of duality



Relationships among different types of duality



Relationships among different types of duality



Tensor representable categories are Grothendieck–Verdier

Theorem ([HZ23, Theorem 3.2])

If for every $x \in \mathcal{C}$, there exist adjunctions

$$- \otimes Lx \dashv - \otimes x \dashv - \otimes Rx,$$

then \mathcal{C} is Grothendieck–Verdier.

Tensor representable categories are Grothendieck–Verdier

Theorem ([HZ23, Theorem 3.2])

If for every $x \in \mathcal{C}$, there exist adjunctions

$$- \otimes Lx \dashv - \otimes x \dashv - \otimes Rx,$$

then \mathcal{C} is Grothendieck–Verdier.

Proof.

The assumption implies that L is a quasi-inverse of R ; for example,

$$\mathcal{C}(LRx, y) \cong \mathcal{C}(1, y \otimes Rx) \cong \mathcal{C}(x, y).$$

Tensor representable categories are Grothendieck–Verdier

Theorem ([HZ23, Theorem 3.2])

If for every $x \in C$, there exist adjunctions

$$- \otimes Lx \dashv - \otimes x \dashv - \otimes Rx,$$

then C is Grothendieck–Verdier.

Proof.

The assumption implies that L is a quasi-inverse of R ; for example,

$$C(LRx, y) \cong C(1, y \otimes Rx) \cong C(x, y).$$

Set $D := R$.

Tensor representable categories are Grothendieck–Verdier

Theorem ([HZ23, Theorem 3.2])

If for every $x \in \mathcal{C}$, there exist adjunctions

$$- \otimes Lx \dashv - \otimes x \dashv - \otimes Rx,$$

then \mathcal{C} is Grothendieck–Verdier.

Proof.

The assumption implies that L is a quasi-inverse of R ; for example,

$$\mathcal{C}(LRx, y) \cong \mathcal{C}(1, y \otimes Rx) \cong \mathcal{C}(x, y).$$

Set $D := R$. In order to show that \mathcal{C} is a Grothendieck–Verdier category, we need to find some $d \in \mathcal{C}$ such that

$$\mathcal{C}(y \otimes x, d) \cong \mathcal{C}(y, Dx).$$



Tensor representable categories are Grothendieck–Verdier

Theorem ([HZ23, Theorem 3.2])

If for every $x \in \mathcal{C}$, there exist adjunctions

$$- \otimes Lx \dashv - \otimes x \dashv - \otimes Rx,$$

then \mathcal{C} is Grothendieck–Verdier.

Proof.

The assumption implies that L is a quasi-inverse of R ; for example,

$$\mathcal{C}(LRx, y) \cong \mathcal{C}(1, y \otimes Rx) \cong \mathcal{C}(x, y).$$

Set $D := R$. In order to show that \mathcal{C} is a Grothendieck–Verdier category, we need to find some $d \in \mathcal{C}$ such that

$$\mathcal{C}(y \otimes x, d) \cong \mathcal{C}(y, Dx).$$



Tensor representable categories are Grothendieck–Verdier

Theorem ([HZ23, Theorem 3.2])

If for every $x \in \mathcal{C}$, there exist adjunctions

$$- \otimes Lx \dashv - \otimes x \dashv - \otimes Rx,$$

then \mathcal{C} is Grothendieck–Verdier.

Proof.

The assumption implies that L is a quasi-inverse of R ; for example,

$$\mathcal{C}(LRx, y) \cong \mathcal{C}(1, y \otimes Rx) \cong \mathcal{C}(x, y).$$

Set $D := R$. In order to show that \mathcal{C} is a Grothendieck–Verdier category, we need to find some $d \in \mathcal{C}$ such that

$$\mathcal{C}(y \otimes x, d) \cong \mathcal{C}(y, d \otimes Dx).$$



Tensor representable categories are Grothendieck–Verdier

Theorem ([HZ23, Theorem 3.2])

If for every $x \in \mathcal{C}$, there exist adjunctions

$$- \otimes Lx \dashv - \otimes x \dashv - \otimes Rx,$$

then \mathcal{C} is Grothendieck–Verdier.

Proof.

The assumption implies that L is a quasi-inverse of R ; for example,

$$\mathcal{C}(LRx, y) \cong \mathcal{C}(1, y \otimes Rx) \cong \mathcal{C}(x, y).$$

Set $D := R$. In order to show that \mathcal{C} is a Grothendieck–Verdier category, we need to find some $d \in \mathcal{C}$ such that


$$\mathcal{C}(y \otimes x, 1) \cong \mathcal{C}(y, Dx).$$



**Is tensor representability and
rigidity the same thing?**

The problem in string diagrams


The rigid case



$$(x \otimes \text{ev}_x) \circ (\text{coev}_x \otimes x) = \text{id}_x$$


The problem in string diagrams

The rigid case



$$(x \otimes \text{ev}_x) \circ (\text{coev}_x \otimes x) = \text{id}_x$$

The tensor representable case



$$\epsilon_x^x \circ (\eta_1^x \otimes x) = \text{id}_x$$

A negative result


Theorem ([HZ23, Theorems 2.5 and 4.15])

There exist categories that are tensor representable, but not rigid.

A negative result

Theorem ([HZ23, Theorems 2.5 and 4.15])


There exist categories that are tensor representable, but not rigid.



A negative result

Theorem ([HZ23, Theorems 2.5 and 4.15])

There exist categories that are tensor representable, but not rigid.



Mackey functors

Definition

Let k be a field.

Mackey functors

Definition

Let k be a field. A
functor $M: \text{Sp}_G \longrightarrow \text{vect}_k$.

Mackey functor M is a coproduct preserving

Mackey functors

Definition

Let k be a field. A *finite-dimensional Mackey functor* M is a coproduct preserving functor $M: \mathbf{Sp}_G \longrightarrow \mathbf{vect}_k$, such that Mx is finite-dimensional for every $x \in \mathbf{Sp}_G$.

Mackey functors

Definition

Let k be a field. A *finite-dimensional Mackey functor* M is a coproduct preserving functor $M: \mathbf{Sp}_G \longrightarrow \mathbf{vect}_k$, such that Mx is finite-dimensional for every $x \in \mathbf{Sp}_G$.

Theorem

The category \mathbf{mky} of finite-dimensional Mackey functors is closed monoidal:

Mackey functors

Definition

Let k be a field. A *finite-dimensional Mackey functor* M is a coproduct preserving functor $M: \mathbf{Sp}_G \rightarrow \mathbf{vect}_k$, such that Mx is finite-dimensional for every $x \in \mathbf{Sp}_G$.

Theorem

The category \mathbf{mky} of finite-dimensional Mackey functors is closed monoidal:

$$(M \star N)x := \int^{a,b} k\mathbf{Sp}_G(a \otimes b, x) \otimes_k Ma \otimes_k Nb,$$

$$[M, N]x := \int_{a,b} \mathbf{vect}_k(k\mathbf{Sp}_G(x \otimes a, b), \mathbf{vect}_k(Ma, Nb)) \cong \mathbf{mky}(M(x^* \otimes -), N).$$

Mackey functors in positive characteristic

Theorem ([HZ23, Theorem 4.15])

The category mky is tensor representable, but not rigid.

Mackey functors in positive characteristic

Theorem ([HZ23, Theorem 4.15])

The category mky is tensor representable, but not rigid.

Proof idea.

1. In vect_k : $\int_y \int^x \cong \int^x \int_y$.

Mackey functors in positive characteristic

Theorem ([HZ23, Theorem 4.15])

The category mky is tensor representable, but not rigid.

Proof idea.

1. In vect_k : $\int_y \int^x \cong \int^x \int_y$.
2. By [Day06]: $\int^x \cong \int_y$.

Mackey functors in positive characteristic

Theorem ([HZ23, Theorem 4.15])

The category mky is tensor representable, but not rigid.

Proof idea.

1. In vect_k : $\int_y \int^x \cong \int^x \int_y$.
2. By [Day06]: $\int^x \cong \int_y$.
3. By [PS07, Section 9]: mky is * -autonomous.

Mackey functors in positive characteristic

Theorem ([HZ23, Theorem 4.15])

The category mky is tensor representable, but not rigid.

Proof idea.

1. In vect_k : $\int_y \int^x \cong \int^x \int_y$.
2. By [Day06]: $\int^x \cong \int_y$.
3. By [PS07, Section 9]: mky is * -autonomous.
4. Calculation: it is even tensor representable.

Mackey functors in positive characteristic

Theorem ([HZ23, Theorem 4.15])

The category mky is tensor representable, but not rigid.

Proof idea.

1. In vect_k : $\int_y \int^x \cong \int^x \int_y$.
2. By [Day06]: $\int^x \cong \int_y$.
3. By [PS07, Section 9]: mky is * -autonomous.
4. Calculation: it is even tensor representable.
5. By [Bou05, Lemma 2.2]: mky is rigid iff all short exact sequences split.

Mackey functors in positive characteristic

Theorem ([HZ23, Theorem 4.15])

The category mky is tensor representable, but not rigid.

Proof idea.

1. In vect_k : $\int_y \int^x \cong \int^x \int_y$.
2. By [Day06]: $\int^x \cong \int_y$.
3. By [PS07, Section 9]: mky is * -autonomous.
4. Calculation: it is even tensor representable.
5. By [Bou05, Lemma 2.2]: mky is rigid iff all short exact sequences split.
6. If the order of G divides the characteristic of k , then mky is not rigid.



Thank you for your attention!

arXiv:2301.03545



- [Bou05] Serge Bouc. “**The Burnside dimension of projective Mackey functors**”. English. In: *RIMS Kôkyûroku* 1440 (2005), pp. 107–120. ISSN: 1880-2818.
- [Day06] Brian J. Day. “**Compact convolution**”. In: *arXiv e-prints* (2006). arXiv: math/0605463 [math.CT].
- [HZ23] Sebastian Halbig and Tony Zorman. “**Duality in Monoidal Categories**”. In: *arXiv e-prints* (2023). arXiv: 2301.03545 [math.CT].
- [PS07] Elango Panchadcharam and Ross Street. “**Mackey functors on compact closed categories**”. English. In: *J. Homotopy Relat. Struct.* 2.2 (2007), pp. 261–293. ISSN: 2193-8407.


Definition

Let G be a finite group, and write $G\text{-set}$ for the category of finite G -sets.

Spans


Definition

Let G be a finite group, and write $G\text{-set}$ for the category of finite G -sets. The category Sp_G of *spans* of finite G -sets has, as objects, finite G -sets, and as morphisms *isomorphism classes of spans*



The explicit construction


Define a category \mathcal{D} whose objects are natural numbers, with morphisms generated by compositions and tensor products of the identities, as well as



subject to appropriate naturality conditions.

The explicit construction


Define a category \mathcal{D} whose objects are natural numbers, with morphisms generated by compositions and tensor products of the identities, as well as



subject to appropriate naturality conditions. Define a *monoidal localisation* \mathcal{C} of \mathcal{D} , in which adjunction properties for ε and η hold.

The explicit construction

Define a category \mathcal{D} whose objects are natural numbers, with morphisms generated by compositions and tensor products of the identities, as well as



subject to appropriate naturality conditions. Define a *monoidal localisation* \mathcal{C} of \mathcal{D} , in which adjunction properties for ε and η hold. By studying strong monoidal functors $\mathcal{C} \rightarrow \mathbf{vect}_k$, we find that the snake equation and the identity must lie in different equivalence classes.