

algebraic topology exercises

Contents

Homework 1	1
0 Preliminaries	1
1 Based spaces and smash product	3
2 Mapping cylinders and Hurewicz cofibrations	4
3 Path spaces and fibrations	6
Homework 1.5	8
1 Exercise on model categories	8
2 Hatcher's exercise on Whitehead's theorem	10
Homework 2	10
cohomology ring of \mathbb{CP}^n	12
References	13

Homework 1

0 Preliminaries

In the category of sets there is a bijection $\text{Hom}(X \times Y, Z) \cong \text{Hom}(X, \text{Hom}(Y, Z))$ that depends naturally on X , Y and Z . The notions related to this bijection are “Cartesian closed category”, “currying” and “internal Hom”.

Definition. A category \mathcal{C} is *Cartesian closed* if:

1. \mathcal{C} has all finite products (Caveat: some require that \mathcal{C} has all finite limits)
2. For any object Y the functor $- \times Y$ has a right adjoint, which we will denote by $\text{Map}(Y, -)$ or by $-^Y$.

Remark. By section 3 [here](#), the second property above implies that we get a functor $\text{Map}(-, -) : \mathcal{C}^{\text{op}} \times \mathcal{C} \rightarrow \mathcal{C}$, and moreover we get natural isomorphisms $\text{Hom}(X, \text{Map}(Y, Z)) \cong \text{Hom}(X \times Y, Z)$ and $\text{Map}(X, \text{Map}(Y, Z)) \cong \text{Map}(X \times Y, Z)$.

Lemma (Yoneda, [wiki](#)). Let F be a functor from a locally small category \mathcal{C} to Set . Then for each object X of \mathcal{C} , the natural transformations $\text{Nat}(\text{Hom}(X, -), F)$ are in one-to-one correspondence with the elements of $F(X)$, that is

$$\text{Nat}(\text{Hom}(X, -), F) \cong F(X)$$

Moreover, this isomorphism is natural in X and F when both sides are regarded as functors from $\mathcal{C} \times \text{Set}^{\mathcal{C}}$ to Set . ($\text{Set}^{\mathcal{C}}$ denotes the category of functors from \mathcal{C} to Set .)

There is a contravariant version of Yoneda lemma asserting that if F is a contravariant functor from \mathcal{C} to \mathbf{Set} ,

$$\mathbf{Nat}(\mathbf{Hom}(-, X), F) \cong F(X).$$

Corollary. $\mathbf{Nat}(\mathbf{Hom}(-, X), \mathbf{Hom}(-, Y)) \cong \mathbf{Hom}(X, Y)$.

Remark. The correspondence $X \mapsto \mathbf{Hom}(-, X)$ is fully faithful, that is, the correspondence $\mathbf{Hom}(X, X') \rightarrow \mathbf{Nat}(\mathbf{Hom}(-, X), \mathbf{Hom}(-, X'))$ is injective and bijective.

Exercise (a). Let \mathcal{C} be any category. Show that if for some objects X and X' we have $\mathbf{Hom}(X, Y) \cong \mathbf{Hom}(X', Y)$ for all objects Y , with isomorphisms being natural in Y , then $X \cong X'$. Dually, if $\mathbf{Hom}(Y, X) \cong \mathbf{Hom}(Y, X')$ naturally in Y , then also $X \cong X'$.

Solution. The latter correspondence sends isomorphisms to isomorphisms. Since we are given a natural isomorphism in the problem, we conclude $X \cong X'$. The dual statement follows from the analogue formulation of Yoneda lemma. \square

Exercise. Let \mathcal{C} be a Cartesian closed category and \mathbf{pt} be the terminal object. Show that for any object X we have $X \cong \mathbf{Map}(\mathbf{pt}, X)$.

Solution. Using item (a) with X and $X' = \mathbf{Map}(\mathbf{pt}, X)$, it suffices to show that

$$\mathbf{Hom}(Y, X) \cong \mathbf{Hom}(Y, \mathbf{Map}(\mathbf{pt}, X))$$

for all objects Y and isomorphisms natural in Y .


Since \mathcal{C} is Cartesian closed, we have isomorphisms **natural** in Y

$$\mathbf{Hom}(Y, \mathbf{Map}(\mathbf{pt}, X)) \cong \mathbf{Hom}(Y \times \mathbf{pt}, X) \cong \mathbf{Hom}(Y, X)$$

since \mathbf{pt} is a terminal object. Indeed:

Claim. In a Cartesian closed category \mathcal{C} with terminal object \mathbf{pt} , we have that $Y \times \mathbf{pt} \cong Y$ for any object Y .

Proof of claim. (**From StackExchange**) The universal property of the product $Y \times \mathbf{pt}$ shows that the maps \mathbf{id}_Y and $t_Y : Y \rightarrow \mathbf{pt}$ must factor through some $u : Y \rightarrow Y \times \mathbf{pt}$, making $\pi_1 \circ u = \mathbf{id}_Y$.




It is also true that $u \circ \pi_1 = \mathbf{id}_{Y \times \mathbf{pt}}$, since

- $\pi_1 \circ u \circ \pi_1 = \mathbf{id}_Y \circ \pi_1 = \pi_1$ and

- $t_{Y \times \text{pt}} \circ u \circ \pi_1 = t_{Y \times \text{pt}}$

so by uniqueness of the universal property we get that $u \circ \pi_1 = \text{id}_{Y \times \text{pt}}$.



□

□

1 Based spaces and smash product

Definition. The appropriate analogue of the Cartesian product in the category of based spaces is the **smash product** $X \wedge Y$ defined by

$$X \wedge Y = X \times Y / X \vee Y.$$

Here $X \vee Y$ is viewed as the subspace of $X \times Y$ consisting of those pairs (x, y) such that either x is the basepoint of X or y is the basepoint of Y .

Exercise. For a based space (X, x_0) let ΣX be $[0, 1] \times X / \{1\} \times X \cup \{0\} \times X \cup [0, 1] \times \{x_0\}$. Check that $\Sigma X \cong S^1 \wedge X$. In particular $S^n \cong S^1 \wedge S^{n-1} \cong (S^1)^{\wedge n}$.

Remark. Another way of defining the reduced suspension ΣX (**I think**) is

$$\Sigma X = (I \times X) / (t, x) \sim (0, y) \sim (1, y) \quad \forall y \in X.$$

Proof. To see that $\Sigma X \cong S^1 \wedge X$ simply notice that "both spaces are the quotient $X \times I$ with $X \times \partial I \cup \{x_0\} \times I$ collapsed to a point" (**Hatcher**, ex. 0.10). This is clear for ΣX . For $X \wedge S^1$, notice that collapsing $X \times \partial I$ to a point in $X \times I$ amounts to taking $X \times S^1$ and collapsing one copy of X to a point. Further, collapsing $x_0 \times I$ to a point amounts to collapsing the copy of S^1 in $X \vee S^1$ to a point.

Let's try induction on n . If $n = 2$, the smash product $S^1 \wedge S^1$ is easily seen to be S^2 since it consists on collapsing the boundary $S^1 \vee S^1$ of the square whose quotient yields $S^1 \times S^1$. For the inductive step **Still incomplete...** □

2 Mapping cylinders and Hurewicz cofibrations

Definition (wikipedia). Let X be a topological space and let $A \subset X$. We say that the pair (X, A) has the *homotopy extension property* if for any space Y , any homotopy $g_\bullet : A \rightarrow Y^I$ and any map $\tilde{g}_0 : X \rightarrow Y$ such that $\tilde{g}_0 \circ \iota = g_0$, there exists an *extension* of f_\bullet to a homotopy $\tilde{g}_\bullet : X \rightarrow Y^I$ such that $\tilde{g}_\bullet \circ \iota = g_\bullet$.

$$\begin{array}{ccc} A & \xrightarrow{g_\bullet} & Y^I \\ \downarrow \iota & \nearrow \tilde{g}_\bullet & \downarrow \pi_0 \\ X & \xrightarrow{\tilde{g}_0} & Y \end{array}$$

A *Hurewicz cofibration* is a map $\iota : A \rightarrow X$ satisfying the homotopy extension property.

Exercise (a). Prove that an inclusion $f : A \rightarrow X$ is a Hurewicz cofibration if and only if $A \times I \cup X \times \{0\}$ is a retract of $X \times I$.

Remark. A little late I noticed the comment on Telegram that we may assume A to be a closed subspace. Maybe I wouldn't have tried the solution following Miller if I had knew this earlier, hehe— still it was nice to see two different solutions.

Solution following Hatcher. (\implies) According to the former definition, choose $Y = (X \times \{0\}) \cup (A \times I)$. The inclusion $A \times I \hookrightarrow Y$ is an homotopy g_\bullet from A to Y . Also, the inclusion $X \times \{0\} \hookrightarrow Y$ is an extension \tilde{g}_0 . Then there exists an extension \tilde{g}_\bullet of the whole homotopy, which is just a map from $X \times I$ to Y . We have thus produced a retraction:

$$\begin{array}{ccc} (X \times \{0\}) \cup (A \times I) & \xrightarrow{\text{id}} & (X \times \{0\}) \cup (A \times I) = Y \\ \downarrow & \nearrow & \\ X \times I & & \end{array}$$

(\impliedby) Now suppose that $(X \times \{0\}) \cup (A \times I)$ is a retract of $X \times I$. Let Y be any space, $g_\bullet : A \rightarrow Y^I$ an homotopy and \tilde{g}_0 a map such that $\tilde{g}_0 = g_0 \circ f$.

The homotopy g_\bullet along with \tilde{g}_0 yield a map $\varphi : (A \times I) \cup (X \times \{0\}) \rightarrow Y \cup (X \times \{0\})$. The key observation is that if A is closed in X , then this map is continuous by the [gluing lemma](#). Then we simply compose the given retraction r with this map to obtain the homotopy extension:


$$\begin{array}{ccc} A \times I & \xrightarrow{g_\bullet} & Y \\ \downarrow & \nearrow \varphi & \\ X \times I & \xrightarrow{\tilde{g}_0} & Y \end{array}$$

$(X \times \{0\}) \cup (A \times I)$


\downarrow
 $X \times I$

A complicated argument in Hatcher's appendix shows that such a function is continuous even without the assumption that A is closed. \square

Solution following Miller. The homotopy extension property may be defined as a map $\iota : A \rightarrow X$ such that for any solid-arrow diagram as below, a dotted blue arrow exists making the whole diagram commute:



Now consider the pushout corresponding to ι and the inclusion $A \rightarrow A \times I$. By the universal property of the pushout, the former diagram must factor by the pushout, and we get the following diagram:



The implication (\implies) of our exercise again follows by choosing $Y = (X \times \{0\}) \cup (A \times I)$. For the implication (\impliedby) **it appears that we have the same problem as before**: we need to construct the blue dashed arrow from the rest of the diagram (using that the black dashed arrow has a left inverse), but it seems that the natural thing to do is defining this function from the two pieces just like before, and we must make sure it is continuous. \square

Definition. Let $f : X \rightarrow Y$ be a map. Let $M_f = X \times [0, 1] \cup_f Y$ be the *mapping cylinder of f* , i.e. the pushout of $X \xrightarrow{\cong} X \times \{0\} \hookrightarrow X \times [0, 1]$ and of $f : X \rightarrow Y$. Let $g : X \rightarrow M_f$ be the map $X \xrightarrow{\cong} X \times \{1\} \rightarrow M_f$. Let $h : M_f \rightarrow Y$ be the map that is induced by $X \times [0, 1] \rightarrow Y : (x, t) \mapsto f(x)$ and $\text{id}_Y : Y \rightarrow Y$. Observe that f is the composition of g and h .

Remark. In both exercises below you might have to use the fact that pushouts are colimits and that colimits commute with products in CGWH, i.e. $(\text{colim } A_i) \times B$ is canonically homeomorphic with $\text{colim}(A_i \times B)$.

Exercise.

- Show that h is a deformation retract, and in particular is a homotopy equivalence.
- Show that $g : X \rightarrow M_f$ is a cofibration. You may use exercise (a), but the direct proof might be simpler.

Solution.

b. We have that

$$\begin{array}{ccc}
 X \times \{0\} & \xrightarrow{f} & Y \\
 \text{id} \times 1 \downarrow & \searrow g & \downarrow \\
 X \times I & \xrightarrow{\quad} & M_f \\
 & \searrow \text{curved} & \downarrow \text{dashed } h \\
 & & Y
 \end{array}
 \quad
 \begin{array}{l}
 \text{curved } (x,t) \mapsto f(x) \\
 \text{curved } \text{id}_Y
 \end{array}$$

We must show that there is a homotopy between the identity map on M_f and a retraction from M_f to Y . So we want $h : M_f \times I \rightarrow M_f$ such that

$$h(-,0) = \text{id}_{M_f}, \quad \text{img } h(-,1) \subset Y \quad \text{and} \quad h(-,1)|_Y = \text{id}_Y$$

Since M_f is a pullback, we can see it as a colimit, that is

$$M_f = \text{colim}(X \times I \leftarrow X \rightarrow Y)$$

and, since colimits commute with products in CGWH, we get

$$M_f \times I = \text{colim}(X \times I \times I \leftarrow X \times I \rightarrow Y \times I)$$

that is,

$$\begin{array}{ccc}
 X \times \{0\} \times I & \longrightarrow & Y \times I \\
 \downarrow & \nearrow \ulcorner & \downarrow ? \\
 X \times I \times I & \xrightarrow{\quad ? \quad} & M_f \times I \\
 & \searrow \text{curved} & \downarrow \text{dashed} \\
 & & M_f
 \end{array}
 \quad
 \begin{array}{l}
 \text{curved } (x,t,s) \mapsto f(x) \\
 \text{curved } (y,s) \mapsto y
 \end{array}$$

[I certainly got stuck in concluding...]

c. [Also in progress...] Consider the following lifting problem:

$$\begin{array}{ccc}
 X & \xrightarrow{H} & Z^I \\
 g \downarrow & \nearrow \text{dashed} & \downarrow \pi_0 \\
 M_f & \xrightarrow{h} & Z
 \end{array}$$

□

3 Path spaces and fibrations

Exercise.

- a. Show that $\text{Map}(I, Y)$ deformation retracts on $\text{Map}(\text{pt}, Y)$. Most likely you'll have to find a correct map $I \times I \rightarrow I$. Also show that $\text{Map}(I, Y) \rightarrow \text{Map}(\text{pt}, Y)$ is a Hurewicz fibration. The key map will be of the form $I \times I \rightarrow I \times I$.

Solution.

- a. (**$\text{Map}(I, Y) \rightarrow \text{Map}(\text{pt}, Y)$ is a Hurewicz fibration.**) Let A be any space. We must show that for any homotopy H and lift h_0 there exists an homotopy \tilde{H} as in the following diagram:

$$\begin{array}{ccc} A \times \{0\} & \xrightarrow{h_0} & \text{Map}(I, Y) \\ \downarrow & \nearrow \tilde{H} & \downarrow p \\ A \times I & \xrightarrow{H} & \text{Map}(0, Y) \end{array}$$

From the isomorphism $\text{Map}(X \times Y, Z) \cong \text{Map}(X, \text{Map}(Y, Z))$ we may rewrite the problem as

$$\begin{array}{ccc} (A \times \{0\}) \times I & & \\ & \searrow H_0 & \\ (A \times I) \times I & \xrightarrow{\tilde{H}} & Y \\ & \nearrow H & \\ (A \times I) \times \{0\} & & \end{array}$$

So we define the dashed arrow by

$$(a, s, t) \mapsto \begin{cases} H_0(a, 0, s - t) & \text{when } s - t \geq 0 \\ H(a, t - s, 0) & \text{when } s - t \leq 0 \end{cases}$$

so that when $s = t$ the functions coincide, when $s = 0$ we get H and when $t = 0$ we get H_0 .

(**$\text{Map}(I, Y)$ deformation retracts on $\text{Map}(\text{pt}, Y)$.**) We must show there is a homotopy

$$h : \text{Map}(I, Y) \times I \longrightarrow \text{Map}(I, Y)$$

such that

$$h(-, 0) = \text{id}_{\text{Map}(I, Y)}, \quad h(-, 1) \subset \text{Map}(\text{pt}, Y)$$

$$\text{and} \quad h(-, 1)|_{\text{Map}(\text{pt}, Y)} = \text{id}_{\text{Map}(\text{pt}, Y)}.$$

Consider the map

$$\begin{aligned} I \times I &\rightarrow I \\ (s, t) &\mapsto s - st \end{aligned}$$

Our deformation retract may be written like

$$h : \text{Map}(I, Y) \times I \longrightarrow \text{Map}(I, Y)$$

$$(f(s), t) \longmapsto f(s - st)$$

Then for $t = 0$ we have the identity on $\text{Map}(I, Y)$, and when $t = 1$ we have ev_0 .

□

Let $f : X \rightarrow Y$ be a map. Let E_f be the pullback of $f : X \rightarrow Y$ and of $\text{ev}_0 : \text{Map}(I, Y) \rightarrow Y$. Let $h : X \rightarrow E_f$ be the map that sends x to $(x, \text{const}(f(x)))$, where $\text{const}(f(x)) : I \rightarrow Y$ is the constant path at $f(x)$. Let $g : E_f \rightarrow Y$ be the composition of projection map $E_f \rightarrow \text{Map}(I, Y)$ with $\text{ev}_1 : \text{Map}(I, Y) \rightarrow Y$.

Homework 1.5

1 Exercise on model categories

Exercise (3.1.8 from Riehl). Verify that the class of morphisms \mathcal{L} characterized by the left lifting property against a fixed class of morphisms \mathcal{R} is closed under coproducts, closed under retracts, and contains the isomorphisms.

Solution. (Coproducts.) Suppose the maps $\ell_i : A_i \rightarrow B_i$ are in \mathcal{L} . Then their coproduct in the arrow category is the obvious map $\coprod A_i \rightarrow \coprod B_i$.

Explicitly, their coproduct is an arrow $\coprod \ell_i$ and a collection of maps $f_i : \ell_i \rightarrow \coprod \ell_i$ such that for any other object $m : A \rightarrow B$ in the arrow category and a map $g : \ell \rightarrow m$, the following diagram is completed uniquely:

$$\begin{array}{ccc} \ell_i & \xrightarrow{f_i} & \coprod \ell_i \xrightarrow{\exists!} m \\ & \searrow g & \uparrow \end{array} \quad \forall i$$

So we conclude that the source of $\coprod \ell_i$ is $\coprod A_i$ and its target $\coprod B_i$. Indeed, we really looking at

$$\begin{array}{ccc} A_i & \xrightarrow{\ell_i} & B_i \\ f_i^! \downarrow & & \downarrow f_i^2 \\ \coprod A_i & \xrightarrow{\coprod \ell_i} & \coprod B_i \\ \exists! \downarrow & & \downarrow \exists! \\ A & \xrightarrow{m} & B \end{array}$$

Now consider the following lifting problem with respect to a morphism $r \in \mathcal{R}$:

$$\begin{array}{ccc} \coprod A_i & \longrightarrow & \bullet \\ \downarrow \coprod \ell_i & & \downarrow r \in \mathcal{R} \\ \coprod B_i & \longrightarrow & \bullet \end{array}$$

Since $\ell_i \in \mathcal{L}$, we have maps

$$\begin{array}{ccccc} A_i & \longrightarrow & \coprod A_i & \longrightarrow & \bullet \\ \mathcal{L} \ni \ell_i \downarrow & & \downarrow & \nearrow & \downarrow r \in \mathcal{R} \\ B_i & \longrightarrow & \coprod B_i & \longrightarrow & \bullet \end{array}$$

which in turn means we have a unique map

$$\begin{array}{ccccc} A_i & \longrightarrow & \coprod A_i & \longrightarrow & \bullet \\ \mathcal{L} \ni \ell_i \downarrow & & \downarrow & \nearrow & \downarrow r \in \mathcal{R} \\ B_i & \longrightarrow & \coprod B_i & \longrightarrow & \bullet \end{array}$$

by the universal property of the coproduct $\coprod B_i$.

To conclude we need to check that the triangles below and above the dashed arrow in the former diagram commute. This follows from the universal property of the coproducts $\coprod A_i$ and $\coprod B_i$ since, [in general](#),

$$\text{Hom}\left(\coprod X_i, Y\right) \cong \prod \text{Hom}(X_i, Y).$$

More explicitly, we now that the red paths in the following diagrams are the same:

$$\begin{array}{ccccc} A_i & \longrightarrow & \coprod A_i & \longrightarrow & \bullet \\ \mathcal{L} \ni \ell_i \downarrow & & \downarrow & \nearrow & \downarrow r \in \mathcal{R} \\ B_i & \longrightarrow & \coprod B_i & \longrightarrow & \bullet \end{array} \quad \text{and} \quad \begin{array}{ccccc} A_i & \longrightarrow & \coprod A_i & \longrightarrow & \bullet \\ \mathcal{L} \ni \ell_i \downarrow & & \downarrow & \nearrow & \downarrow r \in \mathcal{R} \\ B_i & \longrightarrow & \coprod B_i & \longrightarrow & \bullet \end{array}$$

and also

$$\begin{array}{ccccc} A_i & \longrightarrow & \coprod A_i & \longrightarrow & \bullet \\ \mathcal{L} \ni \ell_i \downarrow & & \downarrow & \nearrow & \downarrow r \in \mathcal{R} \\ B_i & \longrightarrow & \coprod B_i & \longrightarrow & \bullet \end{array} \quad \text{and} \quad \begin{array}{ccccc} A_i & \longrightarrow & \coprod A_i & \longrightarrow & \bullet \\ \mathcal{L} \ni \ell_i \downarrow & & \downarrow & \nearrow & \downarrow r \in \mathcal{R} \\ B_i & \longrightarrow & \coprod B_i & \longrightarrow & \bullet \end{array}$$

so the conclusion follows from the former comment.

(Closed under retracts.) Let us at least state what a retract of a morphism g should be in the arrow category. Recall that a retract is just

$$\begin{array}{ccccc} X & \longrightarrow & Y & \longrightarrow & X \\ & \searrow & & \nearrow & \\ & \text{id}_X & & & \end{array}$$

So in the arrow category we get

$$\begin{array}{ccccc} \bullet & \xrightarrow{\quad} & \bullet & \xrightarrow{\quad} & \bullet \\ \downarrow f & & \downarrow g & & \downarrow f \\ \bullet & \xrightarrow{\quad} & \bullet & \xrightarrow{\quad} & \bullet \\ & \searrow & & \nearrow & \\ & \text{id} & & & \end{array}$$

□

2 Hatcher's exercise on Whitehead's theorem

Theorem 1 (Whitehead, May). If X is a CW complex and $e : Y \rightarrow Z$ is an n -equivalence, then $e_* : [X, Y] \rightarrow [X, Z]$ is a bijection if $\dim X < n$ and surjection if $\dim X = n$.

Theorem 2 (Whitehead, May). An n -equivalence between CW complexes of dimension less than n is a homotopy equivalence. A weak equivalence between CW complexes is a homotopy equivalence.

Theorem 3 (Whitehead (4.5), Hatcher). If a map $f : X \rightarrow Y$ between connected CW complexes induces isomorphisms $f_* : \pi_n(X) \rightarrow \pi_n(Y)$ for all n , then f is a homotopy equivalence. In case f is the inclusion of a subcomplex $X \hookrightarrow Y$, the conclusion is stronger: X is a deformation retract of Y .

Exercise (Hatcher 4.1.12). Show that an n -connected, n -dimensional CW complex is contractible.

Solution. Just recall that n -connectedness means that $\pi_i(X) = 0$ for all $i \leq n$, which means that X is contractible by theorem 2. □

Homework 2

Exercise (4.1.3). For an H -space (X, x_0) with multiplication $\mu : X \times X \rightarrow X$, show that the group operation in $\pi_n(X, x_0)$ can also be defined by the rule $(f + g)(x) = \mu(f(x), g(x))$.

Exercise (4.1.19). Consider the equivalence relation \simeq_w generated by weak homotopy equivalence: $X \simeq_w Y$ if there are spaces $X = X_1, X_2, \dots, X_n = Y$ with weak homotopy equivalences $X_i \rightarrow X_{i+1}$ or $X_i \leftarrow X_{i+1}$ for each i . Show that $X \simeq_w Y$ iff X and Y have a common CW approximation.

Solution. (\Leftarrow) Suppose Z is a common CW approximation of X and Y , that is, Z is a CW complex and there are weak homotopy equivalences $Z \rightarrow X$ and $Z \rightarrow Y$. Then the sequence of spaces $X = X_1$, $Z = X_2$ and $Y = X_3$ shows that $X \simeq_w Y$.

(\Rightarrow) Suppose Z is a CW approximation of X and let's show it can be made (somehow) into a CW approximation of Y . There is a weak homotopy equivalence $Z \rightarrow X$, and also a weak homotopy equivalence either $X = X_1 \rightarrow X_2$ or $X = X_1 \leftarrow X_2$. I wonder if this implies that the composition $Z \rightarrow X = X_1 \rightarrow X_2$ is also a weak homotopy equivalence \square

Exercise (4.2.1). Use homotopy groups to show that there is no retraction $\mathbb{R}P^n \rightarrow \mathbb{R}P^k$ for $n > k > 0$.

Solution (in progress...) Suppose there is a retraction

$$\begin{array}{ccccc} \mathbb{R}P^k & \hookrightarrow & \mathbb{R}P^n & \longrightarrow & \mathbb{R}P^k \\ & & \searrow \text{id} & \nearrow & \\ & & & & \end{array}$$

it induces isomorphisms

$$\begin{array}{ccccc} \pi_i(\mathbb{R}P^k) & \longrightarrow & \pi_i(\mathbb{R}P^n) & \longrightarrow & \pi_i(\mathbb{R}P^k) \\ & & \searrow \cong & \nearrow & \\ & & & & \end{array}$$

If we had that this is a Serre fibration, we would get that

content...

\square

Exercise (4.2.13). Show that a map between connected n -dimensional CW complexes is a homotopy equivalence if it induces an isomorphism on π_i for $i \leq n$. [Pass to universal covers and use homology.]

Solution. Let X and Y be n -dimensional CW complexes and $f : X \rightarrow Y$ such that $f_* : \pi_i(X) \rightarrow \pi_i(Y)$ is an isomorphism for $i \leq n$. Let's try to use Hurewicz theorem, which states that a map between simply connected CW complexes is a homotopy equivalence if it induces isomorphisms on all homology groups.

Consider the universal covers \tilde{X} and \tilde{Y} , which are simply connected and also [have CW structures](#). By prop. 4.1, the cover projections induce isomorphisms in the homotopy groups for all $i \geq 2$. By [StackExchange](#) there is a unique lift \tilde{f} to the universal covers making the diagram on the left commute, and by functoriality the diagram on the right also commutes.

$$\begin{array}{ccc}
\tilde{X} & \xrightarrow{\tilde{f}} & \tilde{Y} \\
p \downarrow & & \downarrow q \\
X & \xrightarrow{f} & Y
\end{array}
\qquad
\begin{array}{ccc}
\pi_i(\tilde{X}) & \xrightarrow{\tilde{f}_*} & \pi_i(\tilde{Y}) \\
p_* \downarrow \cong & & \cong \downarrow q_* \\
\pi_i(X) & \xrightarrow{f_*} & \pi_i(Y)
\end{array}
\qquad i \geq 2$$

We conclude that \tilde{f} is a **weak homotopy equivalence**, and by prop. 4.21 it induces isomorphisms on homology groups. Finally, by Hurewicz theorem (coro. 4.33) it is an homotopy equivalence and so is f . \square

Exercise. There is a fiber sequence $U(n) \hookrightarrow U(n+1) \rightarrow U(n+1)/U(n) \cong S^{2n+1}$. Use this to show that $\pi_k(U(n)) \rightarrow \pi_k(U(n+1))$ is isomorphism for $n > k/2$. Compute $\pi_k(U(n))$ for $n \geq 2$ and $k = 1, 2, 3$. In fact, if k is even then $\pi_k(U(N)) = 0$ and if k is odd then $\pi_k(U(N)) = \mathbb{Z}$, where again $N > k/2$. These equalities are known as Bott periodicity.

Solution. The required isomorphisms $\pi_k(U(n)) \rightarrow \pi_k(U(n+1))$ follow simply from the fact that S^{2n+1} is $2n+1$ -connected: in the long homotopy sequence of the fiber bundle we have

$$\pi_{k+1}(S^{2n+1}) \longrightarrow \pi_k(U(n)) \longrightarrow \pi_k(U(n+1)) \longrightarrow \pi_k(S^{2n+1})$$

so when $2n+1 > k+1 \iff n > k/2$ the homotopy groups of the spheres vanish and we have an isomorphism.

The group $\pi_1(U(n))$ is isomorphic to \mathbb{Z} . This follows from the fact that $U(1)$ is homeomorphic to a circle and by induction using the former isomorphism $\pi_1(U(n)) \cong \pi_1(U(n+1))$. We also have $\pi_2(U(1)) = 0$, so that again by induction we get $\pi_2(U(n)) = 0$. Finally, a similar argument shows $\pi_3(U(n)) = 0$. \square

Exercise (4.2.32). Show that if $S^k \rightarrow S^m \rightarrow S^n$ is a fiber bundle, then $k = n-1$ and $m = 2n-1$. [Look at the long exact sequence of homotopy groups.]

Solution (in progress...) The sequence of homotopy groups is

$$\cdots \rightarrow \pi_i(S^k) \rightarrow \pi_i(S^m) \rightarrow \pi_i(S^n) \rightarrow \pi_{i-1}(S^k) \rightarrow \pi_{i-1}(S^m) \rightarrow \cdots$$

so that if $i \leq k$ we have $\pi_i(S^m) = \pi_i(S^n)$, \square

cohomology ring of \mathbb{CP}^n

Exercise. Show that

$$H^\bullet(\mathbb{CP}^n) \cong \mathbb{Z}[\alpha]/(\alpha^{n+1})$$

where α has degree 2.

Proof. The CW structure of \mathbb{CP}^n consists of one cell for every even dimension. This gives us the following chain complex:

$$\mathbb{Z} \longrightarrow 0 \longrightarrow \mathbb{Z} \longrightarrow \dots \longrightarrow \mathbb{Z}, \quad \text{if } n \text{ is even}$$

$$\mathbb{Z} \longrightarrow 0 \longrightarrow \mathbb{Z} \longrightarrow \dots \longrightarrow 0, \quad \text{if } n \text{ is odd}$$

which yields the cohomology

$$H^i(\mathbb{CP}^n) = \begin{cases} \mathbb{Z}, & i = 0, 2, 4, \dots, 2n \\ 0, & i \text{ otherwise} \end{cases}$$

so that

$$H^\bullet(\mathbb{CP}^n) = H^0(\mathbb{CP}^n) \oplus H^2(\mathbb{CP}^n) \oplus \dots \oplus H^{2n}(\mathbb{CP}^n)$$

This means that the underlying group of the cohomology ring is the same as that of

$$\mathbb{Z}[\alpha]/(\alpha^{n+1})$$

where α has degree 2. To show that these groups are also isomorphic as algebras we can use Poincaré duality as follows.

Consider the case $n = 2$, where we may immediately multiply the generator of second cohomology group with itself:

$$\begin{aligned} H^2(\mathbb{CP}^2) \times H^2(\mathbb{CP}^2) &\rightarrow H^4(\mathbb{CP}^2) \\ (\alpha, \alpha) &\mapsto \alpha \smile \alpha = \alpha^2 \end{aligned}$$

By Poincaré duality this map is a nondegenerate symmetric bilinear form, so it must map generator to a generator. **The fact that the product of the generator in degree 2 is the generator of degree 4** yields an homomorphism

$$\begin{aligned} \varphi : \mathbb{Z}[\alpha] &\rightarrow H^\bullet(\mathbb{CP}^n) \\ \alpha &\mapsto \alpha \in H^2(\mathbb{CP}^n) \end{aligned}$$

with kernel (α^{n+1}) as desired.

Now the case of \mathbb{CP}^3 is:

$$\begin{aligned} H^2(\mathbb{CP}^3) \times H^4(\mathbb{CP}^3) &\rightarrow H^6(\mathbb{CP}^3) \\ (\alpha, \alpha^2) &\mapsto \alpha \smile \alpha^2 = \alpha^3 \end{aligned}$$

which also maps generator to generator, producing the desired algebra isomorphism. Notice we have used the group isomorphism $H^4(\mathbb{CP}^3) \approx H^4(\mathbb{CP}^2)$ when denoting the generator of $H^4(\mathbb{CP}^3)$ as α^2 . Such an isomorphism is induced by inclusion $\mathbb{CP}^{n-1} \hookrightarrow \mathbb{CP}^n$ via relative cohomology exact sequence.

The case for dimension n follows by induction. □

References

- [1] A. Hatcher. *Algebraic topology*. Cambridge: Cambridge Univ. Press, 2000 (cit. on pp. 3, 4, 10).
- [2] J.P. May. *A Concise Course in Algebraic Topology*. Chicago Lectures in Mathematics. University of Chicago Press, 1999. ISBN: 9780226511832 (cit. on p. 10).
- [3] H.R. Miller. *Lectures On Algebraic Topology*. World Scientific Publishing Company, 2021. ISBN: 9789811231261. URL: <https://books.google.com.br/books?id=LIZGEAAAQBAJ> (cit. on pp. 4, 5).
- [4] Emily Riehl. *Homotopical categories: from model categories to $(\infty, 1)$ -categories*. 2020. arXiv: 1904.00886 [math.AT] (cit. on p. 8).