Computing Wasserstein Barycenter via Operator Splitting: the Method of Averaged Marginals

Daniel Mimouni[1,2], Paul Malisani[2], Jiamin Zhu[2], Welington de Oliveira[1]

1 **Centre de Mathématiques Appliquées - CMA, Mines Paris PSL**

2 **IFP Energies nouvelles**

25th International Symposium on Mathematical Programming
Advances in Variational Analysis and Nonsmooth Optimization

Overview

I. Some context

II. Navigating the Wasserstein Barycenter Challenge

III. Our Contribution

IV. Applications

V. Conclusion

Some Context

- In applied probability, stochastic optimization, and data science, a crucial aspect is the ability to compare, summarize, and reduce the dimensionality of empirical/discrete measures

- Since these tasks rely heavily on pairwise comparisons of measures, it is essential to use an appropriate metric for accurate data analysis

- Different metrics define different barycenters of a set of measures: a barycenter is a mean element that minimizes the (weighted) sum of all its distances to the set of target measures

- When the chosen metric is the optimal transport one, and there is mass equality between the measures, the underlying barycenter is denoted by Wasserstein Barycenter (WB)

Example extracted from [1]

30 artificial images

Barycenters using

(a) Euclidean distance

(b) Euclidean + re-centering

(c) Jeffrey centroid

(d) RKHS distance

(e) 2-Wasserstein distance:
Wasserstein barycenter



[1] M. Cuturi, A. Doucet. JMLR, 2014

Navigate the Wasserstein Barycenter Challenge

The Wasserstein Distance

The Wasserstein distance - General setting

Let (Ω, P) a probabilty space and two measurable random vectors $\xi, \zeta : \Omega \mapsto \mathbb{R}^d$ such that $\mu := \xi_\# P$ and $\nu := \zeta_\# P$. Their (quadratic) 2-Wasserstein distance is

$$W_2(\xi, \zeta) := \left(\inf_{\pi \in U(\mu,\nu)} \int_{\mathbb{R}^d \times \mathbb{R}^d} \|x - y\|^2 \mathrm{d}\pi(x, y) \right)^{1/2}, \tag{WD}$$

Here $U(\mu, \nu)$ is the set of all probability measures on $\mathbb{R}^d \times \mathbb{R}^d$ having marginals μ and ν.

We focus on discrete/empirical with finite support:

$$\mu = \sum_{r=1}^{R} p_r \delta_{\xi_r} \quad \text{and} \quad \nu = \sum_{s=1}^{S} q_s \delta_{\zeta_s}$$

Wasserstein distance - discrete setting

For discrete probability measures μ and ν, their 2-Wasserstein distance W_2 is:

$$W_2(\mu, \nu) := \left(\min_{\pi \in U(\mu,\nu)} \sum_{r=1}^{R} \sum_{s=1}^{S} \|\xi_r - \zeta_s\|^2 \pi_{rs} \right)^{1/2} \tag{1}$$

Wasserstein Barycenter (WB)

Wasserstein Barycenter problem

Given M measures $\{\nu^{(1)}, \ldots, \nu^{(M)}\} \subset \mathcal{P}(\mathbb{R}^d)$ and $\alpha \in \Delta_M$, an 2-*Wasserstein barycenter* with weights α is a solution to the following optimization problem

$$\min_{\mu \in \mathcal{P}(\mathbb{R}^d)} \sum_{m=1}^{M} \alpha_m W_2^2(\mu, \nu^{(m)}) \,. \tag{2}$$

This is a nonconvex optimization problem: the optimization is performed over the support $\texttt{supp}(\mu)$ and mass p



Huge scale Linear Program (LP)

WB problem

It boils down to solve this huge scale LP :

$$
\begin{cases}
\min\limits_{p,\pi} & \sum_{r=1}^{R}\sum_{s=1}^{S^{(1)}} c_{rs}^{(1)}\pi_{rs}^{(1)} \quad + \cdots + \quad \sum_{r=1}^{R}\sum_{s=1}^{S^{(M)}} c_{rs}^{(M)}\pi_{rs}^{(M)} \\
\text{s.t.} & \sum_{r=1}^{R}\pi_{rs}^{(1)} \qquad\qquad\qquad\qquad = q_s^{(1)}, \qquad s=1,\dots,S^{(1)} \\
& \qquad\qquad \ddots \qquad\qquad\qquad\qquad \vdots \\
& \qquad\qquad\qquad\qquad \sum_{r=1}^{R}\pi_{rs}^{(M)} = q_s^{(M)}, \qquad s=1,\dots,S^{(M)} \\
& \sum_{s=1}^{S^{(1)}}\pi_{rs}^{(1)} \qquad\qquad\qquad\qquad = p_r, \qquad r=1,\dots,R \\
& \qquad\qquad \ddots \qquad\qquad\qquad\qquad \vdots \\
& \qquad\qquad\qquad\qquad \sum_{s=1}^{S^{(M)}}\pi_{rs}^{(M)} = p_r, \qquad r=1,\dots,R \\
& p\in\Delta_R, \pi^{(1)}\geq 0 \quad \cdots \quad \pi^{(M)}\geq 0,
\end{cases}
$$

- If $M = 100$ $R = S^{(m)} = 1600$, $m = 1,\dots,M$ (corresponding to figures with 40×40 pixels), the above LP has 256 001 600 variables.
- This LP may scale exponentially in the number M of measures [2]
- Assume $S^{(m)} = S$ for all $m = 1,\dots,M$: then $R = S^M$ and the LP has $M(S)^{M+1} + (S)^M$ variables and $M(S)^M + MS$ equality constraints

[2] free support setting; S. Borgwardt. Operational Research (2022)

A vast body of the literature deals with inexact WBs

Inexact approaches

- Mostly based on reformulations via an entropic regularization: several papers by M. Cuturi, G. Peyré, G. Carlier and others
- Block-coordinate approach: fix the support and optimize the probability, then fix the probability and optimize the support [3, 4, 5]
- Other approaches [6,7,8]

Exact methods

- Methods for computing exact WBs are based on linear programming techniques [9,10]

[3] M. Cuturi, A. Doucet. JMLR, 2014
[4] J. Ye, J. Li. IEEE ICP (214)
[5] J. Ye et al. IEEE Transactions on Signal Processing (2017)
[6] G. Puccetti, L. Ruschendorf, S. Vanduffe. JMVA (2020)
[7] S. Borgwardt. Operational Research (2022)
[8] J. von Lindheim. COAP (2023)
[9] S. Borgwardt, S. Patterson (2020). INFORM J. Optimization
[10] J. Altschuler, E. Adsera. JMLR (2021)

Our Contribution

Our contribution

We provide an easy-to-implement, memory efficient and parallelizable algorithm based on the Douglas-Rachford splitting scheme to compute a solution to LPs of the form

$$\begin{cases} \min\limits_{p,\pi\geq 0} & \sum\limits_{m=1}^{M}\sum\limits_{r=1}^{R}\sum\limits_{s=1}^{S^{(m)}} d_{rs}^{(m)}\pi_{rs}^{(m)} \\ \text{s.t.} & \sum\limits_{r=1}^{R}\pi_{rs}^{(m)} = q_s^{(m)}, \quad s=1,\dots,S^{(m)},\ m=1,\dots,M \\ & \sum\limits_{s=1}^{S^{(m)}}\pi_{rs}^{(m)} = p_r, \quad r=1,\dots,R,\ m=1,\dots,M \end{cases}$$

with given $d^{(m)} \in \mathbb{R}^{R\times S^{(m)}}$

Reformulation of the LP

Separable sets

$$\Pi^{(m)} := \left\{ \pi^{(m)} \geq 0 : \sum_{r=1}^{R} \pi_{rs}^{(m)} = q_s^{(m)},\ s = 1, \ldots, S^{(m)} \right\},\ m = 1, \ldots, M \tag{3}$$

and the linear subspace

$$\mathcal{B} := \left\{ \pi = (\pi^{(1)}, \ldots, \pi^{(M)}) : \begin{array}{lll} \sum_{s=1}^{S^{(1)}} \pi_{rs}^{(1)} & = \sum_{s=1}^{S^{(2)}} \pi_{rs}^{(2)}, & \forall r \\ \sum_{s=1}^{S^{(2)}} \pi_{rs}^{(2)} & = \sum_{s=1}^{S^{(3)}} \pi_{rs}^{(3)}, & \forall r \\ & \vdots & \\ \sum_{s=1}^{S^{(M-1)}} \pi_{rs}^{(M-1)} & = \sum_{s=1}^{S^{(M)}} \pi_{rs}^{(M)}, & \forall r \end{array} \right\} \tag{4}$$

Douglas-Rachford theory

- Consider the indicator function $\mathbf{i}_C$ of a convex set C, and reformulate the problem:

$$f^{(m)}(\pi^{(m)}) := \sum_{r=1}^{R} \sum_{s=1}^{S^{(m)}} c_{rs}^{(m)} \pi_{rs}^{(m)} + \mathbf{i}_{\Pi^{(m)}}(\pi^{(m)}), \quad m = 1, \ldots, M, \tag{5}$$

- Recast problem:

$$f(\pi) := \sum_{m=1}^{M} f^{(m)}(\pi^{(m)}) \quad \text{and} \quad g(x) := \begin{cases} \mathbf{i}_{\mathcal{B}}(\pi) & \text{if balanced} \\ \gamma\, \mathtt{dist}_{\mathcal{B}}(\pi) & \text{if unbalanced.} \end{cases} \tag{6}$$

- Since f is polyhedral, it follows that computing one of its solutions is equivalent to

$$\text{find} \quad \pi \quad \text{such that} \quad 0 \in \partial f(\pi) + \partial g(\pi). \tag{7}$$

Douglas-Rachford algorithm

Given initial point $\theta^0 = (\theta^{(1),0}, \ldots, \theta^{(M),0})$ and prox-parameter $\rho > 0$:

$$\begin{cases} \pi^{k+1} & = & \mathrm{prox}_{g/\rho}(\theta^k) \\ \hat{\pi}^{k+1} & = & \mathrm{prox}_{f/\rho}(2\pi^{k+1} - \theta^k) \\ \theta^{k+1} & = & \theta^k + \hat{\pi}^{k+1} - \pi^{k+1} \end{cases} \tag{8}$$

The Method of Averaged Marginals (MAM)

The Douglas Rachford and The MAM Algorithm

$$\begin{cases} \pi^{k+1} & = & \texttt{Proj}_{\mathcal{B}}(\theta^k) \xrightarrow{\text{explicit}} \pi_{rs}^{(m)} := \theta_{rs}^{(m)} + \frac{(p_r^k - p_r^{(m)})}{S^{(m)}} \\ \hat{\pi}^{k+1} & := & \begin{pmatrix} \hat{\pi}_{1s}^{(m)} \\ \vdots \\ \hat{\pi}_{Rs}^{(m)} \end{pmatrix} = \texttt{Proj}_{\Delta_R(q_s^{(m)})} \begin{pmatrix} y_{1s} - \frac{1}{\rho} c_{1s}^{(m)} \\ \vdots \\ y_{Rs} - \frac{1}{\rho} c_{Rs}^{(m)} \end{pmatrix}, \quad s = 1, \ldots, S^{(m)} \\ \theta^{k+1} & = & \theta^k + \hat{\pi}^{k+1} - \pi^{k+1} \end{cases}$$

- At every iteration the barycenter approximation p^k is a weighted average of the M marginals $p^{(m)} := \sum_{s=1}^{S^{(m)}} \theta_{rs}^{(m)}$.
- The projection onto the simplex can be accomplished in parallel, exactly and efficiently by specialized algorithms[11].
- The whole sequence converges to an exact barycenter.

[11] see L. Condat 2016; computation in $O(R \log(R))$.

Unbalanced Wasserstein barycenters

Linear subspace of balanced plans:

$$\mathcal{B} = \left\{ \pi : \sum_{s=1}^{S^1} \pi^1_{rs} = \cdots = \sum_{s=1}^{S^M} \pi^M_{rs}, \quad r = 1, \ldots, R \right\}$$

ν^m, $m = 1, \ldots, M$, have equal masses

Balanced WB

$$\begin{cases} \min\limits_{\pi \in \mathcal{B}} & \sum\limits_{m=1}^{M} \langle d^m, \pi^m \rangle \\ \text{s.t.} & \pi^1 \in \Pi^m \\ & \vdots \\ & \pi^M \in \Pi^M \end{cases}$$

ν^m, $m = 1, \ldots, M$, have different masses

Unbalanced WB ($\gamma > 0$)

$$\begin{cases} \min\limits_{\pi} & \sum\limits_{m=1}^{M} \langle d^m, \pi^m \rangle + \gamma \, \mathtt{dist}_{\mathcal{B}}(\pi) \\ \text{s.t.} & \pi^1 \in \Pi^m \\ & \vdots \\ & \pi^M \in \Pi^M \end{cases}$$

MAM can be easily adapted to deal with both balanced or unbalanced WBs[12]

[12] the new proximal is known to be solved analytically; Bauschke and P. L. Combettes, Spinger 2017

Applications

Qualitative comparisons - Fixed support $R = 1\,600$

We benchmark MAM, randomized MAM, and IBP (Iterative Bregman Projection of [13]) on the MNIST database with $M = 100$ images of 40×40 pixels. LP's dimension: 256 001 600 variables and 320 000 constraints

[13] [J.-D. Benamou et al. SIAM Journal on Scientific Computing. (2015)]

Qualitative comparisons - Fixed support $R = 1\,600$

WB

Qualitative comparisons - Fixed support $R = 1\,600$



WB

Quantitative comparisons - Fixed support $R = 1\,600$



Evolution with respect to time of the difference between the Wasserstein barycenter distance of an approximation, $\bar{W}_2^2(p^k)$, and the Wasserstein barycentric distance of the exact solution $\bar{W}_2^2(p_{exact})$ given by the LP. The time step between two points is 30 seconds

Exact free-support resolution

The dataset we use is the one from [14]: $M = 10$ images of 60×60 pixels
LP's dimension: $1.2574 \cdot 10^{10}$ variables and $3.5288 \cdot 10^{6}$ constraints
We compare with the dedicated solver of Altschuler and Boix-Adsera, available at [15]

- All measures share the same finite support: suppose that all measures $\nu^{(m)}$ are supported on a d-dimensional regular grid of integer step sizes in each direction, each coordinate going from 1 to K: $S^{(m)} = S = K^d$.

- The measures are evenly weighted $\alpha_m = \frac{1}{M}$, $m = 1, \ldots, M$



Evolution of the approximated MAM barycenter with time in regards with the exact barycenter of the Altschuler and Bois-Adsera algorithm computed in 3.5 hours [16]

MAM can solve larger problems than the method Altschuler and Boix-Adsera

[14] J. M. Altschuler and E. Boix-Adsera. JMLR (2021)
[15] `https://github.com/eboix/high_precision_barycenters`
[16] S. Borgwardt, S. Patterson (2020). INFORM J. Optimization



The optimal value of the WB problem is 0.2666

After 1 hour of processing, MAM had a barycenter distance of 0.2702, which improved to 0.2667 after 3.5 hours, when the solver of Altschuler and Boix-Adsera halts

Unbalanced WB



$$\begin{cases} \min\limits_{\pi} & \sum_{m=1}^{M} \langle d^{(m)}, \pi^{(m)} \rangle + \gamma \mathtt{dist}_{\mathcal{B}}(\pi) \\ \text{s.t.} & \pi^{(1)} \in \Pi^{(m)}, \ldots, \pi^{(M)} \in \Pi^{(M)} \end{cases}$$



Constrained Wasserstein barycenters

Suppose the probability vector p is constrained to a closed convex set $X \subset \mathbb{R}^R$:

$$\begin{cases} \min\limits_{p,\pi \geq 0} & \sum\limits_{m=1}^{M} \langle d^m, \pi^m \rangle \\ \text{s.t.} & \sum\limits_{r=1}^{R} \pi_{rs}^m = q_s^m, \quad s = 1, \ldots, S^m, \; m = 1, \ldots, M \\ & \sum\limits_{s=1}^{S^m} \pi_{rs}^m = p_r, \quad r = 1, \ldots, R, \; m = 1, \ldots, M \\ & p \in X \end{cases}$$

We can easily adapt MAM to compute constrained WBS

Ajouter application ici





Take-away messages

- New easy-to-implement and memory efficient algorithm for computing WBs, which is parallelizable and can run in a randomized manner if necessary
- It can be applied to both balanced WB and unbalanced WB problems upon setting a single parameter
- It can be applied to the free or fixed-support settings
- The method is submitted (after review) in SIAM Journal on Mathematics and Data Science `https://arxiv.org/pdf/2309.05315.pdf`
- Our Python code is freely available at `https://ifpen-gitlab.appcollaboratif.fr/detocs/mam_wb`

Thank you!

D. Mimouni, P. Malisani, J. Zhu, W. de Oliveira. Computing Wasserstein barycenter via operator splitting: the method of averaged marginals, https://arxiv.org/pdf/2309.05315.pdf, 2023

Contact:

daniel.mimouni@ifpen.fr

https://dan-mim.github.io

Annexes

Special setting: grid-structured data

- All measures share the same finite support: suppose that all measures $\nu^{(m)}$ are supported on a d-dimensional regular grid of integer step sizes in each direction, each coordinate going from 1 to K: $S^{(m)} = S = K^d$, and $\texttt{supp}(\nu^{(m)}) := \{\zeta_1, \ldots, \zeta_S\}$, $m = 1, \ldots, M$
- The measures are evenly weighted $\alpha_m = \frac{1}{M}$, $m = 1, \ldots, M$
- Then $\texttt{supp}(\mu)$ has at most

$$R \leq ((K-1)M + 1)^d$$

points, as the finer grid only runs between the boundary points [17]

This significantly reduces the LP's dimension

[17] S. Borgwardt, S. Patterson (2020). INFORM J. Optimization

Example (LP's dimensions)

Consider the case: $M = 10$, $d = 2$, $K = 40 \Rightarrow S = 1600$

data	$\lvert\mathtt{supp}(\mu)\rvert$ R	# variables $(MS+1)R$	# eq. constraints $(S+R)M$
general	$1.0995 \cdot 10^{32}$	$1.7593 \cdot 10^{36}$	$1.0995 \cdot 10^{33}$
grid-structured	152881	$2.4462 \cdot 10^{9}$	1544810

- In contrast to the worst-case, exponentially sized possible support set, there always exists a WB $\bar{\mu}$ with provably sparse support

$$|\mathtt{supp}(\bar{\mu})| \leq \sum_{m=1}^{M} S^{(m)} - M + 1$$

- For the above example $|\mathtt{supp}(\bar{\mu})| \leq 15991$

- This fact motivates heuristics for computing inexact WBs: fixed-support approaches, which generally fix R to $\sum_{m=1}^{M} S^{(m)} - M + 1$ (or fewer) points

Algorithm

Input: initial plan $\pi = (\pi^1, \dots, \pi^m)$ and parameters $\rho, \gamma > 0$

Define $a_m \leftarrow (\frac{1}{S^m})/(\sum_{j=1}^{M} \frac{1}{S^j})$ and set $p^m \leftarrow \sum_{s=1}^{S^m} \pi^m_{rs}$, $m = 1, \dots, M$

while not converged **do**

  $p \leftarrow \sum_{m=1}^{M} a_m p^m$ ▷ Average the marginals

  Set $t \leftarrow 1$ if $\rho \sqrt{\sum_{m=1}^{M} \frac{\|p - p^m\|^2}{S^m}} \leq \gamma$; else $t \leftarrow \gamma / \left(\rho \sqrt{\sum_{m=1}^{M} \frac{\|p - p^m\|^2}{S^m}} \right)$

  for $m = 1, \dots, M$ **do**

    for $s = 1, \dots, S^m$ **do**

      $\pi^m_{:s} \leftarrow \texttt{Proj}_{\Delta(q^m_s)} \left(\pi^m_{:s} + 2t \frac{p - p^m}{S^m} - \frac{1}{\rho} d^m_{:s} \right) - t \frac{p - p^m}{S^m}$

    end for

    $p^m \leftarrow \sum_{s=1}^{S^m} \theta^m_{rs}$ ▷ Update the mth marginal

  end for

end while

Set $\gamma = \infty$ to compute balanced WB (if the measures are balanced)
Otherwise, choose $\gamma \in (0, \infty)$ to compute unbalanced WB

The Method of Averaged Marginals - MAM

Constrained setting

Algorithm

Input: initial plan $\pi = (\pi^1, \dots, \pi^m)$ and parameter $\rho > 0$

Define $a_m \leftarrow (\frac{1}{S^m})/(\sum_{j=1}^{M} \frac{1}{S^j})$ and set $p^m \leftarrow \sum_{s=1}^{S^m} \pi^m_{rs}$, $m = 1, \dots, M$

while not converged **do**

  $p \leftarrow \texttt{Proj}_X \left(\sum_{m=1}^{M} a_m p^m \right)$ ▷ Average the marginals

  for $m = 1, \dots, M$ **do**

    for $s = 1, \dots, S^m$ **do**

      $\pi^m_{:s} \leftarrow \texttt{Proj}_{\Delta(q^m_s)} \left(\pi^m_{:s} + 2\frac{p - p^m}{S^m} - \frac{1}{\rho} d^m_{:s} \right) - \frac{p - p^m}{S^m}$

    end for

    $p^m \leftarrow \sum_{s=1}^{S^m} \theta^m_{rs}$ ▷ Update the mth marginal

  end for

end while