

# Lecture 1

Dani Nguyen

September 23, 2025

## 1 Invertible Matrix Theorem

### 1.1 Definition

Let  $A$  be a square  $n \times n$  matrix. Then the following statements are equivalent. That is, for a given  $A$ , the statements are either all true or all false.

1.  $A$  is invertible.
2.  $A$  is row equivalent to  $I$ .
3.  $A$  has  $n$  pivot positions.
4. The equation  $A\mathbf{x} = \mathbf{0}$  has only the trivial solution.
5. The columns of  $A$  are linearly independent.
6. The linear transformation  $\mathbf{x} \mapsto A\mathbf{x}$  is one-to-one.
7. The equation  $A\mathbf{x} = \mathbf{b}$  has at least one solution for each  $\mathbf{b}$  in  $\mathbb{R}^n$ .
8. The columns of  $A$  span  $\mathbb{R}^n$ .
9. The linear transformation  $\mathbf{x} \mapsto A\mathbf{x}$  maps  $\mathbb{R}^n$  onto  $\mathbb{R}^n$ .
10. There is an  $n \times n$  matrix  $C$  such that  $CA = I$ .
11. There is an  $n \times n$  matrix  $D$  such that  $AD = I$ .
12.  $A^T$  is invertible.

13. The columns of  $A$  form a basis of  $\mathbb{R}^n$ .
14.  $\text{Col } A = \mathbb{R}^n$ .
15.  $\dim \text{Col } A = n$ .
16.  $\text{rank } A = n$
17.  $\text{Nul } A = \mathbf{0}$
18.  $\dim \text{Nul } A = 0$
19. The number 0 is *not* an eigenvalue of  $A$ .
20.  $\det A \neq 0$ .
21.  $(\text{Col } A)^\perp = \mathbf{0}$
22.  $(\text{Nul } A)^\perp = \mathbb{R}^n$
23.  $\text{Row } A = \mathbb{R}^n$

## Exercise

**Thm 9.1.16** Let  $A \in M_n$  and let  $\lambda \in \mathbb{C}$ . Then the following statements are equivalent:

1.  $\lambda$  is an eigenvalue of  $A$ .
2.  $A\mathbf{x} = \lambda\mathbf{x}$  for some nonzero  $\mathbf{x} \in \mathbb{C}^n$ .
3.  $(A - \lambda I)\mathbf{x} = \mathbf{0}$  has a nontrivial solution, that is,  $\text{nullity}(A - \lambda I) > 0$ .
4.  $\text{rank}(A - \lambda I) < n$ .
5.  $A - \lambda I$  is not invertible.
6.  $A^\top - \lambda I$  is not invertible.
7.  $\lambda$  is an eigenvalue of  $A^\top$ .

## Proof