

MATH 114 - Computational Stochastics

Dani Nguyen

Winter 2025

Contents

1	Overview	1
1.1	Introduction	1
1.2	Sampling	2
1.3	MCMC	2
2	Markov Chains	3
2.1	Lecture 9	3
2.2	Lecture 10	4
2.3	Lecture 11 - Invariant Distribution	6
2.4	Lecture 12 - MCMC	9
2.5	Lecture 13 - Metropolis	12
2.6	Lecture 14	13
2.7	Lecture 15 - Midterm Review	16
2.8	Lecture 17 - MCMC & SDEs	20

1 Overview

1.1 Introduction

Stochastic models describe real life phenomena like motion of the cell, shape of polymers stock prices, and more. What can we compute analytically and numerically from these models?

Key questions:

1. What is the distribution of the system?
2. How can we sample from the distribution?
3. How can we compute mean, variance, and higher moments of these systems?

Main topics:

1. Sampling RVs
2. Markov chain Monte Carlo (MCMC)
3. Stochastic differential equations (numerical & computation)

1.2 Sampling

- Generating "hard" RVs from "easy" RVs.
- Compute "hard" integrals.

ex Area of the unit circle, $A = \pi$

Let $U_1 \sim U(-1, 1)$ and $U_2 \sim U(-1, 1)$

Let $x = \begin{cases} 0 & \text{if } (u_1, u_2) \text{ outside circle} \\ 1 & \text{if } (u_1, u_2) \text{ inside circle} \end{cases}$

$\implies P(X = 1) = \frac{A}{4}, P(X = 0) = 1 - \frac{A}{4}$

$\mathbb{E}(X) = \frac{A}{4}$

After N trials we generate $X_1, X_2, \dots, X_n$ iid

By the strong law of large numbers

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{i=1}^N X_i = \frac{A}{4} = \frac{\pi}{4}$$

1.3 MCMC

Construct Markov chains to generate target distribution.

ex Ising model

2 Markov Chains

2.1 Lecture 9

Def: A stochastic process is a family of RVs, $X_t \in S$, indexed by t , where $S \subseteq \mathbb{R}^d$ is a state space. Typically, $t \in [0, \infty)$, or $t \in [a, b]$, or $t \in \{0, 1, 2, 3, \dots\}$, etc.

- A discrete time stochastic process, is a sequence of RVs, $X_n \in S$, $n = 0, 1, 2, \dots$ or $n = 1, 2, 3, \dots$
- If S is finite or countably infinite then the stochastic process is a discrete state process.

First, we will examine discrete-time, discrete space, stochastic process.

Two basic questions:

- Find the joint distribution

$$P(X_0 = i_0, X_1 = i_1, \dots, X_n = i_n) \\ i_0, i_1, \dots, i_n \in S$$

- Find the long-time behavior of the distribution of X_n :

$$P(x_n = i), i \in S, n \gg 1$$

Def: Markov processes

- A stochastic process $\{X_n\}_{n=0}^\infty$ is a Markov process if

$$P(X_{n+1} = i_{n+1} | x_0 = i_0, \dots, x_n = i_n) \\ = P(X_{n+1} = i_{n+1} | X_n = i_n) \\ \forall n \geq 0, \forall i_0, \dots, i_{n+1} \in S$$

- A Markov process is time homogeneous if:

$$P(x_{n+1} = j | x_n = i) \\ = P(X_1 = j | X_0 = i) \\ \forall n \geq 0, \forall i, j \in S$$

Here we consider discrete time/state, time homogeneous Markov Chains (MC).

Def: Transition probability

$$\begin{aligned} P(i, j) &= P(X_1 = j | X_0 = i) \\ &= P(i \rightarrow j) \quad \forall i, j \in S \end{aligned}$$

Def: Transition matrix

$$P = [p(i, j)] \text{ if } S = \{1, 2, 3, \dots\} \text{ or } \{0, 1, 2, \dots\}$$

Here i, j are states and also are row/column labels

For any $n \geq 1, i_0, \dots, i_n \in S$, we have:

$$\begin{aligned} &P(X_0 = i_0, \dots, X_n = i_n) \\ &= P(X_0 = i_0)P(X_1 = i_1 | X_0 = i_0)P(X_2 = i_2 | X_1 = i_1, X_0 = i_0) \\ &\dots P(X_n = i_n | X_{n-1} = i_{n-1}, \dots, X_0 = i_0) \quad [\text{conditional probability}] \end{aligned}$$

ex Let $X_0, X_1, \dots, X_n \in S$ iid. Then $\{X_n\}_{n=0}^\infty$ is a time homogeneous MC.
Let $S = \{1, 2, \dots, N\}$. Then $p(i, j) = \mathbb{P}(X_1 = j | X_0 = i) = \mathbb{P}(X_1 = j) = p_j$

$$\Rightarrow p = \begin{bmatrix} p_1 & \dots & p_n \\ \vdots & \ddots & \vdots \\ p_1 & \dots & p_n \end{bmatrix}, \quad \sum_{i=1}^n p_i = 1$$

2.2 Lecture 10

Recap:

- Discrete time/state, time homogeneous
- MC: $\{X_n\}_{n=0}^\infty \in S$
- $\mathbb{P}(X_{n+1} = i_{n+1} | X_0 = i_0, \dots, X_n = i_n) = \mathbb{P}(X_{n+1} = i_{n+1} | X_n = i_n)$
 $= p(i_n, i_{n+1})$
- Transition probability $\mathbb{P}(X_1 = j | X_0 = i)$
 $= p(i, j) = p_{ij} \quad \forall i, j \in S$
- Transition matrix $\mathbb{P}[p(i, j)]$

ex Two-state MC, e.g. state of phone $X_n \in S = \{0, 1\}$

$$\begin{cases} X_n = 0 & \text{phone free at time } n \\ X_n = 1 & \text{phone busy at time } n \end{cases}$$

$$\mathbb{P}(\text{free} \rightarrow \text{busy in time step}) = p$$

$$\mathbb{P}(\text{busy} \rightarrow \text{free in time step}) = q$$

$$\mathbb{P}(\text{free} \rightarrow \text{free in time step}) = 1 - p$$

$$\mathbb{P}(\text{busy} \rightarrow \text{busy in time step}) = 1 - q$$

Transition matrix:

$$P = \begin{bmatrix} 1-p & p \\ q & 1-q \end{bmatrix}$$

e.g. $p = \frac{1}{2}, q = \frac{1}{3}$

$$P = \begin{bmatrix} \frac{1}{2} & \frac{1}{3} \\ \frac{1}{3} & \frac{2}{3} \end{bmatrix}$$

Let $\mathbb{P}(X_0 = 0) = \frac{1}{5}, \mathbb{P}(X_0 = 1) = \frac{4}{5}$

Then $\mathbb{P}(X_0 = 0, X_1 = 1, X_2 = 0)$

$$= \mathbb{P}(X_0 = 0) \cdot p(0, 1) \cdot p(1, 0)$$

$$= \frac{1}{5} \cdot \frac{1}{2} \cdot \frac{1}{3} = \frac{1}{30}$$

And $\mathbb{P}(X_0 = 0, X_2 = 0) = \sum_{i=0}^1 \mathbb{P}(X_0 = 0, X_1 = i, X_2 = 0)$

ex $P_0 = I, p_0(i, j) = \delta_{ij} = \begin{cases} 1 & i = j \\ 0 & i \neq j \end{cases}$

$P_1 = P, p_1(i, j) = p(i, j)$

The distribution vector π_n for X_n is

$$\pi_n = (\pi_n(0), \pi_n(1), \dots)$$

where $\pi_n(i) = \mathbb{P}(X_n = i), i \in S$

Note that π_n is a probability vector

Property: Let $X_n \in S = \{1, 2, 3, \dots\}, n \geq 0$ be the time homogeneous Markov Chain with transition matrix P . Then

- P is a stochastic matrix

- $P \cdot \mathbb{I} = \mathbb{I}$ where $\mathbb{I} = \begin{bmatrix} 1 \\ 1 \\ \vdots \\ 1 \end{bmatrix}$
- 1 is an eigenvalue of P with eigenvector $\mathbb{I}$
- $P_{m+n} = P_m \cdot P_n$ (Chapman-Kolmogorov equation)
- $P_n = P^n$
- $\pi_n = \pi_0 P_n = \pi_0 P^n$
- $\pi_{n+1} = \pi_n P, \forall n \geq 0$

Proof: direct verification

e.g. $\mathbb{P}(X_1 = j) = \sum_{i \in S} \mathbb{P}(X_0 = i) \mathbb{P}(X_1 = j | X_0 = i)$

$\pi_1(j) = \sum \pi_0(j) P_{ij}$

Long-run behavior, invariant distribution

Def: Let $P = [p(i, j)]$ be the transition matrix for Markov Chain $\{X_n\}_{n=0}^{\infty}$ with $S = \{1, 2, \dots, N\}$. A probability vector $\pi \in \mathbb{R}^n$ is a stationary or invariant distribution if $\pi^P = \pi$

2.3 Lecture 11 - Invariant Distribution

- Recap: Discrete time/state, time homogeneous Markov Chains. Transition matrix $P = [p(i, j)]$, $p(i, j) = \mathbb{P}(X_1 = j | X_0 = i)$, $p_n(i, j) = \mathbb{P}(X_n = j | X_0 = i)$
- Markov Chains $\rightarrow$ MCMC

ex 3 state MC

$$P = \begin{bmatrix} \frac{3}{4} & \frac{1}{4} & 0 \\ \frac{1}{8} & \frac{2}{3} & \frac{5}{24} \\ 0 & \frac{1}{6} & \frac{1}{6} \end{bmatrix}$$

$$S = \{1, 2, 3\}$$

$$\mathbb{P}(X_1 = 1|X_0 = 1) = \frac{3}{4}, \text{etc.}$$

$$\lim_{n \rightarrow \infty} P^n = \begin{bmatrix} \pi \\ \pi \\ \pi \end{bmatrix}, \quad \pi = \left(\frac{2}{11}, \frac{4}{11}, \frac{5}{11} \right)$$

$$\begin{aligned} \text{Set } \pi_0 &= (\mathbb{P}(X_0 = 1), \mathbb{P}(X_0 = 2), \mathbb{P}(X_0 = 3)) \\ &= \mathbb{P}(X_0 = i) \end{aligned}$$

Then

$$\begin{aligned} \pi_n &= \pi_0 P^n \\ \lim_{n \rightarrow \infty} \pi_n &= \pi_0 \lim_{n \rightarrow \infty} P^n \\ &= \pi_0 \begin{bmatrix} \pi \\ \pi \\ \pi \end{bmatrix} \\ &= \left(\pi(1) \cdot \sum_i \pi_0(i), \pi(2) \cdot \sum_i \pi_0(i), \dots \right) \\ &= \pi \end{aligned}$$

Remarks

- $\lim_{n \rightarrow \infty} \mathbb{P}(X_n = j) = \pi(j) \forall j$
-

$$\begin{aligned} \pi_n &= \pi_{n-1} P \\ \lim_{n \rightarrow \infty} \pi_n &= \lim_{n \rightarrow \infty} \pi_{n-1} P \\ \pi &= \pi P \end{aligned}$$

- π is a left eigenvector of P with eigenvalue 1.

Def: Let $P = [p(i, j)]$ be the transition matrix for MC $\{X_n\}_{n=0}^\infty$ with $S = \{1, \dots, N\}$. A probability vector $\pi \in \mathbb{R}^n$ is a stationary or invariant distribution for X_n (or P) if: $\pi = \pi P$

Remarks

Above example shows $\lim_{n \rightarrow \infty} \pi_n =$ invariant distribution (with some conditions).

Questions: invariant distribution.

- When do they exist?
- Are they unique?
- Roughly: typically have unique invariant distribution unless reducible or periodic.

Def: A finite state time-homogeneous MC $X_n (n \geq 0)$ with transition matrix P is regular if there exists an integer $k \geq 1$, such that $P^k > 0$, (i.e. all entries of P^k are strictly > 0).

Thm

Let $X_n (n \geq 0)$ be a regular N-state Markov Chain with transition matrix P. Then:

1. $\{X_n\}_{n=0}^\infty$ has a unique stationary distribution π , $\pi > 0$
2. $\lim_{n \rightarrow \infty} P^n = \begin{bmatrix} \pi \\ \vdots \\ \pi \end{bmatrix}$, i.e. $\lim_{n \rightarrow \infty} p_n(i, j) = \pi(j)$
3. For any initial distribution π_0 , $\lim_{n \rightarrow \infty} \pi_0 P^n = \pi$ i.e. $\lim_{n \rightarrow \infty} \mathbb{P}(X_n = j) = \pi(j) \forall j$

Remarks

- 2. $\implies \lim_{n \rightarrow \infty} \mathbb{P}(X_n = j | X_0 = i) = \pi(j)$ independent of i

- From 2.:

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=0}^{n-1} p_k(i, j) = \lim_{n \rightarrow \infty} p_n(i, j) = \pi(j)$$

$$\begin{aligned} \text{And } \frac{1}{n} \sum_{k=0}^{n-1} p_k(i, j) &= \frac{1}{n} \sum_{k=0}^{n-1} \mathbb{P}(X_k = j | X_0 = i) \\ &= \frac{1}{n} \sum_{k=0}^{n-1} \mathbb{E}[\mathbb{I}[X_k = j] | X_0 = i] \\ &= \mathbb{E} \left[\frac{1}{n} \sum_{k=0}^{n-1} \mathbb{I}(X_k = j) | X_0 = i \right] \\ &\rightarrow \pi(j) \text{ as } n \rightarrow \infty \end{aligned}$$

Which means the mean fraction of time spent in $J \rightarrow \pi(j)$

How do we find π ?

- Solve $\pi P = \pi$
- Diagonalize $P = U \Lambda U^{-1}$

$$\begin{aligned} \Rightarrow \lim_{n \rightarrow \infty} P^n &= \lim_{n \rightarrow \infty} (U \Lambda U^{-1})^n \\ &= U \left(\lim_{n \rightarrow \infty} \Lambda^n \right) U^{-1} \\ &= \begin{bmatrix} \pi \\ \vdots \\ \pi \end{bmatrix} \end{aligned}$$

2.4 Lecture 12 - MCMC

Recap:

- Discrete time/space MC $\{X_n\}_{n=0}^{\infty}, X_n \in S$
- $P = [p(i, j)], p(i, j) = \mathbb{P}(X_1 = j | X_0 = i)$
- $\pi_0 = \mathbb{P}(X_0 = i), \pi_n = \mathbb{P}(X_n = i)$ (pmf X_n)

- Stationary distribution π s.t. $\pi P = \pi$
- Regular: $\exists k \geq 1$ s.t. $P^k > 0 \Rightarrow$ unique stationary distribution $\pi > 0$
 And $\lim_{n \rightarrow \infty} P^n = \begin{bmatrix} \pi \\ \vdots \\ \pi \end{bmatrix}$, $\lim_{n \rightarrow \infty} \pi_n = \pi$
- Detailed balance: probability vector λ s.t. $\lambda_i p(i, j) = \lambda_j p(j, i) \forall i, j \in S$
 $\Rightarrow \lambda P = \lambda \Rightarrow \lambda$ stationary with respect to P .

Remarks

1. Can view MC as evolution of probability vectors (pmfs): $\pi_0, \pi_1, \dots$ with $\pi_{n+1} = \pi_n P$. This is hard to compute for $|S|$ large.
2. MC is a random sequence of states $X_0, X_1, \dots \in S$ which is often easier to compute.

Recall: Discrete Inversion (Lecture 6)

e.g.: Generate discrete RV X

$$P(X = i) = p_i, \quad i = 1, 2, 3$$

$$p_1 + p_2 + p_3 = 1$$

1. $U \sim U[0, 1]$
2. $k = \min\{m \mid \sum_{i=1}^m p_i \geq U\}$
3. Set $X \leftarrow k$

Algorithm: Generate / simulate MC

Input: transition matrix P , initial distribution π_0

Output: MC $X_0, X_1, \dots \in S, X_n \sim \pi_k = \pi_0 P^k$

1. Generate $X_0 \sim \pi_0$
2. For $k = 1, \dots, n$, generate X_{n+1} from $p(X_k)$. i.e. pmf over S , given by the X_k 'th row of P .
3. Repeat n times to get samples from π_k

MCMC: (R&K ch.6)

- Key idea: generate distribution π by simulating MC with stationary distribution π
- Often π is known up to a normalizing constant.
- $\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n \mathbb{I}\{X_k = i\} = \pi(i) \quad \forall i \in S$
- $\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n g(X_k) = \mathbb{E}_\pi(g(x))$

We assume that

- RV $X \in S$ is finite, e.g. $S = \{1, 2, \dots, N\}$
- Target pmf of X is $\pi = (\pi(1), \dots, \pi_N)$, $\pi(i) = \mathbb{P}(X = i) \quad \forall i \in S$
- π has special form $\pi(i) = \frac{1}{z} b(i) > 0 \quad \forall i \in S$.
Where $b(i) > 0$ are known and $z > 0$ is unknown norm.

$$\sum_{i \in S} \pi(i) = 1 \iff \sum_{i \in S} b(i) = z$$

Remarks

- z is hard to compute, (sum of $N = |S|$ terms)
- z is called a partition function

Algorithm: Metropolis

- First construct proposal transition matrix $Q = [q(i, j)]$, where Q is symmetric.
- Choose initial state $X_0 \in S$ (e.g. randomly).

Given $X_k = i \in S$

1. Sample $Y \in S$ from $q(i, \cdot)$, i.e. $\mathbb{P}(Y = y | X_k = i) = q(i, y) \quad \forall y \in S$
Assume $Y = j$ is observed.
2. Let $\alpha = \alpha_{ij} = \min\left(1, \frac{\pi(j)}{\pi(i)}\right) \in [0, 1]$ and generate $U \sim U[0, 1]$.
If $U \leq \alpha$, accept Y , $X_{k+1} \leftarrow Y$
If $U > \alpha$, reject Y , $X_{k+1} \leftarrow X_k$
(Accept Y with probability α)
Output: $X_0, X_1, X_2, \dots$

Remarks

- $\{X_n\}_{n=0}^\infty$ is MC with $\lim_{n \rightarrow \infty} X_n \sim \pi$
- Can choose $Q = [q(i, j)]$ to be simple, e.g. uniform jumps, $q(i, j) = \frac{1}{N}$
- Only need the ratios $\frac{\pi(j)}{\pi(i)}$, so we don't need z .
- Call Y a proposed state.
- Call α the acceptance probability.
- In practice, $\alpha = a_{ij}$ is easy to compute.
- Can be extended to non-symmetric Q

ex 1D Ising model

$$\begin{aligned} S &= \{\sigma = (\sigma_1, \dots, \sigma_m) \mid \sigma_i = -1 \text{ or } +1, 1 \leq i \leq m\} \\ &= \{-1, +1\}^m \\ |S| &= 2^m \text{ states} \end{aligned}$$

2.5 Lecture 13 - Metropolis

Metropolis Algorithm

$$S = \{1, 2, \dots, N\}$$

Goal: Sample random variable X from some target distribution $\Pi(i) = \mathbb{P}(X = i) \implies \Pi_i(i) = \frac{1}{z}b(i)$

We have $Q = [q(i, j)]$ which is our transition matrix (symmetric).

1. Choose $X_0 \in S$. Given $X_n \in S$, sample $Y \in S$ from $q(X_n, i)$, i.e. $\mathbb{P}(Y = y \mid X_n = i)$
2. Set $\alpha = \min(1, \frac{\pi(y)}{\pi(X_n)}) \in [0, 1]$.
Generate $U \sim U[0, 1]$
If $U \leq \alpha$, accept $Y : X_{n+1} \leftarrow Y$
Otherwise, $U > \alpha$, reject Y , $X_{n+1} \leftarrow X_n$

1D Ising Model

$S = \{\sigma = \{\sigma_1, \sigma_2, \dots, \sigma_n\} : \sigma_j = \pm 1\}$
 $\Rightarrow 2^m$ states.

$$\Pi(\sigma) = \frac{1}{Z} e^{-\beta \mathcal{H}(\sigma)}$$
$$\mathcal{H} = -J \sum_{j=1}^{m-1} \sigma_j \sigma_{j+1} - h \sum_{j=1}^m \sigma_j$$
$$J, h > 0$$

Large $\Pi(\sigma) \longleftrightarrow$ Low energy states.
Calculate physical quantities.

$$m(\sigma) = \langle \sigma \rangle = \frac{1}{m} (N_+ - N_-)$$
$$= \frac{1}{m} \sum (N_+ - N_-) \Pi_p(\sigma)$$

$$\frac{\Pi_\beta(y)}{\Pi_\beta(X_n)} = e^{-\beta(\Delta \mathcal{H})}$$

a. $1 < i < m$

$$\Delta \mathcal{H} = 2J\sigma_i(\sigma_{i+1} + \sigma_{i-1}) + 2h\sigma_i$$

2.6 Lecture 14

Recap: Metropolis Chain

We have

- State space $S = \{1, 2, \dots, N\}$
- Proposed transition matrix $Q = [q(i, j)]$ which is symmetric
- Target distribution $\pi > 0$

Metropolis summary

- Given $X_k \in S$, choose y sampled from $q(X_k, \cdot)$, i.e. X_k 'th row of Q
- Accept Y with probability $\min(1, \frac{\pi(y)}{\pi(X_k)})$

- Output: MC $X_0, X_1, \dots, \in S$ with $X_n \sim \pi$ as $n \rightarrow \infty$

Convergence

- How big should n be to get $X_n \sim \pi$ 'roughly'?
- Hard to compute!
- Heuristic: Consider $R(n) = \text{Cov}(X_n, X_0)$

Thm

1. The transition matrix $P = [p(i, j)]$ of the Metropolis chain is

$$p(i, j) = \begin{cases} q(i, j) \min(1, \frac{\pi(j)}{\pi(i)}) & i \neq j \\ 1 - \sum_{k \in S, k \neq i} q(i, k) \min(\frac{\pi(k)}{\pi(i)}) & i = j \end{cases} \quad \forall i, j \in S$$

2. π and P satisfy detailed balance (DB)

$$\pi(i)p(i, j) = \pi(j)p(j, i) \quad \forall i, j \in S$$

Hence π is stationary

3. If $q(i, j) > 0 \forall i, j \in S$ and $p(i, i) > 0 \quad \forall i \in S$ Then π is a unique stationary distribution and $\lim_{n \rightarrow \infty} \pi_0 P^n = \pi$

Proof

1. For $i \neq j$

$$p(i, j) = \mathbb{P}(\text{propose } j|i) \cdot \mathbb{P}(\text{accept } j) \\ q(i, j) \min(1, \frac{\pi(j)}{\pi(i)})$$

Then

$$\sum_{j \in S} p(i, j) = 1 \quad \forall i \in S \\ \Rightarrow p(i, j) = 1 - \sum_{k \in S, k \neq i} q(i, k) \min(1, \frac{\pi(k)}{\pi(i)})$$

2. Want to show: $\pi(i)p(i, j) = \pi(j)p(j, i) \forall i, j \in S$.

This is always true for $i = j$

For $i \neq j$:

$$\begin{aligned}\pi(i)p(i, j) &= \pi(i)q(i, j) \min(1, \frac{\pi(j)}{\pi(i)}) \\ &= q(i, j) \min(\pi(i), \pi(j)) \\ &= q(j, i) \min(\pi(j), \pi(i)) \\ &= \pi(j)p(j, i) \Rightarrow DB\checkmark \\ \pi &\text{ is stationary (Lecture 11)}\end{aligned}$$

3. Conditions $\Rightarrow p(i, j) > 0 \forall i, j \in S \Rightarrow P$ is regular

$\Rightarrow \pi$ is unique and $\lim_{n \rightarrow \infty} \pi_0 P^n = \pi$

Remarks

- For Metropolis Algorithm applied to Ising model, we had:

$$S = \{\underline{\sigma} = (\sigma_1, \dots, \sigma_n), \sigma_j = \pm 1\}$$

Spin flips $(\sigma_1, \dots, \sigma_j, \dots, \sigma_n) \mapsto (\sigma_1, \dots, \sigma_j', \dots, \sigma_n)$

$$\Rightarrow q(\sigma, \sigma') = \begin{cases} \frac{1}{m} & \text{if } \sigma \mapsto \sigma' \text{ is 1 flip} \\ 0 & \text{otherwise} \end{cases}$$

- WE can show that this q gives a Metropolis chain which is regular
(Hint: any 2 states in S are at most m flips apart, $P^m > 0$)

The Gibbs Sampler

ex: Ising model (Glauber dynamics)

$$\text{States : } S = \{\sigma = (\sigma_1, \dots, \sigma_m), \sigma_j = \pm 1\}$$

$$\text{Target dist : } \pi, \pi(\sigma) = \frac{b(\sigma)}{Z} > 0 \forall \sigma \in S$$

$$\begin{aligned}\text{Notation : } \sigma^{j,+1} &= (\sigma_1, \sigma_2, \dots, \sigma_{j-1}, +1, \sigma_{j+1}, \dots, \sigma_m) \\ \sigma^{j,-1} &= (\sigma_1, \sigma_2, \dots, \sigma_{j-1}, -1, \sigma_{j+1}, \dots, \sigma_m)\end{aligned}$$

Glauber dynamics

(Random-scan Gibbs sampler for Ising model)

Goal: Construct MC, $X_0, X_1, \dots \in S$ s.t. $X_n \sim \pi$ as $n \rightarrow \infty$

Assume $X_k = \sigma \in S$

1. Choose J uniformly at random from $1, 2, \dots, m$

2. Set $\alpha_+ = \frac{\pi(\sigma^{J,+1})}{\pi(\sigma^{J,-1} + \sigma^{J,+1})}$ and $\alpha_- = \frac{\pi(\sigma^{J,-1})}{\pi(\sigma^{J,-1} + \sigma^{J,+1})}$

Note that $\alpha_+ + \alpha_- = 1$

Set $X_{k+1} \leftarrow \sigma^{J,+1}$, with probability α_+

$X_{k+1} \leftarrow \sigma^{J,-1}$, with probability α_-

Remarks

- Each 'spin' is in $\{-1, +1\}$
- $\{\alpha_+, \alpha_-\}$ is pmf over $\{-1, +1\}$
- Set J 'th spin conditional on the value of every other spin.

ex $m = 2$ spins

States : $S = \{(-1, -1), (-1, +1), (+1, -1), (+1, +1), \}$

$$\begin{aligned} \text{Set : } \pi(-1, -1) &= \frac{1}{z}, \pi(-1, +1) = \frac{2}{z} \\ \pi(+1, -1) &= \frac{3}{z}, \pi(+1, +1) = \frac{5}{z} \end{aligned}$$

(Here $z = 11$)

Suppose $X_k = (-1, -1)$, $J = 2$ (Glauber dynamics)

Then $\sigma^{J,+1} = (-1, +1)$, $\sigma^{J,-1} = (-1, -1)$

$$\alpha_+ = \frac{\frac{2}{z}}{\frac{1}{z} + \frac{2}{z}} = \frac{2}{3}, \alpha_- = \frac{1}{1+2} = \frac{1}{3}$$

Then $\mathbb{P}(X_{k+1} = (-1, +1) \mid X_k = (-1, -1)) = \frac{2}{3} \cdot \frac{1}{2} = \alpha_+ \cdot \mathbb{P}(J = 2)$

2.7 Lecture 15 - Midterm Review

Recap:

- Inversion, transformation
- Acceptance-rejection method
- MC integration, crude and importance sampling.
- Variance reduction.

- Markov chains (not MCMC)

MC Integration

Goal: Calculate

$$I = \int_D g(x)f(x)dx \quad D \subseteq \mathbb{R}^d$$

where D is bounded

$$f, g : D \rightarrow \mathbb{R}$$

f is a pdf on D

For $X \sim f$:

$$\begin{aligned} I &= \mathbb{E}_f(g(x)) \\ \text{Var}_f(g(x)) &= \mathbb{E}_f(g(x)^2) - [\mathbb{E}_f(g(x))]^2 \\ &= \int_D g(x)^2 f(x)dx - I^2 \end{aligned}$$

Crude MC estimator:

$$I_N = \frac{1}{N} \sum_{i=1}^N g(x_i) \quad X_1, \dots, X_N \sim f, iid$$

Then (check):

- $\mathbb{E}(I_N) = I$ (unbiased)
- $\text{Var} I_N = \frac{1}{N} \text{Var}_f(g(x_i))$
- $I_N \rightarrow I$ with probability 1

Importance sampling:

The importance function ϕ is a pdf on D .

Then

$$\begin{aligned} I &= \int_D \frac{g(x)f(x)}{\phi(x)} \phi(x)dx \\ &= \mathbb{E}_\phi \left[\frac{g(y)f(y)}{\phi(y)} \right], \quad y \sim \phi \end{aligned}$$

and we have our estimator

$$J_N = \frac{1}{N} \sum_{i=1}^N \frac{g(y_i)f(y_i)}{\phi(y_i)} \quad y_1, \dots, y_N \sim \text{iid}$$

Then (check):

- $\mathbb{E}(J_N) = I$ (unbiased)
- $\text{Var} J_N = \frac{1}{N} \left[\int_D \frac{g(x)^2 f(x)^2}{\phi(x)} dx - I^2 \right]$
- Choose ϕ to be easy to compute and to give small $\text{Var} J_N$

Variance Reduction

Let $U, V \sim U[0, 1]$, $\alpha, \beta : \mathbb{R} \rightarrow \mathbb{R}$

Set $W = \alpha(U) + \beta(V)$, W is an estimator for $\mu = \mathbb{E}(\alpha(U) + \beta(V)) = \mathbb{E}W$

Goal: Find relationship between U, V to reduce $\text{Var} W$

Note: $\text{Var} W = \text{Var}(\alpha(U)) + \text{Var}\beta(V) + 2\text{Cov}(\alpha(U), \beta(V))$

Note: $\text{Cov}(X, Y) = \mathbb{E}(XY) - \mathbb{E}(X)\mathbb{E}(Y)$ which = 0 if X, Y are independent.

1. α, β increasing
 $\rightarrow$ Use antithetic RVs: $V = 1 - U$
 $\text{Cov}(\alpha(U), \beta(1 - U)) \leq 0, \quad \alpha, \beta \uparrow$
2. $\alpha \uparrow, \beta \downarrow$
 $\rightarrow$ Use common RVs: $V = U$
 $\text{Cov}(\alpha(U), \beta(U)) \leq 0, \quad \alpha \uparrow, \beta \downarrow$
 See practice midterm Q4, HW2 Q4.

Markov Chain

- Discrete time/state, time homogeneous MC
 $X_0, X_1, \dots \in S = \{1, 2, 3, \dots, N\}$
- Transition matrix
 $P = [p(i, j)], \quad p(i, j) = \mathbb{P}(X_1 = j | X_0 = i) \quad \forall i, j \in S$
- Distribution vectors: $\pi_n = [\mathbb{P}(X_n = i)] \quad \forall i \in S$, pmf X_n
- $\pi_{n+1} = \pi_n P$ (HW3)

- Stationary/invariant distribution
 π s.t. $\pi P = \pi$
- Regular MC or transition matrix means that $\exists k \geq 1$ s.t. $P^k > 0$
 (Lecture 10, 11)
 Regular $\Rightarrow$ unique stationary distribution $\pi > 0$ and $\pi_n \rightarrow \pi$ as $n \rightarrow \infty$

ex $P = \begin{bmatrix} \frac{1}{2} & \frac{1}{3} \\ \frac{1}{3} & \frac{2}{3} \end{bmatrix}$ Note that this matrix is regular, with a unique stationary distribution $\Rightarrow \pi_0 P^n \rightarrow \pi$ for any π_0 .
 Solve $\pi P = \pi$, let $\pi = (a, b)$ (HW3, practice midterm)

$$\begin{aligned} \Rightarrow \frac{1}{2}a + \frac{1}{3}b &= a \\ \frac{1}{2}a + \frac{2}{3}b &= b \\ a + b &= 1 \end{aligned}$$

Solving this yields $a = \frac{2}{5}, b = \frac{3}{5}$

Acceptance-Rejection

We have

f : target pdf of $Z \in \mathbb{R}^d$
 αg : proposed pdf, $\alpha > 0$ s.t. $f(x) \leq \alpha g(x) \forall x$

Algorithm: sampling $Z \sim f$

1. Generate $X \sim g$
2. Generate $U \sim U[0, 1]$
3. Set $Y = \alpha g(x) \cdot U$
4. If $U \leq \frac{f(x)}{\alpha g(x)}$, then accept X
5. Return $Z \leftarrow X$

Efficiency: $\frac{1}{\alpha} = \frac{\text{vol B}}{\text{Vol A}} = \text{acceptance rate}$

2.8 Lecture 17 - MCMC & SDEs

Recap - MCMC

- Goal: sample from target distribution f on a state space S , by constructing MC $X_0, X_1, \dots \in S$ with $X_n \sim f$ as $n \rightarrow \infty$
- Output: $X_0, X_1, X_2, \dots \in S$
- For large enough n , $X_n, X_{n+1}, X_{n+2}, \dots$ is approximately a sample from f , but $X_n, X_{n+1}, X_{n+2}, \dots$ are dependent, not iid.
- For independent sample, we would need to run MCMC many times.
- Estimator: Suppose $X \sim f$ and $Y = g(X)$, we can use MCMC to estimate $\mathbb{E}Y$

$$\text{Let } Y_n = \frac{1}{n} \sum_{i=1}^n g(X_i)$$

Typically, $Y_n \rightarrow \mathbb{E}Y$ (HW4)

- Y_n can be used to assess convergence (often)
- Can also assess convergence with sample autocovariance. (HW4)

Gibbs Sampler

- Goal: sample a target pdf

$$f(x), \quad x \in \mathbb{R}^n$$

- Conditional distribution:

$$f_j(y|x_1, \dots, x_n) = \frac{f(x_1, \dots, x_{j-1}, y, x_j, \dots, x_{m+1})}{\int dy f(x_1, \dots, x_{j-1}, y, x_j, \dots, x_{m+1})}$$

for $1 \leq j \leq m$

So $f_j(\cdot | x_1, \dots, x_{m-1})$ is a pdf on $\mathbb{R}$

Algorithm: Random-scan Gibbs Sampler

Given $X_k = (X_{k,1}, \dots, X_{k,m}) \in \mathbb{R}^m$

1. Choose J in $\{1, \dots, m\}$ randomly (e.g. from unique distribution on $\{1, \dots, m\}$)
2. Sample Y from distribution:

$$f_J(\cdot \mid \underbrace{X_{k,1}, \dots, X_{k,J-1}, X_{k,J+1}, \dots, X_{k,m}}_{\in \mathbb{R}^{m-1}})$$

Set:

$$X_{k+1} \leftarrow \underbrace{(X_{k,1}, \dots, X_{k,J-1}, Y, X_{k,J+1}, \dots, X_{k,m})}_{\in \mathbb{R}^m}$$

ex

Target: $f(x, y) = c \exp(-(x^2 + 1)(y^2 + 1))$, $(x, y) \in \mathbb{R}^2$
 c is normalization so

$$\int_{\mathbb{R}^2} c \exp(-(x^2 + 1)(y^2 + 1)) dx dy = 1$$

What are the conditional distributions?

$$\begin{aligned} f_1(x|y) &= \frac{1}{\int_{\mathbb{R}} f(x, y) dx} \cdot c \exp(-(x^2 + 1)(y^2 + 1)) \\ &= c_1(y) \cdot \exp(-(y^2 + 1)x^2) \end{aligned}$$

where $\int_{\mathbb{R}} f(x, y) dx$ is a function of y and $c_1(y)$ is such that

$$\int_{-\infty}^{+\infty} c_1(y) \exp(-(y^2 + 1)x^2) dx = 1$$

i.e.: $c_1(y)$ is the normalization of the pdf

$$f_1(x|y) \propto \exp(-(y^2 + 1)x^2)$$

Recall that $W \sim N(0, \sigma^2)$ has pdf $f_w(w) \propto \exp(\frac{-1}{2\sigma^2} w^2)$

$$\Rightarrow f_1(x|y) \text{ is the pdf of } N\left(0, \frac{1}{2(y^2 + 1)}\right)$$

Similarly, $f_2(y|x)$ is pdf of $N(0, \frac{1}{2(x^2 + 1)})$

Algorithm (Random-scan Gibbs)

Given $(x_k, y_k) \in \mathbb{R}^2$

1. Choose J uniformly at random from $\{1, 2\}$
2. Generate $Z \sim N(0, 1)$
 - If $J = 1$
 - Set $W = \frac{1}{\sqrt{2(y_k^2+1)}}z$ (i.e. $W \sim f_1(\cdot | y_k)$)
 - Set $(x_{k+1}, y_{k+1}) \leftarrow (w, y_k)$
 - If $J = 2$ Set $W = \frac{1}{\sqrt{2(x_k^2+1)}}z$ (i.e. $W \sim f_2(\cdot | x_k)$)
 - Set $(x_{k+1}, y_{k+1}) \leftarrow (x_k, w)$
3. Output: $(x_1, y_1), (x_2, y_2), \dots$

Systemic-scan Gibbs Sampler

- Given: $X_k = (X_{k,1}, \dots, X_{k,m}) \in \mathbb{R}^m$
 - Input: target pdf $f(x)$, $x \in \mathbb{R}^m$
1. Sample Y_1 from $f_1(\cdot | X_{k,2}, \dots, X_{k,m})$
 - For $i = 2$ to m
 - Sample Y_i from

$$f_i(\cdot | y_1, \dots, y_{i-1}, x_k)$$