

Question 1

(a) $L = (\partial_x + x\partial_y)$

$$\begin{aligned}
L(u + v) &= (\partial_x + x\partial_y)(u + v) \\
&= u_x + v_x + xu_y + xv_y \\
&= Lu + Lv \\
L(cu) &= (\partial_x + x\partial_y)(cu) \\
&= cu_x + cxu_y \\
&= c(u_x + xu_y) \\
&= cLu
\end{aligned}$$

Therefore L is a linear operator.

(b) $L = (\partial_x + u_y\partial_y)$

$$\begin{aligned}
L(u + v) &= (\partial_x + (u_y + v_y)\partial_y)(u + v) \\
&= u_x + v_x + (u_y + v_y)u_y + (u_y + v_y)v_y \\
&= u_x + v_x + u_y^2 + 2u_yv_y + v_y^2 \\
&\neq u_x + u_y^2 + v_x + v_y^2 = Lu + Lv
\end{aligned}$$

Therefore, L is NOT a linear operator.

(c) $L = (\partial_x + \partial_y + \frac{1}{u})$

$$\begin{aligned}
L(u + v) &= (\partial_x + \partial_y + \frac{1}{u+v})(u + v) \\
&= u_x + v_x + u_y + v_y + 1 \\
&\neq (u_x + u_y + 1) + (v_x + v_y + 1) = Lu + Lv
\end{aligned}$$

Therefore, L is NOT a linear operator.

(d) $L = (\cos(x)\partial_{xy} + \arctan(\frac{y}{x})\partial_y)$

$$\begin{aligned}
 L(u+v) &= (\cos(x)\partial_{xy} + \arctan(\frac{y}{x})\partial_y)(u+v) \\
 &= \cos(x)u_{xy} + \arctan(\frac{y}{x})u_y + \cos(x)v_{xy} + \arctan(\frac{y}{x})v_y \\
 &= Lu + Lv \\
 L(cu) &= (\cos(x)\partial_{xy} + \arctan(\frac{y}{x})\partial_y)(cu) \\
 &= c\cos(x)u_{xy} + c\arctan(\frac{y}{x})u_y \\
 &= c(\cos(x)u_{xy} + \arctan(\frac{y}{x})u_y) \\
 &= cLu
 \end{aligned}$$

Therefore, L is a linear operator.

Question 2

Suppose that v and w are solutions to the inhomogeneous linear PDE $Lu = g$. Then we have

$$Lv = g \qquad Lw = g$$

Take their difference

$$\begin{aligned}
 Lv - Lw &= g - g \\
 L(v - w) &= 0
 \end{aligned}$$

Since the PDE is linear, $v - w$ is a solution to the homogeneous equation $Lu = 0$.

Question 3

(a)

$$\begin{aligned}
 u_x &= 0 \\
 u &= f(y)
 \end{aligned}$$

(b)

$$\begin{aligned}
 u_x &= x^2 + y \\
 u &= \frac{1}{3}x^3 + yx + f(y)
 \end{aligned}$$

(c)

$$\begin{aligned}
 u_{yy} + u &= 0 \\
 u &= -u_{yy} \\
 &= f(x) \cos(y) + g(x) \sin(y)
 \end{aligned}$$


(d)

$$\begin{aligned}
 u_{xy} + u_y &= 0 \\
 u_y &= -u_{xy} \\
 u &= -u_x \\
 u &= \cos(x) + f(y)
 \end{aligned}$$

Question 4


(a)

$$\begin{aligned}
 A(x, y) &= (2, -1) \\
 u(x, y) &= f(-x - 2y)
 \end{aligned}$$



(b)

$$\begin{aligned}\frac{dx}{d\eta} &= 1 + x^2 & \frac{dy}{d\eta} &= 1 \\ \frac{dy}{dx} &= \frac{1}{1 + x^2} \\ y &= \arctan x + \xi \\ u(\xi) &= f(y - \arctan x)\end{aligned}$$



(c)

$$\frac{dx}{d\eta} = -y \quad \frac{dy}{d\eta} = x$$

$$\frac{dy}{dx} = -\frac{x}{y}$$

$$ydy = -xdx$$

$$\frac{1}{2}y^2 = -\frac{1}{2}x^2$$

$$y = \pm ix + \xi$$

$$u(\xi) = f(y \pm ix)$$

Question 5(a) $yu_x + xu_y = 0$

$$\frac{dx}{d\eta} = y \quad \frac{dy}{d\eta} = x$$


$$\frac{dy}{dx} = \frac{x}{y}$$

$$ydy = xdx$$

$$\frac{1}{2}y^2 = \frac{1}{2}x^2 + \xi$$

$$y^2 = x^2 + \xi$$

$$u(\xi) = f(y^2 - x^2)$$



(b) Given $u(0, y) = y^2$

$$u(0, y) = f(y^2) = y^2$$

$$u(x, y) = (y^2 - x^2)^2$$

Question 6

$$u_x + y u_y = -u$$