

Selamat! Anda lulus!

Nilai diterima 100%

Nilai Pengiriman Terbaru 100%

UNTUK LULUS 80% atau lebih tinggi

Pergi ke item berikutnya

1. Which of the following can address overfitting?

1 / 1 poin

- ☐ Remove a random set of training examples
- ☒ Select a subset of the more relevant features.

Benar
If the model trains on the more relevant features, and not on the less useful features, it may generalize better to new examples.

- ☒ Apply regularization


Benar
Regularization is used to reduce overfitting.

- ☒ Collect more training data

Benar
If the model trains on more data, it may generalize better to new examples.

2. You fit logistic regression with polynomial features to a dataset, and your model looks like this.

1 / 1 poin



What would you conclude? (Pick one)

- ☐ The model has high bias (underfit). Thus, adding data is, by itself, unlikely to help much.
- ☒ The model has high variance (overfit). Thus, adding data is likely to help
- ☐ The model has high bias (underfit). Thus, adding data is likely to help
- ☐ The model has high variance (overfit). Thus, adding data is, by itself, unlikely to help much.

Benar
The model has high variance (it overfits the training data). Adding data (more training examples) can help.

3. Regularization

1 / 1 poin

$$\min_{\vec{w}, b} J(\vec{w}, b) = \min_{\vec{w}, b} \left[\underbrace{\frac{1}{2m} \sum_{i=1}^m (f_{\vec{w}, b}(\vec{x}^{(i)}) - y^{(i)})^2}_{\text{mean squared error}} + \underbrace{\frac{\lambda}{2m} \sum_{j=1}^n w_j^2}_{\text{regularization term}} \right]$$

Suppose you have a regularized linear regression model. If you increase the regularization parameter λ , what do you expect to happen to the parameters w_1, w_2, \dots, w_n ?

- ☒ This will reduce the size of the parameters w_1, w_2, \dots, w_n
- ☐ This will increase the size of the parameters w_1, w_2, \dots, w_n

Benar
Regularization reduces overfitting by reducing the size of the parameters w_1, w_2, \dots, w_n .