

Module:
Techniques in Neuroscience

Week 3:
Immunohistochemistry: Preserving and studying cells of the brain

**Carl Hobbs**

Voice over by Dr Brenda Williams

Topic 1:
**An introduction to
immunohistochemistry**
Part 4 of 4

Part 4

Introduction to part 4

Principles of immunotechniques

Enzyme based
detection methodsFluorescence based
detection methods

Principles of immunodetection: terminology


Immunodetection methods

- take advantage of the interaction between antibodies and their molecular targets in tissues, called antigens
- it makes it possible to not only distinguish between cell types but also to visualise the location of proteins


Terminology

Antigen: Molecule which induces an immune response in our bodies, in particular, the production of antibodies.

Epitope: Small sequence or part of the antigen recognised by the antibody.




Antibody production: monoclonal



Michnick & Sidhu (2008)


Antibody production: polyclonal



Memory B cells:


Cells that remember the same antigen enabling faster antibody production if the same antigen is seen again.

Immunoglobulin structure




Antibodies can be divided into five different types: IgG, IgM, IgA, IgD and IgE (most antibody reagents are IgG or occasionally IgM).


Immunohistochemistry protocol



Detection methods

Fluorescence



Used to detect the presence of more than one protein at the same time.

Enzyme

Sequence in immunostaining of sections

- 1) Incorporation of positive and negative controls
- 2) Antigen retrieval
- 3) Blocking of non-specific binding

Incorporation of positive controls



BrdU staining in the hippocampus as a means to assess neural stem cell proliferation.

Positive control



BrdU labelling in the small intestine

Lundgren et al. (2011); Van Praag et al. (1999)

Incorporation of negative controls

Negative controls

- it is mostly sufficient to omit the primary antibody (using the normal serum from the animal that the secondary antibody was raised in instead)
- one then compares the results of the negative control and the positive controls against the test result before drawing a conclusion

Antigen unmasking/retrieval

Fixation procedures can mask or alter epitopes so that they can no longer bind to the primary antibody.

Antigen unmasking or retrieval refers to any technique where the masking of an epitope is reversed so that the antibody can again bind to it.

Antigen unmasking methods

Heat induced epitope retrieval (HIER)

- Citric acid pH6
- Citrate buffer pH6
- Tris pH9
- Tris/EDTA pH9
- EDTA pH8
- Tris pH10

Protease-induced epitope retrieval

- Proteinase K
- Trypsin
- Pepsin



Blocking non-specific binding

Serum:

- contains proteins that will bind to non-specific sites

Protein – BSA (bovine serum albumin):

- compete with antibodies for non-specific binding sites



To consider when immunostaining of sections

- 1) Inclusion of positive and negative controls
- 2) Do I need to use antigen retrieval?
- 3) Blocking of non-specific binding


Example of indirect IF using 2 antibodies

Paraffin wax section from a **mouse model of Alzheimers Disease** showing astrocytes surrounding an amyloid plaque.



Method used:

- double indirect immunofluorescence staining
- incubated simultaneously with two primary antibodies that each recognised a different protein (glial fibrillary acidic protein (GFAP) and beta amyloid)



Nuclei are stained with a fluorescent blue stain called DAPI.


Example of indirect enzyme IHC

Paraffin wax section from a normal mouse brain showing an **area of the Hippocampus**.




Method used:

- stained with an anti-GFAP antibody that is expressed by astrocytes in the section
- the binding of this antibody was detected by indirect immunoperoxidase staining using DAB



Counterstained with Haemalum (a blue dye) to show all nuclei.

Summary



References

Cho, Y. S., Cho, J. H., Shin, B. N., Cho, G. S., Kim, I. H., Park, J. H., ... & Hong, S. (2015). Ischemic preconditioning maintains the immunoreactivities of glucokinase and glucokinase regulatory protein in neurons of the gerbil hippocampal CA1 region following transient cerebral ischemia. *Molecular medicine reports*, 12(4), 4939-4946.

Glantz, L. A., & Lewis, D. A. (2000). Decreased dendritic spine density on prefrontal cortical pyramidal neurons in schizophrenia. *Archives of general psychiatry*, 57(1), 65-73.

Hermes, J., Anliker, B., Heber, S., Ring, S., Fuhrmann, M., Kretzschmar, H., ... & Müller, U. (2004). Cortical dysplasia resembling human type 2 lissencephaly in mice lacking all three APP family members. *The EMBO journal*, 23(20), 4106-4115.

Lacroix, S., Hamilton, L. K., Vaugois, A., Beaudoin, S., Breault-Dugas, C., Pineau, I., ... & Fernandes, K. J. (2014). Central canal ependymal cells proliferate extensively in response to traumatic spinal cord injury but not demyelinating lesions. *PloS one*, 9(1), e85916.

Lavado, A., He, Y., Paré, J., Neale, G., Olson, E. N., Giovannini, M., & Cao, X. (2013). Tumor suppressor Nf2 limits expansion of the neural progenitor pool by inhibiting Yap/Taz transcriptional coactivators. *Development*, 140(16), 3323-3334.

Lundgren, O., Jodal, M., Jansson, M., Ryberg, A. T., & Svensson, L. (2011). Intestinal epithelial stem/progenitor cells are controlled by mucosal afferent nerves. *PloS one*, 6(2), e16296.

Michnick, S. W., & Sidhu, S. S. (2008). Submitting antibodies to binding arbitration. *Nature chemical biology*, 4(6), 326-329.

References

Milatovic, D., Montine, T. J., Zaja-Milatovic, S., Madison, J. L., Bowman, A. B., & Aschner, M. (2010). Morphometric analysis in neurodegenerative disorders. *Current protocols in toxicology*, 12-16.

Van Praag, H., Kempermann, G., & Gage, F. H. (1999). Running increases cell proliferation and neurogenesis in the adult mouse dentate gyrus. *Nat neurosci*, 2(3), 266-270.

Attributions

Anonymous. (circa 1900). Franz Nissl [photograph]. Retrieved from <https://commons.wikimedia.org/w/index.php?curid=4680740>

Mattei G, Cristiani I, Magliaro C, Ahluwalia A. (2014) Profile analysis of hepatic porcine and murine brain tissue slices obtained with a vibratome. PeerJ PrePrints 2:e561v1 <https://doi.org/10.7287/peerj.preprints.561v1>

Mumssygris. (2015). Cresyl violet staining is commonly used in histology to stain nervous tissues [digital reproduction]. Retrieved from <https://commons.wikimedia.org/w/index.php?curid=38220541>

OpenStax. (2016). Figure 2: Polyclonal and Monoclonal Antibody Production [illustration]. Retrieved from <http://cnx.org/contents/55953ebd-7069-45cf-a114-aa7fe120640c@3>.
Download for free at <http://cnx.org/contents/55953ebd-7069-45cf-a114-aa7fe120640c@3>.

Pr495d. (2006). Golgi-stained neurons from somatosensory cortex in the macaque monkey [photograph]. Retrieved from <https://commons.wikimedia.org/w/index.php?curid=2542549>

Ramón y Cajal, S. (1911). Drawing of the neural circuitry of the rodent hippocampus [drawing]. Retrieved from <https://commons.wikimedia.org/w/index.php?curid=612536>

Unknown author. (1899). Santiago Ramón y Cajal. Spanish Nobel laureate in medicine [digital reproduction]. Retrieved from <https://commons.wikimedia.org/w/index.php?curid=12334552>

Attributions

Unknown author. (1906). Camillo Golgi [digital reproduction]. Retrieved from <https://commons.wikimedia.org/w/index.php?curid=10308907>

Uthman, E. (2006). Tissue Processing 06 [photograph]. Retrieved from <https://commons.wikimedia.org/w/index.php?curid=5158960>

End of topic