

Module:
Techniques in neuroscience

Week 4:
Tissue culture: Growing and studying neural cells in a dish




Dr Graham Cocks

Topic 1:
An introduction to tissue culture
Part 2 of 2

Part 2

Antibiotics


Widespread use of the antibiotics, penicillin and streptomycin in the 1940s onwards reduced the problem of microbial contamination of cultures.



These are ineffective against certain common strains of bacteria such as mycoplasma.

Mycoplasma

Mycoplasma are very small
(<1 micron in length)




- can be a significant problem in long-term culture
- some antibiotics are effective against mycoplasmas
- best practice is to prevent contamination by employing an aseptic technique

Biological safety cabinets

Biological safety cabinets are one of the most important developments in improving aseptic techniques.


Class I cabinets



Characteristics:

- protects both the user and the environment from the sample
- does not protect the sample from airborne particles from the environment


Class II cabinets



Characteristics:

- protects sample from outside contamination
- relies on a continuous uniform flow of clean filtered air travelling down over the sample

Class III cabinets




Characteristics:

- completely encloses the sample, which can only be accessed through the gloves integrated into the cabinet

Kruse et al. (1991)

Tissue culture incubators



Tissue culture incubators maintain a **number of critical parameters** to **allow optimal growth and survival**:


- constant levels of temperature, humidity, CO₂ and oxygen
- most media currently in use for tissue culture use buffers that require an atmosphere of 5 per cent CO₂ to maintain a physiological pH
- some cell types grow better under low oxygen conditions – specific incubators can reduce oxygen levels with displacement by nitrogen

Primary cells, cell lines and cell strains



Leonard Hayflick & Paul Moorhead
(1961)

Derived the first strains of human fibroblasts (WI-38)



Young human diploid cells (strain WI-38)

Made the distinction between primary cells, cell lines and cell strains:

- primary cells are derived from normal tissue and grown without passaging
- cell strains are derived from primary cells which have a limited capacity for growth and division, but retain a normal karyotype
- cell lines have the capacity to grow indefinitely and invariably have abnormal karyotypes

Karyotype:

the size, shape, and number of chromosomes in a cell

Hayflick & Moorhead (1961)

Week 4 Tissue culture: Growing and studying neural cells in a dish

Topic 1: An introduction to tissue culture

7 of 17

Immortal human cell lines




George Otto Gey
(1951)

Gey cultured cells from an individual called Henrietta Lacks who had cervical cancer.

Finding:

Cells derived from the cervical tumour could grow and divide indefinitely.



HeLa cells

- aided the development of the first Polio vaccine
- still being used for research today


Scherer et al. (1953)

Week 4 Tissue culture: Growing and studying neural cells in a dish

Topic 1: An introduction to tissue culture

8 of 17

Embryonic stem cells



Martin Evans
(1982)

Established cultures of cells derived from mouse blastocysts:

- can in principle generate any cell type of the body in a cell culture dish

First generated human embryonic stem cells from human blastocysts:

- allowed for the generation of inaccessible cell types, such as neurons, in large numbers for the first time
- ethical issues need to be considered



Jamie Thomson
(1998)

Evans & Kaufman (1981); Thomson et al. (1998)

Week 4 Tissue culture: Growing and studying neural cells in a dish

Topic 1: An introduction to tissue culture 9 of 17


Induced pluripotent stem cells



Shinya Yamanaka
(2006, 2007)

Directly generated embryonic stem cells by directly manipulating fibroblasts in a process called **reprogramming:**

- produced the so-called induced pluripotent stem cells (iPSCs)
- used to study genetic diseases and inaccessible cell types
- no ethical issues as those surrounding embryonic stem cells




Takahashi et al. (2007); Takahashi & Yamanaka (2006)


Week 4 Tissue culture: Growing and studying neural cells in a dish

Topic 1: An introduction to tissue culture 10 of 17


Visual examples of iPS cells, neurons and neural 'rosettes'



iPS cells growing in culture



Neural 'rosettes' surrounded by neurons




neurons fixed and stained with fluorescent markers

The neural rosettes are composed of radial glial cells (green) which in turn generate intermediate progenitors (red) which then in turn generate neurons (blue). These rosettes are considered to be a 2-D cell culture model of cortical neural tube development.

Click **Next** to continue

Regenerative medicine



Following on from the development of human iPSCs is the exciting prospect of using these cells for personalised **regenerative medicine**.

Major advantage:

Transplanted cells will be genetically identical to that individual, eliminating the risk of rejection or the use of immunosuppressive drugs.

References

- Carrel, A., & Burrows, M. T. (1911). Cultivation of tissues in vitro and its technique. *The Journal of experimental medicine*, 13(3), 387.
- Evans, M. J., & Kaufman, M. H. (1981). Establishment in culture of pluripotential cells from mouse embryos. *Nature*, 292(5819), 154-156.
- Harrison, R. G., Greenman, M. J., Mall, F. P., & Jackson, C. M. (1907). Observations of the living developing nerve fiber. *The Anatomical Record*, 1(5), 116-128.
- Hayflick, L., & Moorhead, P. S. (1961). The serial cultivation of human diploid cell strains. *Experimental cell research*, 25(3), 585-621.
- Jedrzejszak-Slicka, M. (2017). History of Cell Culture. In S. J. T., Gowder (Ed.). *New Insights into Cell Culture Technology* (pp1-41). Rijeka, Croatia: InTech Publishers
- Kruse, R. H., Puckett, W. H., & Richardson, J. H. (1991). Biological safety cabinetry. *Clinical Microbiology Reviews*, 4(2), 207-241.
- Lewis, M. R., & Lewis, W. H. (1911). The cultivation of tissues from chick embryos in solutions of NaCl, CaCl₂, KC1 and NaHCO₃. *The Anatomical Record*, 5(6), 277-293.
- Rous, P., & Jones, F. S. (1916). A method for obtaining suspensions of living cells from the fixed tissues, and for the plating out of individual cells. *Journal of Experimental Medicine*, 23(4), 549-555.

References

- Scherer, W. F., Syverton, J. T., Gey, G. O. (1953). Studies on the propagation in vitro of poliomyelitis viruses. IV. Viral multiplication in a stable strain of human malignant epithelial cells (strain HeLa) derived from an epidermoid carcinoma of the cervix. *The Journal of experimental medicine*, 97(5), 695-710.
- Takahashi, K., Tanabe, K., Ohnuki, M., Narita, M., Ichisaka, T., Tomoda, K., & Yamanaka, S. (2007). Induction of pluripotent stem cells from adult human fibroblasts by defined factors. *cell*, 131(5), 861-872.
- Takahashi, K., & Yamanaka, S. (2006). Induction of pluripotent stem cells from mouse embryonic and adult fibroblast cultures by defined factors. *cell*, 126(4), 663-676.
- Thomson, J. A., Itskovitz-Eldor, J., Shapiro, S. S., Waknitz, M. A., Swiergiel, J. J., Marshall, V. S., & Jones, J. M. (1998). Embryonic stem cell lines derived from human blastocysts. *science*, 282(5391), 1145-1147.

Attributions

Ambrose, C. T. (2017). The Tissue Culture Laboratory of Dr. George Otto Gey 60 yrs ago as recalled by a former student. *In Vitro Cellular & Developmental Biology-Animal*, 1-7.

Cardiff University. (2007). *Martin Evans Nobel Prize* [photograph]. Retrieved from <http://www.cardiff.ac.uk>, CC BY 3.0, <https://commons.wikimedia.org/w/index.php?curid=2893679>

Gitschier, J., PLoS Genetics. (2008). *Jamie Thomson* [photograph]. Retrieved from <https://commons.wikimedia.org/w/index.php?curid=22003886>

Marine Biological Laboratory Archives. (1922). *Embryo Project Encyclopedia (1922)* ISSN: 1940-5030. Retrieved from <http://embryo.asu.edu/handle/10776/2505>

National Institutes of Health (NIH). (2011). HeLa-III [digital reproduction]. Retrieved from <https://commons.wikimedia.org/w/index.php?curid=29540593>

National Institutes of Health. (2013). *Dr. Shinya Yamanaka* [photograph]. Retrieved from <https://commons.wikimedia.org/w/index.php?curid=27101450>

Public Domain. (2010). *Wilhelm Roux (9 June 1850 – 15 September 1924) was a German zoologist and pioneer of experimental embryology* [photograph]. Retrieved from <https://commons.wikimedia.org/w/index.php?curid=10859537>

Public Domain. (n.a.) *French surgeon and biologist Alexis Carrel (1873-1944)* [photograph]. Retrieved from <https://commons.wikimedia.org/w/index.php?curid=2837993>

Attributions

Shay, J. W., & Wright, W. E. (2000). Hayflick, his limit, and cellular ageing. *Nature reviews Molecular cell biology*, 1(1), 72-76.

Smithsonian Institution Archives (1920-1970s). *Montrose Thomas Burrows (1884-1947)* [photograph]. Retrieved from https://siarchives.si.edu/collections/siris_arc_290619

Smithsonian Institution from United States. (n.a.). *Margaret Reed Lewis (1881-1970)* [photograph]. Retrieved from <https://commons.wikimedia.org/w/index.php?curid=45232068>

Source. (2013). *Ross Granville Harrison* [photograph]. Retrieved from <https://en.wikipedia.org/w/index.php?curid=17679355>

Wadman, M. (2013). Cell division. *Nature*, 498(7455), 422.

Weiss, R. A., & Vogt, P. K. (2011). 100 years of Rous sarcoma virus. *Journal of Experimental Medicine*, 208(12), 2351-2355.

End of topic