

INSTITUTE OF PSYCHIATRY, PSYCHOLOGY & NEUROSCIENCE



Module:

Techniques in Neuroscience

Week 5:

Molecular biology: Going inside the cell

Dr Elizabeth Glennon

Topic 1:

An introduction to molecular biology methods

Part 3 of 3

Part 3

Wook

Molecular biology: Going inside the cell

Topic 1: An introduction to molecular biology methods

2 of 14

Lecture outline



In this section, we will study the western blotting process:

Western blotting Limitations of western blotting

Examples of western blotting

Click Next to continue

Week!

Molecular biology: Going inside the cell

Topic 1: An introduction to molecular biology methods

3 of 14

Introduction to western blotting




Introduction


- western blotting is used to detect specific proteins from a mixture of proteins
- during western blotting, we are processing further the polyacrylamide gel
- the technique is used to transfer the proteins from the gel onto a membrane


Molecular biology: Going inside the cell


Topic 1: An introduction to molecular biology methods

4 of 14









Why use western blotting?



Advantages:

- takes 11/2 days (can be faster)
- does not require a lot of specialist equipment
- will work for a wide range of proteins
- can be sensitive and specific with good antibodies
- can be applied to many cells/tissues/experimental procedures
- semi-quantitative




Disadvantages:


- only as good as your antibody
- antibodies are expensive
- post-translational modifications may alter binding or mobility
- · does not work as well if your proteins are very large or very small
- reasonably sensitive, but does not work well if protein levels are low
- · semi-quantitative

Molecular biology: Going inside the cell

Topic 1: An introduction to molecular biology methods

Real life examples of western blotting: BIN1 Control **Sporadic AD** = 100 kDa 70 kDa 50 kDa BIN1 levels in people with AD, compared to age-matched controls BIN1 has an unclear role in the development of AD the authors determined how BIN1 is altered in AD Western blotting showed a signal reduction in the AD samples compared to control, thus indicating a reduction of the protein. Glennon et al. (2013) Molecular biology: Going inside the cell 10 of 14 Topic 1: An introduction to molecular biology methods





References

Boxus, M., Twizere, J. C., Legros, S., Dewulf, J. F., Kettmann, R., & Willems, L. (2008). The HTLV-1 tax interactome. Retrovirology, 5(1), 76.

Glennon, E. B., Whitehouse, I. J., Miners, J. S., Kehoe, P. G., Love, S., Kellett, K. A., & Hooper, N. M. (2013). BIN1 is decreased in sporadic but not familial Alzheimer's disease or in aging. PLoS One, 8(10), e78806.

Stoica, R., De Vos, K. J., Paillusson, S., Mueller, S., Sancho, R. M., Lau, K. F., ... & Dickson, D. W. (2014). ER-mitochondria associations are regulated by the VAPB-PTPIP51 interaction and are disrupted by ALS/FTD-associated TDP-43. Nature communications, 5.

Week

Molecular biology: Going inside the cell

Topic 1: An introduction to molecular biology methods

13 of 14

End of topic

Wook

Molecular biology: Going inside the cell

Topic 1: An introduction to molecular biology methods

14 of 14