

Stem Cells and Epilepsy: Modelling the Brain with Organoids

Daniel Burger

King's College London
public@danielburger.online

13. February 2024

Abstract

Table of Contents

List of Figures	II
1 Introduction	1
1.1 Traditional Models for Epilepsy Research	2
1.2 Current State of the Art	2
2 Discussion	3
2.1 Stem-Cell-Derived Models for Epilepsy	3
2.2 Case Studies and Practical Applications	4
2.3 Other Approaches and Advancements	5
2.4 Ethical Considerations	5
3 Conclusion	6
References	7

List of Figures

1.1 EEG Waveforms: Epileptic vs. Normal Brain Activity	1
--	---

1 Introduction

Epilepsy, a neurological disorder afflicting around 50 million individuals worldwide ([World Health Organization, 2019](#)), manifests through recurrent, unprovoked seizures, which can lead to symptoms such as convulsions (i.e. uncontrolled shaking), loss of consciousness, and unusual sensations or behaviours. The development of seizures in epilepsy is mainly due to a complex interaction between the excitatory and inhibitory activities of neurons in the brain. When there is an imbalance that leads to more excitation than inhibition, it can cause sudden and abnormal electrical activity in localised or entire regions of the brain ([Robinson et al., 1997](#)).

Temporal lobe epilepsy (TLE) is the most common form ([Epilepsy Foundation, 2019](#))—yet it is the drug-resistant epilepsies, especially those without an identifiable epileptogenic zone, that present the most significant treatment challenges ([Iwasaki et al., 2016](#); [Guery & Rheims, 2021](#)). Current interventions, such as neuromodulation ([Fisher & Velasco, 2014](#)), offer some hope, but their effectiveness is variable and can be marred by severe side effects, highlighting the urgent need for novel research directions.

The impact of epilepsy extends far beyond its physiological symptoms, affecting every facet of an individual’s life. A large number of people who are diagnosed suffer from uncontrollable seizures, rendering them incapable of performing routine tasks, securing employment, or even driving—a prohibition enforced in many countries, such as e.g. Switzerland, due to safety concerns ([Schweizerische Epilepsie-Liga, 2021](#)). This pervasive uncertainty cultivates a lifestyle fraught with limitations, emphasising the disease’s profound societal and personal toll.




Figure 1.1: EEG Waveforms: Epileptic vs. Normal Brain Activity. Comparative display of EEG waveforms showing brain electrical activity over 10 seconds. The top graph represents EEG data during an epileptic seizure, characterised by high amplitude and frequent spikes, indicative of abnormal neuronal activity. The bottom graph illustrates a normal EEG with regular wave patterns and lower amplitude, reflecting typical brain function. The x-axis measures time in seconds, and the y-axis measures the amplitude of brain waves in microvolts (μV). Image taken from [Espinosa et al. \(2020\)](#).

To fully grasp the essence of epilepsy, one must consider not only its physical manifestations but also the profound psychological toll it exacts on sufferers. The unpredictability of seizures can instill a persistent fear and sense of helplessness, significantly affecting mental health. The origins of epilepsy are diverse, encompassing genetic predispositions (genotype) and observable characteristics (phenotype), including brain injuries and infections, underscoring the complexity of its etiology. The dichotomy between normal and epileptic brain function is starkly illustrated in EEG recordings, as exemplarily shown in [Figure 1.1](#). These visual representations not only highlight the aberrant electrical activity characteristic of seizures but also underscore the critical need for a deeper understanding of the underlying mechanisms of epilepsy. Such insights are essential for paving the way toward more targeted and efficacious treatments, addressing the disease’s genotype and phenotype.

1.1 Traditional Models for Epilepsy Research

Traditional epilepsy research models, such as in vivo animal subjects and in silico simulations, have been invaluable yet present notable limitations. Animal models, often e.g. employing rodents ([Wang et al., 2022](#)), provide insights but may not fully translate to human epilepsy due to differences in brain structure and function ([Kandratavicius et al., 2014](#)). Similarly, in silico models, like those developed by the Blue Brain Project ([Markram, 2006](#)), offer detailed simulations of neural networks but are limited by current computational capabilities and understanding of the brain ([Mirza et al., 2016](#)).

Stem-cell-derived models, mainly from human induced pluripotent stem cells (hiPSCs), emerge as a promising solution, offering a more accurate and ethical approach to studying epilepsy. Unlike animal models, which face translational hurdles due to species-specific differences in brain architecture and function, hiPSCs can be derived from human cells, ensuring a closer representation of human pathophysiology. Furthermore, the use of hiPSCs sidesteps the ethical quandaries associated with embryonic stem cell research, as they can be obtained from adult cells (e.g. skin cells) without harm to the donor, thereby aligning with ethical standards for human research ([Takahashi & Yamanaka, 2006](#)). These models circumvent traditional methods’ limitations by providing a renewable source of human neural tissue and enabling the exploration of epilepsy’s neurodevelopmental and pathophysiological aspects in a patient-specific context while also allowing the precise analysis and image of the tissues in a highly controlled environment.

1.2 Current State of the Art

The advances in stem cell technology have given rise to another groundbreaking approach in the study of epilepsy: the cultivation of three-dimensional brain organoids derived from hiPSCs. Compared to the first approaches where neuronal cells were cultivated in a two-dimensional manner (sometimes called dish brains), researchers can now generate these cerebral organoids that mimic the brain’s complex architectures in terms of tissue structures and the arrangement of cellular types, cell-to-cell interactions, and synaptic connectivity, surpassing two-dimensional dish brains that fall short in replicating the

physiological interactions, regional specificity, and microenvironment gradients observed in vivo (Clevers, 2016; Wang, 2018).

Another benefit of cultivating brain organoids is that it allows researchers to grow or 3D-print neural tissue, e.g. around electrically conductive matrices or electrodes (Yao et al., 2023), engaging with and measuring the organoids in ways reminiscent of living brains. Additionally, combining multiple brain organoids as an assembloid offers a sophisticated method to study the interactions between different brain regions (Sloan et al., 2018). These assembloids recreate the complex neural networks and facilitate investigations into the inter-regional synaptic connections critical for higher-order brain functions, often disrupted in epilepsy.

These state-of-the-art techniques offer a transformative avenue for epilepsy research and beyond. Using hiPSCs to reflect a patient’s unique genetic makeup, brain organoids serve as personalised models to decipher the complex interplay of factors driving epileptogenesis. The insights garnered through such increasingly precise 3D models promise to accelerate the discovery of innovative treatments, aiming to improve the quality of life for individuals with neurological diseases.

2 Discussion

2.1 Stem-Cell-Derived Models for Epilepsy

One of the most compelling aspects of hiPSC technology is its ability to model genetic forms of epilepsy. By deriving iPSCs from patients with known genetic mutations, researchers can observe the direct effects of these mutations on neuronal development, function, and network formation. This approach has been instrumental in elucidating the pathophysiological mechanisms underlying syndromes such as Dravet Syndrome and Tuberous Sclerosis Complex, where specific gene mutations lead to distinct epileptic phenotypes (Jiao et al., 2013; Nadadthur et al., 2019).

Despite these advances, stem-cell-derived models of epilepsy face several challenges. The differentiation of iPSCs into fully mature, functional neurons and glial cells that accurately represent the diversity and complexity of the human brain remains a daunting task. This is particularly pertinent in modelling age-related epilepsies, where the disease phenotype may only manifest or worsen over time. As mentioned earlier, techniques such as 3D brain organoids have emerged as a promising solution to mimic the brain’s architecture and cellular heterogeneity more closely, though they are not without their own set of limitations, including the lack of vascularization and an immune system (Lancaster et al., 2013; Di Lullo & Kriegstein, 2017).

Recent studies, such as those by Thodeson et al. (2017), underscore the potential of neural stem cells in epilepsy research. These investigations reveal that iPSC models can uncover novel insights into diseases with epileptic phenotypes, despite challenges such as variable expression profiles and differentiation potential among iPSC lines. For example, iPSC models of Rett syndrome have illuminated critical aspects of epileptogenesis by demonstrating decreases in neuronal soma size, neurite outgrowth, and synapse formation compared to controls, thereby highlighting the intricate interplay between neuronal and astrocytic contributions to the disease (Marchetto et al., 2010).

The evaluation of stem-cell-derived models' effectiveness and limitations, as discussed by [Kandemir et al. \(2022\)](#), points to the nuanced relationship between neurogenesis and epilepsy. These models not only provide deep insights into the pathological underpinnings of epilepsy but also illuminate potential therapeutic avenues. However, the specificity of neurogenesis markers and the role of mature astrocytes in epilepsy remain areas of ongoing research, emphasizing the need for continued innovation and refinement in stem-cell-derived epilepsy models ([Jessberger & Parent, 2015](#)).

2.2 Case Studies and Practical Applications

Building on the foundational concepts discussed earlier, we now shift our focus to two pivotal case studies and practical examples. These instances illuminate the real-world application of stem-cell-derived models in epilepsy research, showcasing the depth of insights gained and the potential pathways to therapeutic advancements.

- **Samarasinghe et al. (2021)** delved into the complex neural dynamics of brain organoids, uncovering significant epileptiform activities within organoids modeling Rett syndrome, which is a genetic disorder that typically affects females and is characterised by impairments in language and coordination, repetitive movements, slower growth, difficulty walking and so on. Complications can include seizures, scoliosis, and sleeping problems. However, [Samarasinghe et al.](#) groundbreaking work demonstrated not only the presence of sophisticated physiological activities within these organoids but also the potential for therapeutic intervention. Remarkably, they observed a substantial reduction in epileptiform activities upon administration of pifithrin- α , a TP53 inhibitor, which points towards new directions in epilepsy treatment, especially for conditions exhibiting resistance to conventional therapies.
- **Steinberg et al. (2020)** embarked on an ambitious project to model epileptic encephalopathies using a combination of CRISPR-engineered human ES cells and patient-derived iPSCs, with a focus on the devastating WOREE syndrome. Their meticulous approach unveiled significant cellular and molecular CNS abnormalities, including alterations in GABAergic markers which suggest a disruption in the development of normal and balanced neuronal networks. This work not only underscores the intricate pathophysiology of epileptic encephalopathies but also provides a proof-of-concept for potential therapeutic interventions, such as the modulation of GABAergic responses, which are crucial in the dynamics of developmental epilepsies.

These case studies emphasize the critical importance of accurately mimicking human brain tissue in research to provide a platform for studying epilepsy in a controlled, ethical manner. The advancements in stem-cell-derived models, particularly organoids, open up new avenues for direct therapeutic research and personalized treatment strategies. By enabling the study of individualized brain organoids from patients with epilepsy, researchers can explore tailored treatments for specific cases, thereby enhancing the efficacy and precision of epilepsy management. Furthermore, these models hold promise for innovative treatments, such as the potential use of organoids for cell-based therapies in epilepsy, by genetically modifying implants to address the unique needs of each

patient's condition.

2.3 Other Approaches and Advancements

It should have become evident so far that stem cell technology shows promise in epilepsy research, especially what is to come to tackle this disease. Among the most promising developments is the transplantation of hiPSC-derived brain organoids into living beings, a technique that has shown significant potential in increasing inhibition in epileptic brain areas. [Hunt & Baraban](#)'s (2015) pioneering work on interneuron transplantation laid the groundwork for this approach, demonstrating the feasibility and therapeutic potential of grafting new inhibitory neurons into the epileptic brain.

Building on this foundation, [Neurona Therapeutics](#) (2022) has made significant strides with its lead cell therapy candidate, NRTX-1001, which is currently being evaluated in a Phase 1/2 clinical trial for people with drug-resistant mesial temporal lobe epilepsy (mTLE). The therapy involves a one-time administration of fully-differentiated neural cells, called interneurons, derived from human pluripotent stem cells and intended to provide long-term GABAergic inhibition to repair hyperexcitable neural networks. Early results from the clinical trial are promising, with significant seizure suppression and no serious adverse events reported in patients treated with NRTX-1001. These findings suggest that NRTX-1001 could represent a groundbreaking, non-destructive treatment for epilepsy, offering hope for durable seizure reduction and improved quality of life for individuals living with this debilitating condition.

The intersection of neuroprosthetics and epilepsy research presents another frontier. Neuroprosthetics, which include devices capable of interacting with the nervous system, hold the potential not only for treatment but also for a deeper understanding of epilepsy's mechanisms. The concept of augmenting cognition through synthetic biological neural tissue, as discussed in "Augmenting Cognition" by [Segev & Markram](#) (2011), where one of the book's authors, Mijail Demian Serruya, suggests the expansion of neural substrates through artificial models, opens up possibilities for compensating lost brain functions due to epilepsy.

2.4 Ethical Considerations

While working with stem cells, we are nonetheless confronted with many ethical considerations that demand our attention. [Farahany et al.](#) (2018) provide a comprehensive overview of the ethical landscape when experimenting with human brain tissue, raising critical questions about the potential for consciousness in brain organoids and the moral implications of in vitro experimentation.

The ethical discourse surrounding brain organoids primarily hinges on their increasing complexity and the theoretical possibility of these entities attaining a form of consciousness or subjective experiences. This prospect, albeit currently remote, necessitates a careful examination of the moral status of these organoids and the protections they might warrant, akin to those provided to human or animal research subjects. One of the significant ethical challenges lies in the limitations of current brain organoid models, particularly their lack of a comprehensive vascular system, which restricts their growth and complexity, as mentioned in [subsection 2.1](#). Therefore, the potential to

create sentient brain organoids in laboratories emerges as a topic of ongoing debate and research, necessitating a cautious and principled approach to ensure that we do not inadvertently cross ethical boundaries.

In light of these considerations, engaging in a broad and inclusive dialogue involving civil society, researchers, ethicists, and policymakers is paramount. Such discourse should aim to develop robust ethical frameworks and guidelines that can adapt to the rapid advancements in this field, ensuring that the research proceeds in a manner that respects the potential moral status of brain organoids and is mindful of the broader implications for society.

3 Conclusion

TODO: - Summarising the key points, emphasising the impact of stem-cell-derived models compared to other models, especially e.g. in silico or animal testing. - Discussing future prospects of stem-cell-derived models in neuroscience, considering technological advancements and potential breakthroughs. - Addressing remaining challenges, including technical, ethical, and funding-related issues, to give a balanced view of the field's future.

Personal note: Basically conclude and have a key focus on that this field needs to get more funding, as brain organoids coupled with personalised medicine (also sometimes called precision medicine) and gene-editing is a very promising field of research and has the potential to cure epilepsy and other neurological diseases. Also, mention somehow neuroprosthetics again the the whole point from Mijail Demian Serruya's part of the "Augmenting Cognition" book from Markram.

References

- Clevers, H. (2016). Modeling Development and Disease with Organoids. *Cell*, 165(7), 1586–1597. Publisher: Elsevier.
URL: [https://www.cell.com/cell/abstract/S0092-8674\(16\)30729-2](https://www.cell.com/cell/abstract/S0092-8674(16)30729-2) (Accessed at: 2024-02-11)
- Di Lullo, E., & Kriegstein, A. R. (2017). The use of brain organoids to investigate neural development and disease. *Nature Reviews. Neuroscience*, 18(10), 573–584.
- Epilepsy Foundation (2019). Temporal Lobe Epilepsy (TLE).
URL: <https://www.epilepsy.com/what-is-epilepsy/syndromes/temporal-lobe-epilepsy> (Accessed at: 2024-02-11)
- Espinosa, N., Amorim, A., & Huebner, R. (2020). *Feedforward Neural Network with Backpropagation for Epilepsy Seizure Detection*.
- Farahany, N. A., Greely, H. T., Hyman, S., Koch, C., Grady, C., Paşca, S. P., Sestan, N., Arlotta, P., Bernat, J. L., Ting, J., Lunshof, J. E., Iyer, E. P. R., Hyun, I., Capetany, B. H., Church, G. M., Huang, H., & Song, H. (2018). The Ethics of Experimenting With Human Brain Tissue.
- Fisher, R. S., & Velasco, A. L. (2014). Electrical brain stimulation for epilepsy. *Nature Reviews Neurology*, 10(5), 261–270. Number: 5 Publisher: Nature Publishing Group.
URL: <https://www.nature.com/articles/nrneurol.2014.59> (Accessed at: 2024-02-11)
- Guery, D., & Rheims, S. (2021). Clinical Management of Drug Resistant Epilepsy: A Review on Current Strategies. *Neuropsychiatric Disease and Treatment*, 17, 2229–2242. Publisher: Dove Press.
URL: <https://www.dovepress.com/clinical-management-of-drug-resistant-epilepsy-a-review-on-current-str-peer-reviewed-fulltext-article-NDT> (Accessed at: 2024-02-11)
- Hunt, R. F., & Baraban, S. C. (2015). Interneuron Transplantation as a Treatment for Epilepsy.
- Iwasaki, M., Jin, K., Nakasato, N., & Tominaga, T. (2016). Non-invasive Evaluation for Epilepsy Surgery. *Neurologia medico-chirurgica*, 56(10), 632–640.
- Jessberger, S., & Parent, J. M. (2015). Epilepsy and Adult Neurogenesis. *Cold Spring Harbor Perspectives in Biology*, 7(12), a020677.
URL: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4665072/> (Accessed at: 2024-02-12)
- Jiao, J., Yang, Y., Shi, Y., Chen, J., Gao, R., Fan, Y., Yao, H., Liao, W., Sun, X.-F., & Gao, S. (2013). Modeling Dravet syndrome using induced pluripotent stem cells (iPSCs) and directly converted neurons. *Human Molecular Genetics*, 22(21), 4241–4252.

- Kandemir, C., Yavuz, M., Karakaya, F. B., Çilingir Kaya, O. T., Onat, F., & Şirvanci, S. (2022). Investigation of Neurogenesis in Kindled Wistar and Genetic Absence Epilepsy Rats. *Clinical and Experimental Health Sciences*, 12(3), 753–759. Number: 3 Publisher: Marmara University.
URL: <https://dergipark.org.tr/en/pub/clinexphealthsci/issue/72731/1021171> (Accessed at: 2024-02-11)
- Kandratavicius, L., Balista, P. A., Lopes-Aguiar, C., Ruggiero, R. N., Umeoka, E. H., Garcia-Cairasco, N., Bueno-Junior, L. S., & Leite, J. P. (2014). Animal models of epilepsy: use and limitations. *Neuropsychiatric Disease and Treatment*, 10, 1693–1705. Publisher: Dove Press.
URL: <https://www.dovepress.com/animal-models-of-epilepsy-use-and-limitations-peer-reviewed-fulltext-article-NDT> (Accessed at: 2024-02-11)
- Lancaster, M. A., Renner, M., Martin, C.-A., Wenzel, D., Bicknell, L. S., Hurles, M. E., Homfray, T., Penninger, J. M., Jackson, A. P., & Knoblich, J. A. (2013). Cerebral organoids model human brain development and microcephaly. *Nature*, 501(7467), 373–379. Number: 7467 Publisher: Nature Publishing Group.
URL: <https://www.nature.com/articles/nature12517> (Accessed at: 2024-02-12)
- Marchetto, M. C. N., Carromeu, C., Acab, A., Yu, D., Yeo, G. W., Mu, Y., Chen, G., Gage, F. H., & Muotri, A. R. (2010). A model for neural development and treatment of Rett syndrome using human induced pluripotent stem cells. *Cell*, 143(4), 527–539.
- Markram, H. (2006). The Blue Brain Project. *Nature Reviews Neuroscience*, 7(2), 153–160. Number: 2 Publisher: Nature Publishing Group.
URL: <https://www.nature.com/articles/nrn1848> (Accessed at: 2024-02-11)
- Mirza, N., Vasieva, O., Appleton, R., Burn, S., Carr, D., Crooks, D., du Plessis, D., Duncan, R., Farah, J. O., Josan, V., Miyajima, F., Mohanraj, R., Shukralla, A., Sills, G. J., Marson, A. G., & Pirmohamed, M. (2016). An integrative in silico system for predicting dysregulated genes in the human epileptic focus: Application to SLC transporters. *Epilepsia*, 57(9), 1467–1474. _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1111/epi.13473>.
URL: <https://onlinelibrary.wiley.com/doi/abs/10.1111/epi.13473> (Accessed at: 2024-02-11)
- Nadadhur, A. G., Alsaqati, M., Gasparotto, L., Cornelissen-Steijger, P., van Hugte, E., Dooves, S., Harwood, A. J., & Heine, V. M. (2019). Neuron-Glia Interactions Increase Neuronal Phenotypes in Tuberous Sclerosis Complex Patient iPSC-Derived Models. *Stem Cell Reports*, 12(1), 42–56.
- Neurona Therapeutics (2022). Neurona Therapeutics Presents Preclinical Data at the American Academy of Neurology (AAN) Annual Meeting from Lead Cell Therapy Candidate, NRTX-1001, Being Evaluated in a Phase 1/2 Clinical Trial for Chronic Focal Epilepsy.
URL: <https://www.neuronatherapeutics.com/news/press-releases/040122/> (Accessed at: 2024-02-04)

- Robinson, P., Rennie, C., & Wright, J. (1997). Propagation and stability of waves of electrical activity in the cerebral cortex. *Phys. Rev. E*, 56.
- Samarasinghe, R. A., Miranda, O. A., Buth, J. E., Mitchell, S., Ferando, I., Watanabe, M., Allison, T. F., Kurdian, A., Fotion, N. N., Gandal, M. J., Golshani, P., Plath, K., Lowry, W. E., Parent, J. M., Mody, I., & Novitch, B. G. (2021). Identification of Neural Oscillations and Epileptiform Changes in Human Brain Organoids.
- Schweizerische Epilepsie-Liga (2021). Driving with epilepsy.
URL: <https://www.epi.ch/en/about-epilepsy/in-depth-information/driving-with-epilepsy/> (Accessed at: 2024-02-11)
- Segev, I., & Markram, H. (2011). *Augmenting Cognition*. Lausanne (Suisse): EPFL Press, 1st edition ed.
- Sloan, S. A., Andersen, J., Paşca, A. M., Birey, F., & Paşca, S. P. (2018). Generation and Assembly of Human Brain Region-Specific Three-Dimensional Cultures. *Nature protocols*, 13(9), 2062–2085.
URL: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6597009/> (Accessed at: 2024-02-11)
- Takahashi, K., & Yamanaka, S. (2006). Induction of pluripotent stem cells from mouse embryonic and adult fibroblast cultures by defined factors. *Cell*, 126(4), 663–676.
- Thodeson, D. M., Brulet, R., & Hsieh, J. (2017). Neural Stem Cells and Epilepsy: Functional Roles and Disease-in-a-Dish Models.
- Wang, H. (2018). Modeling Neurological Diseases With Human Brain Organoids. *Frontiers in Synaptic Neuroscience*, 10.
URL: <https://www.frontiersin.org/articles/10.3389/fnsyn.2018.00015> (Accessed at: 2024-02-11)
- Wang, Y., Wei, P., Yan, F., Luo, Y., & Zhao, G. (2022). Animal Models of Epilepsy: A Phenotype-oriented Review. *Aging and Disease*, 13(1), 215–231.
URL: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8782545/> (Accessed at: 2024-02-11)
- World Health Organization (2019). Epilepsy: a public health imperative.
URL: <https://apps.who.int/iris/bitstream/handle/10665/325293/9789241515931-eng.pdf> (Accessed at: 2024-02-07)
- Yao, Y., Coleman, H. A., Meagher, L., Forsythe, J. S., & Parkinson, H. C. (2023). 3D Functional Neuronal Networks in Free-Standing Bioprinted Hydrogel Constructs. *Advanced Healthcare Materials*, 12(28), 2300801. _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1002/adhm.202300801>.
URL: <https://onlinelibrary.wiley.com/doi/abs/10.1002/adhm.202300801> (Accessed at: 2024-02-11)