

Two Closed Negatives in the MODFORM Axis

A hexa cusp-form-dimension definition gap, and the non-lift of the $n = 6$ arithmetic identity to the modular-curve tower

TECS-L domain, `hexa-lang`
dancinlab

github.com/dancinlab/hexa-lang

May 26, 2026

Abstract

The MODFORM axis of the TECS-L domain re-grounds the classical invariants of the congruence subgroup $\Gamma_0(N)$ — the index $\psi(N)$, the cusp count, the genus of $X_0(N)$, the cusp-form weight, and the Atkin–Lehner group order — as machine-checked closed-form recomputations under a single verification discipline (`hexa verify`, tier rubric §3). Against that verified backdrop we pre-register two falsifiers and report two *closed-negative* findings. *Finding 1 (MF4)*. We pre-registered that the `hexa` builtin `dim_cusp_forms(N,2)` realizes the standard theorem $\dim S_2(\Gamma_0(N)) = \text{genus}(X_0(N))$. Sweeping $N \in [1, 30]$ the two quantities agree for $N \leq 7$ but disagree for 20 of the 30 levels (e.g. $N = 11$: `hexa` = 0 vs. classical = 1; $N = 30$: `hexa` = 6 vs. classical = 3). The falsifier is ∇ FALSIFIED: the builtin has a *definition gap*, not the mathematics. *Finding 2 (F7)*. We pre-registered that the $\{1, 6\}$ -type identity $\sigma(n)\phi(n) = n\tau(n)$ — which is exactly the MODFORM-axis $n = 6$ specialness — lifts to a $\{1, 6\}$ -type index identity at the $\Gamma_1(N)/X(N)$ level of the modular-curve tower. The $\Gamma_1(N)$ and $X(N) = \Gamma(N)$ indices are smooth (multiplicative) in N with no $n = 6$ peak; the falsifier is ∇ FALSIFIED: the specialness is a Γ_0 -level arithmetic-function phenomenon that does *not* lift. Both negatives are pre-registered, measured, and deterministic. They rule out two distinct axes — a trusted machine builtin, and a tower-level continuation of an arithmetic accident — and the $\Gamma_0(N)$ -level $n = 6$ bridge survives both intact.

1 Introduction

The modular curve $X_0(N) = \Gamma_0(N) \backslash \mathbb{H}^*$ carries a short list of arithmetic invariants whose closed forms are classical [2, 3]: the index $\psi(N) = [\text{SL}_2(\mathbb{Z}) : \Gamma_0(N)] = N \prod_{p|N} (1 + 1/p)$, the cusp count $\sum_{d|N} \phi(\gcd(d, N/d))$, the genus of $X_0(N)$, the dimension of the weight-2 cusp-form space $S_2(\Gamma_0(N))$, and the Atkin–Lehner involution group of order $2^{\omega(N)}$ [1]. A theme that runs through the TECS-L domain is that the smallest non-trivial level $N = 6$ is special: each of these invariants of $X_0(6)$ reduces to a standard $n = 6$ arithmetic-function value — $\psi(6) = 12 = \sigma(6)$, four cusps = $\tau(6)$, genus 0, first weight 4 = $\tau(6)$, $|\text{AL}| = 4 = 2^{\omega(6)}$ — and the perfect-number identity $\sigma(n)\phi(n) = n\tau(n)$ holds exactly at $n \in \{1, 6\}$ (the “identity locus,” the subject of a companion paper [5]).

A claim that “ $N = 6$ is special” is scientific only to the extent that it comes with a *falsifier* fixed before the computation. This paper isolates and resolves two such falsifiers, both of which turn out negative — and both of which are, in the TECS-L discipline, valuable precisely because they are closed negatives [6].

The first falsifier is internal to the tooling. The TECS-L verification discipline trusts that a `hexa` builtin named after a classical quantity *computes* that quantity. For weight-2 cusp forms the textbook fact is $\dim S_2(\Gamma_0(N)) = \text{genus}(X_0(N))$ [2]. We pre-registered that the builtin `dim_cusp_forms(N,2)` realizes this identity, sweeping it against the independently

trusted `gamma0_genus(N)` (which MF3 had already validated against 22 classical genus values). The sweep falsifies the assumption: the two functions diverge for 20 of the 30 levels $N \in [1, 30]$. This is a *definition gap* in the builtin, not a mathematical falsehood; the gap was filed upstream (PR #1083).

The second falsifier asks whether the $n = 6$ specialness is a property of the *number* 6 that propagates up the modular-curve tower $\Gamma_0(N) \rightarrow \Gamma_1(N) \rightarrow \Gamma(N)$, or whether it is confined to the Γ_0 -level arithmetic-function identities. We pre-registered that a $\{1, 6\}$ -type index identity — a closed-form singularity, peak, or arithmetic collapse at $N = 6$ — exists at the $\Gamma_1(N)/X(N)$ level. The index formulas $[\mathrm{SL}_2(\mathbb{Z}) : \Gamma_1(N)] = \psi(N)\phi(N)/2$ and $[\mathrm{SL}_2(\mathbb{Z}) : \Gamma(N)] = N \cdot [\mathrm{SL}_2(\mathbb{Z}) : \Gamma_1(N)]$ are smooth multiplicative functions of N with no distinguished behaviour at 6; the falsifier is closed-negative. The $n = 6$ specialness is real, but it lives at the Γ_0 -level and at the arithmetic-function level — it does not lift.

The contribution of this paper is therefore (i) a verified $\Gamma_0(N)$ backdrop (§3, §4); (ii) two pre-registered falsifiers, each resolved to a deterministic closed negative (§5); and (iii) an honest delineation of what each negative does and does not rule out (§6). Every numeric quantity in the paper traces to a verbatim `hexa verify` verdict persisted under `.verdicts/` and indexed in `CLAIMS.tape` [4].

2 Background

The $\Gamma_0(N)$ invariants. For a positive integer N , the congruence subgroup $\Gamma_0(N) \subset \mathrm{SL}_2(\mathbb{Z})$ consists of matrices that are upper-triangular modulo N . Its index is the Dedekind psi function $\psi(N) = N \prod_{p|N} (1 + 1/p)$. The quotient $X_0(N)$ is a compact Riemann surface; the number of cusps is $\sum_{d|N} \phi(\gcd(d, N/d))$ and the genus g follows from the Riemann–Hurwitz formula. The space $S_2(\Gamma_0(N))$ of weight-2 cusp forms has dimension equal to the genus, $\dim S_2(\Gamma_0(N)) = g(X_0(N))$, by the standard correspondence between weight-2 cusp forms and holomorphic differentials [2]. The Atkin–Lehner involutions W_Q for $Q \parallel N$ generate an abelian 2-group of order $2^{\omega(N)}$, where $\omega(N)$ is the number of distinct prime factors [1].

The modular-curve tower. Above $\Gamma_0(N)$ sit the finer congruence subgroups $\Gamma_1(N)$ (lower-left entry $\equiv 0$, lower-right $\equiv 1 \pmod{N}$) and the principal congruence subgroup $\Gamma(N)$ (full reduction to the identity mod N), with $\Gamma(N) \subset \Gamma_1(N) \subset \Gamma_0(N)$. For $N > 2$ their indices are

$$[\mathrm{SL}_2(\mathbb{Z}) : \Gamma_1(N)] = \frac{1}{2} \psi(N) \phi(N) = \frac{N^2}{2} \prod_{p|N} (1 - \frac{1}{p^2}), \quad [\mathrm{SL}_2(\mathbb{Z}) : \Gamma(N)] = N \cdot [\mathrm{SL}_2(\mathbb{Z}) : \Gamma_1(N)],$$

both multiplicative and smoothly increasing in N . The curve $X(N)$ associated to $\Gamma(N)$ has index N times that of $\Gamma_1(N)$.

The $n = 6$ arithmetic identity. The companion TECS-L result [5] establishes that $\sigma(n)\phi(n) = n\tau(n)$ holds at exactly $n \in \{1, 6\}$ over a tested range, where σ is the divisor sum, ϕ the totient, and τ the divisor count. At $n = 6$ all of $\sigma(6) = 12$, $\phi(6) = 2$, $\tau(6) = 4$ conspire so that $\sigma(6)\phi(6) = 24 = 6 \cdot \tau(6)$. This is the arithmetic-function “specialness” whose lift to the modular-curve tower Finding 2 tests.

3 Method: the verification discipline

Every numeric claim in this paper is a closed-form recomputation issued through the `hexa verify` command and persisted verbatim. We use four tiers:

- **◆ formal** — a `hexa`-native closed-form recomputation that equals its expected value exactly (`hexa verify --expr <fn> <n> <v>` reporting `SUPPORTED-FORMAL`). These are the verified $\Gamma_0(N)$ invariants.
- **■ numerical** — a `libm`-class value to a fixed tolerance. None appear here; the `MOD-FORM` axis is integer-exact.
- **▽ closed-negative** — a pre-registered falsifier that a recomputation deterministically refutes, ruling out one axis.
- *citation / deferred* — residuals deliberately gated *out* of any terminal claim (e.g. the assembled Γ_1/X indices, which combine **◆** integer components with a cited closed form, are marked citation and never used as a headline).

Falsifier 1 (MF4 — dim vs. genus).

Falsifier 1. The `hexa` builtin `dim_cusp_forms(N,2)` realizes the standard theorem $\dim S_2(\Gamma_0(N)) = \text{genus}(X_0(N))$; i.e. for every N , `dim_cusp_forms(N,2) = gamma0_genus(N)`.

Method: sweep $N \in [1, 30]$, recomputing both builtins, and record per- N agreement. The reference `gamma0_genus` was independently validated in MF3 against 22 classical genus values (15 classical genus-0 levels plus a genus- ≥ 1 boundary), so a disagreement localizes the fault to `dim_cusp_forms`.

Falsifier 2 (F7 — $n = 6$ non-lift).

Falsifier 2. A $\{1, 6\}$ -type index identity — a closed-form singularity, peak, or arithmetic-function collapse distinguishing $N = 6$ from its neighbours, echoing $\sigma(n)\phi(n) = n\tau(n) \iff n \in \{1, 6\}$ — exists at the $\Gamma_1(N)/X(N)$ level of the modular-curve tower.

Method: assemble the $\Gamma_1(N)$ and $X(N)$ indices from **◆** integer components ($\psi(N) = \text{gamma0_index}$, $\phi(N)$) via the cited closed forms, for $N \in \{5, 6, 7, 11, 12\}$ spanning 6 and its neighbours, and inspect whether $N = 6$ sits at a distinguished value. Smoothness (the indices being exactly the generic multiplicative formula output at 6) refutes the falsifier.

The full driver is the `HEAD hexa verify --expr` engine (PR #1153). Raw standard output of every run is persisted verbatim under `.verdicts/tecs-1-modform-*/` and indexed in `CLAIMS.tape` [4]; representative transcripts are reproduced in Appendix C.

4 Verification: the real runs

4.1 The verified $\Gamma_0(N)$ backdrop

The `MODFORM` axis first recomputes the $\Gamma_0(N)$ invariants on $N \in [1, 30]$, all **◆**. The headline counts (verbatim from the persisted verdicts) are:

invariant	sweep	tier count
index $\psi(N)$ (<code>gamma0_index</code>)	$N \in [1, 30]$	30/30 ◆
cusps $\sum \phi(\gcd(d, N/d))$ (<code>gamma0_cusps</code>)	$N \in [1, 30]$	30/30 ◆
genus $g(X_0(N))$ (<code>gamma0_genus</code>)	22 classical	22/22 ◆
quadratic symbols (<code>jacobi/kronecker</code>)	15 instances	15/15 ◆
first cusp-form weight (<code>first_cusp_form_weight</code>)	$N \in [1, 30]$	30/30 ◆

Crucially, `gamma0_genus` matches all 22 classical/known genus values with *zero* mismatches — 15 classical genus-0 levels $N \in \{1-10, 12, 13, 16, 18, 25\}$ and the genus- ≥ 1 boundary ($N = 11, 14, 15, 17, 19 \rightarrow 1$; $N = 22, 23 \rightarrow 2$). This makes `gamma0_genus` a *trusted* reference, which is exactly what Falsifier 1 leans on. The $n = 6$ bridge over this backdrop is $\psi(6) = 12 = \sigma(6)$, four cusps $= \tau(6)$, $g(X_0(6)) = 0$, first weight $4 = \tau(6)$, $|\text{AL}(6)| = 4 = 2^{\omega(6)}$ — all $\blacklozenge$ (the AL row citation).

4.2 Finding-1 run: `dim_cusp_forms` vs. `gamma0_genus`

Sweeping $N \in [1, 30]$ and comparing `dim_cusp_forms(N, 2)` against the trusted `gamma0_genus(N)` gives exactly **20** MISMATCH and **10** MATCH (full table in Appendix A). The match region is the small- N coincidence $N \leq 7$ (plus 9, 13, 25); the divergence begins at $N = 8$ (`hexa` = 1, classical = 0) and persists. Two boundary instances are independently $\blacklozenge$ -verified:

N	<code>dim_cusp_forms(N, 2)</code>	<code>gamma0_genus(N)</code>	verdict
6	0 $\blacklozenge$	0 $\blacklozenge$	MATCH (coincidence)
11	0 $\blacklozenge$	1 $\blacklozenge$	MISMATCH
30	6 $\blacklozenge$	3 $\blacklozenge$	MISMATCH

Both individual values are themselves $\blacklozenge$ FORMAL (the builtins *run* and *report* exact integers); the negative is in their *relationship*. Since `gamma0_genus` is the trusted reference, Falsifier 1 is $\blacktriangledown$ FALSIFIED.

4.3 Finding-2 run: $\Gamma_1(N)/X(N)$ index smoothness

The 10 integer components are $\blacklozenge$: $\psi \in \{6, 12, 8, 12, 24\}$ and $\phi \in \{4, 2, 6, 10, 4\}$ for $N \in \{5, 6, 7, 11, 12\}$ (verbatim `gamma0_index / phi` verdicts). Assembling the cited closed forms gives the indices (citation tier — gated out of any headline):

N	$\Gamma_1(N)$ idx = $\psi\phi/2$	$X(N)$ idx = $N \cdot \Gamma_1$	note
5	$6 \cdot 4/2 = 12$	$5 \cdot 12 = 60$	
6	$12 \cdot 2/2 = 12$	$6 \cdot 12 = 72$	focal
7	$8 \cdot 6/2 = 24$	$7 \cdot 24 = 168$	
11	$12 \cdot 10/2 = 60$	$11 \cdot 60 = 660$	
12	$24 \cdot 4/2 = 48$	$12 \cdot 48 = 576$	

The $N = 6$ values (12, 72) are exactly the generic multiplicative-formula output. A cross-check on the equivalent closed form confirms $X(6) = N^3/2 \cdot \prod_{p|N} (1 - 1/p^2) = 108 \cdot (3/4)(8/9) = 72$, matching the relation $X = N \cdot \Gamma_1$. There is *no* peak, singularity, or arithmetic collapse at $N = 6$: the indices are smooth and multiplicative in N , so $N = 6$ is not distinguished from its neighbours the way $\{1, 6\}$ is distinguished for $\sigma\phi = n\tau$. Falsifier 2 is $\blacktriangledown$ FALSIFIED.

5 Finding

This paper reports two closed-negative findings, each resolving a pre-registered falsifier deterministically.

Finding 1 (MF4 — the `dim_cusp_forms` definition gap). The `hexa` builtin `dim_cusp_forms(N, 2)` does *not* realize the standard theorem $\dim S_2(\Gamma_0(N)) = \text{genus}(X_0(N))$: against the trusted `gamma0_genus`, it disagrees on 20 of the 30 levels $N \in [1, 30]$ (matching only by coincidence for small N). The ruled-out axis is *trust in the builtin's name*: `dim_cusp_forms` carries a definition gap and cannot be used as a weight-2 cusp-form-dimension oracle without the fix filed upstream as PR #1083. Falsifier 1: $\blacktriangledown$ FALSIFIED.

Finding 2 (F7 — the $n = 6$ non-lift). The $\{1, 6\}$ specialness $\sigma(n)\phi(n) = n\tau(n)$ is a Γ_0 -level / arithmetic-function-identity phenomenon that does *not* lift to the modular-curve tower: the $\Gamma_1(N)$ and $X(N) = \Gamma(N)$ indices are smooth multiplicative functions of N with no $n = 6$ peak, singularity, or collapse. The ruled-out axis is *the hypothesis that the $\{1, 6\}$ distinction propagates to higher modular levels*: it does not. Falsifier 2: ∇ FALSIFIED.




Figure 1: The two closed negatives. *Top:* the modular-curve tower $\Gamma_0(N) \rightarrow \Gamma_1(N) \rightarrow \Gamma(N) = X(N)$; the $n = 6$ specialness ($\sigma\phi = n\tau$ at the Γ_0 level) fades out moving up the tower because the Γ_1/X indices are smooth multiplicative functions of N with no $n = 6$ peak (Finding 2). *Bottom:* the **hexa** builtin `dim_cusp_forms(N,2)` agrees with the classical genus $\dim S_2(\Gamma_0(N)) = g(X_0(N))$ only for small N and diverges for $N \geq 11$, 20/30 mismatch (Finding 1, a builtin definition gap not a mathematical falsehood).

Why two negatives strengthen, not weaken, the MODFORM picture. The verified backdrop (§4.1) shows that the $\Gamma_0(N)$ invariants of $X_0(6)$ each reduce to a standard $n = 6$ arithmetic value, and the companion locus result [5] shows the $\sigma\phi = n\tau$ identity is genuinely $\{1, 6\}$ -confined. The two negatives now bound that picture from above and from the side. Finding 2 shows the specialness does not escape *upward* into the tower: it is a property of the smallest level and of the arithmetic functions, not of the integer 6 percolating through every finer congruence structure. Finding 1 shows that the one MODFORM quantity whose **hexa** builtin disagrees with theory does so for a *tooling* reason (a definition gap), leaving the genus, index, cusp, weight, and Atkin–Lehner results — all $\blacklozenge$ against classical values — untouched. The two negatives are therefore the precise boundary of the verified region: everything inside is $\blacklozenge$, the two things outside are a named tool bug and a refuted lift hypothesis.

6 Limitations and honest caveats

- **Finding 1 is a tool gap, not a mathematical falsehood.** The theorem $\dim S_2(\Gamma_0(N)) = \text{genus}(X_0(N))$ is true. What is falsified is the assumption that the builtin `dim_cusp_forms(N,2)` computes it. The small- N matches ($N \leq 7$) are coincidences of small dimension, not evidence of correctness; they are exactly the regime in which both quantities are 0. We report `dim_cusp_forms` values as $\blacklozenge$ *closed-form recomputations of whatever the builtin defines*, and the genus values as independently trusted; the negative lives only in their equality. The fix is filed upstream (PR #1083) and is not claimed as part of this paper.

- **Finding 2 does not deny the $n = 6$ specialness — it locates it.** $N = 6$ is special at the Γ_0 level (every $X_0(6)$ invariant reduces to an $n = 6$ arithmetic value) and $\{1, 6\}$ is the exact locus of $\sigma\phi = n\tau$ in the tested range [5]. The negative is narrow: the specialness does not *lift* to the Γ_1/X index magnitudes, which are smooth. A mild, *non-unique* feature does survive: $N = 6$ belongs to the 3-element set $\{3, 4, 6\}$ where $\Gamma_1(N) \text{ idx} = \Gamma_0(N) \text{ idx}$ (because $\phi(N) = 2$ makes the $\phi/2$ factor unity, and $\phi(N) = 2$ only for $N \in \{3, 4, 6\}$). $N = 6$ is the largest such level, with bridge $\Gamma_1(6) = \Gamma_0(6) = 12 = \sigma(6)$. This is a $\phi(N) = 2$ coincidence, citation-grade, *not* an $n = 6$ -unique identity, and it is gated out of the headline.
- **Assembled Γ_1/X indices are citation tier.** The integer components (ψ, ϕ) are $\blacklozenge$; the assembled indices combine them with a cited closed form and are therefore citation tier, used only to demonstrate *smoothness*, never as a standalone formal result.
- **Shimura-curve direction is not computed.** Compact Shimura curves $X^D(N)$ (indefinite quaternion algebra of discriminant $D > 1$, no cusps) would test the non-lift in a non-cuspidal setting via the Eichler mass formula. The `hexa` stack has no quaternion-discriminant / Eichler-mass builtin, so this is not integer-component verifiable here. It is flagged as a future *capability* gap (a missing function), distinct from Finding 1’s definition *bug*, and was deliberately not filed upstream.
- **Finite range.** Both findings are established on $N \in [1, 30]$ (Finding 1 sweep) and $N \in \{5, 6, 7, 11, 12\}$ (Finding 2 focal set). The `dim` $\neq$ genus divergence is monotone and structural beyond the small- N coincidence, and the Γ_1/X index smoothness is a closed-form (multiplicative) property that holds for all $N > 2$, but neither is here promoted to an unbounded $\forall N$ proof.

7 Related framings

The classical theory of modular forms and the genus of $X_0(N)$ is standard [2, 3]; the Atkin–Lehner involutions are from [1]. The $n = 6$ arithmetic-function identity and its $\{1, 6\}$ locus are the subject of the companion TECS-L paper [5], against which this paper is the modular-geometry continuation: where the companion paper asks “does the identity extend to the second perfect number?” (closed-negative at $n = 28$), this paper asks “does the identity lift to the modular-curve tower?” (closed-negative at the Γ_1/X level). The two are independent negatives about the same specialness from different directions. The discipline of treating a deterministically refuted falsifier as a publishable finding is the TECS-L *closed-negative* policy [6]; the claim-to-verdict audit chain is `CLAIMS.tape` [4]. We note explicitly that this work is *unrelated* to integrated-information-theory (Φ/IIT) measurements that share the TECS-L repository — the overlap is only the verification harness, not the subject matter.

Reproducibility

Every numeric quantity in this paper is a closed-form recomputation issued via `hexa verify --expr` and persisted verbatim under `.verdicts/tecs-l-modform-*/` (backdrop: `index`, `cusps`, `genus`, `symbols`, `weight-al`, `n6-bridge`; Finding 1: `dim-genus`; Finding 2: `other-curves`). Each is indexed in `CLAIMS.tape` (group TECS-L) with a one-to-one pointer to its verdict file. The full 30- N `dim` vs. genus sweep is Appendix A; the Γ_1/X index assembly is Appendix B; representative raw verdict transcripts (ASCII sanitized) are Appendix C.

References

- [1] A. O. L. Atkin and J. Lehner. Hecke operators on $\gamma_0(m)$. *Mathematische Annalen*, 185: 134–160, 1970. doi: 10.1007/BF01359701. Atkin–Lehner involutions W_Q ; group order $2^{\omega(N)}$.
- [2] Fred Diamond and Jerry Shurman. *A First Course in Modular Forms*, volume 228 of *Graduate Texts in Mathematics*. Springer, 2005. doi: 10.1007/978-0-387-27226-9. Genus of $X_0(N)$; $\dim S_2(\Gamma_0(N)) = \text{genus correspondence}$.
- [3] Goro Shimura. *Introduction to the Arithmetic Theory of Automorphic Functions*. Princeton University Press, 1971. Congruence subgroups, indices of $\Gamma_0(N)$, $\Gamma_1(N)$, $\Gamma(N)$.
- [4] TECS-L domain, hexa-lang. Claims.tape — single audit index of verifiable hexa-lang claims (group tecs-l). <https://github.com/dancinlab/hexa-lang> (CLAIMS.tape; verdicts under .verdicts/), 2026.
- [5] TECS-L domain, hexa-lang. The $\{1, 6\}$ identity locus: five arithmetic-function identities of the first perfect number, each closed-negative at the second. <https://github.com/dancinlab/hexa-lang> (PAPER/tecs-l-n6-identity-locus), 2026.
- [6] TECS-L domain, hexa-lang. Closed-negative findings are publishable (paper_negative_ok governance directive). <https://github.com/dancinlab/hexa-lang> (CLAUDE.md / project.tape), 2026.

A Appendix A: full dim vs. genus sweep, $N \in [1, 30]$

The complete sweep underlying Finding 1 (§4.2), recomputed by `hexa verify` and persisted verbatim at `.verdicts/tecs-l-modform-dim-genus/dim_vs_genus_sweep.txt`. Of the 30 levels, 10 match (small- N coincidence, in **bold**) and 20 mismatch. The divergence begins at $N = 8$ and persists.

N	<code>dim_cusp_forms(N,2)</code>	<code>gamma0_genus(N)</code>	verdict
1	0	0	MATCH
2	0	0	MATCH
3	0	0	MATCH
4	0	0	MATCH
5	0	0	MATCH
6	0	0	MATCH
7	0	0	MATCH
8	1	0	MISMATCH
9	0	0	MATCH
10	1	0	MISMATCH
11	0	1	MISMATCH
12	2	0	MISMATCH
13	0	0	MATCH
14	2	1	MISMATCH
15	0	1	MISMATCH
16	3	0	MISMATCH
17	0	1	MISMATCH
18	3	0	MISMATCH
19	0	1	MISMATCH

N	<code>dim_cusp_forms(N,2)</code>	<code>gamma0_genus(N)</code>	verdict
20	4	1	MISMATCH
21	0	1	MISMATCH
22	4	2	MISMATCH
23	0	2	MISMATCH
24	5	1	MISMATCH
25	0	0	MATCH
26	5	2	MISMATCH
27	0	1	MISMATCH
28	6	2	MISMATCH
29	0	2	MISMATCH
30	6	3	MISMATCH
<i>Totals: 10 MATCH, 20 MISMATCH ($N \in [1, 30]$).</i>			

B Appendix B: $\Gamma_1(N)/X(N)$ index table and $n = 6$ focal assembly

The $\Gamma_1(N)$ and $X(N) = \Gamma(N)$ indices underlying Finding 2 (§4.3), assembled from $\blacklozenge$ integer components via the cited closed forms (`.verdicts/tecs-1-modform-other-curves/`). The $\psi = \text{gamma0_index}$ and ϕ rows are $\blacklozenge$; the index columns are citation tier (cited closed form over verified components).

N	$\psi(N)$ ($\blacklozenge$)	$\phi(N)$ ($\blacklozenge$)	$\Gamma_1(N)$ idx = $\psi\phi/2$	$X(N)$ idx = $N \cdot \Gamma_1$
5	6	4	12	60
6	12	2	12	72 (focal)
7	8	6	24	168
11	12	10	60	660
12	24	4	48	576

Cross-check at $N = 6$: $X(6) = N^3/2 \cdot \prod_{p|N} (1 - 1/p^2) = 108 \cdot (3/4)(8/9) = 72 = 6 \cdot 12$.

Smoothness: Γ_1/X indices are multiplicative in N ; $N = 6$ sits at the generic value, no peak.

Mild non-unique feature: $\Gamma_1(6) = \Gamma_0(6) = 12$ since $\phi(6) = 2$ (also holds for $N \in \{3, 4\}$, citation tier).

Atkin–Lehner backdrop (MF7, citation). For completeness, the Atkin–Lehner involution count $|\text{AL}(\Gamma_0(N))| = 2^{\omega(N)}$ over a sample of levels, with $\omega(N)$ the number of distinct prime factors:

N	$\omega(N)$	$ \text{AL} = 2^{\omega}$
1	0	1
6	2	4
12	2	4
30	3	8
2310	5	32
30030	6	64

C Appendix C: representative raw verdict transcripts

Verbatim standard output (ASCII sanitized; tier glyphs rendered as their TIER names) of representative hexa verify runs underlying the two findings. The complete set is under `.verdicts/tecs-l-modform-*/`.

Finding 1 — the two boundary mismatches and the $n = 6$ match.

```
verify --expr dim_cusp_forms(11,2)=0
  calc   = 0 == expected 0
  tier    = [BLUE] SUPPORTED-FORMAL (hexa-native closed-form, TECS-L Tier1)
```

```
verify --expr gamma0_genus(11)=1
  calc   = 1 == expected 1
  tier    = [BLUE] SUPPORTED-FORMAL (hexa-native closed-form, TECS-L Tier1)
=> N=11: dim_cusp_forms=0 != gamma0_genus=1 MISMATCH
```

```
verify --expr dim_cusp_forms(30,2)=6
  calc   = 6 == expected 6
  tier    = [BLUE] SUPPORTED-FORMAL (hexa-native closed-form, TECS-L Tier1)
=> N=30: dim_cusp_forms=6 != gamma0_genus=3 MISMATCH
```

```
verify --expr dim_cusp_forms(6,2)=0
  calc   = 0 == expected 0
  tier    = [BLUE] SUPPORTED-FORMAL (hexa-native closed-form, TECS-L Tier1)
=> N=6: dim_cusp_forms=0 == gamma0_genus=0 MATCH (coincidence)
```

Finding 2 — the $n = 6$ focal components (smooth, no peak).

```
verify --expr gamma0_index(6)=12
  calc   = 12 == expected 12
  tier    = [BLUE] SUPPORTED-FORMAL (hexa-native closed-form, TECS-L Tier1)
```

```
verify --expr phi(6)=2
  calc   = 2 == expected 2
  tier    = [BLUE] SUPPORTED-FORMAL (hexa-native closed-form, TECS-L Tier1)
```

```
assembled (citation, exact integer arithmetic):
  Gamma_1(6) idx = psi(6)*phi(6)/2 = 12*2/2 = 12
  X(6) = Gamma(6) idx = 6 * 12 = 72 (relation X = N*Gamma_1, holds all N>2)
=> index is the generic multiplicative-formula output; no n=6 peak. NON-LIFT.
```

Backdrop — trusted gamma0_genus reference (Finding 1 leans on).

```
verify --expr gamma0_genus(6)=0      [BLUE] SUPPORTED-FORMAL (classical genus=0)
verify --expr gamma0_genus(11)=1     [BLUE] SUPPORTED-FORMAL (genus>=1 boundary)
verify --expr gamma0_genus(22)=2     [BLUE] SUPPORTED-FORMAL
verify --expr gamma0_index(6)=12     [BLUE] SUPPORTED-FORMAL (= sigma(6), n=6 bridge)
verify --expr gamma0_cusps(6)=4      [BLUE] SUPPORTED-FORMAL (= tau(6), n=6 bridge)
-> 22/22 classical genus values matched, 0 mismatch: trusted reference.
```

D Appendix D: the full verified $\Gamma_0(N)$ backdrop sweeps

The two closed negatives sit against a fully verified $\Gamma_0(N)$ backdrop. For completeness and reproducibility we reproduce the three integer sweeps on $N \in [1, 30]$, each 30/30 $\blacklozenge$ FORMAL, re-computed by `hexa verify --expr` and persisted verbatim under `.verdicts/tecs-l-modform-*/`. These are the trusted invariants whose internal consistency makes Falsifier 1 (which leans on `gamma0_genus`) and the smoothness assessment of Falsifier 2 (which leans on `gamma0_index`) credible.

D.1 — Index $\psi(N) = \text{gamma0_index}$, **cusps** $= \text{gamma0_cusps}$, **first weight** $= \text{first_cusp_form_weight}$. The $n = 6$ row is highlighted: $\psi(6) = 12 = \sigma(6)$ and four cusps $= \tau(6)$ are the two arithmetic bridges that Finding 2 then shows do *not* continue up the tower.

N	$\psi(N)$ ($\blacklozenge$)	cusps ($\blacklozenge$)	first weight ($\blacklozenge$)
1	1	1	—
2	3	2	—
3	4	2	—
4	6	3	—
5	6	2	—
6	12 $= \sigma(6)$	4 $= \tau(6)$	4 $= \tau(6)$
7	8	2	—
8	12	4	—
9	12	4	—
10	18	4	—
11	12	2	—
12	24	6	—
13	14	2	—
14	24	4	—
15	24	4	—
16	24	6	—
17	18	2	—
18	36	8	—
19	20	2	—
20	36	6	—
21	32	4	—
22	36	4	—
23	24	2	—
24	48	8	—
25	30	6	—
26	42	4	—
27	36	6	—
28	48	6	—
29	30	2	—
30	72	8	—

All 30/30 $\blacklozenge$ for ψ and cusps; first weight 30/30 $\blacklozenge$ ($1 \rightarrow 12$, $6 \rightarrow 4$, $30 \rightarrow 2$, monotone trend).

D.2 — The genus sweep (22 classical/known values, 0 mismatch). This is the trusted reference for Finding 1. The 15 classical genus-0 levels are $N \in \{1-10, 12, 13, 16, 18, 25\}$; the

genus- ≥ 1 boundary is $g(X_0(N)) = 1$ for $N \in \{11, 14, 15, 17, 19\}$ and $g = 2$ for $N \in \{22, 23\}$. Every one of the 22 values matched its classical value with verdict $\blacklozenge$ FORMAL and *zero* mismatches — which is precisely why a disagreement between `dim_cusp_forms` and `gamma0_genus` (Appendix A) localizes the fault to `dim_cusp_forms`.

D.3 — The $n = 6$ bridge synthesis (MF6). Aggregating four $\blacklozenge$ atoms (plus one citation-tier Atkin–Lehner row), every core invariant of the modular curve $X_0(6)$ reduces to a standard $n = 6$ arithmetic-function value:

invariant of $X_0(6)$	value	$n = 6$ reduction
index $\psi(6)$	12	$= \sigma(6)$
cusps $c(6)$	4	$= \tau(6)$
genus $g(X_0(6))$	0	(classical genus-0)
first cusp-form weight	4	$= \tau(6)$
$ \text{AL}(\Gamma_0(6)) $	4	$= 2^{\omega(6)}$ (citation)

This is the synthesis-reasoned bridge ($\blacksquare$ -class, composed entirely from terminal $\blacklozenge$ atoms) that the two closed negatives bound. Finding 2 shows the covering ($\psi \leftrightarrow \sigma$) and cusp ($c \leftrightarrow \tau$) bridges are Γ_0 -level only — the Γ_1/X index does not inherit them — while Finding 1 shows the one quantity ($\dim S_2$) whose builtin name promises a genus link instead carries a definition gap. The bridge is real at $N = 6$; its closure boundary is exactly the two negatives reported in this paper.