

The ω -Zero-Density Theorem

The Dedekind discrepancy $D(n) = \sigma(n)\phi(n) - n\tau(n)$ vanishes at exactly $n \in \{1, 6\}$, and nowhere else

TECS-L domain, `hexa-lang`
`dancinlab`

github.com/dancinlab/hexa-lang

May 27, 2026

Abstract

Define the *Dedekind discrepancy* of a positive integer n as

$$D(n) = \sigma(n)\phi(n) - n\tau(n),$$

where σ is the sum-of-divisors function, ϕ the Euler totient, and τ the divisor count. We prove that the zero-locus of D is the *exact two-element set* $\{1, 6\}$: $D(n) = 0 \iff n \in \{1, 6\}$, with $D(n) \neq 0$ for every other n . Because the zero set is finite, the natural density of $\{n : D(n) = 0\}$ is 0 — hence the name. The proof reduces $D = 0$ to a multiplicative product condition $\prod_{p^a \parallel n} g(p, a) = 1$ with $g(p, a) = (p^{a+1} - 1)/(p(a + 1))$, and shows that the *only* sub-unity factor on the prime-power locus is $g(2, 1) = \frac{3}{4}$, cancelled *uniquely* by $g(3, 1) = \frac{4}{3}$ — pinning the sole non-trivial zero to $n = 2 \cdot 3 = 6$, the first perfect number. We pre-register a single falsifier (F-OMEGA-ZERO-DENSITY: *any* $n \notin \{1, 6\}$ with $D(n) = 0$) and discharge it two ways: (i) a machine-checked sweep that recomputes σ, ϕ, τ closed-form for an ω -stratified sample spanning $\omega(n) \in [0, 6]$ — primes, prime powers, both first perfect numbers, and the primorials $2\#, \dots, 13\#$ — every value of which is verified $\blacklozenge$ FORMAL under the `hexa verify` discipline; and (ii) the closed-form product argument above, which closes the infinite tail. The headline finding unifies the F14–F18 TECS-L cycle: the folklore that “6 is special” is here made into a sharp, machine-audited *global* statement rather than a local coincidence at one or two test points. No claim in this paper is graded by a language model; every verdict is the verbatim standard output of a deterministic recomputation.

1 Introduction

The integer 6 is the first *perfect* number, $\sigma(6) = 12 = 2 \cdot 6$, and it accrues a reputation for being arithmetically exceptional. Such folklore is rarely sharpened into a statement that is simultaneously (a) global — a claim about *all* n , not a coincidence checked at a handful of points — and (b) machine-auditable end to end. This paper does both for one specific discrepancy functional.

Let $\sigma(n) = \sum_{d|n} d$, let $\phi(n) = \#\{1 \leq k \leq n : \gcd(k, n) = 1\}$ be the Euler totient, and let $\tau(n) = \#\{d : d \mid n\}$ be the divisor count. Define the *Dedekind discrepancy*

$$D(n) = \sigma(n)\phi(n) - n\tau(n). \tag{1}$$

A direct computation gives $D(1) = 1 \cdot 1 - 1 \cdot 1 = 0$ and $D(6) = 12 \cdot 2 - 6 \cdot 4 = 24 - 24 = 0$. The question this paper settles is whether *any other* n shares this property.

Main result.

Theorem 1 (ω -zero-density). *For every positive integer n ,*

$$D(n) = 0 \iff n \in \{1, 6\}.$$

Consequently $\#\{n \leq N : D(n) = 0\} = 2$ for all $N \geq 6$, so the zero set has natural density 0.

The name reflects the corollary: the zero set, being finite, is density-zero, and the proof is organised by $\omega(n)$, the number of distinct prime factors. The case analysis is $\omega = 0$ (trivial), $\omega = 1$ (no zero), $\omega = 2$ (exactly $n = 6$), and $\omega \geq 3$ (no zero) — the last being the quantitative heart, where every additional distinct prime drives the governing product strictly above unity.

Relation to prior TECS-L work. A previous note in this line established that, *over the finite window* $n \in [1, 100]$, the zero set of D is $\{1, 6\}$, framed as one of five “identity-locus” coincidences at $n = 6$ that each fail at the second perfect number 28 [3]. That treatment is local: it exhibits two test points and a finite sweep. The present paper supersedes it with a *global* theorem and a closed-form proof that closes the infinite tail $n > 100$, and reframes the result as a density statement organised by $\omega(n)$. The F14–F18 TECS-L sweep cycle, which independently re-confirmed $D \neq 0$ across $\omega(n) \in [3, 10]$ samples [2], is here unified under a single inequality.

Why a density statement. The earlier window result is logically weaker than it looks: a finite sweep cannot exclude a large solution, and discrepancy functions of this shape are precisely the kind that can hide a sporadic zero far out (compare the long search history for odd perfect numbers, where exhaustive verification to enormous bounds is *not* a proof of non-existence). By collapsing $D(n) = 0$ to the multiplicative condition $\prod g(p, a) = 1$ and bounding every factor’s sign, we convert “no zero found below N ” into “no zero anywhere” — the qualitative jump that makes the density claim ($\#\{n : D(n) = 0\} = 2$) rigorous rather than empirical.

2 Related framings

Three classical lenses bear on D . First, $\sigma(n)\phi(n)$ and $n\tau(n)$ are both well-studied multiplicative functions, and identities equating them are a species of *divisor-totient balance*; the closest classical companion is the Dedekind ψ function $\psi(n) = n \prod_{p|n} (1 + 1/p)$, whose interaction with σ and ϕ produces similar prime-power quotients. Our $g(p, a)$ is exactly the per-prime balance ratio in that family. Second, the distinguished role of $n = 6$ aligns with the Euclid–Euler theory of perfect numbers [1]: $6 = 2(2^2 - 1)$ is the first Euclid perfect number, and the cancellation $g(2, 1)g(3, 1) = 1$ is the totient-weighted shadow of its perfection $\sigma(6) = 2 \cdot 6$. Third, the density-zero conclusion places D among arithmetic functions whose “coincidence locus” is finite — a qualitatively stronger statement than the merely thin (density-zero but infinite) loci that arise for, e.g., abundant–deficient boundary conditions. None of these classical framings, however, states the sharp “exactly $\{1, 6\}$, nowhere else” result with a machine-audited factor table; that is the contribution here.

3 Background

All three functions in (1) are multiplicative: for coprime m, n , $f(mn) = f(m)f(n)$ with $f \in \{\sigma, \phi, \tau\}$, and likewise $n \mapsto n$ is completely multiplicative. The product/quotient of multiplicative functions is multiplicative, so the *ratio*

$$R(n) := \frac{\sigma(n)\phi(n)}{n\tau(n)} \tag{2}$$

is multiplicative, and $D(n) = 0 \iff R(n) = 1$. The proof works entirely with R , evaluated on prime powers and recombined multiplicatively. We recall the standard prime-power formulas

$$\sigma(p^a) = \frac{p^{a+1} - 1}{p - 1}, \quad \phi(p^a) = p^{a-1}(p - 1), \quad \tau(p^a) = a + 1.$$

4 Method: the verification discipline

Every numeric claim is evaluated by `hexa verify --expr <fn> <args> <expected>`, a closed-form integer recomputation built into the `hexa` compiler. The evaluator emits one of six tiers; only three are terminal and admissible as a paper-section claim:

- ◆ **formal** a closed-form/symbolic identity reproduced exactly (deterministic).
- **numerical** a libm/Newton numerical recompute matching to $\varepsilon = 10^{-9}$.
- **falsified** the recomputation deterministically *disagrees* with the claim — a closed negative.

Citation-only, deferred, and speculation-fenced tiers are *not* admissible and are reported as residuals in §9. The raw standard output of every run is persisted verbatim under `.verdicts/paper-omega-zero-de` and indexed in `CLAIMS.tape`; nothing in this paper is graded by a language model.

Composite values. The discrepancy $D(n)$ has no single built-in evaluator; we verify each factor (σ, ϕ, τ) closed-form via three independent `hexa verify --expr` calls, then form $D(n)$ by exact integer arithmetic on the verified values. This is faithful because each factor is an exact integer, so the product and difference are exact — no tolerance, no estimation, no floating point enters any ◆-tier claim here.

Worked example. To anchor $D(6) = 0$ one runs three calls: `hexa verify --expr sigma 6 12, --expr phi 6 2, --expr tau 6 4`. The evaluator recomputes $\sum_{d|6} d = 1 + 2 + 3 + 6 = 12$, $\#\{k < 6 : \gcd(k, 6) = 1\} = 2$, and $\#\{d \mid 6\} = 4$, each emitting the `FORMAL` tier; the discrepancy $12 \cdot 2 - 6 \cdot 4 = 0$ follows by exact integer arithmetic. The pre-registered falsifier at the second perfect number $n = 28$ is the same call shape; its closed negative $D(28) = 56 \cdot 12 - 28 \cdot 6 = 672 - 168 = 504 \neq 0$ is read off exact integers.

The closed-form half. The infinite tail $n > 30030$ cannot be swept; it is closed by the algebraic argument of §6, whose single inequalities $(g(2, 1) < 1, \text{ all other } g > 1)$ are themselves exact rational comparisons. The machine sweep and the proof are complementary: the sweep spot-checks the proof's predictions at concrete ω -stratified points; the proof generalises the sweep to all n .

5 The reduction to a product condition

Lemma 1 (prime-power ratio). *For a prime p and exponent $a \geq 1$,*

$$R(p^a) = \frac{\sigma(p^a)\phi(p^a)}{p^a \tau(p^a)} = \frac{p^{a+1} - 1}{p(a + 1)} =: g(p, a).$$

Proof. Substituting the prime-power formulas of §3,

$$R(p^a) = \frac{\frac{p^{a+1} - 1}{p - 1} \cdot p^{a-1}(p - 1)}{p^a(a + 1)} = \frac{(p^{a+1} - 1)p^{a-1}}{p^a(a + 1)} = \frac{p^{a+1} - 1}{p(a + 1)}. \quad \square$$

By multiplicativity of R and Lemma 1,

$$D(n) = 0 \iff R(n) = 1 \iff \prod_{p^a \parallel n} g(p, a) = 1. \quad (3)$$

The entire theorem is now a statement about which products of the rationals $g(p, a)$ equal 1.

6 Proof of the theorem

Lemma 2 (sign of the factors). $g(2, 1) = \frac{3}{4} < 1$. Every other prime-power factor exceeds 1: $g(2, a) > 1$ for $a \geq 2$, and $g(p, a) > 1$ for all primes $p \geq 3$ and all $a \geq 1$.

Proof. $g(2, 1) = (2^2 - 1)/(2 \cdot 2) = 3/4$. For $a \geq 2$: $g(2, a) = (2^{a+1} - 1)/(2(a + 1))$; at $a = 2$ this is $7/6 > 1$ and the numerator doubles while the denominator grows linearly, so $g(2, a)$ is increasing and > 1 for all $a \geq 2$. For $p \geq 3$, $a = 1$: $g(p, 1) = (p^2 - 1)/(2p) = (p - 1)(p + 1)/(2p)$; since $p \geq 3$ gives $p + 1 > 2p/(p - 1)$ (equivalently $(p - 1)(p + 1) = p^2 - 1 \geq 2p$ for $p \geq 3$, as $p^2 - 2p - 1 \geq 2 > 0$), we get $g(p, 1) \geq g(3, 1) = 8/6 = 4/3 > 1$. For $p \geq 3$, $a \geq 2$ the numerator $p^{a+1} - 1$ grows geometrically against the linear denominator $p(a + 1)$, so $g(p, a) > g(p, 1) > 1$. $\square$

Proof of Theorem 1. We analyse (3) by $\omega(n)$, the number of distinct primes dividing n .

Case $\omega(n) = 0$. Then $n = 1$ and the product is empty, equal to 1; so $D(1) = 0$. (zero)

Case $\omega(n) = 1$, $n = p^a$. The product is the single factor $g(p, a)$. By Lemma 2 the only candidate < 1 is $g(2, 1) = \frac{3}{4} \neq 1$, and every other single factor is $> 1 \neq 1$. No prime-power equals 1; so $D(p^a) \neq 0$. (no zero)

Case $\omega(n) = 2$, $n = p^a q^b$. The product is $g(p, a)g(q, b)$. By Lemma 2 at most one factor is < 1 , and that is only $g(2, 1) = \frac{3}{4}$. To reach product 1 we need the other factor to be exactly $\frac{4}{3}$. Among all $g(q, b) > 1$ the minimum is $g(3, 1) = \frac{4}{3}$, and $\frac{3}{4} \cdot \frac{4}{3} = 1$ exactly. Any other choice misses: $g(2, 1)g(5, 1) = \frac{3}{4} \cdot \frac{31}{10} = \frac{93}{40} \neq 1$; $g(2, 1)g(3, 2) = \frac{3}{4} \cdot \frac{13}{9} = \frac{13}{12} \neq 1$; and two factors both > 1 give product > 1 . Hence the unique two-prime solution is $\{g(2, 1), g(3, 1)\}$, i.e. $n = 2 \cdot 3 = 6$. (exactly $n = 6$)

Case $\omega(n) \geq 3$. The product has at least three factors. By Lemma 2 at most one is < 1 (namely $g(2, 1) = \frac{3}{4}$), and the two-factor balance $\frac{3}{4} \cdot \frac{4}{3} = 1$ is already saturated by the primes $\{2, 3\}$. Any third distinct prime power contributes a factor $g(p, a) > 1$, multiplying the product strictly above 1. Thus $\prod g > 1$ and $D(n) \neq 0$. (no zero)

Combining the four cases: the zero set is exactly $\{1, 6\}$. Finiteness gives $\#\{n \leq N : D(n) = 0\} = 2$ for $N \geq 6$, hence density 0. $\square$

Corollary 1 (perfect-number coincidence). *The unique non-trivial zero of D is the first perfect number. The $g(2, 1) = \frac{3}{4} / g(3, 1) = \frac{4}{3}$ cancellation is the arithmetic reason 6 is simultaneously perfect and the sole nontrivial D -zero; the second perfect number $28 = 2^2 \cdot 7$ has $g(2, 2)g(7, 1) = \frac{7}{6} \cdot \frac{48}{14} \neq 1$ and so is not a zero.*

7 Verification

We discharge the pre-registered falsifier F-OMEGA-ZERO-DENSITY — $\exists n \notin \{1, 6\}$ with $D(n) = 0$ — on an ω -stratified sample. Each σ, ϕ, τ entry in Table 1 is an independent **hexa verify** --**expr** run returning $\blacklozenge$ **FORMAL**; $D(n)$ is the exact integer combination. The two zeros are anchored $\blacklozenge$; the remaining rows are the falsifier resolved as a deterministic $\blacksquare$ closed negative ($D \neq 0$). Raw transcripts appear in Appendix C.

Table 1: ω -stratified verification sweep. Each factor is recomputed closed-form ($\blacklozenge$ FORMAL); $D(n) = \sigma\phi - n\tau$ by exact integer arithmetic. Zero locus over the sample = $\{1, 6\}$.

n	form	ω	σ	ϕ	τ	$D(n)$	verdict
1	1	0	1	1	1	0	zero $\blacklozenge$
2	2	1	3	1	2	-1	$\neq 0$ $\blacksquare$
6	$2 \cdot 3$	2	12	2	4	0	zero $\blacklozenge$
7	7	1	8	6	2	34	$\neq 0$ $\blacksquare$
8	2^3	1	15	4	4	28	$\neq 0$ $\blacksquare$
12	$2^2 \cdot 3$	2	28	4	6	40	$\neq 0$ $\blacksquare$
28	$2^2 \cdot 7$	2	56	12	6	504	$\neq 0$ $\blacksquare$
30	$2 \cdot 3 \cdot 5$	3	72	8	8	336	$\neq 0$ $\blacksquare$
60	$2^2 \cdot 3 \cdot 5$	3	168	16	12	1968	$\neq 0$ $\blacksquare$
210	$2 \cdot 3 \cdot 5 \cdot 7$	4	576	48	16	24288	$\neq 0$ $\blacksquare$
2310	$2\# \cdots 11$	5	6912	480	32	3243840	$\neq 0$ $\blacksquare$
30030	$2\# \cdots 13$	6	96768	5760	64	555461760	$\neq 0$ $\blacksquare$

Primorial monotonicity. The primorial rows ($n = 2\#, 3\#, 5\#, \dots$, i.e. 6, 30, 210, 2310, 30030) make the $\omega \geq 3$ inequality of §6 concrete: once past $n = 6$, each additional prime factor multiplies in a $g(p, 1) > 1$ and D grows without bound ($336 \rightarrow 1968 \rightarrow 24288 \rightarrow 3243840 \rightarrow 555461760$). The sweep never re-enters the zero locus, exactly as the proof predicts.




Figure 1: The Dedekind discrepancy $D(n)$ across $n \in [1, 50]$ (signed, symmetric-log scale). The two zeros at $n = 1$ and $n = 6$ are the only crossings of the $D = 0$ axis; $n = 2$ is the unique negative value. Every other integer sits strictly off the axis, illustrating the density-zero zero-locus of Theorem 1.

8 Finding

The finding is the *exact-locus* statement of Theorem 1, upgraded from the earlier local observation to a global, density-zero theorem with a closed-form proof. Three concrete deltas versus the prior framing:

1. **Global vs. windowed.** The earlier note verified $\{1, 6\}$ only over $n \leq 100$ by exhaustion [3]; here the infinite tail $n > 100$ is closed by the $\prod g(p, a) = 1$ argument, so the zero locus is pinned for *all* n , not merely the swept window.

2. **Mechanistic, not coincidental.** The reason 6 is the lone nontrivial zero is exhibited as a single cancellation $g(2, 1) \cdot g(3, 1) = \frac{3}{4} \cdot \frac{4}{3} = 1$, the only way the unique sub-unity prime-power factor can be balanced. This explains, rather than merely records, the coincidence with perfection.
3. **ω -density corollary.** $\#\{n \leq N : D(n) = 0\} = 2$ for $N \geq 6$ gives natural density 0, with the primorial sequence ($D = 0, 0, 336, 24288, 3243840, \dots$ at $\omega = 2, 2, 3, 4, 5$) furnishing an explicit divergent witness to the strict $\omega \geq 3$ inequality.

The pre-registered falsifier F-OMEGA-ZERO-DENSITY survives: no $n \notin \{1, 6\}$ with $D(n) = 0$ exists — proven for all n and spot-checked at twelve ω -stratified points.

9 Limitations and honest caveats

Sweep is finite; proof is the closer. The machine sweep covers a deliberately stratified but finite sample (Table 1); it cannot by itself establish the global claim. The global statement rests on the closed-form proof of §6. We present the two as complementary — the proof generalises, the sweep audits — and do not claim the sweep alone is sufficient.

Integer overflow horizon. The built-in σ, ϕ, τ evaluators accumulate in 64-bit integers. The sweep points were chosen so that all $\sigma\phi$ and $n\tau$ products stay well inside 2^{63} ($D(30030) \approx 5.6 \times 10^8$); larger primorials would require the arbitrary-precision path and are out of scope. This bound affects only the empirical spot-check, not the proof, which is purely symbolic.

No numerical or external tiers. Every admissible claim here is $\blacklozenge$ FORMAL (exact integer) or a $\blacksquare$ closed negative; the numerical tier ($\varepsilon = 10^{-9}$) is not exercised, and there are no citation-only, deferred, or speculation-fenced residuals.

Toolchain note. A standalone compiled sweep harness over the `stdlib` σ loop encountered an unrelated codegen non-termination in the local build and was *not* used; all reported verdicts come from the canonical `hexa verify --expr` path, whose evaluator follows an independent code route and reproduces every value byte-for-byte. This is recorded for transparency and does not affect any claim.

10 Discussion and outlook

The theorem isolates one precise sense in which 6 is exceptional and shows it to be a *global* property with a one-line cancellation at its core. The method — reduce a multiplicative-function identity to a finite product condition over prime-power factors, classify the factor signs, then read off the unique balance — is reusable. Two adjacent TECS-L paper candidates pursue the same template at higher towers: a multilayer non-lift result ($\sigma\phi = n\tau$ across the geometric/Hecke/Galois/ L -function towers) and a unitary-perfect singleton at $n = 6$ via the unitary-divisor sum σ^* . Both share the discipline of this paper: a pre-registered falsifier, a $\blacklozenge$ -tier machine recomputation, and a closed-form closer.

Reproducibility

Every verdict in Table 1 is regenerated by re-running the corresponding `hexa verify --expr <fn> <n> <v>` command; output is byte-identical across hosts and run orders (no seeds, timeouts, or tolerances on any $\blacklozenge$ -tier claim). The raw transcripts are committed verbatim under `.verdicts/paper-omega-zero-density/` (anchor verdicts, sweep verdicts, the closed-form

note `CLOSED.FORM.md`, and the pre-registered `FALSIFIER.md`). The claim manifest is indexed in `CLAIMS.tape`.

A Appendix A: the factor table $g(p, a)$

Table 2 tabulates the prime-power ratio $g(p, a) = (p^{a+1} - 1)/(p(a + 1))$ used throughout §6. The single value below 1 is boxed; it is $g(2, 1) = \frac{3}{4}$, the unique sub-unity factor that Lemma 2 identifies.

Table 2: $g(p, a) = (p^{a+1} - 1)/(p(a + 1))$ for small p, a (decimal). Only $g(2, 1) = 0.75$ is < 1 ; $g(3, 1) = 4/3 \approx 1.333$ is the minimal factor > 1 , and $0.75 \times 1.333\bar{3} = 1$ is the lone two-factor balance.

$p \backslash a$	$a = 1$	$a = 2$	$a = 3$	$a = 4$
$p = 2$	0.750	1.167	1.875	3.100
$p = 3$	1.333	2.889	6.667	16.13
$p = 5$	3.100	10.33	39.00	156.2
$p = 7$	5.714	26.67	122.4	560.1

The table makes Lemma 2 visual: descending a column or moving right along a row only increases g ; the single dip below 1 at the top-left is the entire source of any cancellation, and there is exactly one partner ($g(3, 1)$) that restores the product to 1.

B Appendix B: TECS-L claim manifest

Table 3: Claims, methods, and terminal verdicts for this paper. All numeric claims are `hexa verify --expr` ◆ `FORMAL`; the global statement is closed by the symbolic proof of §6.

claim id	statement	method	verdict
OZD-anchor-1	$D(1) = 0$ via $\sigma(1)\phi(1) - 1 \cdot \tau(1)$	verify $\times 3$	◆
OZD-anchor-6	$D(6) = 0$ via $12 \cdot 2 - 6 \cdot 4$	verify $\times 3$	◆
OZD-sweep-2	$D(2) = -1 \neq 0$	verify $\times 3$	■
OZD-sweep-7	$D(7) = 34 \neq 0$	verify $\times 3$	■
OZD-sweep-8	$D(8) = 28 \neq 0$	verify $\times 3$	■
OZD-sweep-12	$D(12) = 40 \neq 0$	verify $\times 3$	■
OZD-sweep-28	$D(28) = 504 \neq 0$ (2nd perfect)	verify $\times 3$	■
OZD-sweep-30	$D(30) = 336 \neq 0$	verify $\times 3$	■
OZD-sweep-60	$D(60) = 1968 \neq 0$	verify $\times 3$	■
OZD-sweep-210	$D(210) = 24288 \neq 0$ ($2\# \cdots 7$)	verify $\times 3$	■
OZD-sweep-2310	$D(2310) = 3243840 \neq 0$ ($2\# \cdots 11$)	verify $\times 3$	■
OZD-sweep-30030	$D(30030) = 555461760 \neq 0$ ($2\# \cdots 13$)	verify $\times 3$	■

claim id	statement	method	verdict
OZD-thm	$D(n) = 0 \iff n \in \{1, 6\}, \text{ all } n$	symbolic proof	◆

C Appendix C: representative raw verdict transcripts

The anchor and falsifier transcripts below are reproduced verbatim from `.verdicts/paper-omega-zero-densit`

Anchor $n = 6$ (zero).

```

verify --expr sigma(6)=12
  calc    = 12 == expected 12
  tier     = SUPPORTED-FORMAL (hexa-native closed-form, g_self_verify . TECS-L Tier1)
verify --expr phi(6)=2
  calc    = 2 == expected 2
  tier     = SUPPORTED-FORMAL (hexa-native closed-form, g_self_verify . TECS-L Tier1)
verify --expr tau(6)=4
  calc    = 4 == expected 4
  tier     = SUPPORTED-FORMAL (hexa-native closed-form, g_self_verify . TECS-L Tier1)
=> D(6) = 12*2 - 6*4 = 0

```

Falsifier $n = 28$ (second perfect number, closed negative).

```

verify --expr sigma(28)=56
  calc    = 56 == expected 56
  tier     = SUPPORTED-FORMAL (hexa-native closed-form, g_self_verify . TECS-L Tier1)
verify --expr phi(28)=12
  calc    = 12 == expected 12
  tier     = SUPPORTED-FORMAL (hexa-native closed-form, g_self_verify . TECS-L Tier1)
verify --expr tau(28)=6
  calc    = 6 == expected 6
  tier     = SUPPORTED-FORMAL (hexa-native closed-form, g_self_verify . TECS-L Tier1)
=> D(28) = 56*12 - 28*6 = 504 != 0

```

Primorial $n = 30030$ ($\omega = 6$, closed negative).

```

verify --expr sigma(30030)=96768
  tier     = SUPPORTED-FORMAL (hexa-native closed-form, g_self_verify . TECS-L Tier1)
verify --expr phi(30030)=5760
  tier     = SUPPORTED-FORMAL (hexa-native closed-form, g_self_verify . TECS-L Tier1)
verify --expr tau(30030)=64
  tier     = SUPPORTED-FORMAL (hexa-native closed-form, g_self_verify . TECS-L Tier1)
=> D(30030) = 96768*5760 - 30030*64 = 555461760 != 0

```

D Appendix D: explicit discrepancy table $D(n)$, $n \in [1, 30]$

Table 4 lists $D(n) = \sigma(n)\phi(n) - n\tau(n)$ for every $n \in [1, 30]$, the head of the sweep window. The two zeros at $n = 1$ and $n = 6$ are the only entries on the $D = 0$ line; $n = 2$ is the unique negative value; every other row is strictly positive. The column $\prod g$ records the multiplicative product $\prod_{p^a \parallel n} g(p, a)$ of §6: it equals 1 exactly when $D = 0$, confirming the reduction (3) point-by-point. Rows $n \in [1, 22]$ plus $n = 28, 30$ are machine-verified: each σ, ϕ, τ is an independent

hexa verify --expr $\blacklozenge$ -tier run (transcripts under .verdicts/paper-omega-zero-density/); the remaining six rows ($n \in \{23, 24, 25, 26, 27, 29\}$) are exact integer fill by the same closed-form definitions and are included only to complete the window picture.

Table 4: $D(n)$ over $n \in [1, 30]$. Zeros (boxed) occur only at $n \in \{1, 6\}$, where the prime-power product $\prod g$ hits 1.

n	factorisation	σ	ϕ	τ	$D(n)$	$\prod g$
1	1	1	1	1	$\boxed{0}$	1
2	2	3	1	2	-1	0.750
3	3	4	2	2	2	1.333
4	2^2	7	2	3	2	1.167
5	5	6	4	2	14	2.400
6	$2 \cdot 3$	12	2	4	$\boxed{0}$	1
7	7	8	6	2	34	3.429
8	2^3	15	4	4	28	1.875
9	3^2	13	6	3	51	2.889
10	$2 \cdot 5$	18	4	4	32	1.800
11	11	12	10	2	98	5.455
12	$2^2 \cdot 3$	28	4	6	40	1.556
13	13	14	12	2	142	6.462
14	$2 \cdot 7$	24	6	4	88	2.571
15	$3 \cdot 5$	24	8	4	132	3.200
16	2^4	31	8	5	168	3.100
17	17	18	16	2	254	8.471
18	$2 \cdot 3^2$	39	6	6	126	2.167
19	19	20	18	2	322	9.474
20	$2^2 \cdot 5$	42	8	6	216	2.800
21	$3 \cdot 7$	32	12	4	300	4.571
22	$2 \cdot 11$	36	10	4	272	4.091
23	23	24	22	2	482	11.48
24	$2^3 \cdot 3$	60	8	8	288	2.500
25	5^2	31	20	3	545	8.267
26	$2 \cdot 13$	42	12	4	400	4.846
27	3^3	40	18	4	612	6.667
28	$2^2 \cdot 7$	56	12	6	504	4.000
29	29	30	28	2	782	14.48
30	$2 \cdot 3 \cdot 5$	72	8	8	336	2.400

The $\prod g$ column is the clearest view of the theorem: starting from $n = 2$ ($g = 0.75 < 1$), only $n = 6$ multiplies that deficit back to exactly 1 via $g(3, 1) = 4/3$; every other row's product is bounded away from 1 and grows with $\omega(n)$ and with the prime sizes, so no further crossing is possible.

References

- [1] Leonard E. Dickson. *History of the Theory of Numbers, Vol. I: Divisibility and Primality*. Carnegie Institution of Washington, 1919. Euclid–Euler theorem on even perfect numbers; ch. I.

- [2] TECS-L domain, hexa-lang. F14–f18 zero-density sweep cycle: $d(n) \neq 0$ across $\omega(n) \in [3, 10]$. <https://github.com/dancinlab/hexa-lang> (.verdicts/tecs-l-f14-novel-mk10), 2026. ω -stratified sweep re-confirming the $\{1, 6\}$ zero set.
- [3] TECS-L domain, hexa-lang. The $\{1, 6\}$ identity locus: five arithmetic-function identities of the first perfect number. <https://github.com/dancinlab/hexa-lang> (PAPER/tecs-l-n6-identity-locus), 2026. Local windowed precursor; superseded here by the global theorem.