

Reconfiguration of Colorings and List Colorings: Proofs and Conjectures

Daniel W. Cranston
dcransto@gmail.com

CanaDAM
21 May 2025

What is Coloring Reconfiguration?

What is Coloring Reconfiguration?


The purpose of life is to prove and conjecture...

–Paul Erdős

What is Coloring Reconfiguration?

The purpose of life is to prove and conjecture...


–Paul Erdős



What is Coloring Reconfiguration?


The purpose of life is to prove and conjecture...

–Paul Erdős




What is Coloring Reconfiguration?


The purpose of life is to prove and conjecture...
–Paul Erdős



What is Coloring Reconfiguration?




What is Coloring Reconfiguration?




And another isomorphic component.


What is Coloring Reconfiguration?




And another isomorphic component.



What is Coloring Reconfiguration?





And another isomorphic component.



- “Reconfiguration graphs” of 3-colorings of 5-cycle and 4-cycle.



What is Coloring Reconfiguration?



And another isomorphic component.

- ▶ “Reconfiguration graphs” of 3-colorings of 5-cycle and 4-cycle.
- ▶ Is the reconfiguration graph connected?

What is Coloring Reconfiguration?



And another isomorphic component.

- ▶ “Reconfiguration graphs” of 3-colorings of 5-cycle and 4-cycle.
- ▶ Is the reconfiguration graph connected? What is its diameter?

What is List Coloring Reconfiguration?



- ▶ **list-assignment** L : each vertex v gets allowable colors $L(v)$

What is List Coloring Reconfiguration?

1 2 ○ — ○ 1 3

1 2 ○ — ○ 1 3

1 2 ○ — ○ 1 3

- ▶ **list-assignment** L : each vertex v gets allowable colors $L(v)$
- ▶ **L -coloring** φ : φ is proper and $\varphi(v) \in L(v)$ for all v

What is List Coloring Reconfiguration?

1 2 ○ — ○ 1 3

1 2 ○ — ○ 1 3

1 2 ○ — ○ 1 3

- ▶ **list-assignment** L : each vertex v gets allowable colors $L(v)$
- ▶ **L -coloring** φ : φ is proper and $\varphi(v) \in L(v)$ for all v

Main Questions:

- ▶ Given L -colorings α and β , can we change α to β by recoloring single vertices, keeping L -coloring at each step?

What is List Coloring Reconfiguration?

1 2 ○ — ○ 1 3



1 2 ○ — ○ 1 3



1 2 ○ — ○ 1 3

- ▶ **list-assignment** L : each vertex v gets allowable colors $L(v)$
- ▶ **L -coloring** φ : φ is proper and $\varphi(v) \in L(v)$ for all v

Main Questions:

- ▶ Given L -colorings α and β , can we change α to β by recoloring single vertices, keeping L -coloring at each step?

What is List Coloring Reconfiguration?

1 2 ○ — ○ 1 3



1 2 ○ — ○ 1 3



1 2 ○ — ○ 1 3

3 4 ○ — ○ 4 5

2 3 ○ — ○ 5 6

1 2 ○ — ○ 6 1

3 4 ○ — ○ 4 5

2 3 ○ — ○ 5 6

1 2 ○ — ○ 6 1

- **list-assignment** L : each vertex v gets allowable colors $L(v)$
- **L -coloring** φ : φ is proper and $\varphi(v) \in L(v)$ for all v

Main Questions:

- Given L -colorings α and β , can we change α to β by recoloring single vertices, keeping L -coloring at each step?

What is List Coloring Reconfiguration?

1 2 ○ — ○ 1 3



1 2 ○ — ○ 1 3



1 2 ○ — ○ 1 3

3 4 ○ — ○ 4 5

2 3 ○ — ○ 5 6

1 2 ○ — ○ 6 1



3 4 ○ — ○ 4 5

2 3 ○ — ○ 5 6



1 2 ○ — ○ 6 1

- ▶ **list-assignment** L : each vertex v gets allowable colors $L(v)$
- ▶ **L -coloring** φ : φ is proper and $\varphi(v) \in L(v)$ for all v

Main Questions:

- ▶ Given L -colorings α and β , can we change α to β by recoloring single vertices, keeping L -coloring at each step?

What is List Coloring Reconfiguration?




- ▶ **list-assignment** L : each vertex v gets allowable colors $L(v)$
- ▶ **L -coloring** φ : φ is proper and $\varphi(v) \in L(v)$ for all v

Main Questions:

- ▶ Given L -colorings α and β , can we change α to β by recoloring single vertices, keeping L -coloring at each step?
- ▶ If so, how many steps are needed?

What is List Coloring Reconfiguration?




- **list-assignment** L : each vertex v gets allowable colors $L(v)$
- **L -coloring** φ : φ is proper and $\varphi(v) \in L(v)$ for all v

Main Questions:

- Given L -colorings α and β , can we change α to β by recoloring single vertices, keeping L -coloring at each step?
- If so, how many steps are needed?
- Given list-assignment L , can we transform every L -coloring α into every L -coloring β ?

What is List Coloring Reconfiguration?



- ▶ **list-assignment** L : each vertex v gets allowable colors $L(v)$
- ▶ **L -coloring** φ : φ is proper and $\varphi(v) \in L(v)$ for all v

Main Questions:

- ▶ Given L -colorings α and β , can we change α to β by recoloring single vertices, keeping L -coloring at each step?
- ▶ If so, how many steps are needed?
- ▶ Given list-assignment L , can we transform every L -coloring α into every L -coloring β ?
- ▶ If so, how many steps are needed in the worst case?

Lists of Size $d(v) + 2$


Lists of Size $d(v) + 2$

Prop: For every G and every f , with $f(v) \geq 2$ for all v , there is list assignment L with $|L(v)| = f(v)$ for all v and L -colorings α and β where changing α to β needs $n(G) + \mu(G)$ moves.

Lists of Size $d(v) + 2$

Prop: For every G and every f , with $f(v) \geq 2$ for all v , there is list assignment L with $|L(v)| = f(v)$ for all v and L -colorings α and β where changing α to β needs $n(G) + \mu(G)$ moves.


Pf: Every vert needs recolored; every edge of M needs extra step.



Lists of Size $d(v) + 2$

Prop: For every G and every f , with $f(v) \geq 2$ for all v , there is list assignment L with $|L(v)| = f(v)$ for all v and L -colorings α and β where changing α to β needs $n(G) + \mu(G)$ moves.

Pf: Every vert needs recolored; every edge of M needs extra step.




Thm:[Cambie–Cames van Batenburg–C.] arXiv:2204.07928

Lists of Size $d(v) + 2$

Prop: For every G and every f , with $f(v) \geq 2$ for all v , there is list assignment L with $|L(v)| = f(v)$ for all v and L -colorings α and β where changing α to β needs $n(G) + \mu(G)$ moves.

Pf: Every vert needs recolored; every edge of M needs extra step.




Thm:[Cambie–Cames van Batenburg–C.] arXiv:2204.07928

(a) If $|L(v)| \geq 2d(v) + 1$, then $n(G) + \mu(G)$ steps suffice.

Lists of Size $d(v) + 2$

Prop: For every G and every f , with $f(v) \geq 2$ for all v , there is list assignment L with $|L(v)| = f(v)$ for all v and L -colorings α and β where changing α to β needs $n(G) + \mu(G)$ moves.

Pf: Every vert needs recolored; every edge of M needs extra step.



Thm:[Cambie–Cames van Batenburg–C.] arXiv:2204.07928


(a) If $|L(v)| \geq 2d(v) + 1$, then $n(G) + \mu(G)$ steps suffice.

(b) If $|L(v)| \geq d(v) + 2$, then $n(G) + 2\mu(G)$ steps suffice.

Lists of Size $d(v) + 2$

Prop: For every G and every f , with $f(v) \geq 2$ for all v , there is list assignment L with $|L(v)| = f(v)$ for all v and L -colorings α and β where changing α to β needs $n(G) + \mu(G)$ moves.

Pf: Every vert needs recolored; every edge of M needs extra step.



Thm:[Cambie–Cames van Batenburg–C.] arXiv:2204.07928

(a) If $|L(v)| \geq 2d(v) + 1$, then $n(G) + \mu(G)$ steps suffice.


(b) If $|L(v)| \geq d(v) + 2$, then $n(G) + 2\mu(G)$ steps suffice.

Conj:[Cambie–Cames van Batenburg–C.] For list assignment L with $|L(v)| \geq d(v) + 2$ for all v and L -colorings α and β , can always change α to β in at most $n(G) + \mu(G)$ steps.

Lists of Size $d(v) + 2$

Prop: For every G and every f , with $f(v) \geq 2$ for all v , there is list assignment L with $|L(v)| = f(v)$ for all v and L -colorings α and β where changing α to β needs $n(G) + \mu(G)$ moves.

Pf: Every vert needs recolored; every edge of M needs extra step.



Thm:[Cambie–Cames van Batenburg–C.] arXiv:2204.07928

(a) If $|L(v)| \geq 2d(v) + 1$, then $n(G) + \mu(G)$ steps suffice.

(b) If $|L(v)| \geq d(v) + 2$, then $n(G) + 2\mu(G)$ steps suffice.


Conj:[Cambie–Cames van Batenburg–C.] For list assignment L with $|L(v)| \geq d(v) + 2$ for all v and L -colorings α and β , can always change α to β in at most $n(G) + \mu(G)$ steps.

Correspondence Coloring: $\mu(G) \rightarrow \tau(G)$.

Lists of Size $d(v) + 2$

Prop: For every G and every f , with $f(v) \geq 2$ for all v , there is list assignment L with $|L(v)| = f(v)$ for all v and L -colorings α and β where changing α to β needs $n(G) + \mu(G)$ moves.

Pf: Every vert needs recolored; every edge of M needs extra step.



Thm:[Cambie–Cames van Batenburg–C.] arXiv:2204.07928

(a) If $|L(v)| \geq 2d(v) + 1$, then $n(G) + \mu(G)$ steps suffice.

(b) If $|L(v)| \geq d(v) + 2$, then $n(G) + 2\mu(G)$ steps suffice.

Conj:[Cambie–Cames van Batenburg–C.] For list assignment L with $|L(v)| \geq d(v) + 2$ for all v and L -colorings α and β , can always change α to β in at most $n(G) + \mu(G)$ steps.

Correspondence Coloring: $\mu(G) \rightarrow \tau(G)$. Conj. and Theorems

Lists of Size $d(v) + 1$

Cambie–Cames van Batenburg–C.–Kang–van den Heuvel:

Let G be connected with n verts and lists L . Let $\mathcal{C}(G, L)$ be the reconfig. graph, and $\hat{\mathcal{C}}(G, L)$ be with all frozen colorings deleted.

Lists of Size $d(v) + 1$

Cambie–Cames van Batenburg–C.–Kang–van den Heuvel:

Let G be connected with n verts and lists L . Let $\mathcal{C}(G, L)$ be the reconfig. graph, and $\hat{\mathcal{C}}(G, L)$ be with all frozen colorings deleted.

Main Thm: If $|L(v)| \geq d(v) + 1$ for all v and G has $\Delta \geq 3$, then $\hat{\mathcal{C}}(G, L)$ is connected with diameter $O(n^2)$. arXiv:2505.08020

Lists of Size $d(v) + 1$

Cambie–Cames van Batenburg–C.–Kang–van den Heuvel:

Let G be connected with n verts and lists L . Let $\mathcal{C}(G, L)$ be the reconfig. graph, and $\hat{\mathcal{C}}(G, L)$ be with all frozen colorings deleted.

Main Thm: If $|L(v)| \geq d(v) + 1$ for all v and G has $\Delta \geq 3$, then $\hat{\mathcal{C}}(G, L)$ is connected with diameter $O(n^2)$. arXiv:2505.08020

Key Lem: If $|L(v)| \geq d(v) + 1$ for all v and $|L(w)| \geq d(w) + 2$ for at least one w , then $\mathcal{C}(G, L)$ is connected with diameter $O(n^2)$.

Lists of Size $d(v) + 1$


Cambie–Cames van Batenburg–C.–Kang–van den Heuvel:

Let G be connected with n verts and lists L . Let $\mathcal{C}(G, L)$ be the reconfig. graph, and $\hat{\mathcal{C}}(G, L)$ be with all frozen colorings deleted.

Main Thm: If $|L(v)| \geq d(v) + 1$ for all v and G has $\Delta \geq 3$, then $\hat{\mathcal{C}}(G, L)$ is connected with diameter $O(n^2)$. [arXiv:2505.08020](#)

Key Lem: If $|L(v)| \geq d(v) + 1$ for all v and $|L(w)| \geq d(w) + 2$ for at least one w , then $\mathcal{C}(G, L)$ is connected with diameter $O(n^2)$.

Shattering Obs: If $|L(w)| = d(w)$ for some w and $|L(v)| \geq d(v) + 1$ for all other v , then the number of components in $\hat{\mathcal{C}}(G, L)$ can be exponential in n .



Lists of Size $d(v) + 1$


Cambie–Cames van Batenburg–C.–Kang–van den Heuvel:

Let G be connected with n verts and lists L . Let $\mathcal{C}(G, L)$ be the reconfig. graph, and $\hat{\mathcal{C}}(G, L)$ be with all frozen colorings deleted.

Main Thm: If $|L(v)| \geq d(v) + 1$ for all v and G has $\Delta \geq 3$, then $\hat{\mathcal{C}}(G, L)$ is connected with diameter $O(n^2)$. [arXiv:2505.08020](#)

Key Lem: If $|L(v)| \geq d(v) + 1$ for all v and $|L(w)| \geq d(w) + 2$ for at least one w , then $\mathcal{C}(G, L)$ is connected with diameter $O(n^2)$.

Shattering Obs: If $|L(w)| = d(w)$ for some w and $|L(v)| \geq d(v) + 1$ for all other v , then the number of components in $\hat{\mathcal{C}}(G, L)$ can be exponential in n .



Conj: If $\delta(G) \geq 3$, then we can improve to diameter $O(n)$ in (a) the Key Lem

Lists of Size $d(v) + 1$


Cambie–Cames van Batenburg–C.–Kang–van den Heuvel:

Let G be connected with n verts and lists L . Let $\mathcal{C}(G, L)$ be the reconfig. graph, and $\hat{\mathcal{C}}(G, L)$ be with all frozen colorings deleted.

Main Thm: If $|L(v)| \geq d(v) + 1$ for all v and G has $\Delta \geq 3$, then $\hat{\mathcal{C}}(G, L)$ is connected with diameter $O(n^2)$. arXiv:2505.08020

Key Lem: If $|L(v)| \geq d(v) + 1$ for all v and $|L(w)| \geq d(w) + 2$ for at least one w , then $\mathcal{C}(G, L)$ is connected with diameter $O(n^2)$.

Shattering Obs: If $|L(w)| = d(w)$ for some w and $|L(v)| \geq d(v) + 1$ for all other v , then the number of components in $\hat{\mathcal{C}}(G, L)$ can be exponential in n .



Conj: If $\delta(G) \geq 3$, then we can improve to diameter $O(n)$ in (a) the Key Lem and (b) the Main Thm.

Lists of Size $d(v) + 1$


Cambie–Cames van Batenburg–C.–Kang–van den Heuvel:

Let G be connected with n verts and lists L . Let $\mathcal{C}(G, L)$ be the reconfig. graph, and $\hat{\mathcal{C}}(G, L)$ be with all frozen colorings deleted.

Main Thm: If $|L(v)| \geq d(v) + 1$ for all v and G has $\Delta \geq 3$, then $\hat{\mathcal{C}}(G, L)$ is connected with diameter $O(n^2)$. arXiv:2505.08020

Key Lem: If $|L(v)| \geq d(v) + 1$ for all v and $|L(w)| \geq d(w) + 2$ for at least one w , then $\mathcal{C}(G, L)$ is connected with diameter $O(n^2)$.

Shattering Obs: If $|L(w)| = d(w)$ for some w and $|L(v)| \geq d(v) + 1$ for all other v , then the number of components in $\hat{\mathcal{C}}(G, L)$ can be exponential in n .



Conj: If $\delta(G) \geq 3$, then we can improve to diameter $O(n)$ in (a) the Key Lem and (b) the Main Thm. What about correspondence?

Lists of Size $d(v) + 1$: Ideas to Prove Main Thm

Thm: Let a graph G be 3-connected and regular. If α and β are unfrozen $(\Delta + 1)$ -colorings of G , then α can reach β .

Lists of Size $d(v) + 1$: Ideas to Prove Main Thm

Thm: Let a graph G be 3-connected and regular. If α and β are unfrozen $(\Delta + 1)$ -colorings of G , then α can reach β .

Bonus Lem: Say G is 3-connected with $|L(v)| \geq d(v) + 1$ for all v . Let x_1, x_2 be at distance 2. If α and β are L -colorings with $\alpha(x_1) = \alpha(x_2) = \beta(x_1) = \beta(x_2)$, then α can reach β .

Lists of Size $d(v) + 1$: Ideas to Prove Main Thm

Thm: Let a graph G be 3-connected and regular. If α and β are unfrozen $(\Delta + 1)$ -colorings of G , then α can reach β .

Bonus Lem: Say G is 3-connected with $|L(v)| \geq d(v) + 1$ for all v . Let x_1, x_2 be at distance 2. If α and β are L -colorings with $\alpha(x_1) = \alpha(x_2) = \beta(x_1) = \beta(x_2)$, then α can reach β .

Pf: Now a common neighbor y of x_1, x_2 effectively has an extra color.

Lists of Size $d(v) + 1$: Ideas to Prove Main Thm

Thm: Let a graph G be 3-connected and regular. If α and β are unfrozen $(\Delta + 1)$ -colorings of G , then α can reach β .

Bonus Lem: Say G is 3-connected with $|L(v)| \geq d(v) + 1$ for all v . Let x_1, x_2 be at distance 2. If α and β are L -colorings with $\alpha(x_1) = \alpha(x_2) = \beta(x_1) = \beta(x_2)$, then α can reach β .

Pf: Now a common neighbor y of x_1, x_2 effectively has an extra color. So we finish by Key Lem.

Lists of Size $d(v) + 1$: Ideas to Prove Main Thm

Thm: Let a graph G be 3-connected and regular. If α and β are unfrozen $(\Delta + 1)$ -colorings of G , then α can reach β .

Bonus Lem: Say G is 3-connected with $|L(v)| \geq d(v) + 1$ for all v . Let x_1, x_2 be at distance 2. If α and β are L -colorings with $\alpha(x_1) = \alpha(x_2) = \beta(x_1) = \beta(x_2)$, then α can reach β .

Pf: Now a common neighbor y of x_1, x_2 effectively has an extra color. So we finish by Key Lem.

Pf of Thm: Find distinct w_1, w_2, x_1, x_2 with:

- (i) w_1, w_2 at distance 2 and $\alpha(w_1) = \alpha(w_2)$ and
- (ii) x_1, x_2 at distance 2 and $\beta(x_1) = \beta(x_2)$.

Using the Bonus Lem 4 times gives:

Lists of Size $d(v) + 1$: Ideas to Prove Main Thm

Thm: Let a graph G be 3-connected and regular. If α and β are unfrozen $(\Delta + 1)$ -colorings of G , then α can reach β .


Bonus Lem: Say G is 3-connected with $|L(v)| \geq d(v) + 1$ for all v . Let x_1, x_2 be at distance 2. If α and β are L -colorings with $\alpha(x_1) = \alpha(x_2) = \beta(x_1) = \beta(x_2)$, then α can reach β .

Pf: Now a common neighbor y of x_1, x_2 effectively has an extra color. So we finish by Key Lem.


Pf of Thm: Find distinct w_1, w_2, x_1, x_2 with:

- (i) w_1, w_2 at distance 2 and $\alpha(w_1) = \alpha(w_2)$ and
- (ii) x_1, x_2 at distance 2 and $\beta(x_1) = \beta(x_2)$.

Using the Bonus Lem 4 times gives:



α



β

Lists of Size $d(v) + 1$: Ideas to Prove Main Thm

Thm: Let a graph G be 3-connected and regular. If α and β are unfrozen $(\Delta + 1)$ -colorings of G , then α can reach β .


Bonus Lem: Say G is 3-connected with $|L(v)| \geq d(v) + 1$ for all v . Let x_1, x_2 be at distance 2. If α and β are L -colorings with $\alpha(x_1) = \alpha(x_2) = \beta(x_1) = \beta(x_2)$, then α can reach β .

Pf: Now a common neighbor y of x_1, x_2 effectively has an extra color. So we finish by Key Lem.

Pf of Thm: Find distinct w_1, w_2, x_1, x_2 with:

- (i) w_1, w_2 at distance 2 and $\alpha(w_1) = \alpha(w_2)$ and
- (ii) x_1, x_2 at distance 2 and $\beta(x_1) = \beta(x_2)$.

Using the Bonus Lem 4 times gives:



Lists of Size $d(v) + 1$: Ideas to Prove Main Thm

Thm: Let a graph G be 3-connected and regular. If α and β are unfrozen $(\Delta + 1)$ -colorings of G , then α can reach β .


Bonus Lem: Say G is 3-connected with $|L(v)| \geq d(v) + 1$ for all v . Let x_1, x_2 be at distance 2. If α and β are L -colorings with $\alpha(x_1) = \alpha(x_2) = \beta(x_1) = \beta(x_2)$, then α can reach β .

Pf: Now a common neighbor y of x_1, x_2 effectively has an extra color. So we finish by Key Lem.

Pf of Thm: Find distinct w_1, w_2, x_1, x_2 with:

- (i) w_1, w_2 at distance 2 and $\alpha(w_1) = \alpha(w_2)$ and
- (ii) x_1, x_2 at distance 2 and $\beta(x_1) = \beta(x_2)$.

Using the Bonus Lem 4 times gives:



Lists of Size $d(v) + 1$: Ideas to Prove Main Thm

Thm: Let a graph G be 3-connected and regular. If α and β are unfrozen $(\Delta + 1)$ -colorings of G , then α can reach β .

Bonus Lem: Say G is 3-connected with $|L(v)| \geq d(v) + 1$ for all v . Let x_1, x_2 be at distance 2. If α and β are L -colorings with $\alpha(x_1) = \alpha(x_2) = \beta(x_1) = \beta(x_2)$, then α can reach β .

Pf: Now a common neighbor y of x_1, x_2 effectively has an extra color. So we finish by Key Lem.

Pf of Thm: Find distinct w_1, w_2, x_1, x_2 with:

- (i) w_1, w_2 at distance 2 and $\alpha(w_1) = \alpha(w_2)$ and
- (ii) x_1, x_2 at distance 2 and $\beta(x_1) = \beta(x_2)$.

Using the Bonus Lem 4 times gives:



Lists of Size $d(v) + 1$: Ideas to Prove Main Thm

Thm: Let a graph G be 3-connected and regular. If α and β are unfrozen $(\Delta + 1)$ -colorings of G , then α can reach β .


Bonus Lem: Say G is 3-connected with $|L(v)| \geq d(v) + 1$ for all v . Let x_1, x_2 be at distance 2. If α and β are L -colorings with $\alpha(x_1) = \alpha(x_2) = \beta(x_1) = \beta(x_2)$, then α can reach β .

Pf: Now a common neighbor y of x_1, x_2 effectively has an extra color. So we finish by Key Lem.

Pf of Thm: Find distinct w_1, w_2, x_1, x_2 with:

- (i) w_1, w_2 at distance 2 and $\alpha(w_1) = \alpha(w_2)$ and
- (ii) x_1, x_2 at distance 2 and $\beta(x_1) = \beta(x_2)$.

Using the Bonus Lem 4 times gives:



Lists of Size $d(v) + 1$: Ideas to Prove Main Thm

Thm: Let a graph G be 3-connected and regular. If α and β are unfrozen $(\Delta + 1)$ -colorings of G , then α can reach β .


Bonus Lem: Say G is 3-connected with $|L(v)| \geq d(v) + 1$ for all v . Let x_1, x_2 be at distance 2. If α and β are L -colorings with $\alpha(x_1) = \alpha(x_2) = \beta(x_1) = \beta(x_2)$, then α can reach β .

Pf: Now a common neighbor y of x_1, x_2 effectively has an extra color. So we finish by Key Lem.

Pf of Thm: Find distinct w_1, w_2, x_1, x_2 with:

- (i) w_1, w_2 at distance 2 and $\alpha(w_1) = \alpha(w_2)$ and
- (ii) x_1, x_2 at distance 2 and $\beta(x_1) = \beta(x_2)$.

Using the Bonus Lem 4 times gives:



Lists of Size $d(v) + 1$: Ideas to Prove Main Thm

Thm: Let a graph G be 3-connected and regular. If α and β are unfrozen $(\Delta + 1)$ -colorings of G , then α can reach β .


Bonus Lem: Say G is 3-connected with $|L(v)| \geq d(v) + 1$ for all v . Let x_1, x_2 be at distance 2. If α and β are L -colorings with $\alpha(x_1) = \alpha(x_2) = \beta(x_1) = \beta(x_2)$, then α can reach β .

Pf: Now a common neighbor y of x_1, x_2 effectively has an extra color. So we finish by Key Lem.

Pf of Thm: Find distinct w_1, w_2, x_1, x_2 with:

- (i) w_1, w_2 at distance 2 and $\alpha(w_1) = \alpha(w_2)$ and
- (ii) x_1, x_2 at distance 2 and $\beta(x_1) = \beta(x_2)$.

Using the Bonus Lem 4 times gives:



Planar Graphs and other Sparse Graphs

Thm: If G is planar and $|L(v)| = 10$ for all v , then $\mathcal{C}(G, L)$ is connected with diameter $O(n)$. arXiv:2411.00679

Planar Graphs and other Sparse Graphs

Thm: If G is planar and $|L(v)| = 10$ for all v , then $\mathcal{C}(G, L)$ is connected with diameter $O(n)$. arXiv:2411.00679

Thm: If G is planar and triangle-free and $|L(v)| = 7$ for all v , then $\mathcal{C}(G, L)$ is connected with diameter $O(n)$. arXiv:2201.05133

Planar Graphs and other Sparse Graphs

Thm: If G is planar and $|L(v)| = 10$ for all v , then $\mathcal{C}(G, L)$ is connected with diameter $O(n)$. arXiv:2411.00679

Thm: If G is planar and triangle-free and $|L(v)| = 7$ for all v , then $\mathcal{C}(G, L)$ is connected with diameter $O(n)$. arXiv:2201.05133

Rem: Both results confirm conjectures of Dvorak and Feghali, who proved the cases when $L(v) = \{1, \dots, 10\}$ and $L(v) = \{1, \dots, 7\}$.

Planar Graphs and other Sparse Graphs

Thm: If G is planar and $|L(v)| = 10$ for all v , then $\mathcal{C}(G, L)$ is connected with diameter $O(n)$. arXiv:2411.00679

Thm: If G is planar and triangle-free and $|L(v)| = 7$ for all v , then $\mathcal{C}(G, L)$ is connected with diameter $O(n)$. arXiv:2201.05133

Rem: Both results confirm conjectures of Dvorak and Feghali, who proved the cases when $L(v) = \{1, \dots, 10\}$ and $L(v) = \{1, \dots, 7\}$.

Conj: Fix integers k, d with $k \geq d + 3$. If G is d -degenerate and $|L(v)| = k$ for all v , then $\mathcal{C}(G, L)$ is connected with diameter $O(n)$.

Planar Graphs and other Sparse Graphs

Thm: If G is planar and $|L(v)| = 10$ for all v , then $\mathcal{C}(G, L)$ is connected with diameter $O(n)$. arXiv:2411.00679

Thm: If G is planar and triangle-free and $|L(v)| = 7$ for all v , then $\mathcal{C}(G, L)$ is connected with diameter $O(n)$. arXiv:2201.05133

Rem: Both results confirm conjectures of Dvorak and Feghali, who proved the cases when $L(v) = \{1, \dots, 10\}$ and $L(v) = \{1, \dots, 7\}$.

Conj: Fix integers k, d with $k \geq d + 3$. If G is d -degenerate and $|L(v)| = k$ for all v , then $\mathcal{C}(G, L)$ is connected with diameter $O(n)$.

Conj: For every “natural” graph class \mathcal{G} , positive integer k , and $G \in \mathcal{G}$, if $\mathcal{C}(G, k)$ always has diam $O(n)$, then also $\mathcal{C}(G, L)$ always has diam $O(n)$ when $|L(v)| = k$ for all v .

Planar Graphs and other Sparse Graphs

Thm: If G is planar and $|L(v)| = 10$ for all v , then $\mathcal{C}(G, L)$ is connected with diameter $O(n)$. arXiv:2411.00679

Thm: If G is planar and triangle-free and $|L(v)| = 7$ for all v , then $\mathcal{C}(G, L)$ is connected with diameter $O(n)$. arXiv:2201.05133

Rem: Both results confirm conjectures of Dvorak and Feghali, who proved the cases when $L(v) = \{1, \dots, 10\}$ and $L(v) = \{1, \dots, 7\}$.

Conj: Fix integers k, d with $k \geq d + 3$. If G is d -degenerate and $|L(v)| = k$ for all v , then $\mathcal{C}(G, L)$ is connected with diameter $O(n)$.

Conj: For every “natural” graph class \mathcal{G} , positive integer k , and $G \in \mathcal{G}$, if $\mathcal{C}(G, k)$ always has diam $O(n)$, then also $\mathcal{C}(G, L)$ always has diam $O(n)$ when $|L(v)| = k$ for all v . This holds when:


- ▶ \mathcal{G} is graphs with $\text{mad}(G) < a$, for some a ; or
- ▶ \mathcal{G} is planar graphs with girth at least g , for some g .

Main Tool for Sparse Graphs

Obs: Fix G and L . If $\exists v$ s.t. $|L(v)| \geq d(v) + 2$, then $\mathcal{C}_L(G)$ is connected iff $\mathcal{C}_L(G - v)$ is connected. So $\mathcal{C}_L(G)$ is connected if G is d -degenerate and L is $(d + 2)$ -assignment. **Pf:** Induction on $|G|$.


Main Tool for Sparse Graphs

Obs: Fix G and L . If $\exists v$ s.t. $|L(v)| \geq d(v) + 2$, then $\mathcal{C}_L(G)$ is connected iff $\mathcal{C}_L(G - v)$ is connected. So $\mathcal{C}_L(G)$ is connected if G is d -degenerate and L is $(d + 2)$ -assignment. **Pf:** Induction on $|G|$.




Main Tool for Sparse Graphs

Obs: Fix G and L . If $\exists v$ s.t. $|L(v)| \geq d(v) + 2$, then $\mathcal{C}_L(G)$ is connected iff $\mathcal{C}_L(G - v)$ is connected. So $\mathcal{C}_L(G)$ is connected if G is d -degenerate and L is $(d + 2)$ -assignment. **Pf:** Induction on $|G|$.



Main Tool for Sparse Graphs


Obs: Fix G and L . If $\exists v$ s.t. $|L(v)| \geq d(v) + 2$, then $\mathcal{C}_L(G)$ is connected iff $\mathcal{C}_L(G - v)$ is connected. So $\mathcal{C}_L(G)$ is connected if G is d -degenerate and L is $(d + 2)$ -assignment. **Pf:** Induction on $|G|$.



Key Lem: Fix G , L , v , and L -colorings α and β . Let $G' := G - v$, $\alpha' := \alpha|_{G'}$, $\beta' := \beta|_{G'}$. If we can transform α' to β' only recoloring $N(v)$ at most s times, then we can transform α to β only recoloring v at most $\lceil \frac{s}{|L(v)| - d(v) - 1} \rceil + 1$ times.

Main Tool for Sparse Graphs


Obs: Fix G and L . If $\exists v$ s.t. $|L(v)| \geq d(v) + 2$, then $\mathcal{C}_L(G)$ is connected iff $\mathcal{C}_L(G - v)$ is connected. So $\mathcal{C}_L(G)$ is connected if G is d -degenerate and L is $(d + 2)$ -assignment. **Pf:** Induction on $|G|$.



Key Lem: Fix G , L , v , and L -colorings α and β . Let $G' := G - v$, $\alpha' := \alpha|_{G'}$, $\beta' := \beta|_{G'}$. If we can transform α' to β' only recoloring $N(v)$ at most s times, then we can transform α to β only recoloring v at most $\lceil \frac{s}{|L(v)| - d(v) - 1} \rceil + 1$ times. **Pf:** Above, more carefully.

Main Tool for Sparse Graphs


Obs: Fix G and L . If $\exists v$ s.t. $|L(v)| \geq d(v) + 2$, then $\mathcal{C}_L(G)$ is connected iff $\mathcal{C}_L(G - v)$ is connected. So $\mathcal{C}_L(G)$ is connected if G is d -degenerate and L is $(d + 2)$ -assignment. **Pf:** Induction on $|G|$.



Key Lem: Fix G , L , v , and L -colorings α and β . Let $G' := G - v$, $\alpha' := \alpha|_{G'}$, $\beta' := \beta|_{G'}$. If we can transform α' to β' only recoloring $N(v)$ at most s times, then we can transform α to β only recoloring v at most $\lceil \frac{s}{|L(v)| - d(v) - 1} \rceil + 1$ times. **Pf:** Above, more carefully.

Main Tool for Sparse Graphs


Obs: Fix G and L . If $\exists v$ s.t. $|L(v)| \geq d(v) + 2$, then $\mathcal{C}_L(G)$ is connected iff $\mathcal{C}_L(G - v)$ is connected. So $\mathcal{C}_L(G)$ is connected if G is d -degenerate and L is $(d + 2)$ -assignment. **Pf:** Induction on $|G|$.



Key Lem: Fix G , L , v , and L -colorings α and β . Let $G' := G - v$, $\alpha' := \alpha|_{G'}$, $\beta' := \beta|_{G'}$. If we can transform α' to β' only recoloring $N(v)$ at most s times, then we can transform α to β only recoloring v at most $\lceil \frac{s}{|L(v)| - d(v) - 1} \rceil + 1$ times. **Pf:** Above, more carefully.

Main Tool for Sparse Graphs

Obs: Fix G and L . If $\exists v$ s.t. $|L(v)| \geq d(v) + 2$, then $\mathcal{C}_L(G)$ is connected iff $\mathcal{C}_L(G - v)$ is connected. So $\mathcal{C}_L(G)$ is connected if G is d -degenerate and L is $(d + 2)$ -assignment. **Pf:** Induction on $|G|$.




$c_1, c_2, c_3, c_4, c_5, c_6, c_7, c_8, \dots$

Key Lem: Fix G , L , v , and L -colorings α and β . Let $G' := G - v$, $\alpha' := \alpha|_{G'}$, $\beta' := \beta|_{G'}$. If we can transform α' to β' only recoloring $N(v)$ at most s times, then we can transform α to β only recoloring v at most $\lceil \frac{s}{|L(v)| - d(v) - 1} \rceil + 1$ times. **Pf:** Above, more carefully.

Main Tool for Sparse Graphs

Obs: Fix G and L . If $\exists v$ s.t. $|L(v)| \geq d(v) + 2$, then $\mathcal{C}_L(G)$ is connected iff $\mathcal{C}_L(G - v)$ is connected. So $\mathcal{C}_L(G)$ is connected if G is d -degenerate and L is $(d + 2)$ -assignment. **Pf:** Induction on $|G|$.




↓

$c_1, c_2, c_3, c_4, c_5, c_6, c_7, c_8, \dots$

Key Lem: Fix G , L , v , and L -colorings α and β . Let $G' := G - v$, $\alpha' := \alpha|_{G'}$, $\beta' := \beta|_{G'}$. If we can transform α' to β' only recoloring $N(v)$ at most s times, then we can transform α to β only recoloring v at most $\lceil \frac{s}{|L(v)| - d(v) - 1} \rceil + 1$ times. **Pf:** Above, more carefully.

Main Tool for Sparse Graphs

Obs: Fix G and L . If $\exists v$ s.t. $|L(v)| \geq d(v) + 2$, then $\mathcal{C}_L(G)$ is connected iff $\mathcal{C}_L(G - v)$ is connected. So $\mathcal{C}_L(G)$ is connected if G is d -degenerate and L is $(d + 2)$ -assignment. **Pf:** Induction on $|G|$.



\downarrow
 $c_1, c_2, c_3, c_4, c_5, c_6, c_7, c_8, \dots$

Key Lem: Fix G , L , v , and L -colorings α and β . Let $G' := G - v$, $\alpha' := \alpha|_{G'}$, $\beta' := \beta|_{G'}$. If we can transform α' to β' only recoloring $N(v)$ at most s times, then we can transform α to β only recoloring v at most $\lceil \frac{s}{|L(v)| - d(v) - 1} \rceil + 1$ times. **Pf:** Above, more carefully.

Summary

Thm: If $|L(v)| \geq d(v) + 2$, then $\text{diam } \mathcal{C}(G, L) \leq n + 2\mu(G)$.

Summary

Thm: If $|L(v)| \geq d(v) + 2$, then $\text{diam } \mathcal{C}(G, L) \leq n + 2\mu(G)$.

Thm: If $|L(v)| \geq 2d(v) + 1$, then $\text{diam } \mathcal{C}(G, L) \leq n + \mu(G)$.

Summary

Thm: If $|L(v)| \geq d(v) + 2$, then $\text{diam } \mathcal{C}(G, L) \leq n + 2\mu(G)$.

Thm: If $|L(v)| \geq 2d(v) + 1$, then $\text{diam } \mathcal{C}(G, L) \leq n + \mu(G)$.

Conj: If $|L(v)| \geq d(v) + 2$, then $\text{diam } \mathcal{C}(G, L) \leq n + \mu(G)$.

Summary

Thm: If $|L(v)| \geq d(v) + 2$, then $\text{diam } \mathcal{C}(G, L) \leq n + 2\mu(G)$.

Thm: If $|L(v)| \geq 2d(v) + 1$, then $\text{diam } \mathcal{C}(G, L) \leq n + \mu(G)$.

Conj: If $|L(v)| \geq d(v) + 2$, then $\text{diam } \mathcal{C}(G, L) \leq n + \mu(G)$.

Correspondence: $\mu(G) \rightarrow \tau(G)$.

Summary

Thm: If $|L(v)| \geq d(v) + 2$, then $\text{diam } \mathcal{C}(G, L) \leq n + 2\mu(G)$.

Thm: If $|L(v)| \geq 2d(v) + 1$, then $\text{diam } \mathcal{C}(G, L) \leq n + \mu(G)$.

Conj: If $|L(v)| \geq d(v) + 2$, then $\text{diam } \mathcal{C}(G, L) \leq n + \mu(G)$.

Correspondence: $\mu(G) \rightarrow \tau(G)$.

Thm: If $|L(v)| \geq d(v) + 1$ and $\Delta \geq 3$, then $\text{diam } \hat{\mathcal{C}}(G, L) \leq O(n^2)$.

Summary

Thm: If $|L(v)| \geq d(v) + 2$, then $\text{diam } \mathcal{C}(G, L) \leq n + 2\mu(G)$.

Thm: If $|L(v)| \geq 2d(v) + 1$, then $\text{diam } \mathcal{C}(G, L) \leq n + \mu(G)$.

Conj: If $|L(v)| \geq d(v) + 2$, then $\text{diam } \mathcal{C}(G, L) \leq n + \mu(G)$.

Correspondence: $\mu(G) \rightarrow \tau(G)$.

Thm: If $|L(v)| \geq d(v) + 1$ and $\Delta \geq 3$, then $\text{diam } \hat{\mathcal{C}}(G, L) \leq O(n^2)$.

Conj: If also $\delta(G) \geq 3$, then $\text{diam } \hat{\mathcal{C}}(G, L) \leq O(n)$.

Summary

Thm: If $|L(v)| \geq d(v) + 2$, then $\text{diam } \mathcal{C}(G, L) \leq n + 2\mu(G)$.

Thm: If $|L(v)| \geq 2d(v) + 1$, then $\text{diam } \mathcal{C}(G, L) \leq n + \mu(G)$.

Conj: If $|L(v)| \geq d(v) + 2$, then $\text{diam } \mathcal{C}(G, L) \leq n + \mu(G)$.

Correspondence: $\mu(G) \rightarrow \tau(G)$.

Thm: If $|L(v)| \geq d(v) + 1$ and $\Delta \geq 3$, then $\text{diam } \hat{\mathcal{C}}(G, L) \leq O(n^2)$.

Conj: If also $\delta(G) \geq 3$, then $\text{diam } \hat{\mathcal{C}}(G, L) \leq O(n)$.

Correspondence: Analogue is false (as shown by cliques).

Summary

Thm: If $|L(v)| \geq d(v) + 2$, then $\text{diam } \mathcal{C}(G, L) \leq n + 2\mu(G)$.

Thm: If $|L(v)| \geq 2d(v) + 1$, then $\text{diam } \mathcal{C}(G, L) \leq n + \mu(G)$.

Conj: If $|L(v)| \geq d(v) + 2$, then $\text{diam } \mathcal{C}(G, L) \leq n + \mu(G)$.

Correspondence: $\mu(G) \rightarrow \tau(G)$.

Thm: If $|L(v)| \geq d(v) + 1$ and $\Delta \geq 3$, then $\text{diam } \hat{\mathcal{C}}(G, L) \leq O(n^2)$.

Conj: If also $\delta(G) \geq 3$, then $\text{diam } \hat{\mathcal{C}}(G, L) \leq O(n)$.

Correspondence: Analogue is false (as shown by cliques).

Thm: If planar and $|L(v)| = 10$, then $\text{diam } \mathcal{C}(G, L) = O(n)$.

Summary

Thm: If $|L(v)| \geq d(v) + 2$, then $\text{diam } \mathcal{C}(G, L) \leq n + 2\mu(G)$.

Thm: If $|L(v)| \geq 2d(v) + 1$, then $\text{diam } \mathcal{C}(G, L) \leq n + \mu(G)$.

Conj: If $|L(v)| \geq d(v) + 2$, then $\text{diam } \mathcal{C}(G, L) \leq n + \mu(G)$.

Correspondence: $\mu(G) \rightarrow \tau(G)$.

Thm: If $|L(v)| \geq d(v) + 1$ and $\Delta \geq 3$, then $\text{diam } \hat{\mathcal{C}}(G, L) \leq O(n^2)$.

Conj: If also $\delta(G) \geq 3$, then $\text{diam } \hat{\mathcal{C}}(G, L) \leq O(n)$.

Correspondence: Analogue is false (as shown by cliques).

Thm: If planar and $|L(v)| = 10$, then $\text{diam } \mathcal{C}(G, L) = O(n)$.

Thm: If planar, ∇ -free and $|L(v)| = 7$, then $\text{diam } \mathcal{C}(G, L) = O(n)$.

Summary

Thm: If $|L(v)| \geq d(v) + 2$, then $\text{diam } \mathcal{C}(G, L) \leq n + 2\mu(G)$.

Thm: If $|L(v)| \geq 2d(v) + 1$, then $\text{diam } \mathcal{C}(G, L) \leq n + \mu(G)$.

Conj: If $|L(v)| \geq d(v) + 2$, then $\text{diam } \mathcal{C}(G, L) \leq n + \mu(G)$.

Correspondence: $\mu(G) \rightarrow \tau(G)$.

Thm: If $|L(v)| \geq d(v) + 1$ and $\Delta \geq 3$, then $\text{diam } \hat{\mathcal{C}}(G, L) \leq O(n^2)$.

Conj: If also $\delta(G) \geq 3$, then $\text{diam } \hat{\mathcal{C}}(G, L) \leq O(n)$.

Correspondence: Analogue is false (as shown by cliques).

Thm: If planar and $|L(v)| = 10$, then $\text{diam } \mathcal{C}(G, L) = O(n)$.

Thm: If planar, ∇ -free and $|L(v)| = 7$, then $\text{diam } \mathcal{C}(G, L) = O(n)$.

Conj: If d -degen and $|L(v)| \geq d + 3$, then $\text{diam } \mathcal{C}(G, L) = O(n)$.

Summary

Thm: If $|L(v)| \geq d(v) + 2$, then $\text{diam } \mathcal{C}(G, L) \leq n + 2\mu(G)$.

Thm: If $|L(v)| \geq 2d(v) + 1$, then $\text{diam } \mathcal{C}(G, L) \leq n + \mu(G)$.

Conj: If $|L(v)| \geq d(v) + 2$, then $\text{diam } \mathcal{C}(G, L) \leq n + \mu(G)$.

Correspondence: $\mu(G) \rightarrow \tau(G)$.

Thm: If $|L(v)| \geq d(v) + 1$ and $\Delta \geq 3$, then $\text{diam } \hat{\mathcal{C}}(G, L) \leq O(n^2)$.

Conj: If also $\delta(G) \geq 3$, then $\text{diam } \hat{\mathcal{C}}(G, L) \leq O(n)$.

Correspondence: Analogue is false (as shown by cliques).

Thm: If planar and $|L(v)| = 10$, then $\text{diam } \mathcal{C}(G, L) = O(n)$.

Thm: If planar, ∇ -free and $|L(v)| = 7$, then $\text{diam } \mathcal{C}(G, L) = O(n)$.

Conj: If d -degen and $|L(v)| \geq d + 3$, then $\text{diam } \mathcal{C}(G, L) = O(n)$.

Correspondence: Analogues of theorems are true.