

Rosenfeld Counting: Proper Conflict-free Coloring of Graphs with Large Maximum Degree

Daniel W. Cranston

Virginia Commonwealth University

dcranston@vcu.edu

Joint with Chun-Hung Liu

URI Discrete Math Seminar (virtual)

1 November 2024

Nonrepetitive List-coloring of Paths

Ex: A 3-coloring of P_8 with a **square** and a **square-free** 3-coloring.



Nonrepetitive List-coloring of Paths

Ex: A 3-coloring of P_8 with a **square** and a **square-free** 3-coloring.



Fact: Thue found a square-free 3-coloring of the infinite path.

Nonrepetitive List-coloring of Paths

Ex: A 3-coloring of P_8 with a **square** and a **square-free** 3-coloring.



Fact: Thue found a square-free 3-coloring of the infinite path.

Defn: A coloring is **nonrepetitive** if each path is square-free.

Nonrepetitive List-coloring of Paths

Ex: A 3-coloring of P_8 with a **square** and a **square-free** 3-coloring.



Fact: Thue found a square-free 3-coloring of the infinite path.

Defn: A coloring is **nonrepetitive** if each path is square-free.

Conj: For each 3-assignment L to the verts of P_n , there is a nonrepetitive L -coloring φ (with $\varphi(v) \in L(v)$ for all v).

Nonrepetitive List-coloring of Paths

Ex: A 3-coloring of P_8 with a **square** and a **square-free** 3-coloring.



Fact: Thue found a square-free 3-coloring of the infinite path.

Defn: A coloring is **nonrepetitive** if each path is square-free.

Conj: For each 3-assignment L to the verts of P_n , there is a nonrepetitive L -coloring φ (with $\varphi(v) \in L(v)$ for all v).

Lem: Let L be a 4-assignment to $V(P_n)$. For each $i \geq 1$, let \mathcal{C}_i be set of nonrepetitive L -colorings of first i verts of P_n . For all $i < n$,

$$|\mathcal{C}_{i+1}| \geq 2|\mathcal{C}_i|.$$

Nonrepetitive List-coloring of Paths

Ex: A 3-coloring of P_8 with a **square** and a **square-free** 3-coloring.



Fact: Thue found a square-free 3-coloring of the infinite path.

Defn: A coloring is **nonrepetitive** if each path is square-free.

Conj: For each 3-assignment L to the verts of P_n , there is a nonrepetitive L -coloring φ (with $\varphi(v) \in L(v)$ for all v).

Lem: Let L be a 4-assignment to $V(P_n)$. For each $i \geq 1$, let \mathcal{C}_i be set of nonrepetitive L -colorings of first i verts of P_n . For all $i < n$,

$$|\mathcal{C}_{i+1}| \geq 2|\mathcal{C}_i|.$$

Since $|\mathcal{C}_1| = 4$, path P_n has more than 2^n nonrepetitive L -colorings.

Nonrepetitive List-coloring of Paths (the proof)

Lem: Let L be a 4-assignment to $V(P_n)$. For each $i \geq 1$, let \mathcal{C}_i be set of nonrepetitive L -colorings of first i verts of P_n . For all $i < n$,

$$|\mathcal{C}_{i+1}| \geq 2|\mathcal{C}_i|.$$

Nonrepetitive List-coloring of Paths (the proof)

Lem: Let L be a 4-assignment to $V(P_n)$. For each $i \geq 1$, let \mathcal{C}_i be set of nonrepetitive L -colorings of first i verts of P_n . For all $i < n$,

$$|\mathcal{C}_{i+1}| \geq 2|\mathcal{C}_i|.$$

Pf: Induction on i . Let \mathcal{F} be the set of L -colorings of v_1, \dots, v_{i+1} that are nonrepetitive on v_1, \dots, v_i but have a square with v_{i+1} .

Nonrepetitive List-coloring of Paths (the proof)

Lem: Let L be a 4-assignment to $V(P_n)$. For each $i \geq 1$, let \mathcal{C}_i be set of nonrepetitive L -colorings of first i verts of P_n . For all $i < n$,

$$|\mathcal{C}_{i+1}| \geq 2|\mathcal{C}_i|.$$

Pf: Induction on i . Let \mathcal{F} be the set of L -colorings of v_1, \dots, v_{i+1} that are nonrepetitive on v_1, \dots, v_i but have a square with v_{i+1} . Clearly, $|\mathcal{C}_{i+1}| = 4|\mathcal{C}_i| - |\mathcal{F}|$. Let \mathcal{F}_j be subset of \mathcal{F} with a square of length $2j$. So $\mathcal{F} = \cup_{j \geq 1} \mathcal{F}_j$.

Nonrepetitive List-coloring of Paths (the proof)

Lem: Let L be a 4-assignment to $V(P_n)$. For each $i \geq 1$, let \mathcal{C}_i be set of nonrepetitive L -colorings of first i verts of P_n . For all $i < n$,

$$|\mathcal{C}_{i+1}| \geq 2|\mathcal{C}_i|.$$

Pf: Induction on i . Let \mathcal{F} be the set of L -colorings of v_1, \dots, v_{i+1} that are nonrepetitive on v_1, \dots, v_i but have a square with v_{i+1} . Clearly, $|\mathcal{C}_{i+1}| = 4|\mathcal{C}_i| - |\mathcal{F}|$. Let \mathcal{F}_j be subset of \mathcal{F} with a square of length $2j$. So $\mathcal{F} = \cup_{j \geq 1} \mathcal{F}_j$.

Each $\varphi \in \mathcal{F}_j$ restricts to a nonrepetitive L -coloring φ' of v_1, \dots, v_{i+1-j} .

Nonrepetitive List-coloring of Paths (the proof)

Lem: Let L be a 4-assignment to $V(P_n)$. For each $i \geq 1$, let \mathcal{C}_i be set of nonrepetitive L -colorings of first i verts of P_n . For all $i < n$,

$$|\mathcal{C}_{i+1}| \geq 2|\mathcal{C}_i|.$$

Pf: Induction on i . Let \mathcal{F} be the set of L -colorings of v_1, \dots, v_{i+1} that are nonrepetitive on v_1, \dots, v_i but have a square with v_{i+1} . Clearly, $|\mathcal{C}_{i+1}| = 4|\mathcal{C}_i| - |\mathcal{F}|$. Let \mathcal{F}_j be subset of \mathcal{F} with a square of length $2j$. So $\mathcal{F} = \cup_{j \geq 1} \mathcal{F}_j$.

Each $\varphi \in \mathcal{F}_j$ restricts to a nonrepetitive L -coloring φ' of v_1, \dots, v_{i+1-j} . And φ' uniquely determines φ .

Nonrepetitive List-coloring of Paths (the proof)

Lem: Let L be a 4-assignment to $V(P_n)$. For each $i \geq 1$, let \mathcal{C}_i be set of nonrepetitive L -colorings of first i verts of P_n . For all $i < n$,

$$|\mathcal{C}_{i+1}| \geq 2|\mathcal{C}_i|.$$

Pf: Induction on i . Let \mathcal{F} be the set of L -colorings of v_1, \dots, v_{i+1} that are nonrepetitive on v_1, \dots, v_i but have a square with v_{i+1} . Clearly, $|\mathcal{C}_{i+1}| = 4|\mathcal{C}_i| - |\mathcal{F}|$. Let \mathcal{F}_j be subset of \mathcal{F} with a square of length $2j$. So $\mathcal{F} = \cup_{j \geq 1} \mathcal{F}_j$.

Each $\varphi \in \mathcal{F}_j$ restricts to a nonrepetitive L -coloring φ' of v_1, \dots, v_{i+1-j} . And φ' uniquely determines φ . So $|\mathcal{F}_j| \leq |\mathcal{C}_{i+1-j}|$.

Nonrepetitive List-coloring of Paths (the proof)

Lem: Let L be a 4-assignment to $V(P_n)$. For each $i \geq 1$, let \mathcal{C}_i be set of nonrepetitive L -colorings of first i verts of P_n . For all $i < n$,

$$|\mathcal{C}_{i+1}| \geq 2|\mathcal{C}_i|.$$

Pf: Induction on i . Let \mathcal{F} be the set of L -colorings of v_1, \dots, v_{i+1} that are nonrepetitive on v_1, \dots, v_i but have a square with v_{i+1} . Clearly, $|\mathcal{C}_{i+1}| = 4|\mathcal{C}_i| - |\mathcal{F}|$. Let \mathcal{F}_j be subset of \mathcal{F} with a square of length $2j$. So $\mathcal{F} = \cup_{j \geq 1} \mathcal{F}_j$.

Each $\varphi \in \mathcal{F}_j$ restricts to a nonrepetitive L -coloring φ' of v_1, \dots, v_{i+1-j} . And φ' uniquely determines φ . So $|\mathcal{F}_j| \leq |\mathcal{C}_{i+1-j}|$. By induction, $|\mathcal{C}_{i+1-j}| \leq 2^{-j+1}|\mathcal{C}_i|$, for each $j \geq 1$.

Nonrepetitive List-coloring of Paths (the proof)

Lem: Let L be a 4-assignment to $V(P_n)$. For each $i \geq 1$, let \mathcal{C}_i be set of nonrepetitive L -colorings of first i verts of P_n . For all $i < n$,

$$|\mathcal{C}_{i+1}| \geq 2|\mathcal{C}_i|.$$

Pf: Induction on i . Let \mathcal{F} be the set of L -colorings of v_1, \dots, v_{i+1} that are nonrepetitive on v_1, \dots, v_i but have a square with v_{i+1} . Clearly, $|\mathcal{C}_{i+1}| = 4|\mathcal{C}_i| - |\mathcal{F}|$. Let \mathcal{F}_j be subset of \mathcal{F} with a square of length $2j$. So $\mathcal{F} = \cup_{j \geq 1} \mathcal{F}_j$.

Each $\varphi \in \mathcal{F}_j$ restricts to a nonrepetitive L -coloring φ' of v_1, \dots, v_{i+1-j} . And φ' uniquely determines φ . So $|\mathcal{F}_j| \leq |\mathcal{C}_{i+1-j}|$. By induction, $|\mathcal{C}_{i+1-j}| \leq 2^{-j+1}|\mathcal{C}_i|$, for each $j \geq 1$. Thus,

$$|\mathcal{C}_{i+1}| = 4|\mathcal{C}_i| - |\mathcal{F}|$$

Nonrepetitive List-coloring of Paths (the proof)

Lem: Let L be a 4-assignment to $V(P_n)$. For each $i \geq 1$, let \mathcal{C}_i be set of nonrepetitive L -colorings of first i verts of P_n . For all $i < n$,

$$|\mathcal{C}_{i+1}| \geq 2|\mathcal{C}_i|.$$

Pf: Induction on i . Let \mathcal{F} be the set of L -colorings of v_1, \dots, v_{i+1} that are nonrepetitive on v_1, \dots, v_i but have a square with v_{i+1} . Clearly, $|\mathcal{C}_{i+1}| = 4|\mathcal{C}_i| - |\mathcal{F}|$. Let \mathcal{F}_j be subset of \mathcal{F} with a square of length $2j$. So $\mathcal{F} = \cup_{j \geq 1} \mathcal{F}_j$.

Each $\varphi \in \mathcal{F}_j$ restricts to a nonrepetitive L -coloring φ' of v_1, \dots, v_{i+1-j} . And φ' uniquely determines φ . So $|\mathcal{F}_j| \leq |\mathcal{C}_{i+1-j}|$. By induction, $|\mathcal{C}_{i+1-j}| \leq 2^{-j+1}|\mathcal{C}_i|$, for each $j \geq 1$. Thus,

$$|\mathcal{C}_{i+1}| = 4|\mathcal{C}_i| - |\mathcal{F}| \geq 4|\mathcal{C}_i| - \sum |\mathcal{F}_j|$$

Nonrepetitive List-coloring of Paths (the proof)

Lem: Let L be a 4-assignment to $V(P_n)$. For each $i \geq 1$, let \mathcal{C}_i be set of nonrepetitive L -colorings of first i verts of P_n . For all $i < n$,

$$|\mathcal{C}_{i+1}| \geq 2|\mathcal{C}_i|.$$

Pf: Induction on i . Let \mathcal{F} be the set of L -colorings of v_1, \dots, v_{i+1} that are nonrepetitive on v_1, \dots, v_i but have a square with v_{i+1} . Clearly, $|\mathcal{C}_{i+1}| = 4|\mathcal{C}_i| - |\mathcal{F}|$. Let \mathcal{F}_j be subset of \mathcal{F} with a square of length $2j$. So $\mathcal{F} = \cup_{j \geq 1} \mathcal{F}_j$.

Each $\varphi \in \mathcal{F}_j$ restricts to a nonrepetitive L -coloring φ' of v_1, \dots, v_{i+1-j} . And φ' uniquely determines φ . So $|\mathcal{F}_j| \leq |\mathcal{C}_{i+1-j}|$. By induction, $|\mathcal{C}_{i+1-j}| \leq 2^{-j+1}|\mathcal{C}_i|$, for each $j \geq 1$. Thus,

$$\begin{aligned} |\mathcal{C}_{i+1}| &= 4|\mathcal{C}_i| - |\mathcal{F}| \geq 4|\mathcal{C}_i| - \sum |\mathcal{F}_j| \\ &\geq 4|\mathcal{C}_i| - \sum |\mathcal{C}_{i+1-j}| \end{aligned}$$

Nonrepetitive List-coloring of Paths (the proof)

Lem: Let L be a 4-assignment to $V(P_n)$. For each $i \geq 1$, let \mathcal{C}_i be set of nonrepetitive L -colorings of first i verts of P_n . For all $i < n$,

$$|\mathcal{C}_{i+1}| \geq 2|\mathcal{C}_i|.$$

Pf: Induction on i . Let \mathcal{F} be the set of L -colorings of v_1, \dots, v_{i+1} that are nonrepetitive on v_1, \dots, v_i but have a square with v_{i+1} . Clearly, $|\mathcal{C}_{i+1}| = 4|\mathcal{C}_i| - |\mathcal{F}|$. Let \mathcal{F}_j be subset of \mathcal{F} with a square of length $2j$. So $\mathcal{F} = \cup_{j \geq 1} \mathcal{F}_j$.

Each $\varphi \in \mathcal{F}_j$ restricts to a nonrepetitive L -coloring φ' of v_1, \dots, v_{i+1-j} . And φ' uniquely determines φ . So $|\mathcal{F}_j| \leq |\mathcal{C}_{i+1-j}|$. By induction, $|\mathcal{C}_{i+1-j}| \leq 2^{-j+1}|\mathcal{C}_i|$, for each $j \geq 1$. Thus,

$$\begin{aligned} |\mathcal{C}_{i+1}| &= 4|\mathcal{C}_i| - |\mathcal{F}| \geq 4|\mathcal{C}_i| - \sum |\mathcal{F}_j| \\ &\geq 4|\mathcal{C}_i| - \sum |\mathcal{C}_{i+1-j}| \\ &\geq 4|\mathcal{C}_i| - \sum 2^{-j+1}|\mathcal{C}_i| \end{aligned}$$

Nonrepetitive List-coloring of Paths (the proof)

Lem: Let L be a 4-assignment to $V(P_n)$. For each $i \geq 1$, let \mathcal{C}_i be set of nonrepetitive L -colorings of first i verts of P_n . For all $i < n$,

$$|\mathcal{C}_{i+1}| \geq 2|\mathcal{C}_i|.$$


Pf: Induction on i . Let \mathcal{F} be the set of L -colorings of v_1, \dots, v_{i+1} that are nonrepetitive on v_1, \dots, v_i but have a square with v_{i+1} . Clearly, $|\mathcal{C}_{i+1}| = 4|\mathcal{C}_i| - |\mathcal{F}|$. Let \mathcal{F}_j be subset of \mathcal{F} with a square of length $2j$. So $\mathcal{F} = \cup_{j \geq 1} \mathcal{F}_j$.

Each $\varphi \in \mathcal{F}_j$ restricts to a nonrepetitive L -coloring φ' of v_1, \dots, v_{i+1-j} . And φ' uniquely determines φ . So $|\mathcal{F}_j| \leq |\mathcal{C}_{i+1-j}|$. By induction, $|\mathcal{C}_{i+1-j}| \leq 2^{-j+1}|\mathcal{C}_i|$, for each $j \geq 1$. Thus,

$$\begin{aligned} |\mathcal{C}_{i+1}| &= 4|\mathcal{C}_i| - |\mathcal{F}| \geq 4|\mathcal{C}_i| - \sum |\mathcal{F}_j| \\ &\geq 4|\mathcal{C}_i| - \sum |\mathcal{C}_{i+1-j}| \\ &\geq 4|\mathcal{C}_i| - \sum 2^{-j+1}|\mathcal{C}_i| \geq 2|\mathcal{C}_i|. \end{aligned}$$


Proper Conflict-free Coloring

Defn: A proper coloring of G is **conflict-free** if every non-isolated vertex of G has some color appearing exactly once on its open neighborhood.



Proper Conflict-free Coloring


Defn: A proper coloring of G is **conflict-free** if every non-isolated vertex of G has some color appearing exactly once on its open neighborhood.



Conj: [CPS] $\chi_{pcf}(G) \leq \Delta + 1$ for all connected G with $\Delta \geq 3$.

Proper Conflict-free Coloring

Defn: A proper coloring of G is **conflict-free** if every non-isolated vertex of G has some color appearing exactly once on its open neighborhood.




Conj: [CPS] $\chi_{pcf}(G) \leq \Delta + 1$ for all connected G with $\Delta \geq 3$.

Thm: Fix a positive integer $\Delta \geq 6.5 \cdot 10^7$, fix a real number β with $\Delta \geq \beta \geq 0.6550826\Delta$, and let $a := \left\lceil \Delta + \beta + \sqrt{\Delta} \right\rceil$.

Proper Conflict-free Coloring

Defn: A proper coloring of G is **conflict-free** if every non-isolated vertex of G has some color appearing exactly once on its open neighborhood.




Conj: [CPS] $\chi_{pcf}(G) \leq \Delta + 1$ for all connected G with $\Delta \geq 3$.

Thm: Fix a positive integer $\Delta \geq 6.5 \cdot 10^7$, fix a real number β with $\Delta \geq \beta \geq 0.6550826\Delta$, and let $a := \left\lceil \Delta + \beta + \sqrt{\Delta} \right\rceil$. If G has max degree at most Δ and L is an a -assignment for G , then there are at least $\beta^{|V(G)|}$ proper conflict-free L -colorings of G .

Proper Conflict-free Coloring

Defn: A proper coloring of G is **conflict-free** if every non-isolated vertex of G has some color appearing exactly once on its open neighborhood.




Conj: [CPS] $\chi_{pcf}(G) \leq \Delta + 1$ for all connected G with $\Delta \geq 3$.

Thm: Fix a positive integer $\Delta \geq 6.5 \cdot 10^7$, fix a real number β with $\Delta \geq \beta \geq 0.6550826\Delta$, and let $a := \left\lceil \Delta + \beta + \sqrt{\Delta} \right\rceil$. If G has max degree at most Δ and L is an a -assignment for G , then there are at least $\beta^{|V(G)|}$ proper conflict-free L -colorings of G . Analogous statements hold when $\Delta \geq 4000$ and $\Delta \geq \beta \geq 0.6\Delta$ and when $\Delta \geq 750$ and $\Delta \geq \beta \geq 0.8\Delta$.

Proper Conflict-free Coloring

Defn: A proper coloring of G is **conflict-free** if every non-isolated vertex of G has some color appearing exactly once on its open neighborhood.




Conj: [CPS] $\chi_{pcf}(G) \leq \Delta + 1$ for all connected G with $\Delta \geq 3$.

Thm: Fix a positive integer $\Delta \geq 6.5 \cdot 10^7$, fix a real number β with $\Delta \geq \beta \geq 0.6550826\Delta$, and let $a := \left\lceil \Delta + \beta + \sqrt{\Delta} \right\rceil$. If G has max degree at most Δ and L is an a -assignment for G , then there are at least $\beta^{|V(G)|}$ proper conflict-free L -colorings of G . Analogous statements hold when $\Delta \geq 4000$ and $\Delta \geq \beta \geq 0.6\Delta$ and when $\Delta \geq 750$ and $\Delta \geq \beta \geq 0.8\Delta$.

Cor: So $\chi_{pcf}^\ell(G) \leq 1.6551\Delta(1 + o(1))$.

Proper Conflict-free Coloring

Defn: A proper coloring of G is **conflict-free** if every non-isolated vertex of G has some color appearing exactly once on its open neighborhood.



Conj: [CPS] $\chi_{pcf}(G) \leq \Delta + 1$ for all connected G with $\Delta \geq 3$.


Thm: Fix a positive integer $\Delta \geq 6.5 \cdot 10^7$, fix a real number β with $\Delta \geq \beta \geq 0.6550826\Delta$, and let $a := \left\lceil \Delta + \beta + \sqrt{\Delta} \right\rceil$. If G has max degree at most Δ and L is an a -assignment for G , then there are at least $\beta^{|V(G)|}$ proper conflict-free L -colorings of G . Analogous statements hold when $\Delta \geq 4000$ and $\Delta \geq \beta \geq 0.6\Delta$ and when $\Delta \geq 750$ and $\Delta \geq \beta \geq 0.8\Delta$.

Cor: So $\chi_{pcf}^\ell(G) \leq 1.6551\Delta(1 + o(1))$.

Rem: Liu and Reed showed that $\chi_{pcf}(G) \leq \Delta(1 + o(1))$.

Proper Conflict-free Coloring

Defn: A proper coloring of G is **conflict-free** if every non-isolated vertex of G has some color appearing exactly once on its open neighborhood.



Conj: [CPS] $\chi_{pcf}(G) \leq \Delta + 1$ for all connected G with $\Delta \geq 3$.

Thm: Fix a positive integer $\Delta \geq 6.5 \cdot 10^7$, fix a real number β with $\Delta \geq \beta \geq 0.6550826\Delta$, and let $a := \left\lceil \Delta + \beta + \sqrt{\Delta} \right\rceil$. If G has max degree at most Δ and L is an a -assignment for G , then there are at least $\beta^{|V(G)|}$ proper conflict-free L -colorings of G . Analogous statements hold when $\Delta \geq 4000$ and $\Delta \geq \beta \geq 0.6\Delta$ and when $\Delta \geq 750$ and $\Delta \geq \beta \geq 0.8\Delta$.

Cor: So $\chi_{pcf}^\ell(G) \leq 1.6551\Delta(1 + o(1))$.

Rem: Liu and Reed showed that $\chi_{pcf}(G) \leq \Delta(1 + o(1))$. This bound is stronger than ours, but much less general.

Key Rosenfeld Counting Lemma

Defn: For an integer t , a graph G , and a hypergraph \mathcal{H} with $V(\mathcal{H}) = V(G)$, a coloring φ is a **proper t -conflict-free coloring** of (G, \mathcal{H}) if φ is a proper coloring of G such that for every $f \in E(\mathcal{H})$, some color is used k times by φ on f for some $k \in \{1, \dots, t\}$.

Key Rosenfeld Counting Lemma

Defn: For an integer t , a graph G , and a hypergraph \mathcal{H} with $V(\mathcal{H}) = V(G)$, a coloring φ is a **proper t -conflict-free coloring** of (G, \mathcal{H}) if φ is a proper coloring of G such that for every $f \in E(\mathcal{H})$, some color is used k times by φ on f for some $k \in \{1, \dots, t\}$.

Defn: Fix $t, i, d \in \mathbb{Z}^+$.

Key Rosenfeld Counting Lemma

Defn: For an integer t , a graph G , and a hypergraph \mathcal{H} with $V(\mathcal{H}) = V(G)$, a coloring φ is a **proper t -conflict-free coloring** of (G, \mathcal{H}) if φ is a proper coloring of G such that for every $f \in E(\mathcal{H})$, some color is used k times by φ on f for some $k \in \{1, \dots, t\}$.

Defn: Fix $t, i, d \in \mathbb{Z}^+$. The **t -associated Stirling number of second kind**, $S_t(d, i)$, is the number of partitions of the set $\{1, \dots, d\}$ into i parts, each of size at least t .

Key Rosenfeld Counting Lemma

Defn: For an integer t , a graph G , and a hypergraph \mathcal{H} with $V(\mathcal{H}) = V(G)$, a coloring φ is a **proper t -conflict-free coloring** of (G, \mathcal{H}) if φ is a proper coloring of G such that for every $f \in E(\mathcal{H})$, some color is used k times by φ on f for some $k \in \{1, \dots, t\}$.

Defn: Fix $t, i, d \in \mathbb{Z}^+$. The **t -associated Stirling number of second kind**, $S_t(d, i)$, is the number of partitions of the set $\{1, \dots, d\}$ into i parts, each of size at least t . E.g. $S_2(2i, i) = (2i)!/(i!2^i)$.

Key Rosenfeld Counting Lemma

Defn: For an integer t , a graph G , and a hypergraph \mathcal{H} with $V(\mathcal{H}) = V(G)$, a coloring φ is a **proper t -conflict-free coloring** of (G, \mathcal{H}) if φ is a proper coloring of G such that for every $f \in E(\mathcal{H})$, some color is used k times by φ on f for some $k \in \{1, \dots, t\}$.

Defn: Fix $t, i, d \in \mathbb{Z}^+$. The **t -associated Stirling number of second kind**, $S_t(d, i)$, is the number of partitions of the set $\{1, \dots, d\}$ into i parts, each of size at least t . E.g. $S_2(2i, i) = (2i)!/(i!2^i)$.

Key Lem: Fix G, \mathcal{H}, t as above. Let β be a real number. If a is a real number such that

$$a \geq \Delta(G) + \beta + \sum_{f \in E(\mathcal{H}), f \ni v} \sum_{i=1}^{\lfloor |f|/(t+1) \rfloor} S_{t+1}(|f|, i) \cdot \beta^{i-|f|+1}$$

for every $v \in V(G)$, then for every a -assignment L of G , there are at least $\beta^{|V(G)|}$ proper t -conflict-free L -colorings of (G, \mathcal{H}) .

Bounding $S_2(d, i)$

Bounding $S_2(d, i)$

Helper Lem: Fix $i, d \in \mathbb{Z}^+$ with $d \geq 110$. If $0.3d \leq i \leq d/2$, then

$$S_2(d, i) \leq 8i(0.6251d)^{d-i}.$$

Bounding $S_2(d, i)$

Helper Lem: Fix $i, d \in \mathbb{Z}^+$ with $d \geq 110$. If $0.3d \leq i \leq d/2$, then

$$S_2(d, i) \leq 8i(0.6251d)^{d-i}.$$

Lem: Fix $d, R \in \mathbb{Z}^+$ with $110 \leq d \leq R$. If $\epsilon, c, \beta \in \mathbb{R}^+$ s.t.
 $0.6251 \leq \epsilon < 1$, $0.3 \leq c < \epsilon/2$, $\epsilon R \leq \beta \leq R$, and $d \geq f(c, \epsilon, R)$,
then

$$\sum_{i=1}^{d/2} S_2(d, i) \beta^{i-d+1} \leq R^{-1/2}.$$

Bounding $S_2(d, i)$

Helper Lem: Fix $i, d \in \mathbb{Z}^+$ with $d \geq 110$. If $0.3d \leq i \leq d/2$, then

$$S_2(d, i) \leq 8i(0.6251d)^{d-i}.$$

Lem: Fix $d, R \in \mathbb{Z}^+$ with $110 \leq d \leq R$. If $\epsilon, c, \beta \in \mathbb{R}^+$ s.t.
 $0.6251 \leq \epsilon < 1$, $0.3 \leq c < \epsilon/2$, $\epsilon R \leq \beta \leq R$, and $d \geq f(c, \epsilon, R)$,
then

$$\sum_{i=1}^{d/2} S_2(d, i) \beta^{i-d+1} \leq R^{-1/2}.$$

Pf Sketch:

Bounding $S_2(d, i)$

Helper Lem: Fix $i, d \in \mathbb{Z}^+$ with $d \geq 110$. If $0.3d \leq i \leq d/2$, then

$$S_2(d, i) \leq 8i(0.6251d)^{d-i}.$$

Lem: Fix $d, R \in \mathbb{Z}^+$ with $110 \leq d \leq R$. If $\epsilon, c, \beta \in \mathbb{R}^+$ s.t.
 $0.6251 \leq \epsilon < 1$, $0.3 \leq c < \epsilon/2$, $\epsilon R \leq \beta \leq R$, and $d \geq f(c, \epsilon, R)$,
then

$$\sum_{i=1}^{d/2} S_2(d, i) \beta^{i-d+1} \leq R^{-1/2}.$$

Pf Sketch:

$$\sum_{i=1}^{cd} S_2(d, i) \beta^{i-d+1} \leq \sum_{i=1}^{cd} \binom{d}{i} i^{d-i} 2^{-i} \beta^{i-d+1}$$

Bounding $S_2(d, i)$

Helper Lem: Fix $i, d \in \mathbb{Z}^+$ with $d \geq 110$. If $0.3d \leq i \leq d/2$, then

$$S_2(d, i) \leq 8i(0.6251d)^{d-i}.$$

Lem: Fix $d, R \in \mathbb{Z}^+$ with $110 \leq d \leq R$. If $\epsilon, c, \beta \in \mathbb{R}^+$ s.t.
 $0.6251 \leq \epsilon < 1$, $0.3 \leq c < \epsilon/2$, $\epsilon R \leq \beta \leq R$, and $d \geq f(c, \epsilon, R)$,
then

$$\sum_{i=1}^{d/2} S_2(d, i) \beta^{i-d+1} \leq R^{-1/2}.$$

Pf Sketch:

$$\sum_{i=1}^{cd} S_2(d, i) \beta^{i-d+1} \leq \sum_{i=1}^{cd} \binom{d}{i} i^{d-i} 2^{-i} \beta^{i-d+1} \leq \dots \leq \frac{1}{2} R^{-1/2}$$

Bounding $S_2(d, i)$

Helper Lem: Fix $i, d \in \mathbb{Z}^+$ with $d \geq 110$. If $0.3d \leq i \leq d/2$, then

$$S_2(d, i) \leq 8i(0.6251d)^{d-i}.$$

Lem: Fix $d, R \in \mathbb{Z}^+$ with $110 \leq d \leq R$. If $\epsilon, c, \beta \in \mathbb{R}^+$ s.t. $0.6251 \leq \epsilon < 1$, $0.3 \leq c < \epsilon/2$, $\epsilon R \leq \beta \leq R$, and $d \geq f(c, \epsilon, R)$, then

$$\sum_{i=1}^{d/2} S_2(d, i) \beta^{i-d+1} \leq R^{-1/2}.$$

Pf Sketch:

$$\begin{aligned} \sum_{i=1}^{cd} S_2(d, i) \beta^{i-d+1} &\leq \sum_{i=1}^{cd} \binom{d}{i} i^{d-i} 2^{-i} \beta^{i-d+1} \leq \dots \leq \frac{1}{2} R^{-1/2} \\ \sum_{i=cd}^{d/2} S_2(d, i) \beta^{i-d+1} &\leq \sum_{i=cd}^{d/2} 8i(0.6251d)^{d-i} \beta^{i-d+1} \end{aligned}$$

Bounding $S_2(d, i)$

Helper Lem: Fix $i, d \in \mathbb{Z}^+$ with $d \geq 110$. If $0.3d \leq i \leq d/2$, then

$$S_2(d, i) \leq 8i(0.6251d)^{d-i}.$$

Lem: Fix $d, R \in \mathbb{Z}^+$ with $110 \leq d \leq R$. If $\epsilon, c, \beta \in \mathbb{R}^+$ s.t. $0.6251 \leq \epsilon < 1$, $0.3 \leq c < \epsilon/2$, $\epsilon R \leq \beta \leq R$, and $d \geq f(c, \epsilon, R)$, then

$$\sum_{i=1}^{d/2} S_2(d, i) \beta^{i-d+1} \leq R^{-1/2}.$$

Pf Sketch:


$$\begin{aligned} \sum_{i=1}^{cd} S_2(d, i) \beta^{i-d+1} &\leq \sum_{i=1}^{cd} \binom{d}{i} i^{d-i} 2^{-i} \beta^{i-d+1} \leq \dots \leq \frac{1}{2} R^{-1/2} \\ \sum_{i=cd}^{d/2} S_2(d, i) \beta^{i-d+1} &\leq \sum_{i=cd}^{d/2} 8i(0.6251d)^{d-i} \beta^{i-d+1} \leq \dots \leq \frac{1}{2} R^{-1/2} \end{aligned}$$

Recap

- ▶ Rosenfeld Counting


Recap

- ▶ Rosenfeld Counting
 - ▶ Nonrepetitive 4-list-coloring of paths




Recap

- ▶ Rosenfeld Counting
 - ▶ Nonrepetitive 4-list-coloring of paths
 - ▶ Color iteratively




Recap

- ▶ Rosenfeld Counting
 - ▶ Nonrepetitive 4-list-coloring of paths
 - ▶ Color iteratively
 - ▶ # partial colorings exponential in $|\text{subgraph}|$




Recap

- ▶ Rosenfeld Counting
 - ▶ Nonrepetitive 4-list-coloring of paths
 - ▶ Color iteratively
 - ▶ # partial colorings exponential in $|\text{subgraph}|$
 - ▶ Bad colorings \rightarrow good colorings of subgraph





Recap

- ▶ Rosenfeld Counting
 - ▶ Nonrepetitive 4-list-coloring of paths
 - ▶ Color iteratively
 - ▶ # partial colorings exponential in $|\text{subgraph}|$
 - ▶ Bad colorings \rightarrow good colorings of subgraph
 - ▶ Exponentially many good colorings remain





Recap

- ▶ Rosenfeld Counting
 - ▶ Nonrepetitive 4-list-coloring of paths
 - ▶ Color iteratively
 - ▶ # partial colorings exponential in $|\text{subgraph}|$
 - ▶ Bad colorings \rightarrow good colorings of subgraph
 - ▶ Exponentially many good colorings remain
- ▶ Proper Conflict-free Coloring




Recap

- ▶ Rosenfeld Counting
 - ▶ Nonrepetitive 4-list-coloring of paths
 - ▶ Color iteratively
 - ▶ # partial colorings exponential in |subgraph|
 - ▶ Bad colorings \rightarrow good colorings of subgraph
 - ▶ Exponentially many good colorings remain
- ▶ Proper Conflict-free Coloring
 - ▶ CPS conjectured $\chi_{pcf}(G) \leq \Delta + 1$




Recap

- ▶ Rosenfeld Counting
 - ▶ Nonrepetitive 4-list-coloring of paths
 - ▶ Color iteratively
 - ▶ # partial colorings exponential in |subgraph|
 - ▶ Bad colorings \rightarrow good colorings of subgraph
 - ▶ Exponentially many good colorings remain




- ▶ Proper Conflict-free Coloring
 - ▶ CPS conjectured $\chi_{pcf}(G) \leq \Delta + 1$
 - ▶ We proved $\chi_{pcf}^\ell(G) \leq 1.6551\Delta(1 + o(1))$



Recap


▶ Rosenfeld Counting

- ▶ Nonrepetitive 4-list-coloring of paths
- ▶ Color iteratively
- ▶ # partial colorings exponential in |subgraph|
 - ▶ Bad colorings \rightarrow good colorings of subgraph
 - ▶ Exponentially many good colorings remain



▶ Proper Conflict-free Coloring


- ▶ CPS conjectured $\chi_{pcf}(G) \leq \Delta + 1$
- ▶ We proved $\chi_{pcf}^\ell(G) \leq 1.6551\Delta(1 + o(1))$
- ▶ Corollary of general hypergraph framework



Recap


▶ Rosenfeld Counting

- ▶ Nonrepetitive 4-list-coloring of paths
- ▶ Color iteratively
- ▶ # partial colorings exponential in |subgraph|
 - ▶ Bad colorings \rightarrow good colorings of subgraph
 - ▶ Exponentially many good colorings remain



▶ Proper Conflict-free Coloring


- ▶ CPS conjectured $\chi_{pcf}(G) \leq \Delta + 1$
- ▶ We proved $\chi_{pcf}^\ell(G) \leq 1.6551\Delta(1 + o(1))$
- ▶ Corollary of general hypergraph framework
- ▶ Key step is bounding $S_2(d, i)$



Learn More about Rosenfeld Counting

Graph Coloring Methods


Daniel W. Cranston



Learn More about Rosenfeld Counting

Graph Coloring Methods


Daniel W. Cranston




<https://graphcoloringmethods.com>

Recap

- ▶ Rosenfeld Counting
 - ▶ Nonrepetitive 4-list-coloring of paths
 - ▶ Color iteratively
 - ▶ # partial colorings exponential in |subgraph|
 - ▶ Bad colorings \rightarrow good colorings of subgraph
 - ▶ Exponentially many good colorings remain



- ▶ Proper Conflict-free Coloring
 - ▶ CPS conjectured $\chi_{pcf}(G) \leq \Delta + 1$
 - ▶ We proved $\chi_{pcf}^{\ell}(G) \leq 1.6551\Delta(1 + o(1))$
 - ▶ Corollary of General hypergraph framework
 - ▶ Key step is bounding $S_2(d, i)$




- ▶ Graph Coloring Methods




Recap

- ▶ Rosenfeld Counting
 - ▶ Nonrepetitive 4-list-coloring of paths
 - ▶ Color iteratively
 - ▶ # partial colorings exponential in |subgraph|
 - ▶ Bad colorings \rightarrow good colorings of subgraph
 - ▶ Exponentially many good colorings remain



- ▶ Proper Conflict-free Coloring
 - ▶ CPS conjectured $\chi_{pcf}(G) \leq \Delta + 1$
 - ▶ We proved $\chi_{pcf}^\ell(G) \leq 1.6551\Delta(1 + o(1))$
 - ▶ Corollary of General hypergraph framework
 - ▶ Key step is bounding $S_2(d, i)$




- ▶ Graph Coloring Methods
 - ▶ Graduate Textbook (450 pages)




Recap

- ▶ Rosenfeld Counting
 - ▶ Nonrepetitive 4-list-coloring of paths
 - ▶ Color iteratively
 - ▶ # partial colorings exponential in |subgraph|
 - ▶ Bad colorings \rightarrow good colorings of subgraph
 - ▶ Exponentially many good colorings remain



- ▶ Proper Conflict-free Coloring
 - ▶ CPS conjectured $\chi_{pcf}(G) \leq \Delta + 1$
 - ▶ We proved $\chi_{pcf}^{\ell}(G) \leq 1.6551\Delta(1 + o(1))$
 - ▶ Corollary of General hypergraph framework
 - ▶ Key step is bounding $S_2(d, i)$




- ▶ Graph Coloring Methods
 - ▶ Graduate Textbook (450 pages)
 - ▶ Chapter on Rosenfeld Counting




Recap

- ▶ Rosenfeld Counting
 - ▶ Nonrepetitive 4-list-coloring of paths
 - ▶ Color iteratively
 - ▶ # partial colorings exponential in |subgraph|
 - ▶ Bad colorings \rightarrow good colorings of subgraph
 - ▶ Exponentially many good colorings remain



- ▶ Proper Conflict-free Coloring
 - ▶ CPS conjectured $\chi_{pcf}(G) \leq \Delta + 1$
 - ▶ We proved $\chi_{pcf}^{\ell}(G) \leq 1.6551\Delta(1 + o(1))$
 - ▶ Corollary of General hypergraph framework
 - ▶ Key step is bounding $S_2(d, i)$




- ▶ Graph Coloring Methods
 - ▶ Graduate Textbook (450 pages)
 - ▶ Chapter on Rosenfeld Counting
 - ▶ 11 other chapters (each on 1 method)




Recap

- ▶ Rosenfeld Counting
 - ▶ Nonrepetitive 4-list-coloring of paths
 - ▶ Color iteratively
 - ▶ # partial colorings exponential in |subgraph|
 - ▶ Bad colorings \rightarrow good colorings of subgraph
 - ▶ Exponentially many good colorings remain



- ▶ Proper Conflict-free Coloring
 - ▶ CPS conjectured $\chi_{pcf}(G) \leq \Delta + 1$
 - ▶ We proved $\chi_{pcf}^\ell(G) \leq 1.6551\Delta(1 + o(1))$
 - ▶ Corollary of General hypergraph framework
 - ▶ Key step is bounding $S_2(d, i)$



- ▶ Graph Coloring Methods
 - ▶ Graduate Textbook (450 pages)
 - ▶ Chapter on Rosenfeld Counting
 - ▶ 11 other chapters (each on 1 method)
 - ▶ Free at graphcoloringmethods.com

