



COMPUTATIONAL FINANCE & RISK MANAGEMENT

UNIVERSITY *of* WASHINGTON

Department of Applied Mathematics

CFRM 410: Probability and Statistics for Computational Finance

Week 7 Multivariate Random Variables

Jake Price

Instructor, Computational Finance and Risk Management

University of Washington

Slides originally produced by Kjell Konis

Outline

Outline

Multivariate Random Variables

An *n-dimensional random vector* is a function from the sample space S into \mathbb{R}^n (*n*-dimensional Euclidean space)

Multivariate Random Variables

An *n-dimensional random vector* is a function from the sample space S into \mathbb{R}^n (*n-dimensional Euclidean space*)

Example

- ▶ Consider the random experiment of rolling two 6-sided dice

Multivariate Random Variables

An *n-dimensional random vector* is a function from the sample space S into \mathbb{R}^n (*n-dimensional Euclidean space*)

Example

- ▶ Consider the random experiment of rolling two 6-sided dice
- ▶ For each of the 36 possible elementary outcomes in S let

$$X = \text{sum of the two dice} \quad \text{and} \quad Y = |\text{difference of the two dice}|$$

Multivariate Random Variables

An n -dimensional random vector is a function from the sample space S into \mathbb{R}^n (n -dimensional Euclidean space)

Example

- ▶ Consider the random experiment of rolling two 6-sided dice
- ▶ For each of the 36 possible elementary outcomes in S let

$$X = \text{sum of the two dice} \quad \text{and} \quad Y = |\text{difference of the two dice}|$$

- ▶ The vector (X, Y) is a *bivariate random vector*

Joint Distribution

Let X and Y be two random variables with supports S_X and S_Y

Joint Distribution

Let X and Y be two random variables with supports S_X and S_Y

The vector (X, Y) is a bivariate random vector with support S_{XY}

Joint Distribution

Let X and Y be two random variables with supports S_X and S_Y

The vector (X, Y) is a bivariate random vector with support S_{XY}

The function

$$F_{X,Y}(x, y) = P(X \leq x, Y \leq y)$$

is called the *joint distribution function* of X and Y

Joint Distribution

Let X and Y be two random variables with supports S_X and S_Y

The vector (X, Y) is a bivariate random vector with support S_{XY}

The function

$$F_{X,Y}(x, y) = P(X \leq x, Y \leq y)$$

is called the *joint distribution function* of X and Y

Discrete Case: The joint probability of X and Y is described by the *joint mass function*

$$f_{X,Y}(x_i, y_j) = P(X = x_i, Y = y_j)$$

for all possible pairs $(x_i, y_j) \in S_{XY}$

Joint Distribution

Let X and Y be two random variables with supports S_X and S_Y

The vector (X, Y) is a bivariate random vector with support S_{XY}

The function

$$F_{X,Y}(x, y) = P(X \leq x, Y \leq y)$$

is called the *joint distribution function* of X and Y

Discrete Case: The joint probability of X and Y is described by the *joint mass function*

$$f_{X,Y}(x_i, y_j) = P(X = x_i, Y = y_j)$$

for all possible pairs $(x_i, y_j) \in S_{XY}$

Continuous Case: The joint density of X and Y is described by the *joint density function*

$$f_{X,Y}(x, y)$$

Joint Density

The joint distribution function is

$$P(X \leq x, Y \leq y) = F_{X,Y}(x, y) = \int_{-\infty}^y \int_{-\infty}^x f_{X,Y}(u, v) du dv$$

Joint Density

The joint distribution function is

$$P(X \leq x, Y \leq y) = F_{X,Y}(x, y) = \int_{-\infty}^y \int_{-\infty}^x f_{X,Y}(u, v) \, du \, dv$$

Properties:

- ▶ $f_{X,Y}(x, y) \geq 0$ for all points $(x, y) \in \mathbb{R}^2$

Joint Density

The joint distribution function is

$$P(X \leq x, Y \leq y) = F_{X,Y}(x, y) = \int_{-\infty}^y \int_{-\infty}^x f_{X,Y}(u, v) \, du \, dv$$

Properties:

- ▶ $f_{X,Y}(x, y) \geq 0$ for all points $(x, y) \in \mathbb{R}^2$
- ▶ $\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_{X,Y}(x, y) \, dx \, dy = 1$

Joint Density

The joint distribution function is

$$P(X \leq x, Y \leq y) = F_{X,Y}(x, y) = \int_{-\infty}^y \int_{-\infty}^x f_{X,Y}(u, v) du dv$$

Properties:

- ▶ $f_{X,Y}(x, y) \geq 0$ for all points $(x, y) \in \mathbb{R}^2$
- ▶ $\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_{X,Y}(x, y) dx dy = 1$
- ▶ $f_{X,Y}(x, y) = \frac{\partial^2}{\partial x \partial y} F_{X,Y}(x, y)$

Joint Density

The joint distribution function is

$$P(X \leq x, Y \leq y) = F_{X,Y}(x, y) = \int_{-\infty}^y \int_{-\infty}^x f_{X,Y}(u, v) du dv$$

Properties:

- ▶ $f_{X,Y}(x, y) \geq 0$ for all points $(x, y) \in \mathbb{R}^2$
- ▶ $\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_{X,Y}(x, y) dx dy = 1$
- ▶ $f_{X,Y}(x, y) = \frac{\partial^2}{\partial x \partial y} F_{X,Y}(x, y)$
- ▶ $P(a_1 \leq X \leq b_1, a_2 \leq Y \leq b_2) = \int_{a_2}^{b_2} \int_{a_1}^{b_1} f_{X,Y}(x, y) dx dy$

Marginal Distribution

Let (X, Y) be a bivariate random vector with joint density (mass) function $f_{X,Y}(x, y)$, then the function

Discrete:
$$f_X(x_i) = \sum_{y_j \in S_Y} f_{X,Y}(x_i, y_j)$$

Continuous:
$$f_X(x) = \int_{-\infty}^{\infty} f_{X,Y}(x, y) dy$$

is called the *marginal density (mass) functions* of X

Marginal Distribution

Let (X, Y) be a bivariate random vector with joint density (mass) function $f_{X,Y}(x, y)$, then the function

Discrete:
$$f_X(x_i) = \sum_{y_j \in S_Y} f_{X,Y}(x_i, y_j)$$

Continuous:
$$f_X(x) = \int_{-\infty}^{\infty} f_{X,Y}(x, y) dy$$

is called the *marginal density (mass) functions* of X

The *marginal cumulative distribution function* of X is

Discrete:
$$F_X(x) = \sum_{x_i \leq x} f_X(x_i)$$

Continuous:
$$F_X(x) = \int_{-\infty}^x f_X(u) du$$

Example

Suppose the random vector (X, Y) takes values in the set

$$\{(1, 2), (1, 4), (2, 3), (3, 2), (3, 4)\}$$

each with equal probability

Example

Suppose the random vector (X, Y) takes values in the set

$$\{(1, 2), (1, 4), (2, 3), (3, 2), (3, 4)\}$$

each with equal probability

What are $f_X(x)$ and $f_Y(y)$?

Example

Suppose the random vector (X, Y) takes values in the set

$$\{(1, 2), (1, 4), (2, 3), (3, 2), (3, 4)\}$$

each with equal probability

What are $f_X(x)$ and $f_Y(y)$?

$$f_X(x) = \begin{cases} 0.4 & x = 1 \\ 0.2 & x = 2 \\ 0.4 & x = 3 \end{cases} \quad f_Y(y) = \begin{cases} 0.4 & y = 2 \\ 0.2 & y = 3 \\ 0.4 & y = 4 \end{cases}$$

Expected Value

Let (X, Y) be a bivariate random vector with joint pdf $f_{X,Y}(x, y)$

Expected Value

Let (X, Y) be a bivariate random vector with joint pdf $f_{X,Y}(x, y)$

The expected value of a function $g(X, Y)$ is

$$E(g(X, Y)) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} g(x, y) f_{X,Y}(x, y) dx dy$$

Expected Value of the Sum of Two Random Variables

Suppose (X, Y) is a bivariate random vector with joint pdf $f_{X,Y}(x, y)$

Expected Value of the Sum of Two Random Variables

Suppose (X, Y) is a bivariate random vector with joint pdf $f_{X,Y}(x, y)$

What is the expected value of $(X + Y)$?

Outline

Independence

Two discrete random variables X and Y are *independent* if

$$P(X = x_i, Y = y_j) = P(X = x_i)P(Y = y_j)$$

holds for all pairs $(x_i, y_j) \in S_{XY}$

Independence

Two discrete random variables X and Y are *independent* if

$$P(X = x_i, Y = y_j) = P(X = x_i)P(Y = y_j)$$

holds for all pairs $(x_i, y_j) \in S_{XY}$

In general, X and Y are *independent* if

$$f_{X,Y}(x, y) = f_X(x)f_Y(y) \quad \text{for all } x, y \in \mathbb{R}$$

Independence

Two discrete random variables X and Y are *independent* if

$$P(X = x_i, Y = y_j) = P(X = x_i)P(Y = y_j)$$

holds for all pairs $(x_i, y_j) \in S_{XY}$

In general, X and Y are *independent* if

$$f_{X,Y}(x, y) = f_X(x)f_Y(y) \quad \text{for all } x, y \in \mathbb{R}$$

Also implies that

$$F_{X,Y}(x, y) = F_X(x)F_Y(y) \quad \text{for all } x, y \in \mathbb{R}$$

Independence

Two discrete random variables X and Y are *independent* if

$$P(X = x_i, Y = y_j) = P(X = x_i)P(Y = y_j)$$

holds for all pairs $(x_i, y_j) \in S_{XY}$

In general, X and Y are *independent* if

$$f_{X,Y}(x, y) = f_X(x)f_Y(y) \quad \text{for all } x, y \in \mathbb{R}$$

Also implies that

$$F_{X,Y}(x, y) = F_X(x)F_Y(y) \quad \text{for all } x, y \in \mathbb{R}$$

The notation $X, Y \stackrel{iid}{\sim} f$ means that X and Y are *independent and identically distributed*, that is $f_X(x) = f_Y(y) = f$

Example

Recall the previous example

$$\{(1, 2), (1, 4), (2, 3), (3, 2), (3, 4)\}$$

$$f_X(x) = \begin{cases} 0.4 & x = 1 \\ 0.2 & x = 2 \\ 0.4 & x = 3 \end{cases}$$

$$f_Y(y) = \begin{cases} 0.4 & y = 2 \\ 0.2 & y = 3 \\ 0.4 & y = 4 \end{cases}$$

Example

Recall the previous example

$$\{(1, 2), (1, 4), (2, 3), (3, 2), (3, 4)\}$$

$$f_X(x) = \begin{cases} 0.4 & x = 1 \\ 0.2 & x = 2 \\ 0.4 & x = 3 \end{cases}$$

$$f_Y(y) = \begin{cases} 0.4 & y = 2 \\ 0.2 & y = 3 \\ 0.4 & y = 4 \end{cases}$$

Are X and Y independent?

Example

Recall the previous example

$$\{(1, 2), (1, 4), (2, 3), (3, 2), (3, 4)\}$$

$$f_X(x) = \begin{cases} 0.4 & x = 1 \\ 0.2 & x = 2 \\ 0.4 & x = 3 \end{cases} \quad f_Y(y) = \begin{cases} 0.4 & y = 2 \\ 0.2 & y = 3 \\ 0.4 & y = 4 \end{cases}$$

Are X and Y independent?

$$P(X = 1, Y = 4) = 0.2 \neq 0.16 = P(X = 1)P(Y = 4)$$

Conditional Density

Let (X, Y) be a bivariate random vector with joint pdf $f_{X,Y}(x, y)$

Conditional Density

Let (X, Y) be a bivariate random vector with joint pdf $f_{X,Y}(x, y)$

The *conditional density* of X given that $Y = y$ such that $f_Y(y) > 0$ is

$$f_{X|Y}(x|y) = \frac{f_{X,Y}(x, y)}{f_Y(y)}$$

Conditional Density

Let (X, Y) be a bivariate random vector with joint pdf $f_{X,Y}(x, y)$

The *conditional density* of X given that $Y = y$ such that $f_Y(y) > 0$ is

$$f_{X|Y}(x|y) = \frac{f_{X,Y}(x, y)}{f_Y(y)}$$

If X and Y are independent

$$f_{X|Y}(x|y) = f_X(x) \quad f_{Y|X}(y|x) = f_Y(y) \quad \text{for all } x, y$$

Example

Let (X, Y) be a bivariate random vector with joint pdf

$$f_{X,Y}(x,y) = \begin{cases} x+y & 0 < x < 1, \ 0 < y < 1 \\ 0 & \text{otherwise} \end{cases}$$

Example

Let (X, Y) be a bivariate random vector with joint pdf

$$f_{X,Y}(x,y) = \begin{cases} x+y & 0 < x < 1, \ 0 < y < 1 \\ 0 & \text{otherwise} \end{cases}$$

Find the conditional density $X|Y$

Example

Let (X, Y) be a bivariate random vector with joint pdf

$$f_{X,Y}(x, y) = \begin{cases} x + y & 0 < x < 1, \ 0 < y < 1 \\ 0 & \text{otherwise} \end{cases}$$

Find the conditional density $X|Y$

First, compute the marginal density of Y

$$f_Y(y) = \int_{-\infty}^{\infty} f_{X,Y}(x, y) dx = \int_0^1 (x + y) dx = \left. \frac{1}{2}x^2 + xy \right|_0^1 = y + \frac{1}{2}$$

Example

Let (X, Y) be a bivariate random vector with joint pdf

$$f_{X,Y}(x,y) = \begin{cases} x+y & 0 < x < 1, \ 0 < y < 1 \\ 0 & \text{otherwise} \end{cases}$$

Find the conditional density $X|Y$

First, compute the marginal density of Y

$$f_Y(y) = \int_{-\infty}^{\infty} f_{X,Y}(x,y) dx = \int_0^1 (x+y) dx = \left. \frac{1}{2}x^2 + xy \right|_0^1 = y + \frac{1}{2}$$

By the definition of conditional probability

$$f_{X|Y}(x|y) = \frac{f_{X,Y}(x,y)}{f_Y(y)} = \frac{x+y}{y + \frac{1}{2}} \quad 0 < x < 1, \ 0 < y < 1$$

Outline

Covariance

The *covariance* is a measure of the (linear) dependence between two random variables X and Y

Covariance

The *covariance* is a measure of the (linear) dependence between two random variables X and Y

$$\text{Cov}(X, Y) = E([X - E(X)][Y - E(Y)])$$

Covariance

The *covariance* is a measure of the (linear) dependence between two random variables X and Y

$$\begin{aligned}\text{Cov}(X, Y) &= E([X - E(X)][Y - E(Y)]) \\ &= E(XY) - E(X)E(Y)\end{aligned}$$

Covariance

The *covariance* is a measure of the (linear) dependence between two random variables X and Y

$$\begin{aligned}\text{Cov}(X, Y) &= E([X - E(X)][Y - E(Y)]) \\ &= E(XY) - E(X)E(Y)\end{aligned}$$

Properties:

- ▶ $\text{Cov}(X, Y) = \text{Cov}(Y, X)$

Covariance

The *covariance* is a measure of the (linear) dependence between two random variables X and Y

$$\begin{aligned}\text{Cov}(X, Y) &= E([X - E(X)][Y - E(Y)]) \\ &= E(XY) - E(X)E(Y)\end{aligned}$$

Properties:

- ▶ $\text{Cov}(X, Y) = \text{Cov}(Y, X)$
- ▶ $\text{Cov}(X, X) = \text{Var}(X)$

Covariance

The *covariance* is a measure of the (linear) dependence between two random variables X and Y

$$\begin{aligned}\text{Cov}(X, Y) &= E([X - E(X)][Y - E(Y)]) \\ &= E(XY) - E(X)E(Y)\end{aligned}$$

Properties:

- ▶ $\text{Cov}(X, Y) = \text{Cov}(Y, X)$
- ▶ $\text{Cov}(X, X) = \text{Var}(X)$
- ▶ $\text{Cov}(X + Y, Z) = \text{Cov}(X, Z) + \text{Cov}(Y, Z)$

Covariance

The *covariance* is a measure of the (linear) dependence between two random variables X and Y

$$\begin{aligned}\text{Cov}(X, Y) &= E([X - E(X)][Y - E(Y)]) \\ &= E(XY) - E(X)E(Y)\end{aligned}$$

Properties:

- ▶ $\text{Cov}(X, Y) = \text{Cov}(Y, X)$
- ▶ $\text{Cov}(X, X) = \text{Var}(X)$
- ▶ $\text{Cov}(X + Y, Z) = \text{Cov}(X, Z) + \text{Cov}(Y, Z)$
- ▶ $\text{Cov}(aX + b, cY + d) = ac \text{Cov}(X, Y)$ *a, b, c, d constant*

Covariance

The *covariance* is a measure of the (linear) dependence between two random variables X and Y

$$\begin{aligned}\text{Cov}(X, Y) &= E([X - E(X)][Y - E(Y)]) \\ &= E(XY) - E(X)E(Y)\end{aligned}$$

Properties:

- ▶ $\text{Cov}(X, Y) = \text{Cov}(Y, X)$
- ▶ $\text{Cov}(X, X) = \text{Var}(X)$
- ▶ $\text{Cov}(X + Y, Z) = \text{Cov}(X, Z) + \text{Cov}(Y, Z)$
- ▶ $\text{Cov}(aX + b, cY + d) = ac \text{Cov}(X, Y)$ *a, b, c, d constant*
- ▶ If X and Y are independent then $\text{Cov}(X, Y) = 0$

Covariance

The *covariance* is a measure of the (linear) dependence between two random variables X and Y

$$\begin{aligned}\text{Cov}(X, Y) &= E([X - E(X)][Y - E(Y)]) \\ &= E(XY) - E(X)E(Y)\end{aligned}$$

Properties:

- ▶ $\text{Cov}(X, Y) = \text{Cov}(Y, X)$
- ▶ $\text{Cov}(X, X) = \text{Var}(X)$
- ▶ $\text{Cov}(X + Y, Z) = \text{Cov}(X, Z) + \text{Cov}(Y, Z)$
- ▶ $\text{Cov}(aX + b, cY + d) = ac \text{Cov}(X, Y)$ a, b, c, d constant
- ▶ If X and Y are independent then $\text{Cov}(X, Y) = 0$
- ▶ $\text{Cov}(X, Y) = 0$ does **NOT** imply X and Y are independent

Example

Let X and Y be random variables with joint density

$$f_{X,Y}(x,y) = \begin{cases} x+y & 0 < x < 1, \ 0 < y < 1 \\ 0 & \text{otherwise} \end{cases}$$

Example

Let X and Y be random variables with joint density

$$f_{X,Y}(x,y) = \begin{cases} x+y & 0 < x < 1, \ 0 < y < 1 \\ 0 & \text{otherwise} \end{cases}$$

Find $\text{Var}(X)$ and $\text{Cov}(X, Y)$

Example

Let X and Y be random variables with joint density

$$f_{X,Y}(x,y) = \begin{cases} x+y & 0 < x < 1, \ 0 < y < 1 \\ 0 & \text{otherwise} \end{cases}$$

Find $\text{Var}(X)$ and $\text{Cov}(X, Y)$

$$E(X) = \int_{-\infty}^{\infty} x f_X(x) dx = \int_0^1 x(x + \frac{1}{2}) dx = \frac{1}{3}x^3 + \frac{1}{4}x^2 \Big|_0^1 = \frac{7}{12}$$

Example

Let X and Y be random variables with joint density

$$f_{X,Y}(x,y) = \begin{cases} x+y & 0 < x < 1, \ 0 < y < 1 \\ 0 & \text{otherwise} \end{cases}$$

Find $\text{Var}(X)$ and $\text{Cov}(X, Y)$

$$E(X) = \int_{-\infty}^{\infty} x f_X(x) dx = \int_0^1 x(x + \frac{1}{2}) dx = \frac{1}{3}x^3 + \frac{1}{4}x^2 \Big|_0^1 = \frac{7}{12}$$

$$E(X^2) = \int_{-\infty}^{\infty} x^2 f_X(x) dx = \int_0^1 x^2(x + \frac{1}{2}) dx = \frac{1}{4}x^4 + \frac{1}{6}x^3 \Big|_0^1 = \frac{5}{12}$$

Example

Let X and Y be random variables with joint density

$$f_{X,Y}(x,y) = \begin{cases} x+y & 0 < x < 1, \ 0 < y < 1 \\ 0 & \text{otherwise} \end{cases}$$

Find $\text{Var}(X)$ and $\text{Cov}(X, Y)$

$$E(X) = \int_{-\infty}^{\infty} x f_X(x) dx = \int_0^1 x(x + \frac{1}{2}) dx = \frac{1}{3}x^3 + \frac{1}{4}x^2 \Big|_0^1 = \frac{7}{12}$$

$$E(X^2) = \int_{-\infty}^{\infty} x^2 f_X(x) dx = \int_0^1 x^2(x + \frac{1}{2}) dx = \frac{1}{4}x^4 + \frac{1}{6}x^3 \Big|_0^1 = \frac{5}{12}$$

$$\text{Var}(X) = E(X^2) - (E(X))^2 = \frac{5}{12} - \left[\frac{7}{12}\right]^2 = \frac{11}{144}$$

Example (continued)

$$E(XY) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} xy f_{X,Y}(x, y) dx dy$$

Example (continued)

$$\begin{aligned} E(XY) &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} xy f_{X,Y}(x,y) dx dy \\ &= \int_0^1 \left[\int_0^1 xy(x+y) dx \right] dy \end{aligned}$$

Example (continued)

$$\begin{aligned} E(XY) &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} xy f_{X,Y}(x, y) dx dy \\ &= \int_0^1 \left[\int_0^1 xy(x+y) dx \right] dy \\ &= \int_0^1 \left[\frac{1}{3}x^3y + \frac{1}{2}x^2y^2 \right]_{x=0}^{x=1} dy \end{aligned}$$

Example (continued)

$$\begin{aligned} E(XY) &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} xy f_{X,Y}(x, y) dx dy \\ &= \int_0^1 \left[\int_0^1 xy(x+y) dx \right] dy \\ &= \int_0^1 \left[\frac{1}{3}x^3y + \frac{1}{2}x^2y^2 \right]_{x=0}^{x=1} dy \\ &= \int_0^1 \frac{1}{3}y + \frac{1}{2}y^2 dy \end{aligned}$$

Example (continued)

$$\begin{aligned} E(XY) &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} xy f_{X,Y}(x, y) dx dy \\ &= \int_0^1 \left[\int_0^1 xy(x+y) dx \right] dy \\ &= \int_0^1 \left[\frac{1}{3}x^3y + \frac{1}{2}x^2y^2 \right]_{x=0}^{x=1} dy \\ &= \int_0^1 \frac{1}{3}y + \frac{1}{2}y^2 dy \\ &= \frac{1}{6}y^2 + \frac{1}{6}y^3 \Big|_{y=0}^{y=1} = \frac{1}{3} \end{aligned}$$

Example (continued)

$$\begin{aligned}E(XY) &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} xy f_{X,Y}(x, y) dx dy \\&= \int_0^1 \left[\int_0^1 xy(x+y) dx \right] dy \\&= \int_0^1 \left[\frac{1}{3}x^3y + \frac{1}{2}x^2y^2 \right]_{x=0}^{x=1} dy \\&= \int_0^1 \frac{1}{3}y + \frac{1}{2}y^2 dy \\&= \frac{1}{6}y^2 + \frac{1}{6}y^3 \Big|_{y=0}^{y=1} = \frac{1}{3}\end{aligned}$$

$$\text{Cov}(X, Y) = E(XY) - E(X)E(Y) = \frac{1}{3} - \left[\frac{7}{12} \right] \left[\frac{7}{12} \right] = -\frac{1}{144}$$

Variance of the Sum of Two Random Variables

Let (X, Y) be a bivariate random vector with joint pdf $f_{X,Y}(x, y)$

Variance of the Sum of Two Random Variables

Let (X, Y) be a bivariate random vector with joint pdf $f_{X,Y}(x, y)$

What is the variance of $(X + Y)$?

Variance of the Sum of Two Random Variables

Let (X, Y) be a bivariate random vector with joint pdf $f_{X,Y}(x, y)$

What is the variance of $(X + Y)$?

$$\text{Var}(X + Y) = E[(X + Y)^2] - [E(X + Y)]^2$$

Variance of the Sum of Two Random Variables

Let (X, Y) be a bivariate random vector with joint pdf $f_{X,Y}(x, y)$

What is the variance of $(X + Y)$?

$$\begin{aligned}\text{Var}(X + Y) &= E[(X + Y)^2] - [E(X + Y)]^2 \\ &= E(X^2 + 2XY + Y^2) - [E(X) + E(Y)]^2\end{aligned}$$

Variance of the Sum of Two Random Variables

Let (X, Y) be a bivariate random vector with joint pdf $f_{X,Y}(x, y)$

What is the variance of $(X + Y)$?

$$\begin{aligned}\text{Var}(X + Y) &= E[(X + Y)^2] - [E(X + Y)]^2 \\&= E(X^2 + 2XY + Y^2) - [E(X) + E(Y)]^2 \\&= E(X^2) + E(2XY) + E(Y^2) \\&\quad - [E(X)]^2 - 2E(X)E(Y) - [E(Y)]^2\end{aligned}$$

Variance of the Sum of Two Random Variables

Let (X, Y) be a bivariate random vector with joint pdf $f_{X,Y}(x, y)$

What is the variance of $(X + Y)$?

$$\begin{aligned}\text{Var}(X + Y) &= E[(X + Y)^2] - [E(X + Y)]^2 \\&= E(X^2 + 2XY + Y^2) - [E(X) + E(Y)]^2 \\&= E(X^2) + E(2XY) + E(Y^2) \\&\quad - [E(X)]^2 - 2E(X)E(Y) - [E(Y)]^2 \\&= E(X^2) - [E(X)]^2 \\&\quad + 2[E(XY) - E(X)E(Y)] \\&\quad + E(Y^2) - [E(Y)]^2\end{aligned}$$

Variance of the Sum of Two Random Variables

$$\text{Var}(X + Y) = \text{Var}(X) + \text{Var}(Y) + 2 \text{Cov}(X, Y)$$

Variance of the Sum of Two Random Variables

$$\text{Var}(X + Y) = \text{Var}(X) + \text{Var}(Y) + 2 \text{Cov}(X, Y)$$

More generally,

$$\text{Var}(aX + bY + c) = a^2 \text{Var}(X) + b^2 \text{Var}(Y) + 2ab \text{Cov}(X, Y)$$

Correlation

The *correlation* is a unit-free measure of the (linear) dependence between two random variables X and Y

Correlation

The *correlation* is a unit-free measure of the (linear) dependence between two random variables X and Y

$$\rho = \rho_{X,Y} = \rho(X, Y) = \text{Corr}(X, Y) = \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X) \text{Var}(Y)}}$$

Correlation

The *correlation* is a unit-free measure of the (linear) dependence between two random variables X and Y

$$\rho = \rho_{X,Y} = \rho(X, Y) = \text{Corr}(X, Y) = \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X) \text{Var}(Y)}}$$

Properties:

- ▶ $\text{Corr}(X, Y) = \text{Corr}(Y, X)$

Correlation

The *correlation* is a unit-free measure of the (linear) dependence between two random variables X and Y

$$\rho = \rho_{X,Y} = \rho(X, Y) = \text{Corr}(X, Y) = \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X) \text{Var}(Y)}}$$

Properties:

- ▶ $\text{Corr}(X, Y) = \text{Corr}(Y, X)$
- ▶ $\text{Corr}(X, X) = 1$

Correlation

The *correlation* is a unit-free measure of the (linear) dependence between two random variables X and Y

$$\rho = \rho_{X,Y} = \rho(X, Y) = \text{Corr}(X, Y) = \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X) \text{Var}(Y)}}$$

Properties:

- ▶ $\text{Corr}(X, Y) = \text{Corr}(Y, X)$
- ▶ $\text{Corr}(X, X) = 1$
- ▶ $\text{Corr}(X, -X) = -1$

Correlation

The *correlation* is a unit-free measure of the (linear) dependence between two random variables X and Y

$$\rho = \rho_{X,Y} = \rho(X, Y) = \text{Corr}(X, Y) = \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X) \text{Var}(Y)}}$$

Properties:

- ▶ $\text{Corr}(X, Y) = \text{Corr}(Y, X)$
- ▶ $\text{Corr}(X, X) = 1$
- ▶ $\text{Corr}(X, -X) = -1$
- ▶ $-1 \leq \text{Corr}(X, Y) \leq 1$ (Cauchy-Schwarz inequality)

Correlation

The *correlation* is a unit-free measure of the (linear) dependence between two random variables X and Y

$$\rho = \rho_{X,Y} = \rho(X, Y) = \text{Corr}(X, Y) = \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X) \text{Var}(Y)}}$$

Properties:

- ▶ $\text{Corr}(X, Y) = \text{Corr}(Y, X)$
- ▶ $\text{Corr}(X, X) = 1$
- ▶ $\text{Corr}(X, -X) = -1$
- ▶ $-1 \leq \text{Corr}(X, Y) \leq 1$ (Cauchy-Schwarz inequality)
- ▶ If X and Y are independent then $\text{Corr}(X, Y) = 0$

Correlation

The *correlation* is a unit-free measure of the (linear) dependence between two random variables X and Y

$$\rho = \rho_{X,Y} = \rho(X, Y) = \text{Corr}(X, Y) = \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X) \text{Var}(Y)}}$$

Properties:

- ▶ $\text{Corr}(X, Y) = \text{Corr}(Y, X)$
- ▶ $\text{Corr}(X, X) = 1$
- ▶ $\text{Corr}(X, -X) = -1$
- ▶ $-1 \leq \text{Corr}(X, Y) \leq 1$ (Cauchy-Schwarz inequality)
- ▶ If X and Y are independent then $\text{Corr}(X, Y) = 0$
- ▶ Correlation \neq Causality!

Example

Let X and Y be random variables with joint density

$$f_{X,Y}(x,y) = \begin{cases} x+y & 0 < x < 1, \ 0 < y < 1 \\ 0 & \text{otherwise} \end{cases}$$

Example

Let X and Y be random variables with joint density

$$f_{X,Y}(x,y) = \begin{cases} x+y & 0 < x < 1, \ 0 < y < 1 \\ 0 & \text{otherwise} \end{cases}$$

Find $\text{Corr}(X, Y)$

Example

Let X and Y be random variables with joint density

$$f_{X,Y}(x,y) = \begin{cases} x+y & 0 < x < 1, \ 0 < y < 1 \\ 0 & \text{otherwise} \end{cases}$$

Find $\text{Corr}(X, Y)$

Already know that $\text{Cov}(X, Y) = -\frac{1}{144}$ $\text{Var}(X) = \text{Var}(Y) = \frac{11}{144}$

Example

Let X and Y be random variables with joint density

$$f_{X,Y}(x,y) = \begin{cases} x+y & 0 < x < 1, \ 0 < y < 1 \\ 0 & \text{otherwise} \end{cases}$$

Find $\text{Corr}(X, Y)$

Already know that $\text{Cov}(X, Y) = -\frac{1}{144}$ $\text{Var}(X) = \text{Var}(Y) = \frac{11}{144}$

$$\text{Corr}(X, Y) = \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X) \text{Var}(Y)}} = \frac{-\frac{1}{144}}{\sqrt{\frac{11}{144} \frac{11}{144}}} = -\frac{1}{11}$$

Autocovariance and Autocorrelation

Let $\{X_t, t \in \mathbb{Z}\}$ be a sequence of time-ordered random variables

Autocovariance and Autocorrelation

Let $\{X_t, t \in \mathbb{Z}\}$ be a sequence of time-ordered random variables
 $\{X_t\}$ is called a *time series model*

Autocovariance and Autocorrelation

Let $\{X_t, t \in \mathbb{Z}\}$ be a sequence of time-ordered random variables
 $\{X_t\}$ is called a *time series model*

The mean and covariance functions of $\{X_t\}$ are given by

$$\mu_X(t) = E(X_t)$$

Autocovariance and Autocorrelation

Let $\{X_t, t \in \mathbb{Z}\}$ be a sequence of time-ordered random variables
 $\{X_t\}$ is called a *time series model*

The mean and covariance functions of $\{X_t\}$ are given by

$$\begin{aligned}\mu_X(t) &= E(X_t) \\ \gamma_X(r, s) &= \text{Cov}(X_r, X_s) = E[(X_r - \mu_X(r))(X_s - \mu_X(s))]\end{aligned}$$

Autocovariance and Autocorrelation

Let $\{X_t, t \in \mathbb{Z}\}$ be a sequence of time-ordered random variables
 $\{X_t\}$ is called a *time series model*

The mean and covariance functions of $\{X_t\}$ are given by

$$\begin{aligned}\mu_X(t) &= E(X_t) \\ \gamma_X(r, s) &= \text{Cov}(X_r, X_s) = E[(X_r - \mu_X(r))(X_s - \mu_X(s))]\end{aligned}$$

A time series model is called *stationary* if $\mu_X(t) = \mu_X$ and if $\gamma(r, s)$ depends only on $h = r - s$

Autocovariance and Autocorrelation

Let $\{X_t, t \in \mathbb{Z}\}$ be a sequence of time-ordered random variables
 $\{X_t\}$ is called a *time series model*

The mean and covariance functions of $\{X_t\}$ are given by

$$\begin{aligned}\mu_X(t) &= E(X_t) \\ \gamma_X(r, s) &= \text{Cov}(X_r, X_s) = E[(X_r - \mu_X(r))(X_s - \mu_X(s))]\end{aligned}$$

A time series model is called *stationary* if $\mu_X(t) = \mu_X$ and if $\gamma(r, s)$ depends only on $h = r - s$

For a stationary times series model, the *autocovariance* function is

$$\gamma_X(h) = \text{Cov}(X_{t+h}, X_t)$$

Autocovariance and Autocorrelation

Let $\{X_t, t \in \mathbb{Z}\}$ be a sequence of time-ordered random variables
 $\{X_t\}$ is called a *time series model*

The mean and covariance functions of $\{X_t\}$ are given by

$$\begin{aligned}\mu_X(t) &= E(X_t) \\ \gamma_X(r, s) &= \text{Cov}(X_r, X_s) = E[(X_r - \mu_X(r))(X_s - \mu_X(s))]\end{aligned}$$

A time series model is called *stationary* if $\mu_X(t) = \mu_X$ and if $\gamma(r, s)$ depends only on $h = r - s$

For a stationary times series model, the *autocovariance* function is

$$\gamma_X(h) = \text{Cov}(X_{t+h}, X_t)$$

For a stationary times series model, the *autocorrelation* function is

$$\rho_X(h) = \frac{\gamma_X(h)}{\gamma_X(0)} = \text{Corr}(X_{t+h}, X_t)$$

Sample Autocovariance and Autocorrelation

A realization $\{x_1, x_2, \dots, x_n\}$ of $\{X_t\}$ is called a *time series*

Sample Autocovariance and Autocorrelation

A realization $\{x_1, x_2, \dots, x_n\}$ of $\{X_t\}$ is called a *time series*

The sample mean of a time series is

$$\bar{x} = \frac{1}{n} \sum_{t=1}^n x_t$$

Sample Autocovariance and Autocorrelation

A realization $\{x_1, x_2, \dots, x_n\}$ of $\{X_t\}$ is called a *time series*

The sample mean of a time series is

$$\bar{x} = \frac{1}{n} \sum_{t=1}^n x_t$$

The *sample autocovariance function* is

$$\hat{\gamma}(h) = \frac{1}{n} \sum_{t=1}^{n-|h|} (x_{t+h} - \bar{x})(x_t - \bar{x}) \quad -n < h < n$$

Sample Autocovariance and Autocorrelation

A realization $\{x_1, x_2, \dots, x_n\}$ of $\{X_t\}$ is called a *time series*

The sample mean of a time series is

$$\bar{x} = \frac{1}{n} \sum_{t=1}^n x_t$$

The *sample autocovariance function* is

$$\hat{\gamma}(h) = \frac{1}{n} \sum_{t=1}^{n-|h|} (x_{t+h} - \bar{x})(x_t - \bar{x}) \quad -n < h < n$$

The *sample autocorrelation function* is

$$\hat{\rho}(h) = \frac{\hat{\gamma}(h)}{\hat{\gamma}(0)} \quad -n < h < n$$

Functions of Independent Random Variables

Let X and Y be independent random variables

Functions of Independent Random Variables

Let X and Y be independent random variables

Let $g(x)$ be a function of only x and $h(y)$ a function of only y

Functions of Independent Random Variables

Let X and Y be independent random variables

Let $g(x)$ be a function of only x and $h(y)$ a function of only y

The random variables $U = g(X)$ and $V = h(Y)$ are independent

Functions of Independent Random Variables


Let X and Y be independent random variables

Let $g(x)$ be a function of only x and $h(y)$ a function of only y



The random variables $U = g(X)$ and $V = h(Y)$ are independent

Recall: independence implies uncorrelated

Dell Returns Example



Dell Returns Example



Dell Returns Example

- ▶ The returns on Dell appear uncorrelated


Dell Returns Example

- ▶ The returns on Dell appear uncorrelated
- ▶ If the returns are independent, then any transformation of the returns should also be uncorrelated

Dell Returns Example

- ▶ The returns on Dell appear uncorrelated
- ▶ If the returns are independent, then any transformation of the returns should also be uncorrelated
- ▶ Consider the Dell returns squared . . .

Dell Returns Example



Outline

Mixture Distributions

A random variable X has a *mixture distribution* if its distribution depends on a quantity that also has a distribution

Mixture Distributions

A random variable X has a *mixture distribution* if its distribution depends on a quantity that also has a distribution

Example: Model for asset returns that allows for extreme events in the lower tail

Mixture Distributions

A random variable X has a *mixture distribution* if its distribution depends on a quantity that also has a distribution

Example: Model for asset returns that allows for extreme events in the lower tail

Let ...

$$Y \sim \text{Binomial}(1, 0.05)$$

Mixture Distributions

A random variable X has a *mixture distribution* if its distribution depends on a quantity that also has a distribution

Example: Model for asset returns that allows for extreme events in the lower tail

Let ...

$$Y \sim \text{Binomial}(1, 0.05)$$

$$X|Y = 1 \sim \mathcal{N}(-12, 6^2)$$

Mixture Distributions

A random variable X has a *mixture distribution* if its distribution depends on a quantity that also has a distribution

Example: Model for asset returns that allows for extreme events in the lower tail

Let ...

$$Y \sim \text{Binomial}(1, 0.05)$$

$$X|Y = 1 \sim \mathcal{N}(-12, 6^2)$$

$$X|Y = 0 \sim \mathcal{N}(3, 3^2)$$

Mixture Distributions

A random variable X has a *mixture distribution* if its distribution depends on a quantity that also has a distribution

Example: Model for asset returns that allows for extreme events in the lower tail

Let ...

$$Y \sim \text{Binomial}(1, 0.05)$$


$$X|Y = 1 \sim \mathcal{N}(-12, 6^2)$$

$$X|Y = 0 \sim \mathcal{N}(3, 3^2)$$

The distribution of X depends on Y and Y has a distribution

$\implies X$ has a mixture distribution

Density of X



Conditional Expectation

Let X and Y be any 2 random variables, then

$$E(X) = E[E(X|Y)]$$

provided that the expectations exist

Conditional Expectation

Let X and Y be any 2 random variables, then

$$E(X) = E[E(X|Y)]$$

provided that the expectations exist

Example: Expected value of the random variable X from the last slide

$$E(X|Y = 1) = -12$$

$$E(X|Y = 0) = 3$$

Conditional Expectation

Let X and Y be any 2 random variables, then

$$E(X) = E[E(X|Y)]$$

provided that the expectations exist

Example: Expected value of the random variable X from the last slide

$$E(X|Y = 1) = -12$$

$$E(X|Y = 0) = 3$$

$$E(X) = E(X|Y = 1)P(Y = 1) + E(X|Y = 0)P(Y = 0)$$

Conditional Expectation

Let X and Y be any 2 random variables, then

$$E(X) = E[E(X|Y)]$$

provided that the expectations exist

Example: Expected value of the random variable X from the last slide

$$E(X|Y = 1) = -12$$

$$E(X|Y = 0) = 3$$

$$\begin{aligned} E(X) &= E(X|Y = 1)P(Y = 1) + E(X|Y = 0)P(Y = 0) \\ &= -12 \times 0.05 + 3 \times 0.95 = 2.25 \end{aligned}$$

Conditional Variance

Let X and Y be any 2 random variables, then

$$\text{Var}(X) = E[\text{Var}(X|Y)] + \text{Var}[E(X|Y)]$$

provided that the expectations exists

Conditional Variance

Let X and Y be any 2 random variables, then

$$\text{Var}(X) = E[\text{Var}(X|Y)] + \text{Var}[E(X|Y)]$$

provided that the expectations exists

Example: Variance of the random variable X

Conditional Variance

Let X and Y be any 2 random variables, then

$$\text{Var}(X) = E[\text{Var}(X|Y)] + \text{Var}[E(X|Y)]$$

provided that the expectations exists

Example: Variance of the random variable X

For the first term:

$$\text{Var}(X|Y = 1) = 6^2$$

$$\text{Var}(X|Y = 0) = 3^2$$

Conditional Variance

Let X and Y be any 2 random variables, then

$$\text{Var}(X) = E[\text{Var}(X|Y)] + \text{Var}[E(X|Y)]$$

provided that the expectations exists

Example: Variance of the random variable X

For the first term:

$$\text{Var}(X|Y = 1) = 6^2$$

$$\text{Var}(X|Y = 0) = 3^2$$

$$E[\text{Var}(X|Y)] = \text{Var}(X|Y = 1)P(Y = 1) + \text{Var}(X|Y = 0)P(Y = 0)$$

Conditional Variance

Let X and Y be any 2 random variables, then

$$\text{Var}(X) = E[\text{Var}(X|Y)] + \text{Var}[E(X|Y)]$$

provided that the expectations exists

Example: Variance of the random variable X

For the first term:

$$\text{Var}(X|Y = 1) = 6^2$$

$$\text{Var}(X|Y = 0) = 3^2$$

$$\begin{aligned} E[\text{Var}(X|Y)] &= \text{Var}(X|Y = 1) P(Y = 1) + \text{Var}(X|Y = 0) P(Y = 0) \\ &= 36 \times 0.05 + 9 \times 0.95 = 10.35 \end{aligned}$$

Example (continued)

For the second term:

- ▶ $E(X|Y = 1) = -12$
- ▶ $E(X|Y = 0) = 3$

Example (continued)

For the second term:

- ▶ $E(X|Y = 1) = -12$
- ▶ $E(X|Y = 0) = 3$

Can write as one equation: $E(X|Y) = 3 - 15Y$

$$\text{Var}(E(X|Y)) = \text{Var}(3 - 15Y)$$

Example (continued)

For the second term:

- ▶ $E(X|Y = 1) = -12$
- ▶ $E(X|Y = 0) = 3$

Can write as one equation: $E(X|Y) = 3 - 15Y$

$$\text{Var}(E(X|Y)) = \text{Var}(3 - 15Y) = (-15)^2 \text{Var}(Y)$$

Example (continued)

For the second term:

- ▶ $E(X|Y = 1) = -12$
- ▶ $E(X|Y = 0) = 3$

Can write as one equation: $E(X|Y) = 3 - 15Y$

$$\begin{aligned}\text{Var}(E(X|Y)) &= \text{Var}(3 - 15Y) = (-15)^2 \text{Var}(Y) \\ &= 15^2 \times 0.05 \times (1 - 0.05) = 10.6875\end{aligned}$$

Example (continued)

For the second term:

- ▶ $E(X|Y = 1) = -12$
- ▶ $E(X|Y = 0) = 3$

Can write as one equation: $E(X|Y) = 3 - 15Y$

$$\begin{aligned}\text{Var}(E(X|Y)) &= \text{Var}(3 - 15Y) = (-15)^2 \text{Var}(Y) \\ &= 15^2 \times 0.05 \times (1 - 0.05) = 10.6875\end{aligned}$$

$$\text{Var}(X) = E[\text{Var}(X|Y)] + \text{Var}[E(X|Y)]$$

Example (continued)

For the second term:

- ▶ $E(X|Y = 1) = -12$
- ▶ $E(X|Y = 0) = 3$

Can write as one equation: $E(X|Y) = 3 - 15Y$

$$\begin{aligned}\text{Var}(E(X|Y)) &= \text{Var}(3 - 15Y) = (-15)^2 \text{Var}(Y) \\ &= 15^2 \times 0.05 \times (1 - 0.05) = 10.6875\end{aligned}$$

$$\begin{aligned}\text{Var}(X) &= E[\text{Var}(X|Y)] + \text{Var}[E(X|Y)] \\ &= 10.85 + 10.6875 = 21.5375\end{aligned}$$

Outline

Bivariate Change of Variables Formula

Let (X, Y) be a bivariate random vector with a known pdf

Bivariate Change of Variables Formula

Let (X, Y) be a bivariate random vector with a known pdf

Define a new random vector (U, V)

$$U = g_1(X, Y)$$

$$V = g_2(X, Y)$$

Bivariate Change of Variables Formula

Let (X, Y) be a bivariate random vector with a known pdf

Define a new random vector (U, V)

$$U = g_1(X, Y)$$

$$V = g_2(X, Y)$$

Let $S_{XY} = \{(x, y) \in \mathbb{R}^2 : f_{X,Y}(x, y) > 0\}$

Bivariate Change of Variables Formula

Let (X, Y) be a bivariate random vector with a known pdf

Define a new random vector (U, V)

$$U = g_1(X, Y)$$

$$V = g_2(X, Y)$$

Let $S_{XY} = \{(x, y) \in \mathbb{R}^2 : f_{X,Y}(x, y) > 0\}$

Let $S_{UV} = \{(u, v) : u = g_1(x, y), v = g_2(x, y) \text{ for } (x, y) \in S_{XY}\}$

Bivariate Change of Variables Formula

Let (X, Y) be a bivariate random vector with a known pdf

Define a new random vector (U, V)

$$U = g_1(X, Y)$$

$$V = g_2(X, Y)$$

Let $S_{XY} = \{(x, y) \in \mathbb{R}^2 : f_{X,Y}(x, y) > 0\}$

Let $S_{UV} = \{(u, v) : u = g_1(x, y), v = g_2(x, y) \text{ for } (x, y) \in S_{XY}\}$

The joint pdf $f_{U,V}(u, v)$ is positive on the set $S_{U,V}$

Bivariate Change of Variables Formula

Let (X, Y) be a bivariate random vector with a known pdf

Define a new random vector (U, V)

$$U = g_1(X, Y)$$

$$V = g_2(X, Y)$$

Let $S_{XY} = \{(x, y) \in \mathbb{R}^2 : f_{X,Y}(x, y) > 0\}$

Let $S_{UV} = \{(u, v) : u = g_1(x, y), v = g_2(x, y) \text{ for } (x, y) \in S_{XY}\}$

The joint pdf $f_{U,V}(u, v)$ is positive on the set $S_{U,V}$

The map from S_{XY} to S_{UV} must be one-to-one and onto, that is

$$(u, v) = (g_1(x, y), g_2(x, y))$$

can be inverted

$$(x, y) = (h_1(u, v), h_2(u, v))$$

Bivariate Change of Variables Formula (continued)

The bivariate change of variables formula is given by

$$f_{U,V}(u, v) = f_{X,Y}[h_1(u, v), h_2(u, v)] |J|$$

Bivariate Change of Variables Formula (continued)

The bivariate change of variables formula is given by

$$f_{U,V}(u, v) = f_{X,Y}[h_1(u, v), h_2(u, v)] |J|$$

The quantity J is called the Jacobian of the transformation

$$J = \frac{\partial x}{\partial u} \frac{\partial y}{\partial v} - \frac{\partial y}{\partial u} \frac{\partial x}{\partial v}$$

Bivariate Change of Variables Formula (continued)

The bivariate change of variables formula is given by

$$f_{U,V}(u, v) = f_{X,Y}[h_1(u, v), h_2(u, v)] |J|$$

The quantity J is called the Jacobian of the transformation

$$J = \frac{\partial x}{\partial u} \frac{\partial y}{\partial v} - \frac{\partial y}{\partial u} \frac{\partial x}{\partial v}$$

The partial derivatives are

$$\frac{\partial x}{\partial u} = \frac{\partial h_1(u, v)}{\partial u} \quad \frac{\partial x}{\partial v} = \frac{\partial h_1(u, v)}{\partial v}$$

$$\frac{\partial y}{\partial u} = \frac{\partial h_2(u, v)}{\partial u} \quad \frac{\partial y}{\partial v} = \frac{\partial h_2(u, v)}{\partial v}$$

Example

Let X and Y be independent standard normal random variables

Example

Let X and Y be independent standard normal random variables

Make the transformation

$$U = X + Y$$

$$V = X - Y$$

Example

Let X and Y be independent standard normal random variables

Make the transformation

$$U = X + Y$$

$$V = X - Y$$

In the notation of the previous slide

$$U = g_1(X, Y) \quad \text{where} \quad g_1(x, y) = x + y$$

$$V = g_2(X, Y) \quad \text{where} \quad g_2(x, y) = x - y$$

Example

Let X and Y be independent standard normal random variables

Make the transformation

$$U = X + Y$$

$$V = X - Y$$

In the notation of the previous slide

$$U = g_1(X, Y) \quad \text{where} \quad g_1(x, y) = x + y$$

$$V = g_2(X, Y) \quad \text{where} \quad g_2(x, y) = x - y$$

These are easily solved for

$$x = h_1(u, v) = \frac{1}{2}(u + v)$$

$$y = h_2(u, v) = \frac{1}{2}(u - v)$$

Example

Let X and Y be independent standard normal random variables

Make the transformation

$$U = X + Y$$

$$V = X - Y$$

In the notation of the previous slide

$$U = g_1(X, Y) \quad \text{where} \quad g_1(x, y) = x + y$$

$$V = g_2(X, Y) \quad \text{where} \quad g_2(x, y) = x - y$$

These are easily solved for

$$x = h_1(u, v) = \frac{1}{2}(u + v)$$

$$y = h_2(u, v) = \frac{1}{2}(u - v)$$

X and Y can take any real values $\implies S_{XY} = \mathbb{R}^2$

Example

Let X and Y be independent standard normal random variables

Make the transformation

$$U = X + Y$$

$$V = X - Y$$

In the notation of the previous slide

$$U = g_1(X, Y) \quad \text{where} \quad g_1(x, y) = x + y$$

$$V = g_2(X, Y) \quad \text{where} \quad g_2(x, y) = x - y$$

These are easily solved for

$$x = h_1(u, v) = \frac{1}{2}(u + v)$$

$$y = h_2(u, v) = \frac{1}{2}(u - v)$$

X and Y can take any real values $\implies S_{XY} = \mathbb{R}^2$

The transformation is one-to-one

Example

Let X and Y be independent standard normal random variables

Make the transformation

$$U = X + Y$$

$$V = X - Y$$

In the notation of the previous slide

$$U = g_1(X, Y) \quad \text{where} \quad g_1(x, y) = x + y$$

$$V = g_2(X, Y) \quad \text{where} \quad g_2(x, y) = x - y$$

These are easily solved for

$$x = h_1(u, v) = \frac{1}{2}(u + v)$$

$$y = h_2(u, v) = \frac{1}{2}(u - v)$$

X and Y can take any real values $\implies S_{XY} = \mathbb{R}^2$

The transformation is one-to-one

$$S_{UV} = \mathbb{R}^2$$

Example (continued)

Have: $x = h_1(u, v) = \frac{1}{2}(u + v)$ and $y = h_2(u, v) = \frac{1}{2}(u - v)$

Example (continued)

Have: $x = h_1(u, v) = \frac{1}{2}(u + v)$ and $y = h_2(u, v) = \frac{1}{2}(u - v)$

The Jacobian of the transformation

$$J = \frac{\partial x}{\partial u} \frac{\partial y}{\partial v} - \frac{\partial y}{\partial u} \frac{\partial x}{\partial v}$$

Example (continued)

Have: $x = h_1(u, v) = \frac{1}{2}(u + v)$ and $y = h_2(u, v) = \frac{1}{2}(u - v)$

The Jacobian of the transformation

$$J = \frac{\partial x}{\partial u} \frac{\partial y}{\partial v} - \frac{\partial y}{\partial u} \frac{\partial x}{\partial v} = \frac{1}{2} \left(-\frac{1}{2} \right) - \frac{1}{2} \frac{1}{2}$$

Example (continued)

Have: $x = h_1(u, v) = \frac{1}{2}(u + v)$ and $y = h_2(u, v) = \frac{1}{2}(u - v)$

The Jacobian of the transformation

$$J = \frac{\partial x}{\partial u} \frac{\partial y}{\partial v} - \frac{\partial y}{\partial u} \frac{\partial x}{\partial v} = \frac{1}{2} \left(-\frac{1}{2} \right) - \frac{1}{2} \frac{1}{2} = -\frac{1}{4} - \frac{1}{4} = -\frac{1}{2}$$

Example (continued)

Have: $x = h_1(u, v) = \frac{1}{2}(u + v)$ and $y = h_2(u, v) = \frac{1}{2}(u - v)$

The Jacobian of the transformation

$$J = \frac{\partial x}{\partial u} \frac{\partial y}{\partial v} - \frac{\partial y}{\partial u} \frac{\partial x}{\partial v} = \frac{1}{2} \left(-\frac{1}{2} \right) - \frac{1}{2} \frac{1}{2} = -\frac{1}{4} - \frac{1}{4} = -\frac{1}{2}$$

The joint density of X and Y

$$f_{X,Y}(x, y) = \frac{1}{2\pi} \exp \left[-\frac{x^2 + y^2}{2} \right]$$

Example (continued)

Have: $x = h_1(u, v) = \frac{1}{2}(u + v)$ and $y = h_2(u, v) = \frac{1}{2}(u - v)$

The Jacobian of the transformation

$$J = \frac{\partial x}{\partial u} \frac{\partial y}{\partial v} - \frac{\partial y}{\partial u} \frac{\partial x}{\partial v} = \frac{1}{2} \left(-\frac{1}{2} \right) - \frac{1}{2} \frac{1}{2} = -\frac{1}{4} - \frac{1}{4} = -\frac{1}{2}$$

The joint density of X and Y

$$f_{X,Y}(x, y) = \frac{1}{2\pi} \exp \left[-\frac{x^2 + y^2}{2} \right]$$

The change of variables formula says that

$$f_{U,V}(u, v) = f_{X,Y}(h_1(u, v), h_2(u, v)) |J|$$

Example (continued)

Have: $x = h_1(u, v) = \frac{1}{2}(u + v)$ and $y = h_2(u, v) = \frac{1}{2}(u - v)$

The Jacobian of the transformation

$$J = \frac{\partial x}{\partial u} \frac{\partial y}{\partial v} - \frac{\partial y}{\partial u} \frac{\partial x}{\partial v} = \frac{1}{2} \left(-\frac{1}{2} \right) - \frac{1}{2} \frac{1}{2} = -\frac{1}{4} - \frac{1}{4} = -\frac{1}{2}$$

The joint density of X and Y

$$f_{X,Y}(x, y) = \frac{1}{2\pi} \exp \left[-\frac{x^2 + y^2}{2} \right]$$

The change of variables formula says that

$$f_{U,V}(u, v) = f_{X,Y}(h_1(u, v), h_2(u, v)) |J|$$

Substitute expressions for h_1 , h_2 , and $J \dots$

Example (continued)

Joint density of U and V

$$f_{U,V}(u, v) = f_{X,Y}(x, y)|J|$$

Example (continued)

Joint density of U and V

$$\begin{aligned} f_{U,V}(u,v) &= f_{X,Y}(x,y)|J| \\ &= \frac{1}{2\pi} \exp \left[-\frac{\left(\frac{1}{2}(u+v)\right)^2 + \left(\frac{1}{2}(u-v)\right)^2}{2} \right] \left| -\frac{1}{2} \right| \end{aligned}$$

Example (continued)

Joint density of U and V

$$\begin{aligned}f_{U,V}(u,v) &= f_{X,Y}(x,y)|J| \\&= \frac{1}{2\pi} \exp \left[-\frac{\left(\frac{1}{2}(u+v)\right)^2 + \left(\frac{1}{2}(u-v)\right)^2}{2} \right] \left| -\frac{1}{2} \right| \\&= \frac{1}{2} \frac{1}{2\pi} \exp \left[-\frac{(u^2 + 2uv + v^2) + (u^2 - 2uv + v^2)}{2^3} \right]\end{aligned}$$

Example (continued)

Joint density of U and V

$$\begin{aligned}f_{U,V}(u,v) &= f_{X,Y}(x,y)|J| \\&= \frac{1}{2\pi} \exp \left[-\frac{\left(\frac{1}{2}(u+v)\right)^2 + \left(\frac{1}{2}(u-v)\right)^2}{2} \right] \left| -\frac{1}{2} \right| \\&= \frac{1}{2} \frac{1}{2\pi} \exp \left[-\frac{(u^2 + 2uv + v^2) + (u^2 - 2uv + v^2)}{2^3} \right] \\&= \frac{1}{2} \frac{1}{2\pi} \exp \left[-\frac{2u^2 + 2v^2}{2^3} \right]\end{aligned}$$

Example (continued)

Joint density of U and V

$$\begin{aligned}f_{U,V}(u,v) &= f_{X,Y}(x,y)|J| \\&= \frac{1}{2\pi} \exp \left[-\frac{\left(\frac{1}{2}(u+v)\right)^2 + \left(\frac{1}{2}(u-v)\right)^2}{2} \right] \left| -\frac{1}{2} \right| \\&= \frac{1}{2} \frac{1}{2\pi} \exp \left[-\frac{(u^2 + 2uv + v^2) + (u^2 - 2uv + v^2)}{2^3} \right] \\&= \frac{1}{2} \frac{1}{2\pi} \exp \left[-\frac{2u^2 + 2v^2}{2^3} \right] \\&= \frac{1}{\sqrt{2\pi}\sqrt{2}} \exp \left[-\frac{u^2}{2(\sqrt{2})^2} \right] \frac{1}{\sqrt{2\pi}\sqrt{2}} \exp \left[-\frac{v^2}{2(\sqrt{2})^2} \right]\end{aligned}$$

Independence Revisited

Random variables U and V are independent if the joint density is the product of the marginal densities

$$f_{U,V}(u, v) = f_U(u) f_V(v)$$

Independence Revisited

Random variables U and V are independent if the joint density is the product of the marginal densities

$$f_{U,V}(u, v) = f_U(u) f_V(v)$$

Condition for independence can be stated more generally ...

Independence Revisited

Random variables U and V are independent if the joint density is the product of the marginal densities

$$f_{U,V}(u, v) = f_U(u) f_V(v)$$

Condition for independence can be stated more generally ...

Let (X, Y) be a bivariate random vector with joint density $f_{X,Y}(x, y)$

Independence Revisited

Random variables U and V are independent if the joint density is the product of the marginal densities

$$f_{U,V}(u, v) = f_U(u) f_V(v)$$

Condition for independence can be stated more generally ...

Let (X, Y) be a bivariate random vector with joint density $f_{X,Y}(x, y)$

X and Y are independent iff there exist $g(x)$ and $h(y)$ such that

$$f_{X,Y}(x, y) = g(x) h(y)$$

for all $x \in \mathbb{R}$ and $y \in \mathbb{R}$

Outline

Bivariate Normal Distribution

The bivariate normal distribution is parameterized by

Bivariate Normal Distribution

The bivariate normal distribution is parameterized by

- ▶ μ_X and μ_Y : the means of the marginal distributions

Bivariate Normal Distribution

The bivariate normal distribution is parameterized by

- ▶ μ_X and μ_Y : the means of the marginal distributions
- ▶ σ_X^2 and σ_Y^2 : the variances of the marginal distributions

Bivariate Normal Distribution

The bivariate normal distribution is parameterized by

- ▶ μ_X and μ_Y : the means of the marginal distributions
- ▶ σ_X^2 and σ_Y^2 : the variances of the marginal distributions
- ▶ ρ : the correlation between X and Y

Bivariate Normal Distribution

The bivariate normal distribution is parameterized by

- ▶ μ_X and μ_Y : the means of the marginal distributions
- ▶ σ_X^2 and σ_Y^2 : the variances of the marginal distributions
- ▶ ρ : the correlation between X and Y

The joint density function for the bivariate normal is

$$f_{X,Y}(x,y) =$$

$$k \exp \left\{ - \frac{\left[\frac{x - \mu_X}{\sigma_X} \right]^2 - 2\rho \left[\frac{x - \mu_X}{\sigma_X} \right] \left[\frac{y - \mu_Y}{\sigma_Y} \right] + \left[\frac{y - \mu_Y}{\sigma_Y} \right]^2}{2(1 - \rho^2)} \right\}$$

where the normalization constant

$$k = \frac{1}{2\pi\sigma_X\sigma_Y\sqrt{1 - \rho^2}}$$

Bivariate Normal Density

The joint density can be written compactly in matrix notation

$$f_{X,Y}(\mathbf{x}) = \frac{1}{(\sqrt{2\pi})^2 |\mathbf{\Sigma}|^{\frac{1}{2}}} \exp \left\{ -\frac{1}{2} [\mathbf{x} - \boldsymbol{\mu}]^T \mathbf{\Sigma}^{-1} [\mathbf{x} - \boldsymbol{\mu}] \right\}$$

where

$$\mathbf{x} = \begin{bmatrix} x \\ y \end{bmatrix} \quad \boldsymbol{\mu} = \begin{bmatrix} \mu_X \\ \mu_Y \end{bmatrix} \quad \text{and} \quad \mathbf{\Sigma} = \begin{bmatrix} \sigma_X^2 & \rho\sigma_X\sigma_Y \\ \rho\sigma_Y\sigma_X & \sigma_Y^2 \end{bmatrix}$$

Sum of Normal Random Variables

Let $X \sim \mathcal{N}(\mu, \sigma^2)$ and $Y \sim \mathcal{N}(\gamma, \tau^2)$ be independent RVs

Sum of Normal Random Variables

Let $X \sim \mathcal{N}(\mu, \sigma^2)$ and $Y \sim \mathcal{N}(\gamma, \tau^2)$ be independent RVs

What is the distribution of $X + Y$?

Sum of Normal Random Variables

Let $X \sim \mathcal{N}(\mu, \sigma^2)$ and $Y \sim \mathcal{N}(\gamma, \tau^2)$ be independent RVs

What is the distribution of $X + Y$?

Let $U = X + Y$ and $V = k(X - Y)$ for some $k \in \mathbb{R}$

Sum of Normal Random Variables

Let $X \sim \mathcal{N}(\mu, \sigma^2)$ and $Y \sim \mathcal{N}(\gamma, \tau^2)$ be independent RVs

What is the distribution of $X + Y$?

Let $U = X + Y$ and $V = k(X - Y)$ for some $k \in \mathbb{R}$

Gives transformation

$$u = g_1(x, y) = x + y \quad v = g_2(x, y) = kx - ky$$

with inverse transformation

$$x = h_1(u, v) = \frac{ku + v}{2k} \quad y = h_2(u, v) = \frac{ku - v}{2k}$$

Sum of Normal Random Variables

Let $X \sim \mathcal{N}(\mu, \sigma^2)$ and $Y \sim \mathcal{N}(\gamma, \tau^2)$ be independent RVs

What is the distribution of $X + Y$?

Let $U = X + Y$ and $V = k(X - Y)$ for some $k \in \mathbb{R}$

Gives transformation

$$u = g_1(x, y) = x + y \quad v = g_2(x, y) = kx - ky$$

with inverse transformation

$$x = h_1(u, v) = \frac{ku + v}{2k} \quad y = h_2(u, v) = \frac{ku - v}{2k}$$

The Jacobian of the transformation is

$$J = \frac{\partial x}{\partial u} \frac{\partial y}{\partial v} - \frac{\partial y}{\partial u} \frac{\partial x}{\partial v} = \frac{1}{2} \left(-\frac{1}{2k} \right) - \frac{1}{2} \frac{1}{2k} = -\frac{1}{2k}$$

Sum of Normal Random Variables

Choose $k = \frac{1}{2}$

Sum of Normal Random Variables

Choose $k = \frac{1}{2}$

Gives $x = h_1(u, v) = \frac{1}{2}u + v$ and $y = h_2(u, v) = \frac{1}{2}u - v$

Sum of Normal Random Variables

Choose $k = \frac{1}{2}$

Gives $x = h_1(u, v) = \frac{1}{2}u + v$ and $y = h_2(u, v) = \frac{1}{2}u - v$

More importantly: $|J| = 1$

Sum of Normal Random Variables

Choose $k = \frac{1}{2}$

Gives $x = h_1(u, v) = \frac{1}{2}u + v$ and $y = h_2(u, v) = \frac{1}{2}u - v$

More importantly: $|J| = 1$

The joint density of X and Y is the product of the marginals

$$f_{X,Y}(x, y) = f_X(x) f_Y(y)$$

Sum of Normal Random Variables

Choose $k = \frac{1}{2}$

Gives $x = h_1(u, v) = \frac{1}{2}u + v$ and $y = h_2(u, v) = \frac{1}{2}u - v$

More importantly: $|J| = 1$

The joint density of X and Y is the product of the marginals

$$\begin{aligned} f_{X,Y}(x, y) &= f_X(x) f_Y(y) \\ &= \frac{1}{\sqrt{2\pi\sigma}} \exp\left[-\frac{(x-\mu)^2}{2\sigma^2}\right] \frac{1}{\sqrt{2\pi\tau}} \exp\left[-\frac{(y-\gamma)^2}{2\tau^2}\right] \end{aligned}$$

Sum of Normal Random Variables

Choose $k = \frac{1}{2}$

Gives $x = h_1(u, v) = \frac{1}{2}u + v$ and $y = h_2(u, v) = \frac{1}{2}u - v$

More importantly: $|J| = 1$

The joint density of X and Y is the product of the marginals

$$\begin{aligned}f_{X,Y}(x, y) &= f_X(x) f_Y(y) \\&= \frac{1}{\sqrt{2\pi}\sigma} \exp\left[-\frac{(x-\mu)^2}{2\sigma^2}\right] \frac{1}{\sqrt{2\pi}\tau} \exp\left[-\frac{(y-\gamma)^2}{2\tau^2}\right] \\&= \frac{1}{2\pi\sigma\tau} \exp\left[-\frac{(x-\mu)^2}{2\sigma^2} - \frac{(y-\gamma)^2}{2\tau^2}\right]\end{aligned}$$

Sum of Normal Random Variables

Choose $k = \frac{1}{2}$

Gives $x = h_1(u, v) = \frac{1}{2}u + v$ and $y = h_2(u, v) = \frac{1}{2}u - v$

More importantly: $|J| = 1$

The joint density of X and Y is the product of the marginals

$$\begin{aligned}f_{X,Y}(x, y) &= f_X(x) f_Y(y) \\&= \frac{1}{\sqrt{2\pi}\sigma} \exp\left[-\frac{(x-\mu)^2}{2\sigma^2}\right] \frac{1}{\sqrt{2\pi}\tau} \exp\left[-\frac{(y-\gamma)^2}{2\tau^2}\right] \\&= \frac{1}{2\pi\sigma\tau} \exp\left[-\frac{(x-\mu)^2}{2\sigma^2} - \frac{(y-\gamma)^2}{2\tau^2}\right]\end{aligned}$$

Recall ρ (correlation coefficient) is zero when X and Y independent

Sum of Normal Random Variables

hmmm ...

Sum of Normal Random Variables

hmmm ...

The rest seems like a good homework question



COMPUTATIONAL FINANCE & RISK MANAGEMENT

UNIVERSITY *of* WASHINGTON

Department of Applied Mathematics

<http://computational-finance.uw.edu>