



COMPUTATIONAL FINANCE & RISK MANAGEMENT

UNIVERSITY *of* WASHINGTON

Department of Applied Mathematics

CFRM 410: Probability and Statistics for Computational Finance

Week 10 Hypothesis Testing

Jake Price

Instructor, Computational Finance and Risk Management

University of Washington

Slides originally produced by Kjell Konis

Outline

Test Statistics

Testing a Hypothesis

Significance

p -Values

R Lab

Outline

Test Statistics

Testing a Hypothesis

Significance

p -Values

R Lab

The Scientific Process

- ▶ The scientific process can be summarized by the following steps

The Scientific Process

- ▶ The scientific process can be summarized by the following steps
 1. Propose a hypothesis that can be tested/rejected

The Scientific Process

- ▶ The scientific process can be summarized by the following steps
 1. Propose a hypothesis that can be tested/rejected
 2. Conduct the experiment

The Scientific Process

- ▶ The scientific process can be summarized by the following steps
 1. Propose a hypothesis that can be tested/rejected
 2. Conduct the experiment
 3. Compare outcome with prediction given the hypothesis

The Scientific Process

- ▶ The scientific process can be summarized by the following steps
 1. Propose a hypothesis that can be tested/rejected
 2. Conduct the experiment
 3. Compare outcome with prediction given the hypothesis
 4. Do not reject hypothesis **OR** reject and/or modify hypothesis

The Scientific Process

- ▶ The scientific process can be summarized by the following steps
 1. Propose a hypothesis that can be tested/rejected
 2. Conduct the experiment
 3. Compare outcome with prediction given the hypothesis
 4. Do not reject hypothesis **OR** reject and/or modify hypothesis
- ▶ In statistical terms

The Scientific Process

- ▶ The scientific process can be summarized by the following steps
 1. Propose a hypothesis that can be tested/rejected
 2. Conduct the experiment
 3. Compare outcome with prediction given the hypothesis
 4. Do not reject hypothesis **OR** reject and/or modify hypothesis
- ▶ In statistical terms
 1. Propose a hypothesis (i.e., make a statement about θ)

The Scientific Process

- ▶ The scientific process can be summarized by the following steps
 1. Propose a hypothesis that can be tested/rejected
 2. Conduct the experiment
 3. Compare outcome with prediction given the hypothesis
 4. Do not reject hypothesis **OR** reject and/or modify hypothesis
- ▶ In statistical terms
 1. Propose a hypothesis (i.e., make a statement about θ)
 - ▶ Must be possible to contradict the statement using the data

The Scientific Process

- ▶ The scientific process can be summarized by the following steps
 1. Propose a hypothesis that can be tested/rejected
 2. Conduct the experiment
 3. Compare outcome with prediction given the hypothesis
 4. Do not reject hypothesis **OR** reject and/or modify hypothesis
- ▶ In statistical terms
 1. Propose a hypothesis (i.e., make a statement about θ)
 - ▶ Must be possible to contradict the statement using the data
 2. Perform random experiment, collect data

The Scientific Process

- ▶ The scientific process can be summarized by the following steps
 1. Propose a hypothesis that can be tested/rejected
 2. Conduct the experiment
 3. Compare outcome with prediction given the hypothesis
 4. Do not reject hypothesis **OR** reject and/or modify hypothesis

- ▶ In statistical terms
 1. Propose a hypothesis (i.e., make a statement about θ)
 - ▶ Must be possible to contradict the statement using the data
 2. Perform random experiment, collect data
 3. Reject/don't reject hypothesis by comparing the observed data with the prediction of the statistical model given the hypothesis

The Scientific Process

- ▶ The scientific process can be summarized by the following steps
 1. Propose a hypothesis that can be tested/rejected
 2. Conduct the experiment
 3. Compare outcome with prediction given the hypothesis
 4. Do not reject hypothesis **OR** reject and/or modify hypothesis
- ▶ In statistical terms
 1. Propose a hypothesis (i.e., make a statement about θ)
 - ▶ Must be possible to contradict the statement using the data
 2. Perform random experiment, collect data
 3. Reject/don't reject hypothesis by comparing the observed data with the prediction of the statistical model given the hypothesis
 - ▶ Is the scale significant?

Hypotheses in Statistical Terms

- ▶ Given a statistical model, want to decide between two competing theories

Hypotheses in Statistical Terms

- ▶ Given a statistical model, want to decide between two competing theories
- ▶ Pair of hypotheses called
 - H_0 : the *null* hypothesis
 - H_1 : the *alternative* hypothesis

Hypotheses in Statistical Terms

- ▶ Given a statistical model, want to decide between two competing theories
- ▶ Pair of hypotheses called
 - H_0 : the *null* hypothesis
 - H_1 : the *alternative* hypothesis
- ▶ The hypotheses are statements about the parameter θ

Hypotheses in Statistical Terms

- ▶ Given a statistical model, want to decide between two competing theories
- ▶ Pair of hypotheses called
 - H_0 : the *null* hypothesis
 - H_1 : the *alternative* hypothesis
- ▶ The hypotheses are statements about the parameter θ
 - ▶ H_0 & H_1 must account for all possible values of θ

Hypotheses in Statistical Terms

- ▶ Given a statistical model, want to decide between two competing theories
- ▶ Pair of hypotheses called
 - H_0 : the *null* hypothesis
 - H_1 : the *alternative* hypothesis
- ▶ The hypotheses are statements about the parameter θ
 - ▶ H_0 & H_1 must account for all possible values of θ
- ▶ **Example** In a population with mean μ , form hypotheses about the mean as follows

$$\underbrace{\left\{ \begin{array}{l} H_0 : \mu = \mu_0 \\ H_1 : \mu \neq \mu_0 \end{array} \right\}}_{\text{two-sided test}}$$

Hypotheses in Statistical Terms

- ▶ Given a statistical model, want to decide between two competing theories
- ▶ Pair of hypotheses called
 - H_0 : the *null* hypothesis
 - H_1 : the *alternative* hypothesis
- ▶ The hypotheses are statements about the parameter θ
 - ▶ H_0 & H_1 must account for all possible values of θ
- ▶ **Example** In a population with mean μ , form hypotheses about the mean as follows

$$\underbrace{\left\{ \begin{array}{l} H_0 : \mu = \mu_0 \\ H_1 : \mu \neq \mu_0 \end{array} \right\}}_{\text{two-sided test}} \quad \text{or} \quad \left\{ \begin{array}{l} H_0 : \mu \leq \mu_0 \\ H_1 : \mu > \mu_0 \end{array} \right\}$$

Hypotheses in Statistical Terms

- ▶ Given a statistical model, want to decide between two competing theories
- ▶ Pair of hypotheses called
 - H_0 : the *null* hypothesis
 - H_1 : the *alternative* hypothesis
- ▶ The hypotheses are statements about the parameter θ
 - ▶ H_0 & H_1 must account for all possible values of θ
- ▶ **Example** In a population with mean μ , form hypotheses about the mean as follows

$$\underbrace{\left\{ \begin{array}{l} H_0 : \mu = \mu_0 \\ H_1 : \mu \neq \mu_0 \end{array} \right\}}_{\text{two-sided test}} \quad \text{or} \quad \underbrace{\left\{ \begin{array}{l} H_0 : \mu \leq \mu_0 \\ H_1 : \mu > \mu_0 \end{array} \right\} \quad \text{or} \quad \left\{ \begin{array}{l} H_0 : \mu \geq \mu_0 \\ H_1 : \mu < \mu_0 \end{array} \right\}}_{\text{one-sided tests}}$$

Test Statistic

- ▶ How to choose between the two hypotheses?

Test Statistic

- ▶ How to choose between the two hypotheses?
 - ▶ Take a random sample X_1, \dots, X_n from the population

$$X_1, \dots, X_n \stackrel{iid}{\sim} F$$

Test Statistic

- ▶ How to choose between the two hypotheses?
 - ▶ Take a random sample X_1, \dots, X_n from the population

$$X_1, \dots, X_n \stackrel{iid}{\sim} F$$

- ▶ F is the true population distribution

Test Statistic

- ▶ How to choose between the two hypotheses?
 - ▶ Take a random sample X_1, \dots, X_n from the population

$$X_1, \dots, X_n \stackrel{iid}{\sim} F$$

- ▶ F is the true population distribution
 - ▶ How to use this sample to make the decision?

Test Statistic

- ▶ How to choose between the two hypotheses?
 - ▶ Take a random sample X_1, \dots, X_n from the population

$$X_1, \dots, X_n \stackrel{iid}{\sim} F$$

- ▶ F is the true population distribution
 - ▶ How to use this sample to make the decision?
- ▶ Let $T = T(X_1, \dots, X_n)$ be a statistic that

Test Statistic

- ▶ How to choose between the two hypotheses?
 - ▶ Take a random sample X_1, \dots, X_n from the population

$$X_1, \dots, X_n \stackrel{iid}{\sim} F$$

- ▶ F is the true population distribution
 - ▶ How to use this sample to make the decision?
- ▶ Let $T = T(X_1, \dots, X_n)$ be a statistic that
 - ▶ under the null hypothesis H_0 , typically takes *small* values

Test Statistic

- ▶ How to choose between the two hypotheses?
 - ▶ Take a random sample X_1, \dots, X_n from the population

$$X_1, \dots, X_n \stackrel{iid}{\sim} F$$

- ▶ F is the true population distribution
 - ▶ How to use this sample to make the decision?
- ▶ Let $T = T(X_1, \dots, X_n)$ be a statistic that
 - ▶ under the null hypothesis H_0 , typically takes *small* values
 - ▶ \implies large values of the statistic provide evidence against H_0

Test Statistic

- ▶ How to choose between the two hypotheses?
 - ▶ Take a random sample X_1, \dots, X_n from the population

$$X_1, \dots, X_n \stackrel{iid}{\sim} F$$

- ▶ F is the true population distribution
 - ▶ How to use this sample to make the decision?
- ▶ Let $T = T(X_1, \dots, X_n)$ be a statistic that
 - ▶ under the null hypothesis H_0 , typically takes *small* values
 - ▶ \implies large values of the statistic provide evidence against H_0
- ▶ Decision rule
 - ▶ Reject H_0 in favor of the alternative H_1 if the observed value t of T is big *enough* (exceeds a critical value)

Test Statistic

- ▶ How to choose between the two hypotheses?
 - ▶ Take a random sample X_1, \dots, X_n from the population

$$X_1, \dots, X_n \stackrel{iid}{\sim} F$$

- ▶ F is the true population distribution
 - ▶ How to use this sample to make the decision?
- ▶ Let $T = T(X_1, \dots, X_n)$ be a statistic that
 - ▶ under the null hypothesis H_0 , typically takes *small* values
 - ▶ \implies large values of the statistic provide evidence against H_0
- ▶ Decision rule
 - ▶ Reject H_0 in favor of the alternative H_1 if the observed value t of T is big *enough* (exceeds a critical value)
 - ▶ Do not reject H_0 if the observed value t of T is typical under H_0

Outline

Test Statistics

Testing a Hypothesis

Significance

p -Values

R Lab

Example: Two-Sided Test

- ▶ Let $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$ be a random sample

Example: Two-Sided Test

- ▶ Let $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$ be a random sample
- ▶ Consider the competing hypotheses

$$\left\{ \begin{array}{l} H_0 : \mu = \mu_0 \\ H_1 : \mu \neq \mu_0 \end{array} \right\}$$

Example: Two-Sided Test

- ▶ Let $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$ be a random sample
- ▶ Consider the competing hypotheses

$$\left\{ \begin{array}{l} H_0 : \mu = \mu_0 \\ H_1 : \mu \neq \mu_0 \end{array} \right\}$$

- ▶ Test statistic: Student's t statistic, $T = \frac{\bar{X} - \mu_0}{S/\sqrt{n}}$

Example: Two-Sided Test

- ▶ Let $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$ be a random sample
- ▶ Consider the competing hypotheses

$$\left\{ \begin{array}{l} H_0 : \mu = \mu_0 \\ H_1 : \mu \neq \mu_0 \end{array} \right\}$$

- ▶ Test statistic: Student's t statistic, $T = \frac{\bar{X} - \mu_0}{S/\sqrt{n}}$
- ▶ If H_0 is true, then $T \stackrel{H_0}{\sim} t_{n-1}$ and $E(T) = 0$

Example: Two-Sided Test

- ▶ Let $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$ be a random sample
- ▶ Consider the competing hypotheses

$$\left\{ \begin{array}{l} H_0 : \mu = \mu_0 \\ H_1 : \mu \neq \mu_0 \end{array} \right\}$$

- ▶ Test statistic: Student's t statistic, $T = \frac{\bar{X} - \mu_0}{S/\sqrt{n}}$
- ▶ If H_0 is true, then $T \stackrel{H_0}{\sim} t_{n-1}$ and $E(T) = 0$
 $\implies T$ typically takes values close to 0

Example: Two-Sided Test

- ▶ Let $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$ be a random sample
- ▶ Consider the competing hypotheses

$$\left\{ \begin{array}{l} H_0 : \mu = \mu_0 \\ H_1 : \mu \neq \mu_0 \end{array} \right\}$$

- ▶ Test statistic: Student's t statistic, $T = \frac{\bar{X} - \mu_0}{S/\sqrt{n}}$
- ▶ If H_0 is true, then $T \stackrel{H_0}{\sim} t_{n-1}$ and $E(T) = 0$
 $\implies T$ typically takes values close to 0
- ▶ Values of T far from 0 are considered *extreme*; occur with low probability if H_0 is true

Example: Two-Sided Test

- ▶ Let $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$ be a random sample
- ▶ Consider the competing hypotheses

$$\left\{ \begin{array}{l} H_0 : \mu = \mu_0 \\ H_1 : \mu \neq \mu_0 \end{array} \right\}$$

- ▶ Test statistic: Student's t statistic, $T = \frac{\bar{X} - \mu_0}{S/\sqrt{n}}$
- ▶ If H_0 is true, then $T \stackrel{H_0}{\sim} t_{n-1}$ and $E(T) = 0$
 $\implies T$ typically takes values close to 0
- ▶ Values of T far from 0 are considered *extreme*; occur with low probability if H_0 is true
- ▶ Reject H_0 (in favor of H_1) when $|T|$ is *big enough*

Example: Two-Sided Test

- ▶ Let $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$ be a random sample
- ▶ Consider the competing hypotheses

$$\left\{ \begin{array}{l} H_0 : \mu = \mu_0 \\ H_1 : \mu \neq \mu_0 \end{array} \right\}$$

- ▶ Test statistic: Student's t statistic, $T = \frac{\bar{X} - \mu_0}{S/\sqrt{n}}$
- ▶ If H_0 is true, then $T \stackrel{H_0}{\sim} t_{n-1}$ and $E(T) = 0$
 $\implies T$ typically takes values close to 0
- ▶ Values of T far from 0 are considered *extreme*; occur with low probability if H_0 is true
- ▶ Reject H_0 (in favor of H_1) when $|T|$ is *big enough*
Symmetric reject if $|T| > t^*$ where t^* is the critical value

Example: Two-Sided Test

- ▶ Let $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$ be a random sample
- ▶ Consider the competing hypotheses

$$\left\{ \begin{array}{l} H_0 : \mu = \mu_0 \\ H_1 : \mu \neq \mu_0 \end{array} \right\}$$

- ▶ Test statistic: Student's t statistic, $T = \frac{\bar{X} - \mu_0}{S/\sqrt{n}}$
- ▶ If H_0 is true, then $T \stackrel{H_0}{\sim} t_{n-1}$ and $E(T) = 0$
 $\implies T$ typically takes values close to 0
- ▶ Values of T far from 0 are considered *extreme*; occur with low probability if H_0 is true
- ▶ Reject H_0 (in favor of H_1) when $|T|$ is *big enough*
 Symmetric reject if $|T| > t^*$ where t^* is the critical value
 Non-symmetric reject if H_0 if $T < t_*$ or $T > t^*$

Example: One-Sided Test

- ▶ Let $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$ be a random sample

Example: One-Sided Test

- ▶ Let $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$ be a random sample
- ▶ Consider the competing hypotheses

$$\left\{ \begin{array}{l} H_0 : \mu \leq \mu_0 \\ H_1 : \mu > \mu_0 \end{array} \right\}$$

Example: One-Sided Test

- ▶ Let $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$ be a random sample
- ▶ Consider the competing hypotheses

$$\left\{ \begin{array}{l} H_0 : \mu \leq \mu_0 \\ H_1 : \mu > \mu_0 \end{array} \right\}$$

- ▶ Test statistic: Student's t statistic, $T = \frac{\bar{X} - \mu_0}{S/\sqrt{n}}$

Example: One-Sided Test

- ▶ Let $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$ be a random sample
- ▶ Consider the competing hypotheses

$$\left\{ \begin{array}{l} H_0 : \mu \leq \mu_0 \\ H_1 : \mu > \mu_0 \end{array} \right\}$$

- ▶ Test statistic: Student's t statistic, $T = \frac{\bar{X} - \mu_0}{S/\sqrt{n}}$
- ▶ Will not reject H_0 if the observed value of the test statistic $t \in H_0$

Example: One-Sided Test

- ▶ Let $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$ be a random sample
- ▶ Consider the competing hypotheses

$$\left\{ \begin{array}{l} H_0 : \mu \leq \mu_0 \\ H_1 : \mu > \mu_0 \end{array} \right\}$$

- ▶ Test statistic: Student's t statistic, $T = \frac{\bar{X} - \mu_0}{S/\sqrt{n}}$
- ▶ Will not reject H_0 if the observed value of the test statistic $t \in H_0$
 \implies Interpret $H_0 : \mu \leq \mu_0$ as $\mu = \mu_0$

Example: One-Sided Test

- ▶ Let $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$ be a random sample
- ▶ Consider the competing hypotheses

$$\left\{ \begin{array}{l} H_0 : \mu \leq \mu_0 \\ H_1 : \mu > \mu_0 \end{array} \right\}$$

- ▶ Test statistic: Student's t statistic, $T = \frac{\bar{X} - \mu_0}{S/\sqrt{n}}$
- ▶ Will not reject H_0 if the observed value of the test statistic $t \in H_0$
 \implies Interpret $H_0 : \mu \leq \mu_0$ as $\mu = \mu_0$
- ▶ Under H_0 , $T \stackrel{H_0}{\sim} t_{n-1}$ and $E(T) = 0$

Example: One-Sided Test

- ▶ Let $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$ be a random sample
- ▶ Consider the competing hypotheses

$$\left\{ \begin{array}{l} H_0 : \mu \leq \mu_0 \\ H_1 : \mu > \mu_0 \end{array} \right\}$$

- ▶ Test statistic: Student's t statistic, $T = \frac{\bar{X} - \mu_0}{S/\sqrt{n}}$
- ▶ Will not reject H_0 if the observed value of the test statistic $t \in H_0$
 \implies Interpret $H_0 : \mu \leq \mu_0$ as $\mu = \mu_0$
- ▶ Under H_0 , $T \stackrel{H_0}{\sim} t_{n-1}$ and $E(T) = 0$
 \implies Under H_0 , T typically takes values close to 0

Example: One-Sided Test

- ▶ Let $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$ be a random sample
- ▶ Consider the competing hypotheses

$$\left\{ \begin{array}{l} H_0 : \mu \leq \mu_0 \\ H_1 : \mu > \mu_0 \end{array} \right\}$$

- ▶ Test statistic: Student's t statistic, $T = \frac{\bar{X} - \mu_0}{S/\sqrt{n}}$
- ▶ Will not reject H_0 if the observed value of the test statistic $t \in H_0$
 \implies Interpret $H_0 : \mu \leq \mu_0$ as $\mu = \mu_0$
- ▶ Under H_0 , $T \stackrel{H_0}{\sim} t_{n-1}$ and $E(T) = 0$
 \implies Under H_0 , T typically takes values close to 0
- ▶ One-sided tests: values of T far from 0 are considered *extreme* only when in the direction of the alternative hypothesis H_1

Example: One-Sided Test

- ▶ Let $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$ be a random sample
- ▶ Consider the competing hypotheses

$$\left\{ \begin{array}{l} H_0 : \mu \leq \mu_0 \\ H_1 : \mu > \mu_0 \end{array} \right\}$$

- ▶ Test statistic: Student's t statistic, $T = \frac{\bar{X} - \mu_0}{S/\sqrt{n}}$
- ▶ Will not reject H_0 if the observed value of the test statistic $t \in H_0$
 \implies Interpret $H_0 : \mu \leq \mu_0$ as $\mu = \mu_0$
- ▶ Under H_0 , $T \stackrel{H_0}{\sim} t_{n-1}$ and $E(T) = 0$
 \implies Under H_0 , T typically takes values close to 0
- ▶ One-sided tests: values of T far from 0 are considered *extreme* only when in the direction of the alternative hypothesis H_1
- ▶ Reject H_0 (in favor of H_1) if $T > t^*$ where t^* is the critical value

Critical Values

- ▶ How to choose the critical value(s)?

Critical Values

- ▶ How to choose the critical value(s)?
 - ▶ What are good definitions for *big enough* and *small enough*?

Critical Values

- ▶ How to choose the critical value(s)?
 - ▶ What are good definitions for *big enough* and *small enough*?
 - ▶ At what size does a quantity become significant?

Critical Values

- ▶ How to choose the critical value(s)?
 - ▶ What are good definitions for *big enough* and *small enough*?
 - ▶ At what size does a quantity become significant?
- ▶ Two types of errors

Critical Values

- ▶ How to choose the critical value(s)?
 - ▶ What are good definitions for *big enough* and *small enough*?
 - ▶ At what size does a quantity become significant?
- ▶ Two types of errors
 - Too conservative reject H_0 only for very extreme values of T

Critical Values

- ▶ How to choose the critical value(s)?
 - ▶ What are good definitions for *big enough* and *small enough*?
 - ▶ At what size does a quantity become significant?
- ▶ Two types of errors
 - Too conservative reject H_0 only for very extreme values of T
 - Too aggressive reject H_0 even for typical values of T

Critical Values

- ▶ How to choose the critical value(s)?
 - ▶ What are good definitions for *big enough* and *small enough*?
 - ▶ At what size does a quantity become significant?
- ▶ Two types of errors

Too conservative reject H_0 only for very extreme values of T

Too aggressive reject H_0 even for typical values of T

Decision \ Truth	H_0	H_1
Do not reject H_0		Type II Error
Reject H_0	Type I Error	

Critical Values

- ▶ How to choose the critical value(s)?
 - ▶ What are good definitions for *big enough* and *small enough*?
 - ▶ At what size does a quantity become significant?

- ▶ Two types of errors

Too conservative reject H_0 only for very extreme values of T

Too aggressive reject H_0 even for typical values of T

Decision \ Truth	H_0	H_1
Do not reject H_0		Type II Error
Reject H_0	Type I Error	

- ▶ Natural asymmetry between the two hypotheses \implies type I error worse than type II

Critical Values

- ▶ How to choose the critical value(s)?
 - ▶ What are good definitions for *big enough* and *small enough*?
 - ▶ At what size does a quantity become significant?

- ▶ Two types of errors

Too conservative reject H_0 only for very extreme values of T

Too aggressive reject H_0 even for typical values of T

Decision \ Truth	H_0	H_1
Do not reject H_0		Type II Error
Reject H_0	Type I Error	

- ▶ Natural asymmetry between the two hypotheses \implies type I error worse than type II
 - ▶ For example, an email SPAM filter

Outline

Test Statistics

Testing a Hypothesis

Significance

p -Values

R Lab

Significance

- ▶ Want decision rule that limits the probability of a type I error

$$P[\text{Reject } H_0 \mid H_0 \text{ is true}] \leq \alpha$$

Significance

- ▶ Want decision rule that limits the probability of a type I error

$$P[\text{Reject } H_0 \mid H_0 \text{ is true}] \leq \alpha$$

- ▶ α is a *small* (e.g., $\alpha = 0.05$) probability

Significance

- ▶ Want decision rule that limits the probability of a type I error

$$P[\text{Reject } H_0 \mid H_0 \text{ is true}] \leq \alpha$$

- ▶ α is a *small* (e.g., $\alpha = 0.05$) probability
- ▶ α is called the *significance level* of the test

Significance

- ▶ Want decision rule that limits the probability of a type I error

$$P[\text{Reject } H_0 \mid H_0 \text{ is true}] \leq \alpha$$

- ▶ α is a *small* (e.g., $\alpha = 0.05$) probability
- ▶ α is called the *significance level* of the test
- ▶ Choose the critical value(s) to match probability of type I error

$$P[T \text{ extreme} \mid H_0] = P[\text{Reject } H_0 \mid H_0]$$

Significance

- ▶ Want decision rule that limits the probability of a type I error

$$P[\text{Reject } H_0 \mid H_0 \text{ is true}] \leq \alpha$$


- ▶ α is a *small* (e.g., $\alpha = 0.05$) probability
- ▶ α is called the *significance level* of the test
- ▶ Choose the critical value(s) to match probability of type I error

$$P[T \text{ extreme} \mid H_0] = P[\text{Reject } H_0 \mid H_0]$$



- ▶ In other words, choose the critical value so that

$$P[T \text{ more extreme than critical value(s)} \mid H_0 \text{ is true}] = \alpha$$

Rejection Regions for One and Two Sided Tests



Rejection Regions for One and Two Sided Tests



Example: Finding the Critical Values

- ▶ Let $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$ be a random sample

Example: Finding the Critical Values

- ▶ Let $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$ be a random sample
- ▶ Consider the competing hypotheses: $H_0 : \mu = \mu_0$ and $H_1 : \mu \neq \mu_0$

Example: Finding the Critical Values

- ▶ Let $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$ be a random sample
- ▶ Consider the competing hypotheses: $H_0 : \mu = \mu_0$ and $H_1 : \mu \neq \mu_0$
- ▶ Reject H_0 if $|T| = \left| \frac{\bar{X} - \mu_0}{S/\sqrt{n}} \right|$ is *big enough*, i.e., if $|T| > t^*$

Example: Finding the Critical Values

- ▶ Let $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$ be a random sample
- ▶ Consider the competing hypotheses: $H_0 : \mu = \mu_0$ and $H_1 : \mu \neq \mu_0$
- ▶ Reject H_0 if $|T| = \left| \frac{\bar{X} - \mu_0}{S/\sqrt{n}} \right|$ is *big enough*, i.e., if $|T| > t^*$
- ▶ What is the critical value with respect to the significance level?

Example: Finding the Critical Values

- ▶ Let $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$ be a random sample
- ▶ Consider the competing hypotheses: $H_0 : \mu = \mu_0$ and $H_1 : \mu \neq \mu_0$
- ▶ Reject H_0 if $|T| = \left| \frac{\bar{X} - \mu_0}{S/\sqrt{n}} \right|$ is *big enough*, i.e., if $|T| > t^*$
- ▶ What is the critical value with respect to the significance level?

$$P[\text{Reject } H_0 \mid H_0 \text{ is true}] = \alpha$$

Example: Finding the Critical Values

- ▶ Let $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$ be a random sample
- ▶ Consider the competing hypotheses: $H_0 : \mu = \mu_0$ and $H_1 : \mu \neq \mu_0$
- ▶ Reject H_0 if $|T| = \left| \frac{\bar{X} - \mu_0}{S/\sqrt{n}} \right|$ is *big enough*, i.e., if $|T| > t^*$
- ▶ What is the critical value with respect to the significance level?

$$\begin{aligned} P[\text{Reject } H_0 \mid H_0 \text{ is true}] &= \alpha \\ \Rightarrow P[|T| > t^* \mid H_0 \text{ is true}] &= \alpha \end{aligned}$$

Example: Finding the Critical Values

- ▶ Let $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$ be a random sample
- ▶ Consider the competing hypotheses: $H_0 : \mu = \mu_0$ and $H_1 : \mu \neq \mu_0$
- ▶ Reject H_0 if $|T| = \left| \frac{\bar{X} - \mu_0}{S/\sqrt{n}} \right|$ is *big enough*, i.e., if $|T| > t^*$
- ▶ What is the critical value with respect to the significance level?

$$P[\text{Reject } H_0 \mid H_0 \text{ is true}] = \alpha$$

$$\Rightarrow P[|T| > t^* \mid H_0 \text{ is true}] = \alpha$$

$$\Rightarrow P[T < t_* \text{ or } T > t^* \mid H_0 \text{ is true}] = \alpha$$

Example: Finding the Critical Values

- ▶ Let $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$ be a random sample
- ▶ Consider the competing hypotheses: $H_0 : \mu = \mu_0$ and $H_1 : \mu \neq \mu_0$
- ▶ Reject H_0 if $|T| = \left| \frac{\bar{X} - \mu_0}{S/\sqrt{n}} \right|$ is *big enough*, i.e., if $|T| > t^*$
- ▶ What is the critical value with respect to the significance level?

$$P[\text{Reject } H_0 \mid H_0 \text{ is true}] = \alpha$$

$$\Rightarrow P[|T| > t^* \mid H_0 \text{ is true}] = \alpha$$

$$\Rightarrow P[T < t_* \text{ or } T > t^* \mid H_0 \text{ is true}] = \alpha$$

$$\Rightarrow t_* = t_{\alpha/2} \quad \& \quad t^* = t_{1-\alpha/2}$$

Example: Finding the Critical Values

- ▶ Let $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$ be a random sample
- ▶ Consider the competing hypotheses: $H_0 : \mu = \mu_0$ and $H_1 : \mu \neq \mu_0$
- ▶ Reject H_0 if $|T| = \left| \frac{\bar{X} - \mu_0}{S/\sqrt{n}} \right|$ is *big enough*, i.e., if $|T| > t^*$
- ▶ What is the critical value with respect to the significance level?

$$P[\text{Reject } H_0 \mid H_0 \text{ is true}] = \alpha$$

$$\Rightarrow P[|T| > t^* \mid H_0 \text{ is true}] = \alpha$$

$$\Rightarrow P[T < t_* \text{ or } T > t^* \mid H_0 \text{ is true}] = \alpha$$

$$\Rightarrow t_* = t_{\alpha/2} \quad \& \quad t^* = t_{1-\alpha/2}$$

- ▶ t_γ is the γ quantile of the t_{n-1} distribution

Significance

- ▶ Given
 1. a pair of complementary hypotheses $\{H_0, H_1\}$

Significance

- ▶ Given
 1. a pair of complementary hypotheses $\{H_0, H_1\}$
 2. a test statistic T

Significance

- ▶ Given
 1. a pair of complementary hypotheses $\{H_0, H_1\}$
 2. a test statistic T
 3. a significance level α

Significance

- ▶ Given
 1. a pair of complementary hypotheses $\{H_0, H_1\}$
 2. a test statistic T
 3. a significance level α
- ▶ Reject the null hypothesis H_0 in favor of the alternative hypothesis H_1 at significance level α if T is more extreme than the critical value(s) computed from the significance level α

Significance

- ▶ Given
 1. a pair of complementary hypotheses $\{H_0, H_1\}$
 2. a test statistic T
 3. a significance level α
- ▶ Reject the null hypothesis H_0 in favor of the alternative hypothesis H_1 at significance level α if T is more extreme than the critical value(s) computed from the significance level α
- ▶ **Example** (Two-sided test for a normal population)

Reject the null hypothesis $H_0 : \mu = \mu_0$ in favor of the alternative hypothesis $H_1 : \mu \neq \mu_0$ at significance level α if the observed value t of T satisfies $t \geq t_{1-\alpha/2}$ or $t \leq t_{\alpha/2}$

Outline

Test Statistics

Testing a Hypothesis

Significance

p -Values

R Lab

p Values

- ▶ Instead of computing critical values for a given significance level α , might ask what is the highest significance level where H_0 would be rejected?

p Values

- ▶ Instead of computing critical values for a given significance level α , might ask what is the highest significance level where H_0 would be rejected?
- ▶ This level is called the observed p value

p Values

- ▶ Instead of computing critical values for a given significance level α , might ask what is the highest significance level where H_0 would be rejected?
- ▶ This level is called the observed p value
- ▶ The p value is the probability that the test statistic T takes a value equal to or more extreme than its observed value t

p Values

- ▶ Instead of computing critical values for a given significance level α , might ask what is the highest significance level where H_0 would be rejected?
- ▶ This level is called the observed p value
- ▶ The p value is the probability that the test statistic T takes a value equal to or more extreme than its observed value t
- ▶ Observe $T = t$
 - ▶ Two-sided test: $p = P[|T| \geq t | H_0]$

p Values

- ▶ Instead of computing critical values for a given significance level α , might ask what is the highest significance level where H_0 would be rejected?
- ▶ This level is called the observed p value
- ▶ The p value is the probability that the test statistic T takes a value equal to or more extreme than its observed value t
- ▶ Observe $T = t$
 - ▶ Two-sided test: $p = P[|T| \geq t | H_0]$
 - ▶ One-sided test: $p = P[T \geq t | H_0]$

p Values

- ▶ Instead of computing critical values for a given significance level α , might ask what is the highest significance level where H_0 would be rejected?
- ▶ This level is called the observed p value
- ▶ The p value is the probability that the test statistic T takes a value equal to or more extreme than its observed value t
- ▶ Observe $T = t$
 - ▶ Two-sided test: $p = P[|T| \geq t | H_0]$
 - ▶ One-sided test: $p = P[T \geq t | H_0]$
- ▶ Small values of p provide evidence against the null hypothesis H_0 (i.e., outcome is unlikely under the null hypothesis)

p Values

- ▶ Instead of computing critical values for a given significance level α , might ask what is the highest significance level where H_0 would be rejected?
- ▶ This level is called the observed p value
- ▶ The p value is the probability that the test statistic T takes a value equal to or more extreme than its observed value t
- ▶ Observe $T = t$
 - ▶ Two-sided test: $p = P[|T| \geq t | H_0]$
 - ▶ One-sided test: $p = P[T \geq t | H_0]$
- ▶ Small values of p provide evidence against the null hypothesis H_0 (i.e., outcome is unlikely under the null hypothesis)
- ▶ Still have to decide what values of p are *small enough*

p Values

- ▶ How to decide whether to reject the null hypothesis H_0 using the observed p value?

p Values

- ▶ How to decide whether to reject the null hypothesis H_0 using the observed p value?
- ▶ Interpret the p value on the same scale as the significance level

p Values

- ▶ How to decide whether to reject the null hypothesis H_0 using the observed p value?
- ▶ Interpret the p value on the same scale as the significance level
- ▶ Let α be the maximum allowable probability of type I error

$$\begin{aligned} P[\text{Reject } H_0 \mid H_0 \text{ is true}] &\leq \alpha \\ p &\leq \alpha \end{aligned}$$

(for example: $\alpha = 0.05$ or $\alpha = 0.01$)

p Values

- ▶ How to decide whether to reject the null hypothesis H_0 using the observed p value?
- ▶ Interpret the p value on the same scale as the significance level
- ▶ Let α be the maximum allowable probability of type I error

$$\begin{aligned} P[\text{Reject } H_0 \mid H_0 \text{ is true}] &\leq \alpha \\ p &\leq \alpha \end{aligned}$$

(for example: $\alpha = 0.05$ or $\alpha = 0.01$)

- ▶ The decision rule becomes: reject H_0 if $p \leq \alpha$

p Values

- ▶ How to decide whether to reject the null hypothesis H_0 using the observed p value?
- ▶ Interpret the p value on the same scale as the significance level
- ▶ Let α be the maximum allowable probability of type I error

$$\begin{aligned} P[\text{Reject } H_0 \mid H_0 \text{ is true}] &\leq \alpha \\ p &\leq \alpha \end{aligned}$$

(for example: $\alpha = 0.05$ or $\alpha = 0.01$)

- ▶ The decision rule becomes: reject H_0 if $p \leq \alpha$
- ▶ The probability of type I error using this rule is exactly α

p Values

- ▶ How to decide whether to reject the null hypothesis H_0 using the observed p value?
- ▶ Interpret the p value on the same scale as the significance level
- ▶ Let α be the maximum allowable probability of type I error

$$\begin{aligned} P[\text{Reject } H_0 \mid H_0 \text{ is true}] &\leq \alpha \\ p &\leq \alpha \end{aligned}$$

(for example: $\alpha = 0.05$ or $\alpha = 0.01$)

- ▶ The decision rule becomes: reject H_0 if $p \leq \alpha$
- ▶ The probability of type I error using this rule is exactly α
- ▶ As a decision rule, equivalent to the critical value(s) decision rule

The Elements of a Hypothesis Test

- ▶ A *null hypothesis* H_0 to test against an *alternative hypothesis* H_1

The Elements of a Hypothesis Test

- ▶ A *null hypothesis* H_0 to test against an *alternative hypothesis* H_1
- ▶ A *test statistic* T chosen so that extreme values of T (in the direction of H_1) suggest that H_0 is false

The Elements of a Hypothesis Test

- ▶ A *null hypothesis* H_0 to test against an *alternative hypothesis* H_1
- ▶ A *test statistic* T chosen so that extreme values of T (in the direction of H_1) suggest that H_0 is false
- ▶ The observed value of the test statistic is $T = t$

The Elements of a Hypothesis Test

- ▶ A *null hypothesis* H_0 to test against an *alternative hypothesis* H_1
- ▶ A *test statistic* T chosen so that extreme values of T (in the direction of H_1) suggest that H_0 is false
- ▶ The observed value of the test statistic is $T = t$
- ▶ A significance level α specifying the maximum probability of type I error (rejecting H_0 when H_0 is true)

The Elements of a Hypothesis Test

- ▶ A *null hypothesis* H_0 to test against an *alternative hypothesis* H_1
- ▶ A *test statistic* T chosen so that extreme values of T (in the direction of H_1) suggest that H_0 is false
- ▶ The observed value of the test statistic is $T = t$
- ▶ A significance level α specifying the maximum probability of type I error (rejecting H_0 when H_0 is true)
- ▶ A decision rule:
Critical value(s) Reject the null hypothesis H_0 in favor of the alternative hypothesis H_1 when T is more extreme.
Critical value(s) chosen with respect to significance level α

The Elements of a Hypothesis Test

- ▶ A *null hypothesis* H_0 to test against an *alternative hypothesis* H_1
- ▶ A *test statistic* T chosen so that extreme values of T (in the direction of H_1) suggest that H_0 is false
- ▶ The observed value of the test statistic is $T = t$
- ▶ A significance level α specifying the maximum probability of type I error (rejecting H_0 when H_0 is true)
- ▶ A decision rule:
 - Critical value(s) Reject the null hypothesis H_0 in favor of the alternative hypothesis H_1 when T is more extreme. Critical value(s) chosen with respect to significance level α
 - p value The probability of observing a value of T as extreme or more extreme than t under the null hypothesis H_0 . Reject the null hypothesis H_0 in favor of the alternative hypothesis H_1 when $p \leq \alpha$

The Test Statistic T

- ▶ Test statistic: free to choose any function subject to the qualitative constraint that it is *extreme* when H_0 is false

The Test Statistic T

- ▶ Test statistic: free to choose any function subject to the qualitative constraint that it is *extreme* when H_0 is false
- ▶ The choice of T depends on the alternative hypothesis H_1 , whatever we imagine possible if H_0 is false

The Test Statistic T

- ▶ Test statistic: free to choose any function subject to the qualitative constraint that it is *extreme* when H_0 is false
- ▶ The choice of T depends on the alternative hypothesis H_1 , whatever we imagine possible if H_0 is false
- ▶ The more precise H_1 can be stated, the more important it is to choose an appropriate test statistic T

The Constant Expected Returns (CER) Model

Assumptions:

- ▶ Let $r_t = \log \left(\frac{P_t}{P_{t-1}} \right)$ denote the continuously compounded return on an asset

The Constant Expected Returns (CER) Model

Assumptions:

- ▶ Let $r_t = \log \left(\frac{P_t}{P_{t-1}} \right)$ denote the continuously compounded return on an asset
- ▶ $E(r_t) = \mu$ and $\text{Var}(r_t) = \sigma^2$ for all t

The Constant Expected Returns (CER) Model

Assumptions:

- ▶ Let $r_t = \log \left(\frac{P_t}{P_{t-1}} \right)$ denote the continuously compounded return on an asset
- ▶ $E(r_t) = \mu$ and $\text{Var}(r_t) = \sigma^2$ for all t
- ▶ Normality of returns: $r_t \sim \mathcal{N}(\mu, \sigma^2)$ for all t

The Constant Expected Returns (CER) Model

Assumptions:

- ▶ Let $r_t = \log \left(\frac{P_t}{P_{t-1}} \right)$ denote the continuously compounded return on an asset
- ▶ $E(r_t) = \mu$ and $\text{Var}(r_t) = \sigma^2$ for all t
- ▶ Normality of returns: $r_t \sim \mathcal{N}(\mu, \sigma^2)$ for all t
- ▶ No serial correlation: $\text{Cov}(r_s, r_t) = 0$ for $t \neq s$
 $\implies R_s$ and R_t are independent

Sampling from Normal Populations

One Sample t -test The hypotheses are statements about the mean of a single normal population

Sampling from Normal Populations

One Sample t -test The hypotheses are statements about the mean of a single normal population

Paired t -test A one sample t -test for the difference between two random variables

- Requires that X_i must somehow correspond to Y_i

Sampling from Normal Populations

One Sample t -test The hypotheses are statements about the mean of a single normal population

Paired t -test A one sample t -test for the difference between two random variables

- ▶ Requires that X_i must somehow correspond to Y_i

Two Sample t -test The hypotheses are statements about the means of two separate normal populations, many variations:

- ▶ Variance known: $\sigma_1^2 = \sigma_2^2$ or $\sigma_1^2 \neq \sigma_2^2$
- ▶ Equal variance (unknown): $\sigma_1^2 = \sigma_2^2$ estimated by pooled S^2
- ▶ Different variances (unknown): σ_1^2 estimated by S_1^2 and σ_2^2 estimated by S_2^2

Outline

Test Statistics

Testing a Hypothesis

Significance

p -Values

R Lab

```
> library(quantmod)
```



```
> library(quantmod)

> getSymbols("^RUT")  #Russell 2000
> getSymbols("IWL")   #Russell Top 200
```

```
> library(quantmod)

> getSymbols("^RUT")    #Russell 2000
> getSymbols("IWL")     #Russell Top 200

> head(RUT)
```

```
> library(quantmod)

> getSymbols("^RUT")  #Russell 2000
> getSymbols("IWL")   #Russell Top 200

> head(RUT)
```

	RUT.Open	RUT.High	RUT.Low	RUT.Close	RUT.Adjusted
2007-01-03	788.31	796.62	779.31	787.42	787.42
2007-01-04	786.42	791.83	779.70	789.95	789.95
2007-01-05	787.70	787.70	774.55	775.87	775.87
2007-01-08	776.20	778.83	769.27	776.99	776.99
2007-01-09	777.10	778.74	768.69	778.33	778.33
2007-01-10	775.78	779.45	773.14	778.87	778.87

```

> library(quantmod)

> getSymbols("^RUT")  #Russell 2000
> getSymbols("IWL")   #Russell Top 200

> head(RUT)

```

	RUT.Open	RUT.High	RUT.Low	RUT.Close	RUT.Adjusted
2007-01-03	788.31	796.62	779.31	787.42	787.42
2007-01-04	786.42	791.83	779.70	789.95	789.95
2007-01-05	787.70	787.70	774.55	775.87	775.87
2007-01-08	776.20	778.83	769.27	776.99	776.99
2007-01-09	777.10	778.74	768.69	778.33	778.33
2007-01-10	775.78	779.45	773.14	778.87	778.87

```

> R2000 <- monthlyReturn(RUT["2013"]$RUT.Adjusted)
> RT200 <- monthlyReturn(IWL["2013"]$IWL.Adjusted)

```

```
> library(quantmod)


> getSymbols("^RUT")    #Russell 2000
> getSymbols("IWL")     #Russell Top 200

> head(RUT)
```

	RUT.Open	RUT.High	RUT.Low	RUT.Close	RUT.Adjusted
2007-01-03	788.31	796.62	779.31	787.42	787.42
2007-01-04	786.42	791.83	779.70	789.95	789.95
2007-01-05	787.70	787.70	774.55	775.87	775.87
2007-01-08	776.20	778.83	769.27	776.99	776.99
2007-01-09	777.10	778.74	768.69	778.33	778.33
2007-01-10	775.78	779.45	773.14	778.87	778.87


```
> R2000 <- monthlyReturn(RUT["2013"]$RUT.Adjusted)
> RT200 <- monthlyReturn(IWL["2013"]$IWL.Adjusted)

> plot(density(R2000))
> lines(density(RT200), lty = 2)
```



```
> plot(density(R2000, bw = 0.015))  
> lines(density(RT200, bw = 0.0175), lty = 2)
```

```
> plot(density(R2000, bw = 0.015))  
> lines(density(RT200, bw = 0.0175), lty = 2)
```



A Paired Two-Sample t Test

- ▶ Plain English: Are small cap returns larger than large cap returns?

A Paired Two-Sample t Test

- ▶ Plain English: Are small cap returns larger than large cap returns?
- ▶ Statistics: Is the mean (μ_s) of the small cap returns distribution greater than the mean (μ_l) of the large cap returns distribution?

A Paired Two-Sample t Test

- ▶ Plain English: Are small cap returns larger than large cap returns?
- ▶ Statistics: Is the mean (μ_s) of the small cap returns distribution greater than the mean (μ_l) of the large cap returns distribution?
- ▶ Hypothesis Test: $H_0: \mu_s \leq \mu_l$ vs. $H_1: \mu_s > \mu_l$?

A Paired Two-Sample t Test

- ▶ Plain English: Are small cap returns larger than large cap returns?
- ▶ Statistics: Is the mean (μ_s) of the small cap returns distribution greater than the mean (μ_l) of the large cap returns distribution?
- ▶ Hypothesis Test: $H_0: \mu_s \leq \mu_l$ vs. $H_1: \mu_s > \mu_l$?
- ▶ Paired Test: $H_0: \delta_\mu = \mu_s - \mu_l \leq 0$ vs. $H_1: \delta_\mu = \mu_s - \mu_l > 0$?

A Paired Two-Sample t Test

- ▶ Plain English: Are small cap returns larger than large cap returns?
 - ▶ Statistics: Is the mean (μ_s) of the small cap returns distribution greater than the mean (μ_l) of the large cap returns distribution?
 - ▶ Hypothesis Test: $H_0: \mu_s \leq \mu_l$ vs. $H_1: \mu_s > \mu_l$?
 - ▶ Paired Test: $H_0: \delta_\mu = \mu_s - \mu_l \leq 0$ vs. $H_1: \delta_\mu = \mu_s - \mu_l > 0$?
- ```
> d <- drop(coredata(R2000 - RT200))
```

# A Paired Two-Sample $t$ Test

- ▶ Plain English: Are small cap returns larger than large cap returns?
  - ▶ Statistics: Is the mean ( $\mu_s$ ) of the small cap returns distribution greater than the mean ( $\mu_l$ ) of the large cap returns distribution?
  - ▶ Hypothesis Test:  $H_0: \mu_s \leq \mu_l$  vs.  $H_1: \mu_s > \mu_l$ ?
  - ▶ Paired Test:  $H_0: \delta_\mu = \mu_s - \mu_l \leq 0$  vs.  $H_1: \delta_\mu = \mu_s - \mu_l > 0$ ?
- ```
> d <- drop(coredata(R2000 - RT200))  
> S2 <- var(d)
```

A Paired Two-Sample t Test

- ▶ Plain English: Are small cap returns larger than large cap returns?
 - ▶ Statistics: Is the mean (μ_s) of the small cap returns distribution greater than the mean (μ_l) of the large cap returns distribution?
 - ▶ Hypothesis Test: $H_0: \mu_s \leq \mu_l$ vs. $H_1: \mu_s > \mu_l$?
 - ▶ Paired Test: $H_0: \delta_\mu = \mu_s - \mu_l \leq 0$ vs. $H_1: \delta_\mu = \mu_s - \mu_l > 0$?
- ```
> d <- drop(coredata(R2000 - RT200))
> S2 <- var(d)
> mean(d) / sqrt(S2 / 12)
```

# A Paired Two-Sample $t$ Test

- ▶ Plain English: Are small cap returns larger than large cap returns?
- ▶ Statistics: Is the mean ( $\mu_s$ ) of the small cap returns distribution greater than the mean ( $\mu_l$ ) of the large cap returns distribution?
- ▶ Hypothesis Test:  $H_0: \mu_s \leq \mu_l$  vs.  $H_1: \mu_s > \mu_l$ ?
- ▶ Paired Test:  $H_0: \delta_\mu = \mu_s - \mu_l \leq 0$  vs.  $H_1: \delta_\mu = \mu_s - \mu_l > 0$ ?

```
> d <- drop(coredata(R2000 - RT200))
```

```
> S2 <- var(d)
```

```
> mean(d) / sqrt(S2 / 12)
```

```
[1] 0.5970462
```



# A Paired Two-Sample $t$ Test

- ▶ Plain English: Are small cap returns larger than large cap returns?
- ▶ Statistics: Is the mean ( $\mu_s$ ) of the small cap returns distribution greater than the mean ( $\mu_l$ ) of the large cap returns distribution?
- ▶ Hypothesis Test:  $H_0: \mu_s \leq \mu_l$  vs.  $H_1: \mu_s > \mu_l$ ?
- ▶ Paired Test:  $H_0: \delta_\mu = \mu_s - \mu_l \leq 0$  vs.  $H_1: \delta_\mu = \mu_s - \mu_l > 0$ ?

```
> d <- drop(coredata(R2000 - RT200))
```

```
> S2 <- var(d)
```

```
> mean(d) / sqrt(S2 / 12)
```

```
[1] 0.5970462
```

```
> qt(0.95, df = 11)
```

# A Paired Two-Sample $t$ Test

- ▶ Plain English: Are small cap returns larger than large cap returns?
- ▶ Statistics: Is the mean ( $\mu_s$ ) of the small cap returns distribution greater than the mean ( $\mu_l$ ) of the large cap returns distribution?
- ▶ Hypothesis Test:  $H_0: \mu_s \leq \mu_l$  vs.  $H_1: \mu_s > \mu_l$ ?
- ▶ Paired Test:  $H_0: \delta_\mu = \mu_s - \mu_l \leq 0$  vs.  $H_1: \delta_\mu = \mu_s - \mu_l > 0$ ?

```
> d <- drop(coredata(R2000 - RT200))
```

```
> S2 <- var(d)
```

```
> mean(d) / sqrt(S2 / 12)
```

```
[1] 0.5970462
```

```
> qt(0.95, df = 11)
```

```
[1] 1.795885
```

# A Paired Two-Sample $t$ Test

- ▶ Plain English: Are small cap returns larger than large cap returns?
- ▶ Statistics: Is the mean ( $\mu_s$ ) of the small cap returns distribution greater than the mean ( $\mu_l$ ) of the large cap returns distribution?
- ▶ Hypothesis Test:  $H_0: \mu_s \leq \mu_l$  vs.  $H_1: \mu_s > \mu_l$ ?
- ▶ Paired Test:  $H_0: \delta_\mu = \mu_s - \mu_l \leq 0$  vs.  $H_1: \delta_\mu = \mu_s - \mu_l > 0$ ?

```
> d <- drop(coredata(R2000 - RT200))
```

```
> S2 <- var(d)
```

```
> mean(d) / sqrt(S2 / 12)
```

```
[1] 0.5970462
```

```
> qt(0.95, df = 11)
```

```
[1] 1.795885
```

```
> 1 - pt(0.5970462, df = 11)
```

# A Paired Two-Sample $t$ Test

- ▶ Plain English: Are small cap returns larger than large cap returns?
- ▶ Statistics: Is the mean ( $\mu_s$ ) of the small cap returns distribution greater than the mean ( $\mu_l$ ) of the large cap returns distribution?
- ▶ Hypothesis Test:  $H_0: \mu_s \leq \mu_l$  vs.  $H_1: \mu_s > \mu_l$ ?
- ▶ Paired Test:  $H_0: \delta_\mu = \mu_s - \mu_l \leq 0$  vs.  $H_1: \delta_\mu = \mu_s - \mu_l > 0$ ?

```
> d <- drop(coredata(R2000 - RT200))
```

```
> S2 <- var(d)
```

```
> mean(d) / sqrt(S2 / 12)
```

```
[1] 0.5970462
```

```
> qt(0.95, df = 11)
```

```
[1] 1.795885
```

```
> 1 - pt(0.5970462, df = 11)
```

```
[1] 0.281282
```

## A Paired Two-Sample $t$ Test in R

```
> t.test(R2000, RT200, alternative = "greater", paired = TRUE)
```

## A Paired Two-Sample $t$ Test in R

```
> t.test(R2000, RT200, alternative = "greater", paired = TRUE)
```

Paired t-test

data: R2000 and RT200

$t = 0.597$ ,  $df = 11$ ,  $p\text{-value} = 0.2813$

alternative hypothesis: true difference in means is  
greater than 0

95 percent confidence interval:

-0.006060183                      Inf

sample estimates:

mean of the differences

0.003018096



# COMPUTATIONAL FINANCE & RISK MANAGEMENT

---

UNIVERSITY *of* WASHINGTON

Department of Applied Mathematics

<http://computational-finance.uw.edu>