



COMPUTATIONAL FINANCE & RISK MANAGEMENT

---

UNIVERSITY *of* WASHINGTON

Department of Applied Mathematics

# CFRM 410: Probability and Statistics for Computational Finance

## Week 9 Estimation

Jake Price

Instructor, Computational Finance and Risk Management

University of Washington

Slides originally produced by Kjell Konis

# Fundamental Ideas of Statistics

Statistical Models

Point Estimation

Evaluating Point Estimators

Interval Estimators

# Outline

## Statistical Models

## Point Estimation

## Evaluating Point Estimators

## Interval Estimators

# Statistical Models

Goal: learn something about a population by looking at a subset of the population called a *sample*

# Statistical Models

Goal: learn something about a population by looking at a subset of the population called a *sample*

**Probability Model** Density  $f_{X|\theta}$  is assumed to be a member of a known parametric family (parameter  $\theta$  is unknown)

# Statistical Models

Goal: learn something about a population by looking at a subset of the population called a *sample*

**Probability Model** Density  $f_{X|\theta}$  is assumed to be a member of a known parametric family (parameter  $\theta$  is unknown)

**Data Set** Data  $x_1, \dots, x_n$  are assumed to be a realization of the random sample  $X_1, \dots, X_n \stackrel{iid}{\sim} f_{X|\theta}$  ( $\Rightarrow$  representative)

# Statistical Models

Goal: learn something about a population by looking at a subset of the population called a *sample*

**Probability Model** Density  $f_{X|\theta}$  is assumed to be a member of a known parametric family (parameter  $\theta$  is unknown)

**Data Set** Data  $x_1, \dots, x_n$  are assumed to be a realization of the random sample  $X_1, \dots, X_n \stackrel{iid}{\sim} f_{X|\theta}$  ( $\Rightarrow$  representative)

**Statistic** Random variable  $T = t(X_1, \dots, X_n)$

# Statistical Models

Goal: learn something about a population by looking at a subset of the population called a *sample*

**Probability Model** Density  $f_{X|\theta}$  is assumed to be a member of a known parametric family (parameter  $\theta$  is unknown)

**Data Set** Data  $x_1, \dots, x_n$  are assumed to be a realization of the random sample  $X_1, \dots, X_n \stackrel{iid}{\sim} f_{X|\theta}$  ( $\Rightarrow$  representative)

**Statistic** Random variable  $T = t(X_1, \dots, X_n)$

**Estimator** Statistic used to estimate the parameter  $\theta$



# Statistical Models

Goal: learn something about a population by looking at a subset of the population called a *sample*

**Probability Model** Density  $f_{X|\theta}$  is assumed to be a member of a known parametric family (parameter  $\theta$  is unknown)

**Data Set** Data  $x_1, \dots, x_n$  are assumed to be a realization of the random sample  $X_1, \dots, X_n \stackrel{iid}{\sim} f_{X|\theta}$  ( $\Rightarrow$  representative)

**Statistic** Random variable  $T = t(X_1, \dots, X_n)$

**Estimator** Statistic used to estimate the parameter  $\theta$

**Notation**  $T = t(X_1, \dots, X_n)$       **statistic** (a random variable)

# Statistical Models

Goal: learn something about a population by looking at a subset of the population called a *sample*

**Probability Model** Density  $f_{X|\theta}$  is assumed to be a member of a known parametric family (parameter  $\theta$  is unknown)

**Data Set** Data  $x_1, \dots, x_n$  are assumed to be a realization of the random sample  $X_1, \dots, X_n \stackrel{iid}{\sim} f_{X|\theta}$  ( $\Rightarrow$  representative)

**Statistic** Random variable  $T = t(X_1, \dots, X_n)$

**Estimator** Statistic used to estimate the parameter  $\theta$

|                 |                          |  |
|-----------------|--------------------------|--|
| <b>Notation</b> | $T = t(X_1, \dots, X_n)$ | <b>statistic</b> (a random variable)       |
|                 | $t = t(x_1, \dots, x_n)$ | <b>observed</b> value of the statistic $T$ |

# Statistical Models

Goal: learn something about a population by looking at a subset of the population called a *sample*

**Probability Model** Density  $f_{X|\theta}$  is assumed to be a member of a known parametric family (parameter  $\theta$  is unknown)

**Data Set** Data  $x_1, \dots, x_n$  are assumed to be a realization of the random sample  $X_1, \dots, X_n \stackrel{iid}{\sim} f_{X|\theta}$  ( $\Rightarrow$  representative)

**Statistic** Random variable  $T = t(X_1, \dots, X_n)$

**Estimator** Statistic used to estimate the parameter  $\theta$

|                 |                                 |  |
|-----------------|---------------------------------|--|
| <b>Notation</b> | $T = t(X_1, \dots, X_n)$        | <b>statistic</b> (a random variable)       |
|                 | $t = t(x_1, \dots, x_n)$        | <b>observed</b> value of the statistic $T$ |
|                 | $\hat{\theta}(X_1, \dots, X_n)$ | <b>estimator</b> of the parameter $\theta$ |

# Statistical Models

Goal: learn something about a population by looking at a subset of the population called a *sample*

**Probability Model** Density  $f_{X|\theta}$  is assumed to be a member of a known parametric family (parameter  $\theta$  is unknown)

**Data Set** Data  $x_1, \dots, x_n$  are assumed to be a realization of the random sample  $X_1, \dots, X_n \stackrel{iid}{\sim} f_{X|\theta}$  ( $\Rightarrow$  representative)

**Statistic** Random variable  $T = t(X_1, \dots, X_n)$

**Estimator** Statistic used to estimate the parameter  $\theta$

|                 |  |  |
|-----------------|--|--|
| <b>Notation</b> | $T = t(X_1, \dots, X_n)$                       | <b>statistic</b> (a random variable)       |
|                 | $t = t(x_1, \dots, x_n)$                       | <b>observed</b> value of the statistic $T$ |
|                 | $\hat{\theta}(X_1, \dots, X_n)$                | <b>estimator</b> of the parameter $\theta$ |
|                 | $\hat{\theta} = \hat{\theta}(x_1, \dots, x_n)$ | <b>estimate</b> of the parameter $\theta$  |

# Statistical Models

**Example** Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$

# Statistical Models

**Example** Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$

- ▶  $\hat{\mu} = \bar{X}$  is an estimator of  $\mu$

# Statistical Models

**Example** Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$

- ▶  $\hat{\mu} = \bar{X}$  is an estimator of  $\mu$
- ▶ The observed value of the estimator  $\hat{\mu}$  is  $\bar{x}$

# Statistical Models

**Example** Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$

- ▶  $\hat{\mu} = \bar{X}$  is an estimator of  $\mu$
- ▶ The observed value of the estimator  $\hat{\mu}$  is  $\bar{x}$
- ▶  $\hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2$  is an estimator of  $\sigma^2$



# Statistical Models

**Example** Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$

- ▶  $\hat{\mu} = \bar{X}$  is an estimator of  $\mu$
- ▶ The observed value of the estimator  $\hat{\mu}$  is  $\bar{x}$
- ▶  $\hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2$  is an estimator of  $\sigma^2$
- ▶ The observed value of the estimator  $\hat{\sigma}^2$  is  $\sum_{i=1}^n (x_i - \bar{x})^2$

# Statistical Models

**Example** Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$

- ▶  $\hat{\mu} = \bar{X}$  is an estimator of  $\mu$
- ▶ The observed value of the estimator  $\hat{\mu}$  is  $\bar{x}$
- ▶  $\hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2$  is an estimator of  $\sigma^2$
- ▶ The observed value of the estimator  $\hat{\sigma}^2$  is  $\sum_{i=1}^n (x_i - \bar{x})^2$

## Remarks

- ▶ Since a statistic  $T$  is a function of the random variables  $X_1, \dots, X_n$ ,  $T$  is itself a random variable

# Statistical Models

**Example** Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$

- ▶  $\hat{\mu} = \bar{X}$  is an estimator of  $\mu$
- ▶ The observed value of the estimator  $\hat{\mu}$  is  $\bar{x}$
- ▶  $\hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2$  is an estimator of  $\sigma^2$
- ▶ The observed value of the estimator  $\hat{\sigma}^2$  is  $\sum_{i=1}^n (x_i - \bar{x})^2$

## Remarks

- ▶ Since a statistic  $T$  is a function of the random variables  $X_1, \dots, X_n$ ,  $T$  is itself a random variable
- ▶ The distribution of  $T$  depends on the density of the  $X_i$  and is called the sampling distribution of  $T$

# Statistical Models

**Example** Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$

- ▶  $\hat{\mu} = \bar{X}$  is an estimator of  $\mu$
- ▶ The observed value of the estimator  $\hat{\mu}$  is  $\bar{x}$
- ▶  $\hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2$  is an estimator of  $\sigma^2$
- ▶ The observed value of the estimator  $\hat{\sigma}^2$  is  $\sum_{i=1}^n (x_i - \bar{x})^2$

## Remarks

- ▶ Since a statistic  $T$  is a function of the random variables  $X_1, \dots, X_n$ ,  $T$  is itself a random variable
- ▶ The distribution of  $T$  depends on the density of the  $X_i$  and is called the sampling distribution of  $T$
- ▶  $E(X)$  and  $\text{Var}(X)$  provide partial information on the distribution of  $T$  and are particularly useful when the distribution of  $T$  can be approximated

# Problems to Study

Given a statistical model, might wish to:

# Problems to Study

Given a statistical model, might wish to:

- ▶ Estimate the parameters of the model

# Problems to Study

Given a statistical model, might wish to:

- ▶ Estimate the parameters of the model
- ▶ Answer questions about the parameters of the model: for example, does  $\theta = 0$ ?

# Problems to Study

Given a statistical model, might wish to:

- ▶ Estimate the parameters of the model
- ▶ Answer questions about the parameters of the model: for example, does  $\theta = 0$ ?
- ▶ Make predictions about future observations



# Problems to Study

Given a statistical model, might wish to:

- ▶ Estimate the parameters of the model
- ▶ Answer questions about the parameters of the model: for example, does  $\theta = 0$ ?
- ▶ Make predictions about future observations

There are several methods to estimate parameters, will discuss:

# Problems to Study

Given a statistical model, might wish to:

- ▶ Estimate the parameters of the model
- ▶ Answer questions about the parameters of the model: for example, does  $\theta = 0$ ?
- ▶ Make predictions about future observations

There are several methods to estimate parameters, will discuss:

- ▶ The method of *moments* (easy - almost always works)

# Problems to Study

Given a statistical model, might wish to:

- ▶ Estimate the parameters of the model
- ▶ Answer questions about the parameters of the model: for example, does  $\theta = 0$ ?
- ▶ Make predictions about future observations

There are several methods to estimate parameters, will discuss:

- ▶ The method of *moments* (easy - almost always works)
- ▶ The method of *least squares* (easy)

# Problems to Study

Given a statistical model, might wish to:

- ▶ Estimate the parameters of the model
- ▶ Answer questions about the parameters of the model: for example, does  $\theta = 0$ ?
- ▶ Make predictions about future observations

There are several methods to estimate parameters, will discuss:

- ▶ The method of *moments* (easy - almost always works)
- ▶ The method of *least squares* (easy)
- ▶ The method of *maximum likelihood* (more general - optimal in many situations)

# Outline

Statistical Models

Point Estimation

Evaluating Point Estimators

Interval Estimators

# Method of Moments

Let  $X_1, \dots, X_n$  be a representative random sample from a population characterized by a distribution with density  $f_{X|\theta}$

# Method of Moments

Let  $X_1, \dots, X_n$  be a representative random sample from a population characterized by a distribution with density  $f_{X|\theta}$

To compute the method of moments estimator, equate the *empirical* moments (calculated from the sample) and the *theoretical* moments (calculated from the population, i.e,  $f_{X|\theta}$ )

# Method of Moments

Let  $X_1, \dots, X_n$  be a representative random sample from a population characterized by a distribution with density  $f_{X|\theta}$

To compute the method of moments estimator, equate the *empirical* moments (calculated from the sample) and the *theoretical* moments (calculated from the population, i.e,  $f_{X|\theta}$ )

The  $k^{th}$  moments are:

Theoretical  $M'_k = E(X^k)$



# Method of Moments

Let  $X_1, \dots, X_n$  be a representative random sample from a population characterized by a distribution with density  $f_{X|\theta}$

To compute the method of moments estimator, equate the *empirical* moments (calculated from the sample) and the *theoretical* moments (calculated from the population, i.e,  $f_{X|\theta}$ )

The  $k^{th}$  moments are:

Theoretical  $M'_k = E(X^k)$

Empirical  $m'_k = \frac{1}{n} \sum_{i=1}^k X_i^k$

# Method of Moments

Let  $X_1, \dots, X_n$  be a representative random sample from a population characterized by a distribution with density  $f_{X|\theta}$

To compute the method of moments estimator, equate the *empirical* moments (calculated from the sample) and the *theoretical* moments (calculated from the population, i.e,  $f_{X|\theta}$ )

The  $k^{th}$  moments are:

Theoretical  $M'_k = E(X^k)$

Empirical  $m'_k = \frac{1}{n} \sum_{i=1}^k X_i^k$

Need one moment for each parameter in the model

## Example: Method of Moments for uniform(0, $\theta$ )

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \text{uniform}(0, \theta)$

## Example: Method of Moments for $\text{uniform}(0, \theta)$

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \text{uniform}(0, \theta)$

Find the method of moments estimator of  $\theta$

## Example: Method of Moments for uniform(0, $\theta$ )

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \text{uniform}(0, \theta)$

Find the method of moments estimator of  $\theta$

The first theoretical moment is:  $M'_1 = E(X) = \frac{\theta+0}{2} = \frac{\theta}{2}$

## Example: Method of Moments for uniform(0, $\theta$ )

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \text{uniform}(0, \theta)$

Find the method of moments estimator of  $\theta$

The first theoretical moment is:  $M'_1 = E(X) = \frac{\theta+0}{2} = \frac{\theta}{2}$

The first empirical moment is  $m'_1 = \frac{1}{n} \sum_{i=1}^n X_i = \bar{X}$

## Example: Method of Moments for uniform(0, $\theta$ )

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \text{uniform}(0, \theta)$

Find the method of moments estimator of  $\theta$

The first theoretical moment is:  $M'_1 = E(X) = \frac{\theta+0}{2} = \frac{\theta}{2}$

The first empirical moment is  $m'_1 = \frac{1}{n} \sum_{i=1}^n X_i = \bar{X}$

The method of moments estimator for  $\theta$  is thus

$$\hat{\theta}(X_1, \dots, X_n) = \frac{2}{n} \sum_{i=1}^n X_i = 2\bar{X}$$

## Example: Method of Moments for uniform(0, $\theta$ )

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \text{uniform}(0, \theta)$

Find the method of moments estimator of  $\theta$

The first theoretical moment is:  $M'_1 = E(X) = \frac{\theta+0}{2} = \frac{\theta}{2}$

The first empirical moment is  $m'_1 = \frac{1}{n} \sum_{i=1}^n X_i = \bar{X}$

The method of moments estimator for  $\theta$  is thus

$$\hat{\theta}(X_1, \dots, X_n) = \frac{2}{n} \sum_{i=1}^n X_i = 2\bar{X}$$

**Caveat:** Suppose  $\theta = 1$  and that  $x_1 = 0.98$ ,  $x_2 = 0.34$ ,  $x_3 = 0.12$ ,  $x_4 = 0.48$  and  $x_5 = 0.08$  is a realization of a random sample. The method of moments estimate of  $\theta$  is

$$\hat{\theta} = 2\bar{x} = 0.8$$

which is clearly not consistent with the model!



## Example: Method of Moments for $\mathcal{N}(\mu, \sigma^2)$

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$

## Example: Method of Moments for $\mathcal{N}(\mu, \sigma^2)$

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$

Find the method of moments estimators of  $\mu$  and  $\sigma^2$

## Example: Method of Moments for $\mathcal{N}(\mu, \sigma^2)$

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$

Find the method of moments estimators of  $\mu$  and  $\sigma^2$

Theoretical moments

- ▶  $M'_1 = E(X) = \mu$
- ▶  $M'_2 = E(X^2) = \text{Var}(X) + [E(X)]^2 = \sigma^2 + \mu^2$

## Example: Method of Moments for $\mathcal{N}(\mu, \sigma^2)$

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$

Find the method of moments estimators of  $\mu$  and  $\sigma^2$

Theoretical moments

- ▶  $M'_1 = E(X) = \mu$
- ▶  $M'_2 = E(X^2) = \text{Var}(X) + [E(X)]^2 = \sigma^2 + \mu^2$

Empirical moments

- ▶  $m'_1 = \frac{1}{n} \sum_{i=1}^n X_i$
- ▶  $m'_2 = \frac{1}{n} \sum_{i=1}^n X_i^2$

## Example: Method of Moments for $\mathcal{N}(\mu, \sigma^2)$

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$

Find the method of moments estimators of  $\mu$  and  $\sigma^2$

Theoretical moments

- ▶  $M'_1 = E(X) = \mu$
- ▶  $M'_2 = E(X^2) = \text{Var}(X) + [E(X)]^2 = \sigma^2 + \mu^2$

Empirical moments

- ▶  $m'_1 = \frac{1}{n} \sum_{i=1}^n X_i$
- ▶  $m'_2 = \frac{1}{n} \sum_{i=1}^n X_i^2$

Method of moments estimators for  $\mu$  and  $\sigma^2$ :

- ▶  $\hat{\mu}(X_1, \dots, X_n) = m'_1 = \frac{1}{n} \sum_{i=1}^n X_i = \bar{X}$
- ▶  $\hat{\sigma}^2(X_1, \dots, X_n) = m'_2 - (m'_1)^2 = \frac{1}{n} \sum_{i=1}^n X_i^2 - \bar{X}^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2$

# Method of Least Squares

Let  $X_1, \dots, X_n$  be a random sample

# Method of Least Squares

Let  $X_1, \dots, X_n$  be a random sample

Want to estimate  $\theta$ , the expected value of  $X_i$

# Method of Least Squares

Let  $X_1, \dots, X_n$  be a random sample

Want to estimate  $\theta$ , the expected value of  $X_i$

Each random variable  $X_i$  should be “close” to  $\theta$



# Method of Least Squares

Let  $X_1, \dots, X_n$  be a random sample

Want to estimate  $\theta$ , the expected value of  $X_i$

Each random variable  $X_i$  should be “close” to  $\theta$

Each difference  $|X_i - \theta|$  should be *as small as possible*

# Method of Least Squares

Let  $X_1, \dots, X_n$  be a random sample

Want to estimate  $\theta$ , the expected value of  $X_i$

Each random variable  $X_i$  should be “close” to  $\theta$

Each difference  $|X_i - \theta|$  should be *as small as possible*

A reasonable estimator for  $\theta$  would be the value minimizing

$$S(\theta) = \sum_{i=1}^n (X_i - \theta)^2$$

## Example: Method of Least Squares

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} f$  such that  $E(X_i) = \theta$

## Example: Method of Least Squares

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} f$  such that  $E(X_i) = \theta$

Find the least squares estimator of  $\theta$

## Example: Method of Least Squares

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} f$  such that  $E(X_i) = \theta$

Find the least squares estimator of  $\theta$

$$S(\theta) = \sum_{i=1}^n (X_i - \theta)^2$$

## Example: Method of Least Squares

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} f$  such that  $E(X_i) = \theta$

Find the least squares estimator of  $\theta$

$$\begin{aligned} S(\theta) &= \sum_{i=1}^n (X_i - \theta)^2 \\ &= \sum_{i=1}^n (X_i - \bar{X} + \bar{X} - \theta)^2 \end{aligned}$$

## Example: Method of Least Squares

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} f$  such that  $E(X_i) = \theta$

Find the least squares estimator of  $\theta$

$$\begin{aligned} S(\theta) &= \sum_{i=1}^n (X_i - \theta)^2 \\ &= \sum_{i=1}^n (X_i - \bar{X} + \bar{X} - \theta)^2 \\ &= \sum_{i=1}^n [(X_i - \bar{X}) + (\bar{X} - \theta)]^2 \end{aligned}$$

## Example: Method of Least Squares

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} f$  such that  $E(X_i) = \theta$

Find the least squares estimator of  $\theta$

$$\begin{aligned} S(\theta) &= \sum_{i=1}^n (X_i - \theta)^2 \\ &= \sum_{i=1}^n (X_i - \bar{X} + \bar{X} - \theta)^2 \\ &= \sum_{i=1}^n [(X_i - \bar{X}) + (\bar{X} - \theta)]^2 \\ &= \sum_{i=1}^n (X_i - \bar{X})^2 + 2 \sum_{i=1}^n (X_i - \bar{X})(\bar{X} - \theta) + \sum_{i=1}^n (\bar{X} - \theta)^2 \end{aligned}$$



## Example: Method of Least Squares (continued)

$$= \sum_{i=1}^n (X_i - \bar{X})^2 + 2 \sum_{i=1}^n (X_i - \bar{X})(\bar{X} - \theta) + \sum_{i=1}^n (\bar{X} - \theta)^2$$

## Example: Method of Least Squares (continued)

$$\begin{aligned} &= \sum_{i=1}^n (X_i - \bar{X})^2 + 2 \sum_{i=1}^n (X_i - \bar{X})(\bar{X} - \theta) + \sum_{i=1}^n (\bar{X} - \theta)^2 \\ &= \sum_{i=1}^n (X_i - \bar{X})^2 + 2(\bar{X} - \theta) \sum_{i=1}^n (X_i - \bar{X}) + \sum_{i=1}^n (\bar{X} - \theta)^2 \end{aligned}$$

## Example: Method of Least Squares (continued)

$$\begin{aligned} &= \sum_{i=1}^n (X_i - \bar{X})^2 + 2 \sum_{i=1}^n (X_i - \bar{X})(\bar{X} - \theta) + \sum_{i=1}^n (\bar{X} - \theta)^2 \\ &= \sum_{i=1}^n (X_i - \bar{X})^2 + 2(\bar{X} - \theta) \sum_{i=1}^n (X_i - \bar{X}) + \sum_{i=1}^n (\bar{X} - \theta)^2 \\ &= \sum_{i=1}^n (X_i - \bar{X})^2 + 2(\bar{X} - \theta) \left[ \left( \sum_{i=1}^n X_i \right) - n\bar{X} \right] + \sum_{i=1}^n (\bar{X} - \theta)^2 \end{aligned}$$

## Example: Method of Least Squares (continued)

$$\begin{aligned} &= \sum_{i=1}^n (X_i - \bar{X})^2 + 2 \sum_{i=1}^n (X_i - \bar{X})(\bar{X} - \theta) + \sum_{i=1}^n (\bar{X} - \theta)^2 \\ &= \sum_{i=1}^n (X_i - \bar{X})^2 + 2(\bar{X} - \theta) \sum_{i=1}^n (X_i - \bar{X}) + \sum_{i=1}^n (\bar{X} - \theta)^2 \\ &= \sum_{i=1}^n (X_i - \bar{X})^2 + 2(\bar{X} - \theta) \left[ \left( \sum_{i=1}^n X_i \right) - n\bar{X} \right] + \sum_{i=1}^n (\bar{X} - \theta)^2 \\ &= \sum_{i=1}^n (X_i - \bar{X})^2 + \sum_{i=1}^n (\bar{X} - \theta)^2 \end{aligned}$$

## Example: Method of Least Squares (continued)

$$\begin{aligned} &= \sum_{i=1}^n (X_i - \bar{X})^2 + 2 \sum_{i=1}^n (X_i - \bar{X})(\bar{X} - \theta) + \sum_{i=1}^n (\bar{X} - \theta)^2 \\ &= \sum_{i=1}^n (X_i - \bar{X})^2 + 2(\bar{X} - \theta) \sum_{i=1}^n (X_i - \bar{X}) + \sum_{i=1}^n (\bar{X} - \theta)^2 \\ &= \sum_{i=1}^n (X_i - \bar{X})^2 + 2(\bar{X} - \theta) \left[ \left( \sum_{i=1}^n X_i \right) - n\bar{X} \right] + \sum_{i=1}^n (\bar{X} - \theta)^2 \\ &= \sum_{i=1}^n (X_i - \bar{X})^2 + \sum_{i=1}^n (\bar{X} - \theta)^2 \end{aligned}$$

$$\implies \hat{\theta}(X_1, \dots, X_n) = \bar{X}$$

# Method of Maximum Likelihood

The joint density of a random sample from a parametric family with parameter  $\theta$  is

$$f_{X_1, \dots, X_n}(x_1, \dots, x_n | \theta) = \prod_{i=1}^n f_X(x_i | \theta)$$

# Method of Maximum Likelihood

The joint density of a random sample from a parametric family with parameter  $\theta$  is

$$f_{X_1, \dots, X_n}(x_1, \dots, x_n | \theta) = \prod_{i=1}^n f_X(x_i | \theta)$$

Let  $x_1, \dots, x_n$  (the data) be the realization of a random sample  
 $X_1, \dots, X_n \stackrel{iid}{\sim} f_X(x | \theta)$

# Method of Maximum Likelihood

The joint density of a random sample from a parametric family with parameter  $\theta$  is

$$f_{X_1, \dots, X_n}(x_1, \dots, x_n | \theta) = \prod_{i=1}^n f_X(x_i | \theta)$$

Let  $x_1, \dots, x_n$  (the data) be the realization of a random sample  
 $X_1, \dots, X_n \stackrel{iid}{\sim} f_X(x | \theta)$

**Definition** The *likelihood* of the parameter  $\theta$  is

$$L(\theta | x_1, \dots, x_n) = f_{X_1, \dots, X_n}(x_1, \dots, x_n | \theta) = \prod_{i=1}^n f_X(x_i | \theta)$$



# Method of Maximum Likelihood

The joint density of a random sample from a parametric family with parameter  $\theta$  is

$$f_{X_1, \dots, X_n}(x_1, \dots, x_n | \theta) = \prod_{i=1}^n f_X(x_i | \theta)$$

Let  $x_1, \dots, x_n$  (the data) be the realization of a random sample  
 $X_1, \dots, X_n \stackrel{iid}{\sim} f_X(x | \theta)$

**Definition** The *likelihood* of the parameter  $\theta$  is

$$L(\theta | x_1, \dots, x_n) = f_{X_1, \dots, X_n}(x_1, \dots, x_n | \theta) = \prod_{i=1}^n f_X(x_i | \theta)$$

**Definition** The *maximum likelihood estimator*  $\hat{\theta}_{ML}$  of a parameter  $\theta$  is a value of  $\theta$  giving the largest likelihood possible:

$$L(\hat{\theta}_{ML} | x_1, \dots, x_n) \geq L(\theta | x_1, \dots, x_n)$$

for all admissible values of the parameter  $\theta$

## Calculation of $\hat{\theta}_{ML}$

Often, maximizing the log of the likelihood is easier than maximizing the likelihood itself

## Calculation of $\hat{\theta}_{ML}$

Often, maximizing the log of the likelihood is easier than maximizing the likelihood itself

General algorithm for finding  $\hat{\theta}$

## Calculation of $\hat{\theta}_{ML}$

Often, maximizing the log of the likelihood is easier than maximizing the likelihood itself

General algorithm for finding  $\hat{\theta}$

1. Write down the likelihood function  $L(\theta|x_1, \dots, x_n)$

## Calculation of $\hat{\theta}_{ML}$

Often, maximizing the log of the likelihood is easier than maximizing the likelihood itself

General algorithm for finding  $\hat{\theta}$

1. Write down the likelihood function  $L(\theta|x_1, \dots, x_n)$
2. Define  $\ell(\theta|x_1, \dots, x_n) = \log L(\theta|x_1, \dots, x_n)$  (the log likelihood of  $\theta$ )

## Calculation of $\hat{\theta}_{ML}$

Often, maximizing the log of the likelihood is easier than maximizing the likelihood itself

General algorithm for finding  $\hat{\theta}$

1. Write down the likelihood function  $L(\theta|x_1, \dots, x_n)$
2. Define  $\ell(\theta|x_1, \dots, x_n) = \log L(\theta|x_1, \dots, x_n)$  (the log likelihood of  $\theta$ )
3. Find  $\hat{\theta}_{ML}$  such that

$$\left. \frac{d}{d\theta} \ell(\theta|x_1, \dots, x_n) \right|_{\hat{\theta}_{ML}} = 0$$

## Calculation of $\hat{\theta}_{ML}$

Often, maximizing the log of the likelihood is easier than maximizing the likelihood itself

General algorithm for finding  $\hat{\theta}$

1. Write down the likelihood function  $L(\theta|x_1, \dots, x_n)$
2. Define  $\ell(\theta|x_1, \dots, x_n) = \log L(\theta|x_1, \dots, x_n)$  (the log likelihood of  $\theta$ )
3. Find  $\hat{\theta}_{ML}$  such that

$$\left. \frac{d}{d\theta} \ell(\theta|x_1, \dots, x_n) \right|_{\hat{\theta}_{ML}} = 0$$

4. Verify that  $\hat{\theta}_{ML}$  is a maximum

## Example: Find $\hat{p}_{ML}$ for a Sequence of Bernoulli Trials

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \text{Bernoulli}(p)$  be a random sample



## Example: Find $\hat{p}_{ML}$ for a Sequence of Bernoulli Trials

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \text{Bernoulli}(p)$  be a random sample

Find the maximum likelihood estimator of  $p$

## Example: Find $\hat{p}_{ML}$ for a Sequence of Bernoulli Trials

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \text{Bernoulli}(p)$  be a random sample

Find the maximum likelihood estimator of  $p$

The likelihood function is

$$L(p|x_1, \dots, x_n) = f_{X_1, \dots, X_n}(x_1, \dots, x_n|p)$$

## Example: Find $\hat{p}_{ML}$ for a Sequence of Bernoulli Trials

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \text{Bernoulli}(p)$  be a random sample

Find the maximum likelihood estimator of  $p$

The likelihood function is

$$L(p|x_1, \dots, x_n) = f_{X_1, \dots, X_n}(x_1, \dots, x_n|p) = \prod_{i=1}^n f_X(x_i|p)$$

## Example: Find $\hat{p}_{ML}$ for a Sequence of Bernoulli Trials

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \text{Bernoulli}(p)$  be a random sample

Find the maximum likelihood estimator of  $p$

The likelihood function is

$$\begin{aligned} L(p|x_1, \dots, x_n) &= f_{X_1, \dots, X_n}(x_1, \dots, x_n|p) = \prod_{i=1}^n f_X(x_i|p) \\ &= \prod_{i=1}^n p^{x_i} (1-p)^{(1-x_i)} \end{aligned}$$

## Example: Find $\hat{p}_{ML}$ for a Sequence of Bernoulli Trials

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \text{Bernoulli}(p)$  be a random sample

Find the maximum likelihood estimator of  $p$

The likelihood function is

$$\begin{aligned} L(p|x_1, \dots, x_n) &= f_{X_1, \dots, X_n}(x_1, \dots, x_n|p) = \prod_{i=1}^n f_X(x_i|p) \\ &= \prod_{i=1}^n p^{x_i} (1-p)^{(1-x_i)} \end{aligned}$$

The log likelihood function is

$$\ell(p|x_1, \dots, x_n) = \log L(p|x_1, \dots, x_n) = \log \left[ \prod_{i=1}^n p^{x_i} (1-p)^{(1-x_i)} \right]$$

Find  $\hat{p}_{ML}$  for a Sequence of Bernoulli Trials (continued)

$$= \sum_{i=1}^n \log \left[ p^{x_i} (1-p)^{(1-x_i)} \right]$$

Find  $\hat{p}_{ML}$  for a Sequence of Bernoulli Trials (continued)

$$\begin{aligned} &= \sum_{i=1}^n \log \left[ p^{x_i} (1-p)^{(1-x_i)} \right] \\ &= \log(p) \sum_{i=1}^n x_i + \log(1-p) \sum_{i=1}^n (1-x_i) \end{aligned}$$

## Find $\hat{p}_{ML}$ for a Sequence of Bernoulli Trials (continued)

$$\begin{aligned} &= \sum_{i=1}^n \log \left[ p^{x_i} (1-p)^{(1-x_i)} \right] \\ &= \log(p) \sum_{i=1}^n x_i + \log(1-p) \sum_{i=1}^n (1-x_i) \\ &= \log(p) \sum_{i=1}^n x_i + \log(1-p) \left[ n - \sum_{i=1}^n x_i \right] \end{aligned}$$



## Find $\hat{p}_{ML}$ for a Sequence of Bernoulli Trials (continued)

$$\begin{aligned} &= \sum_{i=1}^n \log \left[ p^{x_i} (1-p)^{(1-x_i)} \right] \\ &= \log(p) \sum_{i=1}^n x_i + \log(1-p) \sum_{i=1}^n (1-x_i) \\ &= \log(p) \sum_{i=1}^n x_i + \log(1-p) \left[ n - \sum_{i=1}^n x_i \right] \end{aligned}$$

Take the derivative of the log likelihood wrt the parameter

$$\frac{d}{dp} \ell(p|x_1, \dots, x_n) = \frac{1}{p} \sum_{i=1}^n x_i - \frac{1}{1-p} \left[ n - \sum_{i=1}^n x_i \right]$$

## Find $\hat{p}_{ML}$ for a Sequence of Bernoulli Trials (continued)

Set the derivative equal to 0 and solve for  $\hat{p}_{ML}$

$$\frac{1}{\hat{p}_{ML}} \sum_{i=1}^n x_i - \frac{1}{1 - \hat{p}_{ML}} \left[ n - \sum_{i=1}^n x_i \right] \stackrel{\text{set}}{=} 0$$

## Find $\hat{p}_{ML}$ for a Sequence of Bernoulli Trials (continued)

Set the derivative equal to 0 and solve for  $\hat{p}_{ML}$

$$\frac{1}{\hat{p}_{ML}} \sum_{i=1}^n x_i - \frac{1}{1 - \hat{p}_{ML}} \left[ n - \sum_{i=1}^n x_i \right] \stackrel{\text{set}}{=} 0$$
$$\frac{1}{\hat{p}_{ML}} \sum_{i=1}^n x_i = \frac{1}{1 - \hat{p}_{ML}} \left[ n - \sum_{i=1}^n x_i \right]$$

## Find $\hat{p}_{ML}$ for a Sequence of Bernoulli Trials (continued)

Set the derivative equal to 0 and solve for  $\hat{p}_{ML}$

$$\frac{1}{\hat{p}_{ML}} \sum_{i=1}^n x_i - \frac{1}{1 - \hat{p}_{ML}} \left[ n - \sum_{i=1}^n x_i \right] \stackrel{\text{set}}{=} 0$$

$$\frac{1}{\hat{p}_{ML}} \sum_{i=1}^n x_i = \frac{1}{1 - \hat{p}_{ML}} \left[ n - \sum_{i=1}^n x_i \right]$$

$$(1 - \hat{p}_{ML}) \sum_{i=1}^n x_i = \hat{p}_{ML} \left[ n - \sum_{i=1}^n x_i \right]$$

## Find $\hat{p}_{ML}$ for a Sequence of Bernoulli Trials (continued)

Set the derivative equal to 0 and solve for  $\hat{p}_{ML}$

$$\frac{1}{\hat{p}_{ML}} \sum_{i=1}^n x_i - \frac{1}{1 - \hat{p}_{ML}} \left[ n - \sum_{i=1}^n x_i \right] \stackrel{\text{set}}{=} 0$$

$$\frac{1}{\hat{p}_{ML}} \sum_{i=1}^n x_i = \frac{1}{1 - \hat{p}_{ML}} \left[ n - \sum_{i=1}^n x_i \right]$$

$$(1 - \hat{p}_{ML}) \sum_{i=1}^n x_i = \hat{p}_{ML} \left[ n - \sum_{i=1}^n x_i \right]$$

$$\sum_{i=1}^n x_i - \hat{p}_{ML} \sum_{i=1}^n x_i = n\hat{p}_{ML} - \hat{p}_{ML} \sum_{i=1}^n x_i$$

## Find $\hat{p}_{ML}$ for a Sequence of Bernoulli Trials (continued)

Set the derivative equal to 0 and solve for  $\hat{p}_{ML}$

$$\frac{1}{\hat{p}_{ML}} \sum_{i=1}^n x_i - \frac{1}{1 - \hat{p}_{ML}} \left[ n - \sum_{i=1}^n x_i \right] \stackrel{\text{set}}{=} 0$$

$$\frac{1}{\hat{p}_{ML}} \sum_{i=1}^n x_i = \frac{1}{1 - \hat{p}_{ML}} \left[ n - \sum_{i=1}^n x_i \right]$$

$$(1 - \hat{p}_{ML}) \sum_{i=1}^n x_i = \hat{p}_{ML} \left[ n - \sum_{i=1}^n x_i \right]$$

$$\sum_{i=1}^n x_i - \hat{p}_{ML} \sum_{i=1}^n x_i = n\hat{p}_{ML} - \hat{p}_{ML} \sum_{i=1}^n x_i$$

$$\hat{p}_{ML} = \frac{1}{n} \sum_{i=1}^n x_i = \bar{x}$$

## Find $\hat{p}_{ML}$ for a Sequence of Bernoulli Trials (continued)

Finally, verify that  $\hat{p}_{ML}$  is indeed a maximum

## Find $\hat{p}_{ML}$ for a Sequence of Bernoulli Trials (continued)

Finally, verify that  $\hat{p}_{ML}$  is indeed a maximum

$$\frac{d^2}{dp^2} \ell(p|x_1, \dots, x_n) = -\frac{1}{p^2} \sum_{i=1}^n x_i - \frac{1}{(1-p)^2} \left[ n - \sum_{i=1}^n x_i \right]$$



## Find $\hat{p}_{ML}$ for a Sequence of Bernoulli Trials (continued)

Finally, verify that  $\hat{p}_{ML}$  is indeed a maximum

$$\begin{aligned}\frac{d^2}{dp^2}\ell(p|x_1, \dots, x_n) &= -\frac{1}{p^2} \sum_{i=1}^n x_i - \frac{1}{(1-p)^2} \left[ n - \sum_{i=1}^n x_i \right] \\ &< 0 \quad \text{for all } p \in (0, 1)\end{aligned}$$

## Find $\hat{p}_{ML}$ for a Sequence of Bernoulli Trials (continued)

Finally, verify that  $\hat{p}_{ML}$  is indeed a maximum

$$\begin{aligned}\frac{d^2}{dp^2}\ell(p|x_1, \dots, x_n) &= -\frac{1}{p^2} \sum_{i=1}^n x_i - \frac{1}{(1-p)^2} \left[ n - \sum_{i=1}^n x_i \right] \\ &< 0 \quad \text{for all } p \in (0, 1)\end{aligned}$$

$\implies \hat{p}_{ML}$  is indeed a maximum of the log likelihood

## Find $\hat{p}_{ML}$ for a Sequence of Bernoulli Trials (continued)

Finally, verify that  $\hat{p}_{ML}$  is indeed a maximum

$$\begin{aligned}\frac{d^2}{dp^2}\ell(p|x_1, \dots, x_n) &= -\frac{1}{p^2} \sum_{i=1}^n x_i - \frac{1}{(1-p)^2} \left[ n - \sum_{i=1}^n x_i \right] \\ &< 0 \quad \text{for all } p \in (0, 1)\end{aligned}$$

$\implies \hat{p}_{ML}$  is indeed a maximum of the log likelihood

Have shown that for every possible  $(X_1 = x_1, \dots, X_n = x_n)$

$$\hat{p}_{ML} = \bar{x}$$

## Find $\hat{p}_{ML}$ for a Sequence of Bernoulli Trials (continued)

Finally, verify that  $\hat{p}_{ML}$  is indeed a maximum


$$\begin{aligned}\frac{d^2}{dp^2}\ell(p|x_1, \dots, x_n) &= -\frac{1}{p^2} \sum_{i=1}^n x_i - \frac{1}{(1-p)^2} \left[ n - \sum_{i=1}^n x_i \right] \\ &< 0 \quad \text{for all } p \in (0, 1)\end{aligned}$$

$\implies \hat{p}_{ML}$  is indeed a maximum of the log likelihood

Have shown that for every possible  $(X_1 = x_1, \dots, X_n = x_n)$

$$\hat{p}_{ML} = \bar{x} \implies \hat{p}_{ML} = \bar{X}$$

Find  $\hat{p}_{ML}$  for a Sequence of Bernoulli Trials (illustration)



## Example: Find $\hat{\mu}$ for a Normal Population

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$

## Example: Find $\hat{\mu}$ for a Normal Population

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$

Find the maximum likelihood estimator of  $\mu$

## Example: Find $\hat{\mu}$ for a Normal Population

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$

Find the maximum likelihood estimator of  $\mu$

The likelihood function is

$$L(\mu, \sigma^2 | x_1, \dots, x_n) = f_{X_1, \dots, X_n}(x_1, \dots, x_n | \mu, \sigma^2)$$



## Example: Find $\hat{\mu}$ for a Normal Population

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$

Find the maximum likelihood estimator of  $\mu$

The likelihood function is

$$\begin{aligned} L(\mu, \sigma^2 | x_1, \dots, x_n) &= f_{X_1, \dots, X_n}(x_1, \dots, x_n | \mu, \sigma^2) \\ &= \prod_{i=1}^n f_X(x_i | \mu, \sigma^2) \end{aligned}$$

## Example: Find $\hat{\mu}$ for a Normal Population

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$

Find the maximum likelihood estimator of  $\mu$

The likelihood function is

$$\begin{aligned} L(\mu, \sigma^2 | x_1, \dots, x_n) &= f_{X_1, \dots, X_n}(x_1, \dots, x_n | \mu, \sigma^2) \\ &= \prod_{i=1}^n f_X(x_i | \mu, \sigma^2) \\ &= \prod_{i=1}^n \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{(x_i - \mu)^2}{2\sigma^2}\right) \end{aligned}$$

## Example: Find $\hat{\mu}$ for a Normal Population

The log likelihood function is

$$\ell(\mu, \sigma^2 | x_1, \dots, x_n) = \log L(\mu, \sigma^2 | x_1, \dots, x_n)$$

## Example: Find $\hat{\mu}$ for a Normal Population

The log likelihood function is

$$\begin{aligned}\ell(\mu, \sigma^2 | x_1, \dots, x_n) &= \log L(\mu, \sigma^2 | x_1, \dots, x_n) \\ &= \log \left[ \prod_{i=1}^n \frac{1}{\sqrt{2\pi}\sigma} \exp \left( -\frac{(x_i - \mu)^2}{2\sigma^2} \right) \right]\end{aligned}$$

## Example: Find $\hat{\mu}$ for a Normal Population

The log likelihood function is

$$\begin{aligned}\ell(\mu, \sigma^2 | x_1, \dots, x_n) &= \log L(\mu, \sigma^2 | x_1, \dots, x_n) \\ &= \log \left[ \prod_{i=1}^n \frac{1}{\sqrt{2\pi}\sigma} \exp \left( -\frac{(x_i - \mu)^2}{2\sigma^2} \right) \right] \\ &= \sum_{i=1}^n \log \frac{1}{\sqrt{2\pi}\sigma} + \sum_{i=1}^n \left( -\frac{(x_i - \mu)^2}{2\sigma^2} \right)\end{aligned}$$

## Example: Find $\hat{\mu}$ for a Normal Population

The log likelihood function is

$$\begin{aligned}\ell(\mu, \sigma^2 | x_1, \dots, x_n) &= \log L(\mu, \sigma^2 | x_1, \dots, x_n) \\&= \log \left[ \prod_{i=1}^n \frac{1}{\sqrt{2\pi}\sigma} \exp \left( -\frac{(x_i - \mu)^2}{2\sigma^2} \right) \right] \\&= \sum_{i=1}^n \log \frac{1}{\sqrt{2\pi}\sigma} + \sum_{i=1}^n \left( -\frac{(x_i - \mu)^2}{2\sigma^2} \right) \\&= -n \log(\sqrt{2\pi}\sigma) - \frac{1}{2\sigma^2} \sum_{i=1}^n (x_i - \mu)^2\end{aligned}$$

## Example: Find $\hat{\mu}$ for a Normal Population

The log likelihood function is

$$\begin{aligned}\ell(\mu, \sigma^2 | x_1, \dots, x_n) &= \log L(\mu, \sigma^2 | x_1, \dots, x_n) \\&= \log \left[ \prod_{i=1}^n \frac{1}{\sqrt{2\pi}\sigma} \exp \left( -\frac{(x_i - \mu)^2}{2\sigma^2} \right) \right] \\&= \sum_{i=1}^n \log \frac{1}{\sqrt{2\pi}\sigma} + \sum_{i=1}^n \left( -\frac{(x_i - \mu)^2}{2\sigma^2} \right) \\&= -n \log(\sqrt{2\pi}\sigma) - \frac{1}{2\sigma^2} \sum_{i=1}^n (x_i - \mu)^2 \\&\propto -\frac{1}{2\sigma^2} \sum_{i=1}^n (x_i - \mu)^2\end{aligned}$$

## Example: Find $\hat{\mu}$ for a Normal Population

Take the derivative of the log likelihood

$$\frac{d}{d\mu} \ell(\mu, \sigma^2 | x_1, \dots, x_n) = \frac{d}{d\mu} \left[ -\frac{1}{2\sigma^2} \sum_{i=1}^n (x_i - \mu)^2 \right]$$



## Example: Find $\hat{\mu}$ for a Normal Population

Take the derivative of the log likelihood

$$\begin{aligned}\frac{d}{d\mu} \ell(\mu, \sigma^2 | x_1, \dots, x_n) &= \frac{d}{d\mu} \left[ -\frac{1}{2\sigma^2} \sum_{i=1}^n (x_i - \mu)^2 \right] \\ &= \frac{1}{\sigma^2} \sum_{i=1}^n (x_i - \mu)\end{aligned}$$

## Example: Find $\hat{\mu}$ for a Normal Population

Take the derivative of the log likelihood

$$\begin{aligned}\frac{d}{d\mu} \ell(\mu, \sigma^2 | x_1, \dots, x_n) &= \frac{d}{d\mu} \left[ -\frac{1}{2\sigma^2} \sum_{i=1}^n (x_i - \mu)^2 \right] \\ &= \frac{1}{\sigma^2} \sum_{i=1}^n (x_i - \mu)\end{aligned}$$

Set the derivative equal to 0 and solve for  $\hat{\mu}_{ML}$

$$\frac{1}{\sigma^2} \sum_{i=1}^n (x_i - \hat{\mu}_{ML}) \stackrel{\text{set}}{=} 0$$

## Example: Find $\hat{\mu}$ for a Normal Population

Take the derivative of the log likelihood

$$\begin{aligned}\frac{d}{d\mu} \ell(\mu, \sigma^2 | x_1, \dots, x_n) &= \frac{d}{d\mu} \left[ -\frac{1}{2\sigma^2} \sum_{i=1}^n (x_i - \mu)^2 \right] \\ &= \frac{1}{\sigma^2} \sum_{i=1}^n (x_i - \mu)\end{aligned}$$

Set the derivative equal to 0 and solve for  $\hat{\mu}_{ML}$

$$\begin{aligned}\frac{1}{\sigma^2} \sum_{i=1}^n (x_i - \hat{\mu}_{ML}) &\stackrel{\text{set}}{=} 0 \\ \sum_{i=1}^n x_i &= n\hat{\mu}_{ML}\end{aligned}$$

## Example: Find $\hat{\mu}$ for a Normal Population

Take the derivative of the log likelihood

$$\begin{aligned}\frac{d}{d\mu} \ell(\mu, \sigma^2 | x_1, \dots, x_n) &= \frac{d}{d\mu} \left[ -\frac{1}{2\sigma^2} \sum_{i=1}^n (x_i - \mu)^2 \right] \\ &= \frac{1}{\sigma^2} \sum_{i=1}^n (x_i - \mu)\end{aligned}$$

Set the derivative equal to 0 and solve for  $\hat{\mu}_{ML}$

$$\begin{aligned}\frac{1}{\sigma^2} \sum_{i=1}^n (x_i - \hat{\mu}_{ML}) &\stackrel{\text{set}}{=} 0 \\ \sum_{i=1}^n x_i &= n\hat{\mu}_{ML} \implies \hat{\mu}_{ML} = \frac{1}{n} \sum_{i=1}^n x_i = \bar{x}\end{aligned}$$

## Example: Find $\hat{\mu}$ for a Normal Population

Finally, need to verify that  $\hat{\mu}_{ML}$  is indeed a maximum

$$\frac{d^2}{d\mu^2} \ell(\mu, \sigma^2 | x_1, \dots, x_n) = \frac{d}{d\mu} \left[ \frac{1}{\sigma^2} \sum_{i=1}^n (x_i - \mu) \right]$$

## Example: Find $\hat{\mu}$ for a Normal Population

Finally, need to verify that  $\hat{\mu}_{ML}$  is indeed a maximum

$$\begin{aligned}\frac{d^2}{d\mu^2}\ell(\mu, \sigma^2 | x_1, \dots, x_n) &= \frac{d}{d\mu} \left[ \frac{1}{\sigma^2} \sum_{i=1}^n (x_i - \mu) \right] \\ &= -\frac{n}{\sigma^2}\end{aligned}$$

## Example: Find $\hat{\mu}$ for a Normal Population

Finally, need to verify that  $\hat{\mu}_{ML}$  is indeed a maximum

$$\begin{aligned}\frac{d^2}{d\mu^2}\ell(\mu, \sigma^2 | x_1, \dots, x_n) &= \frac{d}{d\mu} \left[ \frac{1}{\sigma^2} \sum_{i=1}^n (x_i - \mu) \right] \\ &= -\frac{n}{\sigma^2} \\ &< 0\end{aligned}$$

## Example: Find $\hat{\mu}$ for a Normal Population

Finally, need to verify that  $\hat{\mu}_{ML}$  is indeed a maximum

$$\begin{aligned}\frac{d^2}{d\mu^2}\ell(\mu, \sigma^2 | x_1, \dots, x_n) &= \frac{d}{d\mu} \left[ \frac{1}{\sigma^2} \sum_{i=1}^n (x_i - \mu) \right] \\ &= -\frac{n}{\sigma^2} \\ &< 0\end{aligned}$$

$\Rightarrow \hat{\mu}_{ML}$  is a local maximum of the log likelihood



## Example: Find $\hat{\mu}$ for a Normal Population

Finally, need to verify that  $\hat{\mu}_{ML}$  is indeed a maximum

$$\begin{aligned}\frac{d^2}{d\mu^2}\ell(\mu, \sigma^2 | x_1, \dots, x_n) &= \frac{d}{d\mu} \left[ \frac{1}{\sigma^2} \sum_{i=1}^n (x_i - \mu) \right] \\ &= -\frac{n}{\sigma^2} \\ &< 0\end{aligned}$$

$\Rightarrow \hat{\mu}_{ML}$  is a local maximum of the log likelihood

The maximum likelihood estimator of  $\mu$  is

$$\hat{\mu}_{ML} = \bar{X}$$

# Outline

Statistical Models

Point Estimation

Evaluating Point Estimators

Interval Estimators

# Bias

The *bias* of an estimator  $\hat{\theta}$  is defined to be

$$\text{Bias}_{\hat{\theta}}(\theta) = E(\hat{\theta}) - \theta$$

# Bias

The *bias* of an estimator  $\hat{\theta}$  is defined to be

$$\text{Bias}_{\hat{\theta}}(\theta) = E(\hat{\theta}) - \theta$$

Interpretation of bias:

# Bias

The *bias* of an estimator  $\hat{\theta}$  is defined to be

$$\text{Bias}_{\hat{\theta}}(\theta) = E(\hat{\theta}) - \theta$$

Interpretation of bias:

- ▶ If  $\text{Bias}_{\hat{\theta}}(\theta) < 0$  then  $\hat{\theta}$  tends to underestimate  $\theta$

# Bias

The *bias* of an estimator  $\hat{\theta}$  is defined to be

$$\text{Bias}_{\hat{\theta}}(\theta) = E(\hat{\theta}) - \theta$$

Interpretation of bias:

- ▶ If  $\text{Bias}_{\hat{\theta}}(\theta) < 0$  then  $\hat{\theta}$  tends to underestimate  $\theta$
- ▶ If  $\text{Bias}_{\hat{\theta}}(\theta) > 0$  then  $\hat{\theta}$  tends to overestimate  $\theta$

# Bias

The *bias* of an estimator  $\hat{\theta}$  is defined to be

$$\text{Bias}_{\hat{\theta}}(\theta) = E(\hat{\theta}) - \theta$$

Interpretation of bias:

- ▶ If  $\text{Bias}_{\hat{\theta}}(\theta) < 0$  then  $\hat{\theta}$  tends to underestimate  $\theta$
- ▶ If  $\text{Bias}_{\hat{\theta}}(\theta) > 0$  then  $\hat{\theta}$  tends to overestimate  $\theta$
- ▶ If  $\text{Bias}_{\hat{\theta}}(\theta) \equiv 0$  then  $\hat{\theta}$  is an *unbiased* estimator of  $\theta$

# Bias

The *bias* of an estimator  $\hat{\theta}$  is defined to be

$$\text{Bias}_{\hat{\theta}}(\theta) = E(\hat{\theta}) - \theta$$

Interpretation of bias:

- ▶ If  $\text{Bias}_{\hat{\theta}}(\theta) < 0$  then  $\hat{\theta}$  tends to underestimate  $\theta$
- ▶ If  $\text{Bias}_{\hat{\theta}}(\theta) > 0$  then  $\hat{\theta}$  tends to overestimate  $\theta$
- ▶ If  $\text{Bias}_{\hat{\theta}}(\theta) \equiv 0$  then  $\hat{\theta}$  is an *unbiased* estimator of  $\theta$

A qualitative measure of an estimator  $\hat{\theta}$  is the absence of a systematic relationship between  $\hat{\theta}$  and  $\theta$



# Bias

The *bias* of an estimator  $\hat{\theta}$  is defined to be

$$\text{Bias}_{\hat{\theta}}(\theta) = E(\hat{\theta}) - \theta$$

Interpretation of bias:

- ▶ If  $\text{Bias}_{\hat{\theta}}(\theta) < 0$  then  $\hat{\theta}$  tends to underestimate  $\theta$
- ▶ If  $\text{Bias}_{\hat{\theta}}(\theta) > 0$  then  $\hat{\theta}$  tends to overestimate  $\theta$
- ▶ If  $\text{Bias}_{\hat{\theta}}(\theta) \equiv 0$  then  $\hat{\theta}$  is an *unbiased* estimator of  $\theta$

A qualitative measure of an estimator  $\hat{\theta}$  is the absence of a systematic relationship between  $\hat{\theta}$  and  $\theta$

Later: better measure is given by the average size of  $(\hat{\theta} - \theta)^2$

## Example: Bias

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$

## Example: Bias

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$

Find the bias and the variance of  $\hat{\mu} = \bar{X}$

## Example: Bias

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$

Find the bias and the variance of  $\hat{\mu} = \bar{X}$

$$\text{Bias}_{\hat{\mu}}(\mu) = E(\bar{X}) - \mu$$

## Example: Bias

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$

Find the bias and the variance of  $\hat{\mu} = \bar{X}$

$$\begin{aligned}\text{Bias}_{\hat{\mu}}(\mu) &= \mathbb{E}(\bar{X}) - \mu \\ &= \mathbb{E}\left(\frac{1}{n} \sum_{i=1}^n X_i\right) - \mu\end{aligned}$$

## Example: Bias

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$

Find the bias and the variance of  $\hat{\mu} = \bar{X}$

$$\begin{aligned}\text{Bias}_{\hat{\mu}}(\mu) &= \mathbb{E}(\bar{X}) - \mu \\ &= \mathbb{E}\left(\frac{1}{n} \sum_{i=1}^n X_i\right) - \mu \\ &= \frac{1}{n} \sum_{i=1}^n \mathbb{E}(X_i) - \mu\end{aligned}$$

## Example: Bias

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$

Find the bias and the variance of  $\hat{\mu} = \bar{X}$

$$\begin{aligned}\text{Bias}_{\hat{\mu}}(\mu) &= \mathbb{E}(\bar{X}) - \mu \\ &= \mathbb{E}\left(\frac{1}{n} \sum_{i=1}^n X_i\right) - \mu \\ &= \frac{1}{n} \sum_{i=1}^n \mathbb{E}(X_i) - \mu \\ &= \frac{1}{n} n \mu - \mu\end{aligned}$$

## Example: Bias

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$

Find the bias and the variance of  $\hat{\mu} = \bar{X}$

$$\begin{aligned}\text{Bias}_{\hat{\mu}}(\mu) &= E(\bar{X}) - \mu \\&= E\left(\frac{1}{n} \sum_{i=1}^n X_i\right) - \mu \\&= \frac{1}{n} \sum_{i=1}^n E(X_i) - \mu \\&= \frac{1}{n} n \mu - \mu \\&= 0 \implies \bar{X} \text{ is an unbiased estimator of } \mu\end{aligned}$$



## Example: Bias (continued)

$$\text{Var}(\bar{X}) = \text{Var}\left(\frac{1}{n} \sum_{i=1}^n X_i\right)$$

## Example: Bias (continued)

$$\begin{aligned}\text{Var}(\bar{X}) &= \text{Var}\left(\frac{1}{n} \sum_{i=1}^n X_i\right) \\ &= \frac{1}{n^2} \text{Var}\left(\sum_{i=1}^n X_i\right)\end{aligned}$$

## Example: Bias (continued)

$$\begin{aligned}\text{Var}(\bar{X}) &= \text{Var}\left(\frac{1}{n} \sum_{i=1}^n X_i\right) \\ &= \frac{1}{n^2} \text{Var}\left(\sum_{i=1}^n X_i\right) \\ &= \frac{1}{n^2} \sum_{i=1}^n \text{Var}(X_i)\end{aligned}$$

(b/c the  $X_i$  are independent)

## Example: Bias (continued)

$$\begin{aligned}\text{Var}(\bar{X}) &= \text{Var}\left(\frac{1}{n} \sum_{i=1}^n X_i\right) \\&= \frac{1}{n^2} \text{Var}\left(\sum_{i=1}^n X_i\right) \\&= \frac{1}{n^2} \sum_{i=1}^n \text{Var}(X_i) && \text{(b/c the } X_i \text{ are independent)} \\&= \frac{1}{n^2} n \sigma^2\end{aligned}$$

## Example: Bias (continued)

$$\begin{aligned}\text{Var}(\bar{X}) &= \text{Var}\left(\frac{1}{n} \sum_{i=1}^n X_i\right) \\&= \frac{1}{n^2} \text{Var}\left(\sum_{i=1}^n X_i\right) \\&= \frac{1}{n^2} \sum_{i=1}^n \text{Var}(X_i) \\&= \frac{1}{n^2} n \sigma^2 \\&= \frac{\sigma^2}{n}\end{aligned}$$

(b/c the  $X_i$  are independent)

## Example: Bias (continued)

Find the bias of  $\hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2$

## Example: Bias (continued)

Find the bias of  $\hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2$

$$\text{Bias}_{\hat{\sigma}^2}(\sigma^2) = E \left( \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2 \right) - \sigma^2$$

## Example: Bias (continued)

Find the bias of  $\hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2$

$$\begin{aligned}\text{Bias}_{\hat{\sigma}^2}(\sigma^2) &= E\left(\frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2\right) - \sigma^2 \\ &= \frac{n-1}{n} E\left(\frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2\right) - \sigma^2\end{aligned}$$



## Example: Bias (continued)

Find the bias of  $\hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2$

$$\begin{aligned}\text{Bias}_{\hat{\sigma}^2}(\sigma^2) &= E\left(\frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2\right) - \sigma^2 \\&= \frac{n-1}{n} E\left(\frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2\right) - \sigma^2 \\&= \frac{n-1}{n} E(S^2) - \sigma^2\end{aligned}$$

## Example: Bias (continued)

Find the bias of  $\hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2$

$$\begin{aligned}\text{Bias}_{\hat{\sigma}^2}(\sigma^2) &= E\left(\frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2\right) - \sigma^2 \\&= \frac{n-1}{n} E\left(\frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2\right) - \sigma^2 \\&= \frac{n-1}{n} E(S^2) - \sigma^2 \\&= \frac{n-1}{n} \sigma^2 - \sigma^2\end{aligned}$$

## Example: Bias (continued)

Find the bias of  $\hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2$

$$\begin{aligned}\text{Bias}_{\hat{\sigma}^2}(\sigma^2) &= E\left(\frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2\right) - \sigma^2 \\&= \frac{n-1}{n} E\left(\frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2\right) - \sigma^2 \\&= \frac{n-1}{n} E(S^2) - \sigma^2 \\&= \frac{n-1}{n} \sigma^2 - \sigma^2 \\&= \frac{-\sigma^2}{n} < 0 \Rightarrow \hat{\sigma}^2 \text{ tends to underestimate } \sigma^2\end{aligned}$$

## Mean Squared Error

The *mean squared error* of an estimator  $\hat{\theta}$  is defined to be

$$\text{MSE}(\hat{\theta}) = \text{E} \left[ (\hat{\theta} - \theta)^2 \right]$$

# Mean Squared Error

The *mean squared error* of an estimator  $\hat{\theta}$  is defined to be

$$\text{MSE}(\hat{\theta}) = \text{E} \left[ (\hat{\theta} - \theta)^2 \right]$$

*Bias-variance tradeoff*: the mean squared error has the following decomposition

# Mean Squared Error

The *mean squared error* of an estimator  $\hat{\theta}$  is defined to be

$$\text{MSE}(\hat{\theta}) = \text{E} \left[ (\hat{\theta} - \theta)^2 \right]$$

*Bias-variance tradeoff:* the mean squared error has the following decomposition

$$\text{MSE}(\hat{\theta}) = \text{E} \left[ (\hat{\theta} - \theta)^2 \right]$$

# Mean Squared Error

The *mean squared error* of an estimator  $\hat{\theta}$  is defined to be

$$\text{MSE}(\hat{\theta}) = \text{E} \left[ (\hat{\theta} - \theta)^2 \right]$$

*Bias-variance tradeoff:* the mean squared error has the following decomposition

$$\text{MSE}(\hat{\theta}) = \text{E} \left[ (\hat{\theta} - \theta)^2 \right] = \text{E} \left[ \left( [\hat{\theta} - \text{E}(\hat{\theta})] + [\text{E}(\hat{\theta}) - \theta] \right)^2 \right]$$

# Mean Squared Error

The *mean squared error* of an estimator  $\hat{\theta}$  is defined to be

$$\text{MSE}(\hat{\theta}) = \text{E} \left[ (\hat{\theta} - \theta)^2 \right]$$

*Bias-variance tradeoff:* the mean squared error has the following decomposition

$$\begin{aligned} \text{MSE}(\hat{\theta}) &= \text{E} \left[ (\hat{\theta} - \theta)^2 \right] = \text{E} \left[ \left( [\hat{\theta} - \text{E}(\hat{\theta})] + [\text{E}(\hat{\theta}) - \theta] \right)^2 \right] \\ &= \text{E}[\hat{\theta} - \text{E}(\hat{\theta})]^2 - 2\text{E} \left[ [\hat{\theta} - \text{E}(\hat{\theta})][\text{E}(\hat{\theta}) - \theta] \right] + \text{E}[\text{E}(\hat{\theta}) - \theta]^2 \end{aligned}$$



# Mean Squared Error

The *mean squared error* of an estimator  $\hat{\theta}$  is defined to be

$$\text{MSE}(\hat{\theta}) = \text{E} \left[ (\hat{\theta} - \theta)^2 \right]$$

*Bias-variance tradeoff:* the mean squared error has the following decomposition

$$\begin{aligned} \text{MSE}(\hat{\theta}) &= \text{E} \left[ (\hat{\theta} - \theta)^2 \right] = \text{E} \left[ \left( [\hat{\theta} - \text{E}(\hat{\theta})] + [\text{E}(\hat{\theta}) - \theta] \right)^2 \right] \\ &= \text{E}[\hat{\theta} - \text{E}(\hat{\theta})]^2 - 2\text{E} \left[ [\hat{\theta} - \text{E}(\hat{\theta})][\text{E}(\hat{\theta}) - \theta] \right] + \text{E}[\text{E}(\hat{\theta}) - \theta]^2 \\ &= \text{Var}(\hat{\theta}) + [\text{Bias}_{\hat{\theta}}(\theta)]^2 \end{aligned}$$

## Bias-Variance Tradeoff (illustration)



# Efficiency

Let  $\hat{\theta}_1$  and  $\hat{\theta}_2$  be two unbiased estimators of same parameter  $\theta$

- ▶  $\text{MSE}(\hat{\theta}_1) = \text{Var}(\hat{\theta}_1) + [\text{Bias}_{\hat{\theta}_1}(\theta)]^2 = \text{Var}(\hat{\theta}_1)$
- ▶  $\text{MSE}(\hat{\theta}_2) = \text{Var}(\hat{\theta}_2) + [\text{Bias}_{\hat{\theta}_2}(\theta)]^2 = \text{Var}(\hat{\theta}_2)$

# Efficiency

Let  $\hat{\theta}_1$  and  $\hat{\theta}_2$  be two unbiased estimators of same parameter  $\theta$

- ▶  $\text{MSE}(\hat{\theta}_1) = \text{Var}(\hat{\theta}_1) + [\text{Bias}_{\hat{\theta}_1}(\theta)]^2 = \text{Var}(\hat{\theta}_1)$
- ▶  $\text{MSE}(\hat{\theta}_2) = \text{Var}(\hat{\theta}_2) + [\text{Bias}_{\hat{\theta}_2}(\theta)]^2 = \text{Var}(\hat{\theta}_2)$

$\hat{\theta}_1$  is more *efficient* than  $\hat{\theta}_2$  if

$$\text{Var}(\hat{\theta}_1) \leq \text{Var}(\hat{\theta}_2)$$

# Efficiency

Let  $\hat{\theta}_1$  and  $\hat{\theta}_2$  be two unbiased estimators of same parameter  $\theta$

- ▶  $\text{MSE}(\hat{\theta}_1) = \text{Var}(\hat{\theta}_1) + [\text{Bias}_{\hat{\theta}_1}(\theta)]^2 = \text{Var}(\hat{\theta}_1)$
- ▶  $\text{MSE}(\hat{\theta}_2) = \text{Var}(\hat{\theta}_2) + [\text{Bias}_{\hat{\theta}_2}(\theta)]^2 = \text{Var}(\hat{\theta}_2)$

$\hat{\theta}_1$  is more *efficient* than  $\hat{\theta}_2$  if

$$\text{Var}(\hat{\theta}_1) \leq \text{Var}(\hat{\theta}_2)$$

In this case, the estimator  $\hat{\theta}_1$  is preferable

# Efficiency

Let  $\hat{\theta}_1$  and  $\hat{\theta}_2$  be two unbiased estimators of same parameter  $\theta$

- ▶  $\text{MSE}(\hat{\theta}_1) = \text{Var}(\hat{\theta}_1) + [\text{Bias}_{\hat{\theta}_1}(\theta)]^2 = \text{Var}(\hat{\theta}_1)$
- ▶  $\text{MSE}(\hat{\theta}_2) = \text{Var}(\hat{\theta}_2) + [\text{Bias}_{\hat{\theta}_2}(\theta)]^2 = \text{Var}(\hat{\theta}_2)$

$\hat{\theta}_1$  is more *efficient* than  $\hat{\theta}_2$  if

$$\text{Var}(\hat{\theta}_1) \leq \text{Var}(\hat{\theta}_2)$$

In this case, the estimator  $\hat{\theta}_1$  is preferable

**Example** Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$

# Efficiency

Let  $\hat{\theta}_1$  and  $\hat{\theta}_2$  be two unbiased estimators of same parameter  $\theta$

- ▶  $\text{MSE}(\hat{\theta}_1) = \text{Var}(\hat{\theta}_1) + [\text{Bias}_{\hat{\theta}_1}(\theta)]^2 = \text{Var}(\hat{\theta}_1)$
- ▶  $\text{MSE}(\hat{\theta}_2) = \text{Var}(\hat{\theta}_2) + [\text{Bias}_{\hat{\theta}_2}(\theta)]^2 = \text{Var}(\hat{\theta}_2)$

$\hat{\theta}_1$  is more *efficient* than  $\hat{\theta}_2$  if

$$\text{Var}(\hat{\theta}_1) \leq \text{Var}(\hat{\theta}_2)$$

In this case, the estimator  $\hat{\theta}_1$  is preferable

**Example** Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$

- ▶ Sampling distribution of the median  $M$  is  $\approx \mathcal{N}(\mu, \frac{\sigma^2 \pi}{2n})$

# Efficiency

Let  $\hat{\theta}_1$  and  $\hat{\theta}_2$  be two unbiased estimators of same parameter  $\theta$

- ▶  $\text{MSE}(\hat{\theta}_1) = \text{Var}(\hat{\theta}_1) + [\text{Bias}_{\hat{\theta}_1}(\theta)]^2 = \text{Var}(\hat{\theta}_1)$
- ▶  $\text{MSE}(\hat{\theta}_2) = \text{Var}(\hat{\theta}_2) + [\text{Bias}_{\hat{\theta}_2}(\theta)]^2 = \text{Var}(\hat{\theta}_2)$

$\hat{\theta}_1$  is more *efficient* than  $\hat{\theta}_2$  if

$$\text{Var}(\hat{\theta}_1) \leq \text{Var}(\hat{\theta}_2)$$

In this case, the estimator  $\hat{\theta}_1$  is preferable

**Example** Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$

- ▶ Sampling distribution of the median  $M$  is  $\approx \mathcal{N}(\mu, \frac{\sigma^2 \pi}{2n})$
- ▶ Sampling distribution of the sample mean  $\bar{X}$  is  $\mathcal{N}(\mu, \frac{\sigma^2}{n})$



# Efficiency

Let  $\hat{\theta}_1$  and  $\hat{\theta}_2$  be two unbiased estimators of same parameter  $\theta$

- ▶  $\text{MSE}(\hat{\theta}_1) = \text{Var}(\hat{\theta}_1) + [\text{Bias}_{\hat{\theta}_1}(\theta)]^2 = \text{Var}(\hat{\theta}_1)$
- ▶  $\text{MSE}(\hat{\theta}_2) = \text{Var}(\hat{\theta}_2) + [\text{Bias}_{\hat{\theta}_2}(\theta)]^2 = \text{Var}(\hat{\theta}_2)$

$\hat{\theta}_1$  is more *efficient* than  $\hat{\theta}_2$  if

$$\text{Var}(\hat{\theta}_1) \leq \text{Var}(\hat{\theta}_2)$$

In this case, the estimator  $\hat{\theta}_1$  is preferable

**Example** Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$

- ▶ Sampling distribution of the median  $M$  is  $\approx \mathcal{N}(\mu, \frac{\sigma^2 \pi}{2n})$
- ▶ Sampling distribution of the sample mean  $\bar{X}$  is  $\mathcal{N}(\mu, \frac{\sigma^2}{n})$
- ▶ Which estimator of  $\mu$  is preferable,  $\bar{X}$  or  $M$ ?

## Consistency

Let  $X_1, X_2, \dots$  be a sequence of random variables

## Consistency

Let  $X_1, X_2, \dots$  be a sequence of random variables

Let  $\bar{X}_1 = X_1$ ,  $\bar{X}_2 = (X_1 + X_2)/2$ ,  $\bar{X}_3 = (X_1 + X_2 + X_3)/3$ ,  $\dots$

## Consistency

Let  $X_1, X_2, \dots$  be a sequence of random variables

Let  $\bar{X}_1 = X_1$ ,  $\bar{X}_2 = (X_1 + X_2)/2$ ,  $\bar{X}_3 = (X_1 + X_2 + X_3)/3$ ,  $\dots$

Call  $\bar{X}_1, \bar{X}_2, \bar{X}_3, \dots$  a sequence of estimators

# Consistency

Let  $X_1, X_2, \dots$  be a sequence of random variables

Let  $\bar{X}_1 = X_1$ ,  $\bar{X}_2 = (X_1 + X_2)/2$ ,  $\bar{X}_3 = (X_1 + X_2 + X_3)/3$ ,  $\dots$

Call  $\bar{X}_1, \bar{X}_2, \bar{X}_3, \dots$  a sequence of estimators

A sequence of estimators  $W_n = W_n(X_1, \dots, X_n)$  is a *consistent sequence of estimators* of the parameter  $\theta$  if for every  $\epsilon > 0$  and every  $\theta$

$$\lim_{n \rightarrow \infty} P(|W_n - \theta| < \epsilon) = 1$$

## Example: Consistency

Let  $X_1, X_2, \dots$  be a sequence of iid  $\mathcal{N}(\theta, 1)$  random variables

## Example: Consistency

Let  $X_1, X_2, \dots$  be a sequence of iid  $\mathcal{N}(\theta, 1)$  random variables

Recall  $\bar{X} \sim \mathcal{N}(\theta, \frac{1}{n})$

## Example: Consistency

Let  $X_1, X_2, \dots$  be a sequence of iid  $\mathcal{N}(\theta, 1)$  random variables

Recall  $\bar{X} \sim \mathcal{N}(\theta, \frac{1}{n})$

$$P(|\bar{X}_n - \theta| < \epsilon) = \int_{\theta - \epsilon}^{\theta + \epsilon} \left(\frac{n}{2\pi}\right)^{\frac{1}{2}} e^{-(n/2)(\bar{x}_n - \theta)^2} d\bar{x}_n$$



## Example: Consistency

Let  $X_1, X_2, \dots$  be a sequence of iid  $\mathcal{N}(\theta, 1)$  random variables

Recall  $\bar{X} \sim \mathcal{N}(\theta, \frac{1}{n})$

$$\begin{aligned} \mathbb{P}(|\bar{X}_n - \theta| < \epsilon) &= \int_{\theta - \epsilon}^{\theta + \epsilon} \left(\frac{n}{2\pi}\right)^{\frac{1}{2}} e^{-(n/2)(\bar{x}_n - \theta)^2} d\bar{x}_n \\ &= \int_{-\epsilon}^{\epsilon} \left(\frac{n}{2\pi}\right)^{\frac{1}{2}} e^{-(n/2)y^2} dy \end{aligned}$$

## Example: Consistency

Let  $X_1, X_2, \dots$  be a sequence of iid  $\mathcal{N}(\theta, 1)$  random variables

Recall  $\bar{X} \sim \mathcal{N}(\theta, \frac{1}{n})$

$$\begin{aligned} \mathbb{P}(|\bar{X}_n - \theta| < \epsilon) &= \int_{\theta - \epsilon}^{\theta + \epsilon} \left(\frac{n}{2\pi}\right)^{\frac{1}{2}} e^{-(n/2)(\bar{x}_n - \theta)^2} d\bar{x}_n \\ &= \int_{-\epsilon}^{\epsilon} \left(\frac{n}{2\pi}\right)^{\frac{1}{2}} e^{-(n/2)y^2} dy \\ &= \int_{-\epsilon\sqrt{n}}^{\epsilon\sqrt{n}} \frac{1}{\sqrt{2\pi}} e^{-t^2/2} dt \end{aligned}$$

## Example: Consistency

Let  $X_1, X_2, \dots$  be a sequence of iid  $\mathcal{N}(\theta, 1)$  random variables

Recall  $\bar{X} \sim \mathcal{N}(\theta, \frac{1}{n})$

$$\begin{aligned} \mathbb{P}(|\bar{X}_n - \theta| < \epsilon) &= \int_{\theta-\epsilon}^{\theta+\epsilon} \left(\frac{n}{2\pi}\right)^{\frac{1}{2}} e^{-(n/2)(\bar{x}_n - \theta)^2} d\bar{x}_n \\ &= \int_{-\epsilon}^{\epsilon} \left(\frac{n}{2\pi}\right)^{\frac{1}{2}} e^{-(n/2)y^2} dy \\ &= \int_{-\epsilon\sqrt{n}}^{\epsilon\sqrt{n}} \frac{1}{\sqrt{2\pi}} e^{-t^2/2} dt \\ &= \mathbb{P}(-\epsilon\sqrt{n} < Z < \epsilon\sqrt{n}) \end{aligned}$$

## Example: Consistency

Let  $X_1, X_2, \dots$  be a sequence of iid  $\mathcal{N}(\theta, 1)$  random variables

Recall  $\bar{X} \sim \mathcal{N}(\theta, \frac{1}{n})$

$$\begin{aligned} P(|\bar{X}_n - \theta| < \epsilon) &= \int_{\theta-\epsilon}^{\theta+\epsilon} \left(\frac{n}{2\pi}\right)^{\frac{1}{2}} e^{-(n/2)(\bar{x}_n - \theta)^2} d\bar{x}_n \\ &= \int_{-\epsilon}^{\epsilon} \left(\frac{n}{2\pi}\right)^{\frac{1}{2}} e^{-(n/2)y^2} dy \\ &= \int_{-\epsilon\sqrt{n}}^{\epsilon\sqrt{n}} \frac{1}{\sqrt{2\pi}} e^{-t^2/2} dt \\ &= P(-\epsilon\sqrt{n} < Z < \epsilon\sqrt{n}) \\ &\rightarrow 1 \text{ as } n \rightarrow \infty \end{aligned}$$

## Example: Consistency

Let  $X_1, X_2, \dots$  be a sequence of iid  $\mathcal{N}(\theta, 1)$  random variables

Recall  $\bar{X} \sim \mathcal{N}(\theta, \frac{1}{n})$

$$\begin{aligned} P(|\bar{X}_n - \theta| < \epsilon) &= \int_{\theta - \epsilon}^{\theta + \epsilon} \left(\frac{n}{2\pi}\right)^{\frac{1}{2}} e^{-(n/2)(\bar{x}_n - \theta)^2} d\bar{x}_n \\ &= \int_{-\epsilon}^{\epsilon} \left(\frac{n}{2\pi}\right)^{\frac{1}{2}} e^{-(n/2)y^2} dy \\ &= \int_{-\epsilon\sqrt{n}}^{\epsilon\sqrt{n}} \frac{1}{\sqrt{2\pi}} e^{-t^2/2} dt \\ &= P(-\epsilon\sqrt{n} < Z < \epsilon\sqrt{n}) \\ &\rightarrow 1 \text{ as } n \rightarrow \infty \end{aligned}$$

$\implies \bar{X}_n$  is a consistent sequence of estimators of  $\theta$

## The Delta Method

Let  $X_1, X_2, \dots$  be iid random variables with  $E(X_i) = \mu$ ,  $\text{Var}(X_i) = \sigma^2$

# The Delta Method

Let  $X_1, X_2, \dots$  be iid random variables with  $E(X_i) = \mu$ ,  $\text{Var}(X_i) = \sigma^2$

Central Limit Theorem says

$$\frac{(\bar{X}_n - \mu)}{\sigma/\sqrt{n}} \rightarrow Z \quad Z \sim \mathcal{N}(0, 1)$$

# The Delta Method

Let  $X_1, X_2, \dots$  be iid random variables with  $E(X_i) = \mu$ ,  $\text{Var}(X_i) = \sigma^2$

Central Limit Theorem says

$$\frac{(\bar{X}_n - \mu)}{\sigma/\sqrt{n}} \rightarrow Z \quad Z \sim \mathcal{N}(0, 1)$$

Let  $g(t) = at + b$  be a linear function



# The Delta Method

Let  $X_1, X_2, \dots$  be iid random variables with  $E(X_i) = \mu$ ,  $\text{Var}(X_i) = \sigma^2$

Central Limit Theorem says

$$\frac{(\bar{X}_n - \mu)}{\sigma/\sqrt{n}} \rightarrow Z \quad Z \sim \mathcal{N}(0, 1)$$

Let  $g(t) = at + b$  be a linear function

►  $E(g(\bar{X}_n)) = E(a\bar{X}_n + b) = aE(\bar{X}_n) + b = a\mu + b$

# The Delta Method

Let  $X_1, X_2, \dots$  be iid random variables with  $E(X_i) = \mu$ ,  $\text{Var}(X_i) = \sigma^2$

Central Limit Theorem says

$$\frac{(\bar{X}_n - \mu)}{\sigma/\sqrt{n}} \rightarrow Z \quad Z \sim \mathcal{N}(0, 1)$$

Let  $g(t) = at + b$  be a linear function

- ▶  $E(g(\bar{X}_n)) = E(a\bar{X}_n + b) = aE(\bar{X}_n) + b = a\mu + b$
- ▶  $\text{Var}(g(\bar{X}_n)) = \text{Var}(a\bar{X}_n + b) = a^2 \text{Var}(\bar{X}_n) = \frac{a^2 \sigma^2}{n}$

# The Delta Method

Let  $X_1, X_2, \dots$  be iid random variables with  $E(X_i) = \mu$ ,  $\text{Var}(X_i) = \sigma^2$

Central Limit Theorem says

$$\frac{(\bar{X}_n - \mu)}{\sigma/\sqrt{n}} \rightarrow Z \quad Z \sim \mathcal{N}(0, 1)$$

Let  $g(t) = at + b$  be a linear function

- ▶  $E(g(\bar{X}_n)) = E(a\bar{X}_n + b) = aE(\bar{X}_n) + b = a\mu + b$
- ▶  $\text{Var}(g(\bar{X}_n)) = \text{Var}(a\bar{X}_n + b) = a^2 \text{Var}(\bar{X}_n) = \frac{a^2 \sigma^2}{n}$

Central Limit Theorem result for  $g(\bar{X}_n)$

$$\frac{(g(\bar{X}_n) - g(\mu))}{a\sigma/\sqrt{n}} \rightarrow Z \quad Z \sim \mathcal{N}(0, 1)$$

# The Delta Method

Let  $X_1, X_2, \dots$  be iid random variables with  $E(X_i) = \mu$ ,  $\text{Var}(X_i) = \sigma^2$

Central Limit Theorem says

$$\frac{(\bar{X}_n - \mu)}{\sigma/\sqrt{n}} \rightarrow Z \quad Z \sim \mathcal{N}(0, 1)$$

Let  $g(t) = at + b$  be a linear function

- ▶  $E(g(\bar{X}_n)) = E(a\bar{X}_n + b) = aE(\bar{X}_n) + b = a\mu + b$
- ▶  $\text{Var}(g(\bar{X}_n)) = \text{Var}(a\bar{X}_n + b) = a^2 \text{Var}(\bar{X}_n) = \frac{a^2 \sigma^2}{n}$

Central Limit Theorem result for  $g(\bar{X}_n)$

$$\frac{(g(\bar{X}_n) - g(\mu))}{a\sigma/\sqrt{n}} \rightarrow Z \quad Z \sim \mathcal{N}(0, 1)$$

Can write as

$$\sqrt{n}(g(\bar{X}_n) - g(\mu)) \rightarrow a\sigma Z$$

# The Delta Method

Now, suppose  $g(t)$  is a nonlinear function and that  $g'(\mu)$  exists

## The Delta Method

Now, suppose  $g(t)$  is a nonlinear function and that  $g'(\mu)$  exists

If  $\bar{X}_n$  is a consistent estimator of  $\mu$ , then  $\bar{X}_n$  close to  $\mu$  for large  $n$

# The Delta Method

Now, suppose  $g(t)$  is a nonlinear function and that  $g'(\mu)$  exists

If  $\bar{X}_n$  is a consistent estimator of  $\mu$ , then  $\bar{X}_n$  close to  $\mu$  for large  $n$

Can approximate  $g(t)$  with a linear function

$$g(t) \approx g(\mu) + g'(\mu)(t - \mu)$$

# The Delta Method

Now, suppose  $g(t)$  is a nonlinear function and that  $g'(\mu)$  exists

If  $\bar{X}_n$  is a consistent estimator of  $\mu$ , then  $\bar{X}_n$  close to  $\mu$  for large  $n$

Can approximate  $g(t)$  with a linear function

$$g(t) \approx g(\mu) + g'(\mu)(t - \mu)$$

$$\sqrt{n}(g(\bar{X}_n) - g(\mu)) \rightarrow g'(\mu)\sigma Z \sim \mathcal{N}(0, \sigma^2[g'(\mu)]^2)$$



## Example: Estimating Binomial Variance

Let  $X_n \sim \text{binomial}(n, p)$  (sum of  $n$  independent Bernoulli( $p$ ) trials)

## Example: Estimating Binomial Variance

Let  $X_n \sim \text{binomial}(n, p)$  (sum of  $n$  independent Bernoulli( $p$ ) trials)

MLE  $\frac{1}{n}X_n$  consistent, unbiased estimator of  $p$

## Example: Estimating Binomial Variance

Let  $X_n \sim \text{binomial}(n, p)$  (sum of  $n$  independent Bernoulli( $p$ ) trials)

MLE  $\frac{1}{n}X_n$  consistent, unbiased estimator of  $p$

Let  $g(t) = t(1 - t)$

$$\implies g\left(\frac{X_n}{n}\right) = \frac{X_n}{n} \left(1 - \frac{X_n}{n}\right) = \frac{X_n(n - X_n)}{n^2}$$

an estimator for the variance  $p(1 - p)$

## Example: Estimating Binomial Variance

Let  $X_n \sim \text{binomial}(n, p)$  (sum of  $n$  independent Bernoulli( $p$ ) trials)

MLE  $\frac{1}{n}X_n$  consistent, unbiased estimator of  $p$

Let  $g(t) = t(1 - t)$

$$\implies g\left(\frac{X_n}{n}\right) = \frac{X_n}{n} \left(1 - \frac{X_n}{n}\right) = \frac{X_n(n - X_n)}{n^2}$$

an estimator for the variance  $p(1 - p)$

Since  $g'(p) = 1 - 2p$

$$\sqrt{n} \left( g\left(\frac{X_n}{n}\right) - p(1 - p) \right) \rightarrow \mathcal{N}(0, p(1 - p)(1 - 2p)^2)$$

# Outline

Statistical Models

Point Estimation

Evaluating Point Estimators

Interval Estimators

# Interval Estimators

Instead of estimating the parameter  $\theta$  with only a number  $\hat{\theta}$ , consider a random interval that covers  $\theta$  with high probability

# Interval Estimators

Instead of estimating the parameter  $\theta$  with only a number  $\hat{\theta}$ , consider a random interval that covers  $\theta$  with high probability

Let  $\hat{\theta}$  be an estimator of  $\theta$  and suppose that there exists a function  $T = q(\hat{\theta}, \theta)$  whose distribution is known and does not depend on  $\theta$

# Interval Estimators

Instead of estimating the parameter  $\theta$  with only a number  $\hat{\theta}$ , consider a random interval that covers  $\theta$  with high probability

Let  $\hat{\theta}$  be an estimator of  $\theta$  and suppose that there exists a function  $T = q(\hat{\theta}, \theta)$  whose distribution is known and does not depend on  $\theta$

Let  $\alpha_1 + \alpha_2 = \alpha < 1$  and let  $a$  and  $b$  be constants such that

$$\alpha_1 = P(T < a) \quad \alpha_2 = P(T > b) \quad (\text{often } \alpha_1 = \alpha_2)$$



# Interval Estimators

Instead of estimating the parameter  $\theta$  with only a number  $\hat{\theta}$ , consider a random interval that covers  $\theta$  with high probability

Let  $\hat{\theta}$  be an estimator of  $\theta$  and suppose that there exists a function  $T = q(\hat{\theta}, \theta)$  whose distribution is known and does not depend on  $\theta$

Let  $\alpha_1 + \alpha_2 = \alpha < 1$  and let  $a$  and  $b$  be constants such that

$$\alpha_1 = P(T < a) \quad \alpha_2 = P(T > b) \quad (\text{often } \alpha_1 = \alpha_2)$$

Then

$$P(a \leq T \leq b) = P(T \leq b) - P(T < a) = (1 - \alpha_2) - \alpha_1 = 1 - \alpha$$

# Interval Estimators

Instead of estimating the parameter  $\theta$  with only a number  $\hat{\theta}$ , consider a random interval that covers  $\theta$  with high probability

Let  $\hat{\theta}$  be an estimator of  $\theta$  and suppose that there exists a function  $T = q(\hat{\theta}, \theta)$  whose distribution is known and does not depend on  $\theta$

Let  $\alpha_1 + \alpha_2 = \alpha < 1$  and let  $a$  and  $b$  be constants such that

$$\alpha_1 = P(T < a) \quad \alpha_2 = P(T > b) \quad (\text{often } \alpha_1 = \alpha_2)$$

Then

$$P(a \leq T \leq b) = P(T \leq b) - P(T < a) = (1 - \alpha_2) - \alpha_1 = 1 - \alpha$$

Isolate  $\theta$  to find random variables  $I$  and  $S$  such that

$$P(I \leq \theta \leq S) = 1 - \alpha$$

# Interval Estimators

Instead of estimating the parameter  $\theta$  with only a number  $\hat{\theta}$ , consider a random interval that covers  $\theta$  with high probability

Let  $\hat{\theta}$  be an estimator of  $\theta$  and suppose that there exists a function  $T = q(\hat{\theta}, \theta)$  whose distribution is known and does not depend on  $\theta$

Let  $\alpha_1 + \alpha_2 = \alpha < 1$  and let  $a$  and  $b$  be constants such that

$$\alpha_1 = P(T < a) \quad \alpha_2 = P(T > b) \quad (\text{often } \alpha_1 = \alpha_2)$$

Then

$$P(a \leq T \leq b) = P(T \leq b) - P(T < a) = (1 - \alpha_2) - \alpha_1 = 1 - \alpha$$

Isolate  $\theta$  to find random variables  $I$  and  $S$  such that

$$P(I \leq \theta \leq S) = 1 - \alpha$$

$[I, S]$  is called a  $100(1 - \alpha)\%$  *confidence interval* for  $\theta$

# Interval Estimators

Instead of estimating the parameter  $\theta$  with only a number  $\hat{\theta}$ , consider a random interval that covers  $\theta$  with high probability

Let  $\hat{\theta}$  be an estimator of  $\theta$  and suppose that there exists a function  $T = q(\hat{\theta}, \theta)$  whose distribution is known and does not depend on  $\theta$

Let  $\alpha_1 + \alpha_2 = \alpha < 1$  and let  $a$  and  $b$  be constants such that

$$\alpha_1 = P(T < a) \quad \alpha_2 = P(T > b) \quad (\text{often } \alpha_1 = \alpha_2)$$

Then

$$P(a \leq T \leq b) = P(T \leq b) - P(T < a) = (1 - \alpha_2) - \alpha_1 = 1 - \alpha$$

Isolate  $\theta$  to find random variables  $I$  and  $S$  such that

$$P(I \leq \theta \leq S) = 1 - \alpha$$

$[I, S]$  is called a  $100(1 - \alpha)\%$  *confidence interval* for  $\theta$

The quantity  $1 - \alpha$  is called the *confidence coefficient*

## Example: Normal Confidence Interval, $\sigma^2$ Known

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, 1^2)$

## Example: Normal Confidence Interval, $\sigma^2$ Known

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, 1^2)$

Find a  $(1 - 0.05)100\%$  symmetric confidence interval for the mean  $\mu$

## Example: Normal Confidence Interval, $\sigma^2$ Known

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, 1^2)$

Find a  $(1 - 0.05)100\%$  symmetric confidence interval for the mean  $\mu$

- ▶  $\alpha = 0.05$ , symmetric  $\implies \alpha_1 = \alpha_2 = 0.025$

## Example: Normal Confidence Interval, $\sigma^2$ Known

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, 1^2)$

Find a  $(1 - 0.05)100\%$  symmetric confidence interval for the mean  $\mu$

- ▶  $\alpha = 0.05$ , symmetric  $\implies \alpha_1 = \alpha_2 = 0.025$
- ▶  $Z = \sqrt{n}(\bar{X} - \mu) \sim \mathcal{N}(0, 1)$



## Example: Normal Confidence Interval, $\sigma^2$ Known

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, 1^2)$

Find a  $(1 - 0.05)100\%$  symmetric confidence interval for the mean  $\mu$

- ▶  $\alpha = 0.05$ , symmetric  $\implies \alpha_1 = \alpha_2 = 0.025$
- ▶  $Z = \sqrt{n}(\bar{X} - \mu) \sim \mathcal{N}(0, 1)$
- ▶  $\{\Phi(a) = 0.025, \Phi(b) = 0.975\} \Rightarrow \{a = z_{0.025}, b = z_{0.975}\}$

## Example: Normal Confidence Interval, $\sigma^2$ Known

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, 1^2)$

Find a  $(1 - 0.05)100\%$  symmetric confidence interval for the mean  $\mu$

- ▶  $\alpha = 0.05$ , symmetric  $\implies \alpha_1 = \alpha_2 = 0.025$
- ▶  $Z = \sqrt{n}(\bar{X} - \mu) \sim \mathcal{N}(0, 1)$
- ▶  $\{\Phi(a) = 0.025, \Phi(b) = 0.975\} \Rightarrow \{a = z_{0.025}, b = z_{0.975}\}$
- ▶  $P(-1.96 \leq \sqrt{n}(\bar{X} - \mu) \leq 1.96) = 0.95$

## Example: Normal Confidence Interval, $\sigma^2$ Known

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, 1^2)$

Find a  $(1 - 0.05)100\%$  symmetric confidence interval for the mean  $\mu$

- ▶  $\alpha = 0.05$ , symmetric  $\implies \alpha_1 = \alpha_2 = 0.025$
- ▶  $Z = \sqrt{n}(\bar{X} - \mu) \sim \mathcal{N}(0, 1)$
- ▶  $\{\Phi(a) = 0.025, \Phi(b) = 0.975\} \Rightarrow \{a = z_{0.025}, b = z_{0.975}\}$
- ▶  $P(-1.96 \leq \sqrt{n}(\bar{X} - \mu) \leq 1.96) = 0.95$
- ▶ Isolate the parameter:

$$\frac{-1.96}{\sqrt{n}} \leq \bar{X} - \mu \leq \frac{1.96}{\sqrt{n}}$$

## Example: Normal Confidence Interval, $\sigma^2$ Known

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, 1^2)$

Find a  $(1 - 0.05)100\%$  symmetric confidence interval for the mean  $\mu$

- ▶  $\alpha = 0.05$ , symmetric  $\implies \alpha_1 = \alpha_2 = 0.025$
- ▶  $Z = \sqrt{n}(\bar{X} - \mu) \sim \mathcal{N}(0, 1)$
- ▶  $\{\Phi(a) = 0.025, \Phi(b) = 0.975\} \Rightarrow \{a = z_{0.025}, b = z_{0.975}\}$
- ▶  $P(-1.96 \leq \sqrt{n}(\bar{X} - \mu) \leq 1.96) = 0.95$
- ▶ Isolate the parameter:

$$\frac{-1.96}{\sqrt{n}} \leq \bar{X} - \mu \leq \frac{1.96}{\sqrt{n}}$$

$$-\bar{X} - \frac{1.96}{\sqrt{n}} \leq -\mu \leq -\bar{X} + \frac{1.96}{\sqrt{n}}$$

## Example: Normal Confidence Interval, $\sigma^2$ Known

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, 1^2)$

Find a  $(1 - 0.05)100\%$  symmetric confidence interval for the mean  $\mu$

- ▶  $\alpha = 0.05$ , symmetric  $\implies \alpha_1 = \alpha_2 = 0.025$
- ▶  $Z = \sqrt{n}(\bar{X} - \mu) \sim \mathcal{N}(0, 1)$
- ▶  $\{\Phi(a) = 0.025, \Phi(b) = 0.975\} \Rightarrow \{a = z_{0.025}, b = z_{0.975}\}$
- ▶  $P(-1.96 \leq \sqrt{n}(\bar{X} - \mu) \leq 1.96) = 0.95$
- ▶ Isolate the parameter:

$$\frac{-1.96}{\sqrt{n}} \leq \bar{X} - \mu \leq \frac{1.96}{\sqrt{n}}$$

$$-\bar{X} - \frac{1.96}{\sqrt{n}} \leq -\mu \leq -\bar{X} + \frac{1.96}{\sqrt{n}}$$

$$\bar{X} - \frac{1.96}{\sqrt{n}} \leq \mu \leq \bar{X} + \frac{1.96}{\sqrt{n}}$$

## Example: Normal Confidence Interval, $\sigma^2$ Known

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, 1^2)$

Find a  $(1 - 0.05)100\%$  symmetric confidence interval for the mean  $\mu$

- ▶  $\alpha = 0.05$ , symmetric  $\implies \alpha_1 = \alpha_2 = 0.025$
- ▶  $Z = \sqrt{n}(\bar{X} - \mu) \sim \mathcal{N}(0, 1)$
- ▶  $\{\Phi(a) = 0.025, \Phi(b) = 0.975\} \Rightarrow \{a = z_{0.025}, b = z_{0.975}\}$
- ▶  $P(-1.96 \leq \sqrt{n}(\bar{X} - \mu) \leq 1.96) = 0.95$
- ▶ Isolate the parameter:

$$\frac{-1.96}{\sqrt{n}} \leq \bar{X} - \mu \leq \frac{1.96}{\sqrt{n}}$$

$$-\bar{X} - \frac{1.96}{\sqrt{n}} \leq -\mu \leq -\bar{X} + \frac{1.96}{\sqrt{n}}$$

$$\bar{X} - \frac{1.96}{\sqrt{n}} \leq \mu \leq \bar{X} + \frac{1.96}{\sqrt{n}}$$

- ▶  $\left[ \frac{\bar{X} - 1.96}{\sqrt{n}}, \frac{\bar{X} + 1.96}{\sqrt{n}} \right]$  is a 95% confidence interval for  $\mu$

## Example: Normal Confidence Interval, Variance Unknown

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$

## Example: Normal Confidence Interval, Variance Unknown

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$

Find a  $(1 - 0.05)100\%$  symmetric confidence interval for the mean  $\mu$



## Example: Normal Confidence Interval, Variance Unknown

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$

Find a  $(1 - 0.05)100\%$  symmetric confidence interval for the mean  $\mu$

- ▶  $\alpha = 0.05$ , symmetric  $\implies \alpha_1 = \alpha_2 = 0.025$

## Example: Normal Confidence Interval, Variance Unknown

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$

Find a  $(1 - 0.05)100\%$  symmetric confidence interval for the mean  $\mu$

►  $\alpha = 0.05$ , symmetric  $\implies \alpha_1 = \alpha_2 = 0.025$

►  $T = \frac{\bar{X} - \mu}{S/\sqrt{n}} \sim t_{n-1}$

## Example: Normal Confidence Interval, Variance Unknown

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$

Find a  $(1 - 0.05)100\%$  symmetric confidence interval for the mean  $\mu$

- ▶  $\alpha = 0.05$ , symmetric  $\implies \alpha_1 = \alpha_2 = 0.025$
- ▶  $T = \frac{\bar{X} - \mu}{S/\sqrt{n}} \sim t_{n-1}$
- ▶  $P(t_{0.025, n-1} \leq \frac{\bar{X} - \mu}{S/\sqrt{n}} \leq t_{0.975, n-1}) = 0.95$

## Example: Normal Confidence Interval, Variance Unknown

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$

Find a  $(1 - 0.05)100\%$  symmetric confidence interval for the mean  $\mu$

►  $\alpha = 0.05$ , symmetric  $\implies \alpha_1 = \alpha_2 = 0.025$

►  $T = \frac{\bar{X} - \mu}{S/\sqrt{n}} \sim t_{n-1}$

►  $P(t_{0.025, n-1} \leq \frac{\bar{X} - \mu}{S/\sqrt{n}} \leq t_{0.975, n-1}) = 0.95$

► Isolate the parameter:

$$\frac{S}{\sqrt{n}} t_{0.025, n-1} \leq \bar{X} - \mu \leq \frac{S}{\sqrt{n}} t_{0.975, n-1}$$

## Example: Normal Confidence Interval, Variance Unknown

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$

Find a  $(1 - 0.05)100\%$  symmetric confidence interval for the mean  $\mu$

►  $\alpha = 0.05$ , symmetric  $\implies \alpha_1 = \alpha_2 = 0.025$

►  $T = \frac{\bar{X} - \mu}{S/\sqrt{n}} \sim t_{n-1}$

►  $P(t_{0.025, n-1} \leq \frac{\bar{X} - \mu}{S/\sqrt{n}} \leq t_{0.975, n-1}) = 0.95$

► Isolate the parameter:

$$\frac{S}{\sqrt{n}} t_{0.025, n-1} \leq \bar{X} - \mu \leq \frac{S}{\sqrt{n}} t_{0.975, n-1}$$

$$-\bar{X} + \frac{S}{\sqrt{n}} t_{0.025, n-1} \leq -\mu \leq -\bar{X} + \frac{S}{\sqrt{n}} t_{0.975, n-1}$$

## Example: Normal Confidence Interval, Variance Unknown

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$

Find a  $(1 - 0.05)100\%$  symmetric confidence interval for the mean  $\mu$

►  $\alpha = 0.05$ , symmetric  $\implies \alpha_1 = \alpha_2 = 0.025$

►  $T = \frac{\bar{X} - \mu}{S/\sqrt{n}} \sim t_{n-1}$

►  $P(t_{0.025, n-1} \leq \frac{\bar{X} - \mu}{S/\sqrt{n}} \leq t_{0.975, n-1}) = 0.95$

► Isolate the parameter:

$$\frac{S}{\sqrt{n}} t_{0.025, n-1} \leq \bar{X} - \mu \leq \frac{S}{\sqrt{n}} t_{0.975, n-1}$$

$$-\bar{X} + \frac{S}{\sqrt{n}} t_{0.025, n-1} \leq -\mu \leq -\bar{X} + \frac{S}{\sqrt{n}} t_{0.975, n-1}$$

$$\bar{X} + \frac{S}{\sqrt{n}} t_{0.025, n-1} \leq \mu \leq \bar{X} + \frac{S}{\sqrt{n}} t_{0.975, n-1}$$

# Confidence Interval Interpretation

The parameter  $\theta$  is unknown but fixed

# Confidence Interval Interpretation

The parameter  $\theta$  is unknown but fixed      It is *NOT* random



# Confidence Interval Interpretation

The parameter  $\theta$  is unknown but fixed      It is *NOT* random

The confidence interval  $[I, S]$  is a random interval that covers  $\theta$  with probability  $(1 - \alpha)$

# Confidence Interval Interpretation

The parameter  $\theta$  is unknown but fixed     It is *NOT* random

The confidence interval  $[I, S]$  is a random interval that covers  $\theta$  with probability  $(1 - \alpha)$

Let  $A$  be the event  $I \leq \theta$  and let  $B$  be the event  $S \geq \theta$

$$P(A \cap B) = (1 - \alpha)$$

# Confidence Interval Interpretation

The parameter  $\theta$  is unknown but fixed     It is *NOT* random

The confidence interval  $[I, S]$  is a random interval that covers  $\theta$  with probability  $(1 - \alpha)$

Let  $A$  be the event  $I \leq \theta$  and let  $B$  be the event  $S \geq \theta$

$$P(A \cap B) = (1 - \alpha)$$

Saying that  $\theta \in [I, S]$  with probability  $(1 - \alpha)$  is incorrect because it is  $I$  and  $S$  that are random (and not  $\theta$ )

# Confidence Interval Interpretation

The parameter  $\theta$  is unknown but fixed      It is *NOT* random

The confidence interval  $[I, S]$  is a random interval that covers  $\theta$  with probability  $(1 - \alpha)$

Let  $A$  be the event  $I \leq \theta$  and let  $B$  be the event  $S \geq \theta$

$$P(A \cap B) = (1 - \alpha)$$

Saying that  $\theta \in [I, S]$  with probability  $(1 - \alpha)$  is incorrect because it is  $I$  and  $S$  that are random (and not  $\theta$ )




Figure: 30 repetitions of the confidence interval calculation

# Confidence Intervals: Remarks

How can we ...

- ▶ ...increase the probability that the confidence interval covers  $\theta$ ?

# Confidence Intervals: Remarks

How can we ...

- ▶ ... increase the probability that the confidence interval covers  $\theta$ ?
  - ▶ Make  $\alpha$  smaller, then  $(1 - \alpha)$  gets bigger making the confidence interval larger

# Confidence Intervals: Remarks

How can we ...

- ▶ ...increase the probability that the confidence interval covers  $\theta$ ?
  - ▶ Make  $\alpha$  smaller, then  $(1 - \alpha)$  gets bigger making the confidence interval larger
- ▶ ...make the confidence interval shorter?

# Confidence Intervals: Remarks

How can we ...

- ▶ ...increase the probability that the confidence interval covers  $\theta$ ?
  - ▶ Make  $\alpha$  smaller, then  $(1 - \alpha)$  gets bigger making the confidence interval larger
- ▶ ...make the confidence interval shorter?
  - ▶ Confidence intervals for the mean are always of the form

$$\bar{x} \pm \frac{c}{\sqrt{n}}$$



# Confidence Intervals: Remarks

How can we ...

- ▶ ...increase the probability that the confidence interval covers  $\theta$ ?
  - ▶ Make  $\alpha$  smaller, then  $(1 - \alpha)$  gets bigger making the confidence interval larger
- ▶ ...make the confidence interval shorter?
  - ▶ Confidence intervals for the mean are always of the form

$$\bar{x} \pm \frac{c}{\sqrt{n}}$$

- ▶ If  $n$  increases, then the confidence interval gets shorter

# Confidence Intervals: Remarks

How can we ...

- ▶ ... increase the probability that the confidence interval covers  $\theta$ ?
  - ▶ Make  $\alpha$  smaller, then  $(1 - \alpha)$  gets bigger making the confidence interval larger
- ▶ ... make the confidence interval shorter?
  - ▶ Confidence intervals for the mean are always of the form

$$\bar{x} \pm \frac{c}{\sqrt{n}}$$

- ▶ If  $n$  increases, then the confidence interval gets shorter

One-sided confidence intervals

- ▶ Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, 1^2)$

# Confidence Intervals: Remarks

How can we ...

- ▶ ...increase the probability that the confidence interval covers  $\theta$ ?
  - ▶ Make  $\alpha$  smaller, then  $(1 - \alpha)$  gets bigger making the confidence interval larger
- ▶ ...make the confidence interval shorter?
  - ▶ Confidence intervals for the mean are always of the form

$$\bar{x} \pm \frac{c}{\sqrt{n}}$$

- ▶ If  $n$  increases, then the confidence interval gets shorter

One-sided confidence intervals

- ▶ Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, 1^2)$
- ▶ A confidence interval of the form  $(-\infty, \bar{X} + z_{(1-\alpha)}/\sqrt{n}]$  is a one-sided confidence interval for  $\mu$

# Confidence Intervals for Maximum Likelihood Estimators

Often construct confidence intervals based on maximum likelihood estimators

# Confidence Intervals for Maximum Likelihood Estimators

Often construct confidence intervals based on maximum likelihood estimators

**Theorem** Let  $\hat{\theta}_{ML}$  the maximum likelihood estimator of  $\theta$ , then for large  $n$ ,

$$\hat{\theta}_{ML} \dot{\sim} \mathcal{N}\left(\theta, J(\hat{\theta}_{ML})^{-1}\right)$$

# Confidence Intervals for Maximum Likelihood Estimators

Often construct confidence intervals based on maximum likelihood estimators

**Theorem** Let  $\hat{\theta}_{ML}$  the maximum likelihood estimator of  $\theta$ , then for large  $n$ ,

$$\hat{\theta}_{ML} \simeq \mathcal{N}\left(\theta, J(\hat{\theta}_{ML})^{-1}\right)$$

$J(\theta) = -\frac{d^2}{d\theta^2}\ell(\theta)$  is the *observed information* for  $\theta$

# Confidence Intervals for Maximum Likelihood Estimators

Often construct confidence intervals based on maximum likelihood estimators

**Theorem** Let  $\hat{\theta}_{ML}$  the maximum likelihood estimator of  $\theta$ , then for large  $n$ ,

$$\hat{\theta}_{ML} \simeq \mathcal{N}\left(\theta, J(\hat{\theta}_{ML})^{-1}\right)$$

$J(\theta) = -\frac{d^2}{d\theta^2}\ell(\theta)$  is the *observed information* for  $\theta$

Can make a symmetric  $(1 - \alpha)100\%$  confidence interval for  $\theta$

$$\left[ \hat{\theta}_{ML} - z_{1-\frac{\alpha}{2}} J(\hat{\theta}_{ML})^{-\frac{1}{2}}, \hat{\theta}_{ML} + z_{1-\frac{\alpha}{2}} J(\hat{\theta}_{ML})^{-\frac{1}{2}} \right]$$



# COMPUTATIONAL FINANCE & RISK MANAGEMENT

---

UNIVERSITY *of* WASHINGTON

Department of Applied Mathematics

<http://computational-finance.uw.edu>