MA 3475 Exam II Review

1. Find the extremal of the functional

$$J[y,z] = \int_0^1 \left((y')^2 - (z')^2 - 8y'y - 4y^2 \right) dx,$$

subject to the boundary conditions y(0) = 1, y(1) = 0, z(0) = 0, z(1) = e.

2. Find the extremal of the functional

$$J[y,z] = \int_{a}^{b} ((y')^{2} + (z')^{2} + yz) dx$$

3. Find the extremal of the functional

$$J[y] = \int_0^{\pi/2} \left((y')^2 - (y'')^2 \right) dx$$

subject to the boundary conditions $y(0) = 0, y'(0) = 0, y\left(\frac{\pi}{2}\right) = 1, y'\left(\frac{\pi}{2}\right) = 1.$

4. Find the extremal of the functional

$$J[y] = \int_{a}^{b} ((y''')^{2} + (y'')^{2}) dx$$

How many boundary conditions do you need to specify at the two endpoints to determine the extremal uniquely?

5. Find the extremal of the functional

$$J[y] = \int_0^1 \left(x^2 + (y')^2\right) dx$$

subject to the conditions y(0) = 0, y(1) = 0 and the constraint $\int_0^1 (x^2 + 2y(x)) dx = 1$.

6. Find the extremal of the functional

$$J[y] = \int_0^1 \frac{1}{2} (y')^2 dx$$

subject to the conditions y(0) = 0, y(1) = 1 and the constraint $\int_0^1 \frac{1}{2} y^2 dx = 1$. For this problem you may stop after expressing y(x) in terms of the multiplier λ but this should be the only unknown constant. For a challenge, determine λ to finish solving the problem.