

Algebraic Geometry

Daniele Faenzi

August 11, 2026



"My art is not abstract, it lives and breathes"

MARK ROTHKO
No. 3 / No. 13, 1949

Contents

1	Basic material	6
I	Introduction: algebraic sets	6
I.A	Ideals and affine algebraic sets	6
I.A.1	From ideals to algebraic sets	6
I.A.2	From subsets to ideals	14
I.B	Morphisms of affine algebraic sets	15
I.B.1	Morphisms of algebraic sets and polynomial mappings	15
I.B.2	Coordinate rings	16
I.C	Radicals and the Nullstellensatz	18
I.C.1	Radicals	18
I.C.2	Integral ring extensions	19
I.C.3	Hilbert's Nullstellensatz	21
I.C.4	Geometrically regular rational functions	23
I.D	Noether normalisation	25
I.D.1	The algebraic statement	25
I.D.2	Geometric version of Noether normalisation	27
II	Projective algebraic sets	29
II.A	Projective space and projective subspaces	29
II.B	Projective equations and homogenisation	31
II.B.1	Homogeneous ideals and projective subsets	31
II.B.2	Dehomogenisation, affine charts and projective closure	32
II.C	Morphisms of projective algebraic sets	34
II.C.1	Rational functions on projective algebraic sets	34
II.C.2	Polynomial maps in local charts	35
III	Localisation	38
III.A	Localisation and inversion of elements	38
III.B	Properties of the localisation	39
III.B.1	Universal property of localisation	39
III.B.2	Pull-back properties of localisation	39
III.C	Localisation and the fraction field	40
III.D	Local rings	41
IV	Sheaves	42
IV.A	Sheaves of abelian groups	42
IV.B	Morphisms	44
IV.C	Stalks, direct and inverse images	44
IV.C.1	Direct limits	45
IV.C.2	The stalk	45

IV.D	Defining a sheaf from compatible data	47
IV.D.1	The sheafification	47
IV.D.2	The étalé space of a presheaf	47
IV.D.3	Sheaves on a basis of the topology	49
IV.E	Properties of sheaves	56
IV.E.1	Kernels, images, cokernels stalks	56
IV.E.2	Injectivity and surjectivity on stalks	58
IV.F	Functorial construction with sheaves	59
IV.F.1	Direct image	59
IV.F.2	Inverse image	60
2	Affine schemes	61
I	Prime spectrum of a ring	61
I.A	Prime ideals	61
I.B	Functoriality of the prime spectrum	62
II	The Zariski topology	62
II.A	Closed subsets of the affine spectrum	62
II.A.1	Closed sets and ideals	62
II.A.2	First properties of closed sets	63
II.B	Intersection of primes and the radical	64
II.C	Dominant morphisms	65
II.D	Open and principal open subsets	67
II.D.1	Localisation and principal open subsets	68
II.D.2	Residue field	69
II.D.3	Localisation and homeomorphism	71
III	The structure sheaf	72
III.A	The principal-open description and associated modules	73
III.B	Equivalent definition by sheafification	74
IV	Affine schemes as locally ringed spaces	78
IV.A	The affine spectrum as a locally ringed space	78
IV.A.1	The stalk of the structure sheaf and the local ring	78
IV.A.2	Ringed spaces and locally ringed spaces	80
IV.B	Morphisms of locally ringed spaces	81
V	Affine subschemes	87
V.A	Open subsets	87
V.B	Closed affine subschemes	87
VI	Subschemes of the affine space	90
VI.A	Subschemes of the affine line over a field	90
VI.B	Points of the affine plane over a field	91
VI.C	Closed subschemes of affine space over a field	92
VI.D	Arithmetic affine schemes	94
VI.E	Rings of integers in number fields	95
VI.E.1	Number fields	95
VI.E.2	Rings of integers of number fields	95

3	Schemes	97
I	Schemes	97
I.A	Schemes as locally ringed spaces	97
I.A.1	Locally ringed spaces	97
I.A.2	Morphisms of schemes	98
I.B	Quasi-coherent sheaves	98
I.B.1	Generating sections	99
I.B.2	Finite type and finite presentation	99
I.B.3	Quasi-coherent sheaves on an affine scheme	100
I.C	Closed subschemes	101
I.D	Gluing schemes	104
I.D.1	The gluing lemma	105
I.D.2	The line with two origins	108
I.D.3	Projective line	109
I.D.4	Higher-dimensional projective space	109
I.E	Affine targets and affinisation	111
I.E.1	The affine-target adjunction	111
I.E.2	Global sections as left adjoint of affine spectrum	114
I.E.3	The affinisation morphism	115
I.E.4	What affinisation forgets	116
II	Schemes over a base	116
II.A	The category of schemes over a base	116
II.A.1	Morphisms of finite type	117
II.A.2	Schemes over a ring	117
II.A.3	The relative adjunction	118
II.A.4	Schemes over a field	119
II.B	The functor of points	120
II.C	Field-valued and rational points	122
III	Projective schemes	123
III.A	Graded rings and the homogeneous spectrum	123
III.B	The Zariski topology and the structure sheaf in the graded setting	125
III.C	Projective schemes and closed subschemes	133
III.D	Graded maps and rational maps	135
IV	Invertible sheaves and projective space	138
IV.A	Invertible sheaves	138
IV.B	Twisting sheaves on projective space	138
IV.C	Morphisms to projective space	139
4	First properties of schemes	140
I	Connected and reduced schemes	140
I.A	Connected schemes	140
I.B	Reduced schemes	141
I.B.1	Reduced structure on a scheme	141
I.B.2	The Zariski tangent space	144
II	Irreducible schemes	147
II.A	Irreducible components	147
II.B	Irreducible affine schemes and minimal primes	148
II.C	Generic points and specialisations	149
III	Integral schemes	151

III.A	Integral schemes and integral domains	151
III.B	The function field	152
IV	Noetherianity and dimension	154
IV.A	Noetherianity	154
IV.B	Dimension	155
IV.B.1	Dimension of schemes and rings	155
IV.B.2	Dimension of algebraic schemes over a field	157
V	Regular schemes	160
V.A	Regular local rings and the cotangent space	160
V.B	The Jacobian criterion for regularity	161
5	Global properties of morphisms	163
I	Fibred products	163
I.A	The universal property of the fibred product	163
I.B	Constructing the fibred product	164
I.C	Properties of the fibred product	166
I.D	Base change	167
II	Smoothness over a field	168
III	Fibres	170
III.A	Fibres and residue fields	170
III.B	Dimension of fibres	172
IV	Separated morphisms	175
IV.A	Separated morphisms and the diagonal	175
IV.B	Affine coverings of separated schemes	177
IV.C	Projective morphisms	179
V	Proper morphisms	180
V.A	Closed and universally closed morphisms	180
V.B	The valuative criterion and projective morphisms	182
V.C	Finite morphisms	184
6	A glimpse towards moduli spaces	188
I	The Hilbert scheme functor	188
I.A	Families and the functor of points	188
I.B	The Hilbert scheme functor of n points of the plane	189
I.C	Commuting matrices and a cyclic vector	190
I.D	Dual numbers and the Barth–Nakajima tangent calculation	191
I.E	Exercises	193
II	What about more general moduli problems?	194
II.A	Hilbert schemes of points on curves, surfaces and threefolds	194
II.B	Beyond Hilbert schemes of points	195

Chapter 1

Basic material

This chapter is partly taken from [Hul03]. For complementary accounts of the classical affine and projective material, see [Hol12] and [Har77, Chapter I]. The scheme-theoretic viewpoint developed later is particularly well motivated in [EH00], while [Liu02] gives a more systematic treatment.

I Introduction: algebraic sets

Further reading. For additional examples and a classical treatment of affine and projective algebraic sets, see [Hul03], [Hol12], and [Har77, Chapter I, §§1–3].

We start by defining algebraic subsets of the affine space over a field. This gives a traditional and basic picture of how to relate algebraic objects such as rings and ideals (actually, polynomial rings over a field) to geometric objects, essentially by looking at the vanishing of polynomial equations.

I.A Ideals and affine algebraic sets

Let $\mathbb{k}$ be a field and consider the polynomial ring $A = \mathbb{k}[x_1, \dots, x_n]$.

I.A.1 From ideals to algebraic sets

Given a subset P of A , we set

$$\mathbf{V}(P) = \{a \in \mathbb{k}^n \mid f(a) = 0, \forall f \in P\}.$$

Remark I.A.1. Subsets of the form $\mathbf{V}(P)$ are naturally related to ideals. A first indication of this is that P defines the same subset as (P) , the ideal generated by P , namely:

$$\mathbf{V}(P) = \mathbf{V}((P)).$$

Indeed, let $a \in \mathbf{V}((P))$. Then, since any element f of P lies in (P) , we have $f(a) = 0$, so $a \in \mathbf{V}(P)$. Conversely, consider $a \in \mathbf{V}(P)$. For any $f \in (P)$, there are $p_1, \dots, p_s \in P$ and $g_1, \dots, g_s \in A$ such that $f = g_1 p_1 + \dots + g_s p_s$. We have $p_1(a) = \dots = p_s(a) = 0$, since $p_1, \dots, p_s \in P$ and $a \in \mathbf{V}(P)$. Therefore, $f(a) = g_1(a)p_1(a) + \dots + g_s(a)p_s(a) = 0$. So $a \in \mathbf{V}((P))$.

According to the previous remark, we will usually replace P by the ideal it generates and write $\mathbf{V}(P)$ without further comment.

Definition I.A.2. A subset of the form $\mathbf{V}(P)$ is called an affine algebraic set. Literally, P can be any subset of $A = \mathbb{k}[x_1, \dots, x_n]$, but as we have just seen, the relevant subsets of A are the ideals of A .

Example I.A.3. In $\mathbb{k}^2$, the algebraic set

$$\mathbf{V}(y - x^2)$$

is the parabola with equation $y = x^2$. The algebraic set

$$\mathbf{V}(xy)$$

is the union of the two coordinate axes, since $xy = 0$ if and only if $x = 0$ or $y = 0$.

Example I.A.4. In $\mathbb{k}^2$, the algebraic set

$$\mathbf{V}(x^2 + y^2 - 1)$$

depends on the field $\mathbb{k}$. For instance, over $\mathbb{R}$ it is the usual circle, while over an algebraically closed field it should be regarded simply as the zero locus of the polynomial $x^2 + y^2 - 1$.

Lemma I.A.5. *Affine algebraic sets satisfy the following properties.*

- i) We have $\mathbf{V}(0) = \mathbb{k}^n$ and $\mathbf{V}(A) = \emptyset$.
- ii) For all ideals $\mathfrak{a} \subset \mathfrak{b}$, we have $\mathbf{V}(\mathfrak{a}) \supset \mathbf{V}(\mathfrak{b})$.
- iii) For all ideals $\mathfrak{a}, \mathfrak{b}$ of A , we have $\mathbf{V}(\mathfrak{a} \cap \mathfrak{b}) = \mathbf{V}(\mathfrak{a}) \cup \mathbf{V}(\mathfrak{b})$.
- iv) Given ideals $(\mathfrak{a}_i \mid i \in I)$, we have $\mathbf{V}(\sum_{i \in I} \mathfrak{a}_i) = \cap_{i \in I} \mathbf{V}(\mathfrak{a}_i)$.

Proof. For i), $\mathbf{V}(0) = \mathbb{k}^n$ is obvious, while $\mathbf{V}(A) = \emptyset$ is clear by noting that $\mathbf{V}(1) = \emptyset$. Item ii) is clear since any $b \in \mathbf{V}(\mathfrak{b})$ satisfies $f(b) = 0$, for all $f \in \mathfrak{a}$, as $\mathfrak{a} \subset \mathfrak{b}$. For iv), let us take $a \in \mathbf{V}(\sum_{i \in I} \mathfrak{a}_i)$. Then $f(a) = 0$ for all $f \in \mathfrak{a}_i$, for all $i \in I$, because $\mathfrak{a}_i \subset \sum_{i \in I} \mathfrak{a}_i$. Hence $a \in \cap_{i \in I} \mathbf{V}(\mathfrak{a}_i)$. Conversely, given $a \in \cap_{i \in I} \mathbf{V}(\mathfrak{a}_i)$, we have $f_i(a) = 0$, for all $i \in I$ and all $f_i \in \mathfrak{a}_i$. Since any $f \in \sum_{i \in I} \mathfrak{a}_i$ is a finite combination of some $(f_i \mid i \in I)$, we have $f(a) = 0$.

In conclusion, the only non-trivial item is iii). The inclusion $\mathbf{V}(\mathfrak{a}) \cup \mathbf{V}(\mathfrak{b}) \subset \mathbf{V}(\mathfrak{a} \cap \mathfrak{b})$ follows from ii), because $\mathfrak{a}$ and $\mathfrak{b}$ both contain $\mathfrak{a} \cap \mathfrak{b}$. Conversely, consider $a \in \mathbb{k}^n$ and assume that $a \notin \mathbf{V}(\mathfrak{a}) \cup \mathbf{V}(\mathfrak{b})$, so that $a \notin \mathbf{V}(\mathfrak{a})$ and $a \notin \mathbf{V}(\mathfrak{b})$. Then, there are $f \in \mathfrak{a}$ and $g \in \mathfrak{b}$ such that $f(a) \neq 0 \neq g(a)$. Hence $fg \in \mathfrak{a} \cap \mathfrak{b}$ gives $fg(a) \neq 0$, so $a \notin \mathbf{V}(\mathfrak{a} \cap \mathfrak{b})$. $\square$

Exercise I.A.6. Draw the following affine algebraic sets in $\mathbb{R}^2$:

$$\mathbf{V}(xy), \quad \mathbf{V}(y - x^2), \quad \mathbf{V}((x - 1)(x + 1), y).$$

Solution. The set $\mathbf{V}(xy)$ is the union of the two coordinate axes. The set $\mathbf{V}(y - x^2)$ is the parabola $y = x^2$. Finally,

$$\mathbf{V}((x - 1)(x + 1), y)$$

is the set of points satisfying $y = 0$ and $(x - 1)(x + 1) = 0$, hence it is the two-point set $\{(-1, 0), (1, 0)\}$.

Exercise I.A.7. Show that every finite subset of $\mathbb{k}^n$ is an affine algebraic set.

Solution. It is enough to show that a single point is algebraic, since finite unions of algebraic sets are algebraic by Lemma iii). If $a = (a_1, \dots, a_n) \in \mathbb{k}^n$, then

$$\{a\} = \mathbf{V}(x_1 - a_1, \dots, x_n - a_n).$$

Thus any finite subset of $\mathbb{k}^n$ is a finite union of algebraic sets, hence is algebraic.

Exercise I.A.8 (Finite sets of points and collision of points). Let

$$A = \mathbb{k}[x_1, x_2]$$

and consider the finite subsets of $\mathbb{k}^2$:

$$\begin{aligned} X_1 &= \{(0, 0)\}, \\ X_2 &= \{(0, 0), (1, 0)\}, \\ X_3 &= \{(0, 0), (1, 0), (0, 1)\}, \\ X_4 &= \{(0, 0), (1, 0), (0, 1), (1, 1)\} \end{aligned}$$

1. Compute the vanishing ideals $\mathbf{I}(X_i)$.
2. Find a basis of the $\mathbb{k}$ -vector space $A/\mathbf{I}(X_i)$ for $1 \leq i \leq 4$.
3. Show that the ideals

$$\mathfrak{a} = (x_1, x_2) \quad \text{and} \quad \mathfrak{b} = (x_1, x_2)^2$$

have the same zero set, but that the quotient algebras $A/\mathfrak{a}$ and $A/\mathfrak{b}$ are not isomorphic.

4. For $a \in \mathbb{k}$, consider the ideals

$$\begin{aligned} \mathfrak{a}_{2,a} &= (x_2, x_1(x_1 - a)), \\ \mathfrak{a}_{3,a} &= (x_1x_2, x_1(x_1 - a), x_2(x_2 - a)), \\ \mathfrak{a}_{4,a} &= (x_1(x_1 - a), x_2(x_2 - a)). \end{aligned}$$

Determine their zero sets and find bases of the quotient algebras $A/\mathfrak{a}_{r,a}$.

5. Find generators for the relations among the displayed generators of $\mathfrak{a}_{2,a}$, $\mathfrak{a}_{3,a}$ and $\mathfrak{a}_{4,a}$.

Solution. For a point $p = (a_1, a_2) \in \mathbb{k}^2$, its vanishing ideal is

$$\mathbf{I}(p) = (x_1 - a_1, x_2 - a_2).$$

Moreover, the vanishing ideal of a finite union is the intersection of the vanishing ideals:

$$\mathbf{I}(Y \cup Z) = \mathbf{I}(Y) \cap \mathbf{I}(Z).$$

1. *Computation of the ideals.* For the first set, we immediately obtain

$$\mathbf{I}(X_1) = (x_1, x_2).$$

For X_2 , the polynomial x_2 vanishes at both points, as does $x_1(x_1 - 1)$. We claim that

$$\mathbf{I}(X_2) = (x_2, x_1(x_1 - 1)).$$

Indeed, modulo the ideal on the right, every polynomial has a unique representative of the form

$$\alpha + \beta x_1, \quad \alpha, \beta \in \mathbb{k}.$$

If such a polynomial vanishes at $(0, 0)$ and $(1, 0)$, then

$$\alpha = 0, \quad \alpha + \beta = 0,$$

so $\alpha = \beta = 0$. Hence the displayed ideal contains every polynomial vanishing on X_2 .

For X_3 , put

$$J_3 = (x_1x_2, x_1(x_1 - 1), x_2(x_2 - 1)).$$

All three generators vanish on X_3 , so

$$J_3 \subset \mathbf{I}(X_3).$$

Modulo J_3 , we have

$$x_1x_2 = 0, \quad x_1^2 = x_1, \quad x_2^2 = x_2.$$

Thus every polynomial is congruent modulo J_3 to a linear combination of

$$1, \quad x_1, \quad x_2.$$

Suppose

$$\alpha + \beta x_1 + \gamma x_2$$

vanishes at the three points of X_3 . Evaluating successively at $(0, 0)$, $(1, 0)$ and $(0, 1)$ gives

$$\alpha = 0, \quad \alpha + \beta = 0, \quad \alpha + \gamma = 0.$$

Hence $\alpha = \beta = \gamma = 0$. Therefore

$$\mathbf{I}(X_3) = (x_1x_2, x_1(x_1 - 1), x_2(x_2 - 1)).$$

Finally, put

$$J_4 = (x_1(x_1 - 1), x_2(x_2 - 1)).$$

Modulo J_4 , we have

$$x_1^2 = x_1, \quad x_2^2 = x_2.$$

Consequently, every polynomial is congruent to a linear combination of

$$1, \quad x_1, \quad x_2, \quad x_1x_2.$$

If

$$\alpha + \beta x_1 + \gamma x_2 + \delta x_1x_2$$

vanishes on X_4 , evaluation at the four points gives

$$\begin{aligned} \alpha &= 0, \\ \alpha + \beta &= 0, \\ \alpha + \gamma &= 0, \\ \alpha + \beta + \gamma + \delta &= 0. \end{aligned}$$

It follows that all four coefficients vanish. Hence

$$\mathbf{I}(X_4) = (x_1(x_1 - 1), x_2(x_2 - 1)).$$

We have therefore proved

$$\begin{aligned} \mathbf{I}(X_1) &= (x_1, x_2), \\ \mathbf{I}(X_2) &= (x_2, x_1(x_1 - 1)), \\ \mathbf{I}(X_3) &= (x_1x_2, x_1(x_1 - 1), x_2(x_2 - 1)), \\ \mathbf{I}(X_4) &= (x_1(x_1 - 1), x_2(x_2 - 1)). \end{aligned}$$

2. *Bases of the quotient algebras.* The computations above show that bases are given by

quotient algebra	basis over $\mathbb{k}$
$A/\mathbf{I}(X_1)$	$\{1\}$
$A/\mathbf{I}(X_2)$	$\{1, x_1\}$
$A/\mathbf{I}(X_3)$	$\{1, x_1, x_2\}$
$A/\mathbf{I}(X_4)$	$\{1, x_1, x_2, x_1x_2\}$.

In particular,

$$\dim_{\mathbb{k}}(A/\mathbf{I}(X_i)) = i.$$

One may also see this directly from the evaluation morphism

$$A \longrightarrow \mathbb{k}^{X_i}, \quad f \longmapsto (f(p))_{p \in X_i}.$$

Its kernel is $\mathbf{I}(X_i)$, and in the present examples it is surjective. Thus

$$A/\mathbf{I}(X_i) \simeq \mathbb{k}^i$$

as $\mathbb{k}$ -algebras.

3. *Two ideals with the same zero set.* Let

$$\mathfrak{a} = (x_1, x_2), \quad \mathfrak{b} = (x_1, x_2)^2 = (x_1^2, x_1x_2, x_2^2).$$

Since $\mathfrak{b} \subset \mathfrak{a}$, we have

$$\mathbf{V}(\mathfrak{a}) \subset \mathbf{V}(\mathfrak{b}).$$

Conversely, if $(a_1, a_2) \in \mathbf{V}(\mathfrak{b})$, then

$$a_1^2 = a_1a_2 = a_2^2 = 0.$$

Since $\mathbb{k}$ is a field, this implies $a_1 = a_2 = 0$. Hence

$$\mathbf{V}(\mathfrak{a}) = \mathbf{V}(\mathfrak{b}) = \{(0, 0)\}.$$

On the other hand,

$$A/\mathfrak{a} \simeq \mathbb{k},$$

whereas

$$A/\mathfrak{b} = \mathbb{k}[x_1, x_2]/(x_1^2, x_1x_2, x_2^2)$$

has basis

$$\{1, x_1, x_2\}.$$

Thus

$$\dim_{\mathbb{k}}(A/\mathfrak{a}) = 1, \quad \dim_{\mathbb{k}}(A/\mathfrak{b}) = 3,$$

so the two quotient algebras cannot be isomorphic.

Moreover, the classes of x_1 and x_2 in $A/\mathfrak{b}$ are nonzero nilpotent elements. For example,

$$\bar{x}_1 \neq 0, \quad \bar{x}_1^2 = 0.$$

4. *Specialisation of the ideals.* Assume first that $a \neq 0$.

For

$$\mathfrak{a}_{2,a} = (x_2, x_1(x_1 - a)),$$

a common zero must satisfy

$$x_2 = 0, \quad x_1(x_1 - a) = 0.$$

Therefore

$$\mathbf{V}(\mathfrak{a}_{2,a}) = \{(0, 0), (a, 0)\}.$$

For

$$\mathfrak{a}_{3,a} = (x_1x_2, x_1(x_1 - a), x_2(x_2 - a)),$$

we have

$$x_1 \in \{0, a\}, \quad x_2 \in \{0, a\}, \quad x_1x_2 = 0.$$

The last condition excludes (a, a) , and hence

$$\mathbf{V}(\mathfrak{a}_{3,a}) = \{(0, 0), (a, 0), (0, a)\}.$$

Finally,

$$\mathfrak{a}_{4,a} = (x_1(x_1 - a), x_2(x_2 - a))$$

gives independently

$$x_1 \in \{0, a\}, \quad x_2 \in \{0, a\}.$$

Thus

$$\mathbf{V}(\mathfrak{a}_{4,a}) = \{(0, 0), (a, 0), (0, a), (a, a)\}.$$

For $a = 0$, the ideals become

$$\begin{aligned} \mathfrak{a}_{2,0} &= (x_2, x_1^2), \\ \mathfrak{a}_{3,0} &= (x_1^2, x_1x_2, x_2^2) = (x_1, x_2)^2, \\ \mathfrak{a}_{4,0} &= (x_1^2, x_2^2). \end{aligned}$$

They all have the same zero set:

$$\mathbf{V}(\mathfrak{a}_{2,0}) = \mathbf{V}(\mathfrak{a}_{3,0}) = \mathbf{V}(\mathfrak{a}_{4,0}) = \{(0, 0)\}.$$

Thus, after the specialisation $a = 0$, respectively two, three and four points have the same underlying zero set.

Let us now compute the quotient algebras. Modulo $\mathfrak{a}_{2,a}$, we have

$$x_2 = 0, \quad x_1^2 = ax_1.$$

Every polynomial therefore reduces to a linear combination of

$$1, \quad x_1.$$

These classes are linearly independent. For $a \neq 0$, this follows by evaluating at $(0, 0)$ and $(a, 0)$; for $a = 0$, it is immediate from

$$A/\mathfrak{a}_{2,0} \simeq \mathbb{k}[x_1]/(x_1^2).$$

Hence

$$\{1, x_1\}$$

is a basis of $A/\mathfrak{a}_{2,a}$ for every $a \in \mathbb{k}$.

Modulo $\mathfrak{a}_{3,a}$, we have

$$x_1x_2 = 0, \quad x_1^2 = ax_1, \quad x_2^2 = ax_2.$$

Thus every polynomial reduces to a linear combination of

$$1, \quad x_1, \quad x_2.$$

For $a \neq 0$, their independence follows by evaluation at

$$(0, 0), \quad (a, 0), \quad (0, a).$$

For $a = 0$, these are the standard monomials modulo $(x_1, x_2)^2$. Consequently,

$$\{1, x_1, x_2\}$$

is a basis of $A/\mathfrak{a}_{3,a}$ for every $a \in \mathbb{k}$.

Finally, modulo $\mathfrak{a}_{4,a}$, we have

$$x_1^2 = ax_1, \quad x_2^2 = ax_2.$$

Every monomial reduces to a linear combination of

$$1, \quad x_1, \quad x_2, \quad x_1x_2.$$

For $a \neq 0$, independence follows by evaluating at the four points of $\mathbf{V}(\mathfrak{a}_{4,a})$. For $a = 0$, these are the standard monomials modulo (x_1^2, x_2^2) . Hence

$$\{1, x_1, x_2, x_1x_2\}$$

is a basis of $A/\mathfrak{a}_{4,a}$.

In conclusion,

$$\dim_{\mathbb{k}}(A/\mathfrak{a}_{r,a}) = r$$

for $r = 2, 3, 4$, independently of a .

5. *Relations among the generators.* For $\mathfrak{a}_{2,a}$, write

$$g_1 = x_2, \quad g_2 = x_1(x_1 - a).$$

We have the relation

$$x_1(x_1 - a)g_1 - x_2g_2 = 0.$$

We claim that this relation generates all the relations between g_1 and g_2 .

Indeed, suppose

$$ug_1 + vg_2 = 0.$$

Then

$$ux_2 = -vx_1(x_1 - a).$$

Since x_2 and $x_1(x_1 - a)$ are coprime, $x_1(x_1 - a)$ divides u . Thus there exists $h \in A$ such that

$$u = hx_1(x_1 - a).$$

Substituting, we obtain

$$v = -hx_2.$$

Hence

$$(u, v) = h(x_1(x_1 - a), -x_2).$$

For $\mathfrak{a}_{3,a}$, put

$$q_1 = x_1x_2, \quad q_2 = x_1(x_1 - a), \quad q_3 = x_2(x_2 - a).$$

We have two relations:

$$(a - x_1)q_1 + x_2q_2 = 0$$

and

$$(a - x_2)q_1 + x_1q_3 = 0.$$

We show that they generate all relations.

Suppose

$$uq_1 + vq_2 + wq_3 = 0.$$

Factoring x_1 from the first two terms gives

$$x_1(ux_2 + v(x_1 - a)) = -wx_2(x_2 - a).$$

Since x_1 is coprime to $x_2(x_2 - a)$, it follows that x_1 divides w . Write

$$w = x_1\beta$$

for some $\beta \in A$. Dividing the equality by x_1 gives

$$ux_2 + v(x_1 - a) + \beta x_2(x_2 - a) = 0,$$

and therefore

$$v(x_1 - a) = -x_2(u + \beta(x_2 - a)).$$

Since $x_1 - a$ and x_2 are coprime, x_2 divides v . Write

$$v = x_2\alpha$$

for some $\alpha \in A$. Substituting, we obtain

$$u + \alpha(x_1 - a) + \beta(x_2 - a) = 0,$$

or equivalently

$$u = \alpha(a - x_1) + \beta(a - x_2).$$

Thus

$$(u, v, w) = \alpha(a - x_1, x_2, 0) + \beta(a - x_2, 0, x_1).$$

Hence all relations are generated by

$$(a - x_1, x_2, 0) \quad \text{and} \quad (a - x_2, 0, x_1).$$

Finally, for $\mathfrak{a}_{4,a}$, put

$$h_1 = x_1(x_1 - a), \quad h_2 = x_2(x_2 - a).$$

The relation

$$x_2(x_2 - a)h_1 - x_1(x_1 - a)h_2 = 0$$

generates all relations. Indeed, h_1 and h_2 are coprime, so the same argument as for $\mathfrak{a}_{2,a}$ applies.

We have therefore obtained the following generators of the modules of relations:

ideal	generating relations
$\mathfrak{a}_{2,a}$	$(x_1(x_1 - a), -x_2)$
$\mathfrak{a}_{3,a}$	$(a - x_1, x_2, 0), (a - x_2, 0, x_1)$
$\mathfrak{a}_{4,a}$	$(x_2(x_2 - a), -x_1(x_1 - a)).$

In particular, the number of generators and the number of generating relations do not change when a is specialised to zero.

Exercise I.A.9. Assume that $\mathbb{k}$ is infinite. Show that an infinite strict subset $X \subsetneq \mathbb{k}$ is not an affine algebraic set.

Solution. The affine algebraic subsets of $\mathbb{k}$ are the subsets of the form $\mathbf{V}(f_1, \dots, f_s)$. If all the f_i are zero, then $\mathbf{V}(f_1, \dots, f_s) = \mathbb{k}$. Otherwise, one of the f_i is a non-zero polynomial, and $\mathbf{V}(f_1, \dots, f_s)$ is contained in the finite set of roots of this polynomial. Hence an algebraic subset of $\mathbb{k}$ is either finite or equal to $\mathbb{k}$. Therefore an infinite strict subset of $\mathbb{k}$ is not algebraic.

I.A.2 From subsets to ideals

Given a subset $X \subset \mathbb{k}^n$, one defines the ideal

$$\mathbf{I}(X) = \{f \in A \mid f(a) = 0, \forall a \in X\}.$$

We call $\mathbf{I}(X)$ the ideal *associated with* X , or the ideal of *functions vanishing at* X .

Lemma I.A.10. *Ideals associated with subsets have the following properties.*

- i) For all subsets $X \subset Y \subset \mathbb{k}^n$, we have $\mathbf{I}(X) \supset \mathbf{I}(Y)$.
- ii) Given $X \subset \mathbb{k}^n$, we have $X \subset \mathbf{V}(\mathbf{I}(X))$. Equality holds if and only if X is an algebraic subset of $\mathbb{k}^n$.
- iii) If $\mathfrak{a}$ is an ideal of A , then $\mathfrak{a} \subset \mathbf{I}(\mathbf{V}(\mathfrak{a}))$.

Proof. Item i) is obvious. For iii), let $f \in \mathfrak{a}$. Then, for any $a \in \mathbf{V}(\mathfrak{a})$, we have $f(a) = 0$. So $f \in \mathbf{I}(\mathbf{V}(\mathfrak{a}))$.

For ii), we let $a \in X$ and take $f \in \mathbf{I}(X)$. Then $f(a) = 0$ by definition. Since this happens for all $f \in \mathbf{I}(X)$, we have $a \in \mathbf{V}(\mathbf{I}(X))$. This shows that $X \subset \mathbf{V}(\mathbf{I}(X))$. If equality is attained, then by definition X is an algebraic set, actually an algebraic subset of $\mathbb{k}^n$. Conversely, if X is an algebraic subset of $\mathbb{k}^n$, then there is an ideal $\mathfrak{b}$ such that $X = \mathbf{V}(\mathfrak{b})$. Hence $\mathfrak{b} \subset \mathbf{I}(\mathbf{V}(\mathfrak{b})) = \mathbf{I}(X)$. So $\mathbf{V}(\mathbf{I}(X)) \subset \mathbf{V}(\mathfrak{b}) = X$. $\square$

Example I.A.11. Let $X = \{0, 1\} \subset \mathbb{k}$. Then

$$\mathbf{I}(X) = (x(x - 1)).$$

Indeed, a polynomial vanishes at both 0 and 1 if and only if it is divisible by x and by $x - 1$, hence by $x(x - 1)$.

Example I.A.12. Let $X = \{(0, 0)\} \subset \mathbb{k}^2$. Then

$$\mathbf{I}(X) = (x, y).$$

Indeed, $f \in \mathbb{k}[x, y]$ vanishes at the origin if and only if its constant term is zero, which is equivalent to saying that $f \in (x, y)$.

Exercise I.A.13. Let $X \subset \mathbb{k}^n$ and $Y \subset \mathbb{k}^n$. Show that

$$\mathbf{I}(X \cup Y) = \mathbf{I}(X) \cap \mathbf{I}(Y).$$

Solution. A polynomial f belongs to $\mathbf{I}(X \cup Y)$ if and only if it vanishes at every point of $X \cup Y$. This is equivalent to saying that f vanishes on X and on Y , namely that $f \in \mathbf{I}(X)$ and $f \in \mathbf{I}(Y)$. Therefore

$$\mathbf{I}(X \cup Y) = \mathbf{I}(X) \cap \mathbf{I}(Y).$$

I.B Morphisms of affine algebraic sets

Here we define morphisms of algebraic subsets as maps from one set to another, such that each coordinate of the target depends in a polynomial way on the coordinates of the source.

I.B.1 Morphisms of algebraic sets and polynomial mappings

Let us consider two affine algebraic sets

$$X \subset \mathbb{k}^n, \quad Y \subset \mathbb{k}^m.$$

Definition I.B.1. A map $\varphi : X \rightarrow Y$ is a *morphism of affine algebraic sets* if there exist polynomials

$$f_1, \dots, f_m \in \mathbb{k}[x_1, \dots, x_n]$$

such that

$$\varphi(a) = (f_1(a), \dots, f_m(a)) \quad \text{for all } a \in X.$$

In other words, a morphism is a map whose coordinate functions are given by polynomials on the ambient affine space.

Example I.B.2. Every polynomial map

$$\mathbb{k}^n \rightarrow \mathbb{k}^m, \quad a \mapsto (f_1(a), \dots, f_m(a)),$$

is a morphism. Its restriction to an affine algebraic subset is a morphism provided that its image is contained in the target algebraic set.

Example I.B.3. Let

$$X = \mathbf{V}(y - x^2) \subset \mathbb{k}^2.$$

The map

$$\alpha : \mathbb{k} \rightarrow X, \quad t \mapsto (t, t^2),$$

is a morphism. The first projection

$$\pi : X \rightarrow \mathbb{k}, \quad (x, y) \mapsto x,$$

is also a morphism, and

$$\pi \circ \alpha = \text{id}_{\mathbb{k}}, \quad \alpha \circ \pi = \text{id}_X.$$

Thus the parabola is isomorphic, as an affine algebraic set, to the affine line.

I.B.2 Coordinate rings

Let $X \subset \mathbb{A}^n$ be an affine algebraic subset. The morphisms of algebraic sets $X \rightarrow \mathbb{A}^1$ are given by polynomial maps $\mathbb{A}^n \rightarrow \mathbb{A}^1$ of the form $(a_1, \dots, a_n) \rightarrow f(a_1, \dots, a_n)$. Replacing f by $f + g$, with $g \in \mathbf{I}(X)$, will not change the value of f at any point of X , so it is natural to associate with a morphism of algebraic sets $X \rightarrow \mathbb{A}^1$ an element of $\mathbb{A}^1[X] = \mathbb{A}^1[x_1, \dots, x_n]/\mathbf{I}(X)$.

Definition I.B.4. Let $X \subset \mathbb{A}^n$ be an affine algebraic set. Its *coordinate ring* is

$$\mathbb{A}^1[X] := \mathbb{A}^1[x_1, \dots, x_n]/\mathbf{I}(X).$$

An element of $\mathbb{A}^1[X]$ is called a *polynomial function* on X .

Proposition I.B.5. Let $X \subset \mathbb{A}^n$ and $Y \subset \mathbb{A}^m$ be affine algebraic sets. Every morphism $\varphi : X \rightarrow Y$ induces a homomorphism of $\mathbb{A}^1$ -algebras

$$\varphi^* : \mathbb{A}^1[Y] \longrightarrow \mathbb{A}^1[X], \quad h \longmapsto h \circ \varphi.$$

Moreover, the assignment $\varphi \mapsto \varphi^*$ gives a bijection

$$\mathrm{Hom}_{(\mathrm{AffAlg})_{\mathbb{A}^1}}(X, Y) \simeq \mathrm{Hom}_{(\mathbb{A}^1\text{-Alg})}(\mathbb{A}^1[Y], \mathbb{A}^1[X]).$$

Proof. Write

$$\varphi(a) = (f_1(a), \dots, f_m(a))$$

with $f_i \in \mathbb{A}^1[x_1, \dots, x_n]$. If $h \in \mathbb{A}^1[y_1, \dots, y_m]$, then

$$h \circ \varphi = h(f_1, \dots, f_m)$$

is a polynomial function on X . If $h \in \mathbf{I}(Y)$, then h vanishes on Y , hence $h \circ \varphi$ vanishes on X . Therefore substitution induces a well-defined homomorphism

$$\varphi^* : \mathbb{A}^1[Y] \rightarrow \mathbb{A}^1[X].$$

Conversely, let

$$\psi : \mathbb{A}^1[Y] \longrightarrow \mathbb{A}^1[X]$$

be a homomorphism of $\mathbb{A}^1$ -algebras. Denote by $\bar{y}_i \in \mathbb{A}^1[Y]$ the class of the coordinate y_i , and choose a polynomial $f_i \in \mathbb{A}^1[x_1, \dots, x_n]$ representing $\psi(\bar{y}_i)$. Define

$$\varphi : X \longrightarrow \mathbb{A}^m, \quad a \longmapsto (f_1(a), \dots, f_m(a)).$$

For every $h \in \mathbf{I}(Y)$, the class of h in $\mathbb{A}^1[Y]$ is zero. Hence

$$h(f_1, \dots, f_m) \in \mathbf{I}(X),$$

so

$$h(\varphi(a)) = 0 \quad \text{for every } a \in X.$$

Thus $\varphi(a) \in Y$, and $\varphi : X \rightarrow Y$ is a morphism. By construction, $\varphi^* = \psi$. The two constructions are inverse to one another. $\square$

Remark I.B.6. The correspondence is contravariant: if

$$X \xrightarrow{\varphi} Y \xrightarrow{\psi} Z$$

are morphisms, then

$$(\psi \circ \varphi)^* = \varphi^* \circ \psi^*.$$

In particular, a morphism $\varphi : X \rightarrow Y$ is an isomorphism if and only if $\varphi^* : \mathbb{A}^1[Y] \rightarrow \mathbb{A}^1[X]$ is an isomorphism of $\mathbb{A}^1$ -algebras.

Exercise I.B.7. Let

$$X = \mathbf{V}(y - x^2) \subset \mathbb{A}^2.$$

Compute the homomorphisms on coordinate rings induced by

$$\alpha : \mathbb{k} \rightarrow X, \quad t \mapsto (t, t^2),$$

and by the projection $\pi : X \rightarrow \mathbb{k}, (x, y) \mapsto x$. Verify algebraically that they are inverse isomorphisms.

Solution. We have

$$\mathbb{k}[X] = \mathbb{k}[x, y]/(y - x^2) \simeq \mathbb{k}[x].$$

The pullback by α is

$$\alpha^* : \mathbb{k}[x, y]/(y - x^2) \longrightarrow \mathbb{k}[t], \quad \bar{x} \longmapsto t, \quad \bar{y} \longmapsto t^2.$$

The pullback by π is

$$\pi^* : \mathbb{k}[t] \longrightarrow \mathbb{k}[x, y]/(y - x^2), \quad t \longmapsto \bar{x}.$$

Since $\bar{y} = \bar{x}^2$ in $\mathbb{k}[X]$, these homomorphisms are inverse to one another.

Exercise I.B.8. Let $X \subset Y \subset \mathbb{A}^n$ be affine algebraic sets. Show that the inclusion $X \hookrightarrow Y$ is a morphism and that the induced homomorphism

$$\mathbb{k}[Y] \longrightarrow \mathbb{k}[X]$$

is surjective.

Solution. The inclusion is the restriction of the identity map of $\mathbb{A}^n$, hence is a morphism. Since $X \subset Y$, every polynomial vanishing on Y also vanishes on X , so

$$\mathbf{I}(Y) \subset \mathbf{I}(X).$$

Therefore the quotient map induces a surjection

$$\mathbb{k}[x_1, \dots, x_n]/\mathbf{I}(Y) \longrightarrow \mathbb{k}[x_1, \dots, x_n]/\mathbf{I}(X).$$

Exercise I.B.9. Consider

$$\nu : \mathbb{k} \longrightarrow \mathbf{V}(y^2 - x^3) \subset \mathbb{A}^2, \quad t \longmapsto (t^2, t^3).$$

Show that ν is a bijective morphism, but that it is not an isomorphism of affine algebraic sets.

Solution. The map is a morphism and its image lies in the cusp because

$$(t^3)^2 = (t^2)^3.$$

It is injective: if $t^2 = u^2$ and $t^3 = u^3$, then either both are zero, or $t = t^3/t^2 = u^3/u^2 = u$. It is also surjective. Indeed, if (x, y) satisfies $y^2 = x^3$ and $x \neq 0$, then $t = y/x$ satisfies $t^2 = x$ and $t^3 = y$; if $x = 0$, then also $y = 0$.

The induced homomorphism is

$$\nu^* : \mathbb{k}[x, y]/(y^2 - x^3) \longrightarrow \mathbb{k}[t], \quad \bar{x} \longmapsto t^2, \quad \bar{y} \longmapsto t^3.$$

Its image is $\mathbb{k}[t^2, t^3]$, which does not contain t . Hence ν^* is not surjective, so ν is not an isomorphism.

I.C Radicals and the Nullstellensatz

Further reading. For the classical affine theory, including the Nullstellensatz, see [Har77, Chapter I, §1]; for a more elementary presentation and further examples, see [Hul03] and [Hol12].

I.C.1 Radicals

Given a ring A , it is important to study the set of nilpotent elements, namely elements a such that $a^m = 0$ for some integer $m \geq 1$. If there is no non-zero nilpotent element, then A is said to be *reduced*.

Definition I.C.1. The *radical* of a ring A , or nilradical, is the set of nilpotent elements of A . It is denoted by $\text{Rad}(A)$. More generally, given an ideal $\mathfrak{a}$ of A , we define the radical of $\mathfrak{a}$ as follows:

$$\sqrt{\mathfrak{a}} = \{a \in A \mid a^n \in \mathfrak{a} \text{ for some integer } n \geq 1\}.$$

An ideal $\mathfrak{a}$ is *radical* if $\mathfrak{a} = \sqrt{\mathfrak{a}}$.

Remark I.C.2. We have the following properties.

- i) For any ideal $\mathfrak{a}$ of A , the radical $\sqrt{\mathfrak{a}}$ is an ideal of A .
- ii) Any ideal $\mathfrak{a}$ of A is contained in its radical, $\mathfrak{a} \subset \sqrt{\mathfrak{a}}$.
- iii) Given $X \subset \mathbb{k}^n$, the ideal $\mathbf{I}(X)$ is radical.

The first item follows by expanding sufficiently high powers of a sum. The second item is obvious. For the third item, given $n \in \mathbb{N}$ and $f \in A$ such that $f^n \in \mathbf{I}(X)$, we must have $f^n(a) = 0$ for all $a \in X$, so $f(a) = 0$ for all $a \in X$, hence $f \in \mathbf{I}(X)$. So $\sqrt{\mathbf{I}(X)} \subset \mathbf{I}(X)$.

Remark I.C.3. We have $\sqrt{(0)} = \text{Rad}(A)$. Given an ideal $\mathfrak{a}$ of A , consider the ring $B = A/\mathfrak{a}$ and the canonical projection $\varphi : A \rightarrow B$. Then $\varphi^{-1}(\text{Rad}(B)) = \sqrt{\mathfrak{a}}$.

Remark I.C.4. We note that, if $\mathfrak{a}$ is a prime ideal, then $\sqrt{\mathfrak{a}} = \mathfrak{a}$, i.e., $\mathfrak{a}$ is radical. Indeed, given $f \in A$, if $f^n \in \mathfrak{a}$ for some $n \geq 1$, then since $\mathfrak{a}$ is prime we have $f \in \mathfrak{a}$ or, when $n > 1$, $f^{n-1} \in \mathfrak{a}$, and continuing this way we finally get $f \in \mathfrak{a}$ anyway, so $\sqrt{\mathfrak{a}} \subset \mathfrak{a}$. Since the inclusion $\mathfrak{a} \subset \sqrt{\mathfrak{a}}$ holds for any ideal, we get $\mathfrak{a} = \sqrt{\mathfrak{a}}$ when $\mathfrak{a}$ is prime.

Example I.C.5. In $\mathbb{k}[x]$, we have

$$\sqrt{(x^m)} = (x)$$

for every $m \geq 1$. More generally, if

$$f = c \prod_{i=1}^r (x - a_i)^{m_i}$$

over an algebraically closed field, then

$$\sqrt{(f)} = \left(\prod_{i=1}^r (x - a_i) \right).$$

Example I.C.6. The so-called *ring of dual numbers* is

$$\mathbb{k}[x]/(x^2).$$

This ring is not reduced: the class of x is non-zero, but its square is zero. On the other hand, the ring $\mathbb{k}[x]/(x)$ is isomorphic to $\mathbb{k}$, hence is reduced.

Exercise I.C.7. Compute the radical of the ideal

$$\mathfrak{a} = (x^2, y^3) \subset \mathbb{k}[x, y].$$

Solution. We have $x^2 \in \mathfrak{a}$ and $y^3 \in \mathfrak{a}$, hence $x, y \in \sqrt{\mathfrak{a}}$. Therefore $(x, y) \subset \sqrt{\mathfrak{a}}$. Conversely, since $\mathfrak{a} \subset (x, y)$ and (x, y) is prime, hence radical, we have $\sqrt{\mathfrak{a}} \subset (x, y)$. Thus

$$\sqrt{(x^2, y^3)} = (x, y).$$

Exercise I.C.8. Decide whether the following rings are reduced:

$$\mathbb{k}[x]/(x^2), \quad \mathbb{k}[x, y]/(xy), \quad \mathbb{k}[x, y]/(x^2, xy).$$

Solution. The ring $\mathbb{k}[x]/(x^2)$ is not reduced because the class of x is nilpotent. The ring $\mathbb{k}[x, y]/(xy)$ is reduced if $\mathbb{k}[x, y]$ is reduced and (xy) is a radical ideal. Indeed,

$$(xy) = (x) \cap (y),$$

so (xy) is radical. Hence $\mathbb{k}[x, y]/(xy)$ is reduced. Finally, $\mathbb{k}[x, y]/(x^2, xy)$ is not reduced because the class of x is non-zero but satisfies $x^2 = 0$.

Exercise I.C.9. Compute $\sqrt{(x^2y^3)}$ in $\mathbb{k}[x, y]$.

Solution. Since $x^2y^3 \in (x^2y^3)$, we have $xy \in \sqrt{(x^2y^3)}$. Thus $(xy) \subset \sqrt{(x^2y^3)}$. Conversely, if $f \in \sqrt{(x^2y^3)}$, then $f^N \in (x^2y^3)$ for some N . Since (xy) is radical and contains (x^2y^3) , we get $f \in (xy)$. Therefore

$$\sqrt{(x^2y^3)} = (xy).$$

I.C.2 Integral ring extensions

Given an inclusion of rings $A \subset B$, an element $b \in B$ is said to be *integral over* A if there is a monic polynomial $f \in A[x]$ such that $f(b) = 0$. If all elements of B are integral over A then B is said to be *integral over* A .

Lemma I.C.10. Let $A \subset B$ be an inclusion of rings.

1. Assume that A and B are domains, with B integral over A . Then A is a field if and only if B is a field.
2. An element $b \in B$ is integral over A if and only if $A[b]$ is finitely generated as an A -module.

Therefore, the A -integral elements of B are a subring of B .

Proof. We first prove the second assertion. If b is integral over A , then there is a monic relation

$$b^d + a_{d-1}b^{d-1} + \cdots + a_1b + a_0 = 0, \quad a_i \in A.$$

Hence every power b^m with $m \geq d$ can be rewritten as an A -linear combination of $1, b, \dots, b^{d-1}$. Therefore $A[b]$ is generated as an A -module by $1, b, \dots, b^{d-1}$.

Conversely, assume that $A[b]$ is finitely generated as an A -module, say by $e_1, \dots, e_r$. For $1 \leq i \leq r$ there are coefficients $a_{i,j} \in A$, with $1 \leq j \leq r$, such that

$$be_i = \sum_{1 \leq j \leq r} a_{i,j}e_j, \quad \text{for } 1 \leq i \leq r.$$

Thus the matrix

$$M = (b\delta_{i,j} - a_{i,j})_{1 \leq i,j \leq r}$$

annihilates the column vector $(e_1, \dots, e_r)^t$. Multiplying by the adjugate matrix, we see that $\det(M)$ annihilates every e_i , with $1 \leq i \leq r$, and hence all of $A[b]$. In particular, $\det(M) \cdot 1 = 0$. Since $\det(M)$ is a monic polynomial in b with coefficients in A , the element b is integral over A .

Now assume that B is integral over A . If A is a field, then every element $b \in B$ is algebraic over A . Since B is an integral domain, the inverse of any non-zero $b \in B$ belongs to the field $A(b)$ and is a polynomial in b with coefficients in A . Hence $b^{-1} \in B$, so B is a field.

Conversely, assume that B is a field and let $a \in A$ be non-zero. Then $a^{-1} \in B$. Since B is integral over A , there is a monic relation

$$(a^{-1})^d + c_{d-1}(a^{-1})^{d-1} + \dots + c_1 a^{-1} + c_0 = 0, \quad c_i \in A.$$

Multiplying by a^{d-1} gives

$$a^{-1} = -(c_{d-1} + c_{d-2}a + \dots + c_0 a^{d-1}) \in A.$$

Thus every non-zero element of A is invertible, and A is a field.

Finally, if $b, c \in B$ are integral over A , then $A[b]$ is finite over A , and $A[b, c]$ is finite over $A[b]$, hence finite over A . Since $b + c$, bc and $-b$ belong to $A[b, c]$, the criterion just proved shows that they are integral over A . Therefore the integral elements form a subring of B . $\square$

Example I.C.11. The element $\sqrt{2} \in \mathbb{Q}(\sqrt{2})$ is integral over $\mathbb{Q}$, since it satisfies the monic equation $T^2 - 2 = 0$. More generally, every element algebraic over a field $\mathbb{k}$ is integral over $\mathbb{k}$.

On the other hand, $x^{-1} \in \mathbb{k}[x, x^{-1}]$ is not integral over $\mathbb{k}[x]$. Indeed, a monic equation for x^{-1} over $\mathbb{k}[x]$, multiplied by a suitable power of x , would imply that $x^{-1} \in \mathbb{k}[x]$, which is impossible.

Exercise I.C.12. Show that if $b \in B$ is integral over A , then the image of b in $B/\mathfrak{b}$ is integral over the image of A , for every ideal $\mathfrak{b} \subset B$.

Solution. If b satisfies a monic equation

$$b^d + a_{d-1}b^{d-1} + \dots + a_0 = 0, \quad a_i \in A,$$

then reducing the same equation modulo $\mathfrak{b}$ gives a monic equation for the image of b in $B/\mathfrak{b}$ with coefficients in the image of A .

Lemma I.C.13. Let κ be a field extension of a field $\mathbb{k}$ which is finitely generated as a $\mathbb{k}$ -algebra. Then κ is a finite field extension of $\mathbb{k}$.

Proof. We prove the statement by induction on the number of algebra generators. Write

$$\kappa = \mathbb{k}[\alpha_1, \dots, \alpha_n].$$

If all the α_i are algebraic over $\mathbb{k}$, then κ is a finite field extension of $\mathbb{k}$, and there is nothing to prove.

Assume, for a contradiction, that one of the generators is transcendental over $\mathbb{k}$; after reordering, say α_n is transcendental. Put

$$K = \mathbb{k}(\alpha_n).$$

Since

$$\kappa = K[\alpha_1, \dots, \alpha_{n-1}],$$

the induction hypothesis, applied over the field K , shows that $\alpha_1, \dots, \alpha_{n-1}$ are algebraic over K . Let $f_i \in K[T]$ be a monic polynomial satisfied by α_i , for $1 \leq i \leq n-1$. Choose a non-zero polynomial $g \in \mathbb{k}[\alpha_n]$ clearing all denominators of the coefficients of these polynomials. Then the α_i are integral over

$$A = \mathbb{k}[\alpha_n, g^{-1}].$$

Hence

$$A[\alpha_1, \dots, \alpha_{n-1}]$$

is integral over A . Moreover

$$\kappa = \mathbb{k}[\alpha_n][\alpha_1, \dots, \alpha_{n-1}] \subset A[\alpha_1, \dots, \alpha_{n-1}] \subset \kappa,$$

so equality holds and κ is integral over A .

Since κ is a field, the previous lemma implies that A is a field. But this is impossible: as α_n is transcendental over $\mathbb{k}$, the ring $\mathbb{k}[\alpha_n, g^{-1}]$ is a localisation of the polynomial ring $\mathbb{k}[\alpha_n]$ obtained by inverting only the powers of g , and it is not a field. Indeed, one can choose an irreducible polynomial in $\mathbb{k}[\alpha_n]$ which does not divide g , and its class is then non-invertible in $\mathbb{k}[\alpha_n, g^{-1}]$.

This contradiction shows that all the α_i are algebraic over $\mathbb{k}$. Therefore κ is finite over $\mathbb{k}$. $\square$

I.C.3 Hilbert's Nullstellensatz

Theorem I.C.14. *Let $\mathbb{k}$ be an algebraically closed field and put $A = \mathbb{k}[x_1, \dots, x_n]$.*

- i) *Any maximal ideal of A is of the form $\mathfrak{m}_a = (x_1 - a_1, \dots, x_n - a_n)$ for some $a = (a_1, \dots, a_n) \in \mathbb{k}^n$.*
- ii) *For any ideal $\mathfrak{a} \neq A$, we have $\mathbf{V}(\mathfrak{a}) \neq \emptyset$.*
- iii) *Then, for any ideal $\mathfrak{a} \subset A$, we have $\mathbf{I}(\mathbf{V}(\mathfrak{a})) = \sqrt{\mathfrak{a}}$.*

Proof. Let us prove i). First, we show that $\mathfrak{m}_a$ is maximal. To see this, note that the evaluation map

$$\varphi_a : A \rightarrow \mathbb{k}, \quad f \mapsto f(a)$$

is a surjective ring homomorphism. So $\ker(\varphi_a)$ is a maximal ideal. It turns out that $\ker(\varphi_a) = \mathfrak{m}_a$. Indeed, it is clear that $\mathfrak{m}_a$ lies in $\ker(\varphi_a)$, for $\varphi_a(x_i - a_i) = 0$ for $1 \leq i \leq n$. Conversely, if $f \in \ker(\varphi_a)$, then $f(a) = 0$, i.e. the polynomial $f(x + a)$ has no constant term. Hence $f(x + a)$ lies in the ideal $(x_1, \dots, x_n)$, and therefore f lies in $(x_1 - a_1, \dots, x_n - a_n) = \mathfrak{m}_a$.

The main point of i) is that any maximal ideal $\mathfrak{m} \subset A$ is of the form $\mathfrak{m} = \mathfrak{m}_a$, for some $a \in \mathbb{k}^n$. For this, let $\kappa = A/\mathfrak{m}$, write $\pi : A \rightarrow \kappa$ for the canonical projection and $\varphi : \mathbb{k} \rightarrow \kappa$ for the map given by the inclusion of $\mathbb{k}$ in A as the set of constant polynomials, composed with π .

The morphism φ is necessarily injective, so κ is a field extension of $\mathbb{k}$. Also, the field κ is finitely generated as a $\mathbb{k}$ -algebra, as A itself is generated by $x_1, \dots, x_n$ as a $\mathbb{k}$ -algebra and κ is a quotient of A . Therefore, by Lemma I.C.13, κ is a finite field extension of $\mathbb{k}$. So, since $\mathbb{k}$ is algebraically closed, $\varphi : \mathbb{k} \rightarrow \kappa$ must be an isomorphism. Hence, denoting by b_i the image of x_i in κ , and $a_i = \varphi^{-1}(b_i)$, for $1 \leq i \leq n$, we have $\pi(x_i - a_i) = 0$ so $x_i - a_i \in \mathfrak{m}$. Hence $\mathfrak{m}_a \subset \mathfrak{m}$. But $\mathfrak{m}_a$ is maximal, and $\mathfrak{m} \neq A$, so $\mathfrak{m} = \mathfrak{m}_a$.

Let us show ii). Consider an ideal $\mathfrak{a} \subsetneq A$. Then, there is a maximal ideal containing $\mathfrak{a}$, by Krull's theorem, based on Zorn's lemma. Therefore, there is $a \in \mathbb{k}^n$ such that $\mathfrak{a} \subset \mathfrak{m}_a$. Hence a lies in $\mathbf{V}(\mathfrak{a})$, which is thus non-empty.

Finally, we prove iii). Let $f \in \sqrt{\mathfrak{a}}$. Then $f^n \in \mathfrak{a}$ for some $n \in \mathbb{N}$. Take $a \in \mathbf{V}(\mathfrak{a})$. Then $f^n(a) = 0$, so $f(a) = 0$. This happens for all $a \in \mathbf{V}(\mathfrak{a})$, so $f \in \mathbf{I}(\mathbf{V}(\mathfrak{a}))$.

To prove the other inclusion, we use Rabinowitsch's trick. Consider $f \in \mathbf{I}(\mathbf{V}(\mathfrak{a}))$, so $f(a) = 0$, for all $a \in \mathbf{V}(\mathfrak{a})$. Consider the ideal

$$\mathfrak{b} = \mathfrak{a} + (ft - 1) \subset A[t] = \mathbb{k}[x_1, \dots, x_n, t].$$

We claim that $\mathbf{V}(\mathfrak{b}) = \emptyset$. Indeed, if $(a, \lambda) \in \mathbb{k}^{n+1}$ were a common zero of $\mathfrak{b}$, then $a \in \mathbf{V}(\mathfrak{a})$ and $f(a)\lambda - 1 = 0$. But f vanishes on $\mathbf{V}(\mathfrak{a})$ by assumption, so $f(a) = 0$, a contradiction.

By ii), applied to the polynomial ring $A[t]$, since $\mathbf{V}(\mathfrak{b}) = \emptyset$, the ideal $\mathfrak{b}$ must be the whole ring $A[t]$. Hence there exist $h_1, \dots, h_s \in \mathfrak{a}$ and $q_1, \dots, q_s, q \in A[t]$ such that

$$1 = q_1 h_1 + \dots + q_s h_s + q(ft - 1).$$

Passing to the quotient ring $(A/\mathfrak{a})[t]$, the first terms disappear. If we denote by $\bar{f}$ the image of f in $A/\mathfrak{a}$, we obtain

$$1 = \bar{q}(\bar{f}t - 1) \quad \text{in } (A/\mathfrak{a})[t].$$

Write

$$\bar{q} = a_0 + a_1 t + \dots + a_N t^N, \quad a_i \in A/\mathfrak{a},$$

with $a_N \neq 0$ if $\bar{q} \neq 0$. Then

$$1 = (a_0 + a_1 t + \dots + a_N t^N)(\bar{f}t - 1).$$

Comparing the coefficient of t^0 , we get

$$-a_0 = 1,$$

hence $a_0 = -1$. Comparing the coefficient of t^1 , we get

$$a_0 \bar{f} - a_1 = 0,$$

hence $a_1 = a_0 \bar{f} = -\bar{f}$. Comparing the coefficient of t^2 , we get

$$a_1 \bar{f} - a_2 = 0,$$

hence $a_2 = -\bar{f}^2$. Continuing in the same way, we obtain

$$a_i = -\bar{f}^i \quad \text{for } 0 \leq i \leq N.$$

Finally, comparing the coefficient of t^{N+1} gives

$$a_N \bar{f} = 0.$$

Since $a_N = -\bar{f}^N$, we get

$$\bar{f}^{N+1} = 0$$

in $A/\mathfrak{a}$. Therefore $f^{N+1} \in \mathfrak{a}$, and hence $f \in \sqrt{\mathfrak{a}}$. □

Example I.C.15. Assume that $\mathbb{k}$ is algebraically closed. In $\mathbb{k}[x]$, the ideal

$$(x^2(x-1)^3)$$

defines the same algebraic set as its radical:

$$\mathbf{V}(x^2(x-1)^3) = \{0, 1\} = \mathbf{V}(x(x-1)).$$

The Nullstellensatz says precisely that

$$\mathbf{I}(\{0, 1\}) = (x(x-1)) = \sqrt{(x^2(x-1)^3)}.$$

Example I.C.16. In $\mathbb{k}[x, y]$, with $\mathbb{k}$ algebraically closed, the ideals

$$(x^2, y) \quad \text{and} \quad (x, y)$$

define the same algebraic set, namely the origin. The first ideal is not radical, and its radical is (x, y) .

Exercise I.C.17. Let $\mathbb{k}$ be algebraically closed. Compute $\mathbf{I}(\mathbf{V}(\mathfrak{a}))$ for

$$\mathfrak{a} = (x^2, y^3) \subset \mathbb{k}[x, y].$$

Solution. We have

$$\mathbf{V}(\mathfrak{a}) = \{(0, 0)\}.$$

Hence $\mathbf{I}(\mathbf{V}(\mathfrak{a})) = \mathbf{I}(\{(0, 0)\}) = (x, y)$. Equivalently, by the Nullstellensatz,

$$\mathbf{I}(\mathbf{V}(\mathfrak{a})) = \sqrt{(x^2, y^3)} = (x, y).$$

Exercise I.C.18. Let $\mathbb{k}$ be algebraically closed and let

$$\mathfrak{a} = (xy, xz) \subset \mathbb{k}[x, y, z].$$

Describe $\mathbf{V}(\mathfrak{a})$ geometrically and compute $\sqrt{\mathfrak{a}}$.

Solution. The equations are $xy = 0$ and $xz = 0$, i.e. $xy = xz = 0$. Thus either $x = 0$, or both $y = 0$ and $z = 0$. Therefore

$$\mathbf{V}(\mathfrak{a}) = \mathbf{V}(x) \cup \mathbf{V}(y, z),$$

the union of the plane $x = 0$ and the x -axis. Correspondingly,

$$\sqrt{\mathfrak{a}} = (x) \cap (y, z).$$

Since $(x) \cap (y, z) = (xy, xz)$, the ideal $\mathfrak{a}$ is already radical.

I.C.4 Geometrically regular rational functions

We introduce the field of rational functions in n variables

$$K = \mathbb{k}(x_1, \dots, x_n).$$

It consists of equivalence classes of fractions u/v , where $u, v \in A = \mathbb{k}[x_1, \dots, x_n]$ and $v \neq 0$, with

$$\frac{u}{v} = \frac{u'}{v'} \iff uv' = u'v.$$

A chosen fraction u/v defines a $\mathbb{k}$ -valued function away from $\mathbf{V}(v)$. To indicate that it need not be defined everywhere, we sometimes write

$$\frac{u}{v} : \mathbb{k}^n \dashrightarrow \mathbb{k}.$$

We shall also consider the same fraction over an algebraic closure $\overline{\mathbb{k}}$.

Definition I.C.19. Let $P \subset \overline{\mathbb{k}}^n$. A rational function $g \in K$ is *geometrically regular on P* if it admits a representation

$$g = \frac{u}{v}, \quad u, v \in A,$$

such that $v(a) \neq 0$ for every $a \in P$. Note that the condition concerns the rational function and not a particular fraction representing it. For instance, $x_1/x_1 = 1$ is geometrically regular everywhere, although the displayed fraction is not defined along $\mathbf{V}(x_1)$.

For $f \in A$, put

$$D(f) = \overline{\mathbb{k}}^n \setminus \mathbf{V}(f).$$

Every fraction of the form u/f^m , with $u \in A$ and $m \geq 0$, is geometrically regular on $D(f)$. The converse is also true.

Lemma I.C.20. *Let $f \in A$. A rational function $g \in K$ is geometrically regular on $D(f)$ if and only if*

$$g = \frac{u}{f^m}$$

for some $u \in A$ and $m \geq 0$.

Proof. One implication is immediate. Conversely, write $g = a/b$, with $a, b \in A$, $b \neq 0$, and with a and b having no common irreducible factor. Since g is geometrically regular on $D(f)$, we may choose this representation so that b has no zero on $D(f)$. Thus

$$\mathbf{V}(b) \subset \mathbf{V}(f)$$

in $\overline{\mathbb{k}}^n$. By the Nullstellensatz, there is an integer $N \geq 1$ such that $f^N \in (b)$. Consequently $f^N = bc$ for some $c \in A$, and hence

$$g = \frac{a}{b} = \frac{ac}{f^N}.$$

□

Taking $f = 1$, we obtain that the rational functions geometrically regular on all of $\overline{\mathbb{k}}^n$ are precisely the polynomial functions. We now pass from affine space to an affine algebraic set. Let $X = \mathbf{V}(\mathfrak{a}) \subset \overline{\mathbb{k}}^n$. For a reason that will be apparent later, we assume that $\mathfrak{a} \subset A$ is a prime ideal. The immediate consequence is that the coordinate ring

$$\mathbb{k}[X] = A/\mathfrak{a}$$

is an integral domain.

Definition I.C.21. The fraction field

$$K(X) = \text{Frac}(\mathbb{k}[X])$$

is called the *field of rational functions on X* . A rational function $g \in K(X)$ is geometrically regular on a subset $P \subset X$ if it admits a representation

$$g = \frac{u}{v}, \quad u, v \in \mathbb{k}[X],$$

such that $v(p) \neq 0$ for every $p \in P$.

If $f \in \mathbb{k}[X]$, we write

$$D_X(f) = \{p \in X \mid f(p) \neq 0\}.$$

Choosing a polynomial representative $\tilde{f} \in A$, this is simply

$$D_X(f) = X \cap D(\tilde{f}),$$

and the right-hand side does not depend on the chosen representative.

Exactly as above, a rational function on X is geometrically regular on $D_X(f)$ if and only if it can be written

$$\frac{u}{f^m}, \quad u \in \mathbb{k}[X], \quad m \geq 0.$$

Later, this construction will be formulated systematically using localisation. For the moment, the important point is geometric: allowing powers of f in the denominator gives precisely the rational functions which remain regular wherever f does not vanish.

I.D Noether normalisation

Further reading. See [Har77, Chapter I, §1] for the classical geometric use of Noether normalisation, and [Liu02] for a systematic algebraic treatment.

Let $A \rightarrow B$ be a ring homomorphism. Then B is equipped with an A -module structure. We say that B is *finite* over A if B is a finitely generated A -module. We will use the notion of *algebraically independent elements* $f_1, \dots, f_n$ of a $\mathbb{k}$ -algebra A , namely, there is no non-zero polynomial $g \in \mathbb{k}[x_1, \dots, x_n]$ such that $g(f_1, \dots, f_n) = 0$.

In this case, the subring $\mathbb{k}[f_1, \dots, f_n] \subset A$ generated by these elements is isomorphic to $\mathbb{k}[x_1, \dots, x_n]$, where $x_1, \dots, x_n$ are indeterminates. Indeed, any element in the kernel of the map $\mathbb{k}[x_1, \dots, x_n] \rightarrow A$ sending x_i to f_i for $1 \leq i \leq n$ is a polynomial $g \in \mathbb{k}[x_1, \dots, x_n]$ such that $g(f_1, \dots, f_n) = 0$.

I.D.1 The algebraic statement

Theorem I.D.1. *Let $\mathbb{k}$ be a field and let $B \neq 0$ be a finitely generated $\mathbb{k}$ -algebra. Then there exist an integer m and algebraically independent elements $y_1, \dots, y_m \in B$ such that B is finite over $\mathbb{k}[y_1, \dots, y_m]$.*

Proof. Since B is a finitely generated $\mathbb{k}$ -algebra, there is a surjection $\varphi : \mathbb{k}[x_1, \dots, x_n] \rightarrow B$, where n is the number of generators of B as a $\mathbb{k}$ -algebra and $x_1, \dots, x_n$ are indeterminates. We write simply $\mathbb{k}[x] = \mathbb{k}[x_1, \dots, x_n]$. Then we have $B \simeq \mathbb{k}[x]/\mathfrak{a}$, where $\mathfrak{a} = \ker(\varphi)$. We replace B with $\mathbb{k}[x]/\mathfrak{a}$ and we prove the statement by induction on n . If $n = 0$, then $B \simeq \mathbb{k}$ and there is nothing to prove. Also, if $\mathfrak{a} = 0$, then $B \simeq \mathbb{k}[x_1, \dots, x_n]$ and the statement is clear.

So let $n \geq 1$, $\mathfrak{a} \neq 0$. We may assume that the statement holds for $\mathbb{k}$ -algebras generated by $n - 1$ elements. Consider a non-zero element $f \in \mathfrak{a}$ and write

$$f = \sum_{\mu \in \mathbb{N}^n} a_\mu x^\mu, \quad \text{for some } a_\mu \in \mathbb{k}, \text{ for all } \mu \in \mathbb{N}^n, \text{ with } x^\mu = x_1^{\mu_1} \cdots x_n^{\mu_n}.$$

For every $(\nu_1, \dots, \nu_{n-1}) \in \mathbb{N}^{n-1}$, we consider $\nu = (\nu_1, \dots, \nu_{n-1}, 1) \in \mathbb{N}^n$ and we introduce:

$$s_i = x_i - x_n^{\nu_i}, \quad \text{for } 1 \leq i \leq n-1.$$

We write $s = (s_1, \dots, s_{n-1})$. Since for $1 \leq i \leq n-1$, x_i can be expressed in terms of s_i and x_n , we have

$$\mathbb{k}[x] = \mathbb{k}[s][x_n].$$

The point is that, for a suitable choice of ν , writing f in the variables $s_1, \dots, s_{n-1}, x_n$ results in a monic polynomial in x_n , up to a non-zero scalar. Indeed, we consider the lexicographic order on $\mathbb{N}^n$, so that for any $\mu, \mu' \in \mathbb{N}^n$, we let $\mu < \mu'$ if there is $1 \leq j \leq n$ such that $\mu_i = \mu'_i$ for $1 \leq i \leq j-1$ and $\mu_j < \mu'_j$. Then we put $d = \deg(f)$, $N = d + 1$ and we let

$$\mu^0 = \max\{\mu \in \mathbb{N}^n \mid a_\mu \neq 0\}, \quad \nu_i = N^{n-i}, \quad 1 \leq i \leq n.$$

We claim that, after the change of variables

$$x_i = s_i + x_n^{\nu_i}, \quad 1 \leq i \leq n-1,$$

the polynomial $f(x_1, \dots, x_n)$, viewed as an element of $\mathbb{k}[s_1, \dots, s_{n-1}][x_n]$, is monic in x_n , up to the coefficient a_{μ^0} . More precisely,

$$f(x_1, \dots, x_n) = f(s_1 + x_n^{\nu_1}, \dots, s_{n-1} + x_n^{\nu_{n-1}}, x_n) = \sum_{\mu \in \mathbb{N}^n} a_\mu \prod_{1 \leq i \leq n-1} (s_i + x_n^{\nu_i})^{\mu_i} x_n^{\mu_n}.$$

In this expression, the largest power of x_n contributed by the monomial x^μ is $x_n^{\langle \nu | \mu \rangle}$, where

$$\langle \nu | \mu \rangle = \nu_1 \mu_1 + \cdots + \nu_{n-1} \mu_{n-1} + \mu_n.$$

Now, take $\mu \neq \mu^0$ with $a_\mu \neq 0$. We have $\mu < \mu^0$, so there is $1 \leq j \leq n$ such that $\mu_i = \mu_i^0$ for $1 \leq i \leq j-1$ and $\mu_j < \mu_j^0$. Then our choice of ν gives

$$\langle \nu | \mu^0 - \mu \rangle = (\mu_j^0 - \mu_j) N^{n-j} + \sum_{j+1 \leq i \leq n} (\mu_i^0 - \mu_i) N^{n-i}.$$

The first term is at least N^{n-j} . The second term has absolute value at most

$$d(N^{n-j-1} + \cdots + 1) = d \frac{N^{n-j} - 1}{N - 1} = N^{n-j} - 1,$$

because $d = N - 1$. Therefore

$$\langle \nu | \mu^0 - \mu \rangle > 0,$$

hence $\langle \nu | \mu \rangle < \langle \nu | \mu^0 \rangle$. In conclusion, only μ^0 contributes to the coefficient of the highest power of x_n , namely $x_n^{\langle \nu | \mu^0 \rangle}$, and the coefficient of this power in the expression of $f(x_1, \dots, x_n)$ in the variables $s_1, \dots, s_{n-1}, x_n$ is a_{μ^0} . Dividing by this non-zero scalar, we obtain a monic polynomial in x_n .

Having set up this expression, we consider $\mathfrak{b} = \mathfrak{a} \cap \mathbb{k}[s]$ and $C = \mathbb{k}[s]/\mathfrak{b}$. Since C is generated by $n - 1$ elements, by induction, there are algebraically independent elements $y_1, \dots, y_m$ of C such that C is finite over $\mathbb{k}[y] = \mathbb{k}[y_1, \dots, y_m]$. By construction, we have an injection $C \hookrightarrow B$, and B is generated by the image b_n of x_n in B , as a C -algebra. Since the equation $f = 0$ gives a monic equation in x_n , the element b_n is integral over C , so B is finitely generated as a C -module by Lemma I.C.10. Also, C is finitely generated as a $\mathbb{k}[y]$ -module, so B is finitely generated as a $\mathbb{k}[y]$ -module too. $\square$

Example I.D.2. Let

$$B = \mathbb{k}[x, y]/(y^2 - x^3).$$

The image of x is algebraically independent over $\mathbb{k}$, and the image of y is integral over $\mathbb{k}[x]$, since it satisfies $T^2 - x^3 = 0$. Hence B is finite over $\mathbb{k}[x]$.

Example I.D.3. Let

$$B = \mathbb{k}[x, y]/(xy - 1).$$

This algebra is isomorphic to $\mathbb{k}[x, x^{-1}]$. It is not finite over $\mathbb{k}[x]$, since x^{-1} is not integral over $\mathbb{k}[x]$. However, Noether normalisation says that, after choosing a better coordinate, B is finite over a polynomial subring in one variable.

Exercise I.D.4. Show that $\mathbb{k}[x, y]/(y^2 - x)$ is finite over $\mathbb{k}[y]$ and also finite over $\mathbb{k}[x]$. Compare the two module structures.

Solution. Let $B = \mathbb{k}[x, y]/(y^2 - x)$. Since $x = y^2$ in B , we have $B = \mathbb{k}[y]$. Therefore B is finite over $\mathbb{k}[y]$, in fact free of rank 1. It is also finite over $\mathbb{k}[x]$: the element y satisfies the monic equation $T^2 - x = 0$, so B is generated by $1, y$ as a $\mathbb{k}[x]$ -module. Thus the first module structure has rank 1, while the second one has rank 2.

I.D.2 Geometric version of Noether normalisation

Consider an ideal $\mathfrak{a} \subset \mathbb{k}[x_1, \dots, x_n]$ and the algebraic set $X = \mathbf{V}(\mathfrak{a}) \subset \mathbb{k}^n$. Put

$$B = \mathbb{k}[x_1, \dots, x_n]/\mathfrak{a}.$$

By Noether normalisation, there are algebraically independent elements $y_1, \dots, y_m \in B$ such that, if

$$\mathbb{k}[y_1, \dots, y_m] \subset B,$$

then B is finite as an A -module. In other words, setting $A = \mathbb{k}[t_1, \dots, t_m]$, where $t_1, \dots, t_m$ are indeterminates, we have an injective morphism of rings $A \rightarrow B$ defined by sending t_i to y_i for $1 \leq i \leq m$, and B is a finitely generated A -module.

Choose polynomial representatives $f_1, \dots, f_m \in \mathbb{k}[x_1, \dots, x_n]$ of $y_1, \dots, y_m$. These functions induce a map of algebraic sets

$$f : X \rightarrow \mathbb{k}^m, \quad b \mapsto (f_1(b), \dots, f_m(b)).$$

Proposition I.D.5. *For all $a \in \mathbb{k}^m$, the fibre $f^{-1}(a)$ is finite. If the field $\mathbb{k}$ is algebraically closed, this fibre is also non-empty.*

Proof. Let $a = (a_1, \dots, a_m) \in \mathbb{k}^m$. We know that B is finitely generated as an A -module. For $1 \leq j \leq n$, the image $\bar{x}_j$ of x_j in B is therefore integral over A . Hence, for each j , there is a monic polynomial

$$h_j = h_j(t_1, \dots, t_m)(T) \in A[T]$$

such that

$$h_j(y_1, \dots, y_m)(\bar{x}_j) = 0$$

in B . By mapping t_i to f_i , for $1 \leq i \leq m$, we get a polynomial

$$H_j = h_j(f_1, \dots, f_m)(T) \in \mathbb{k}[x_1, \dots, x_n, T]$$

which is monic in T and satisfies

$$H_j(x_1, \dots, x_n)(x_j) \in \mathfrak{a}.$$

Now let $b = (b_1, \dots, b_n) \in f^{-1}(a)$. Then $b \in X$, so $g(b) = 0$ for all $g \in \mathfrak{a}$, and also

$$f_i(b) = a_i \quad \text{for } 1 \leq i \leq m.$$

Since $H_j(x_1, \dots, x_n)(x_j) \in \mathfrak{a}$, evaluating at b gives

$$H_j(b_1, \dots, b_n)(b_j) = 0, \quad \text{for } 1 \leq j \leq n.$$

In other words, if we set $\ell_j(T) = h_j(a_1, \dots, a_m)(T) \in \mathbb{k}[T]$, for $1 \leq j \leq n$, then

$$\ell_j(b_j) = H_j(b_1, \dots, b_n)(b_j) = h_j(f_1(b), \dots, f_m(b))(b_j) = 0, \quad \text{for } 1 \leq j \leq n.$$

Since, for $1 \leq j \leq n$, each $\ell_j(T) = h_j(a_1, \dots, a_m)(T)$ is a monic polynomial in $\mathbb{k}[T]$, it has only finitely many roots. Hence each coordinate b_j can take only finitely many values. Therefore there are only finitely many possible points $b = (b_1, \dots, b_n)$ in the fibre $f^{-1}(a)$.

Let us now prove non-emptiness, assuming that $\mathbb{k}$ is algebraically closed. By the Nullstellensatz, it is enough to prove that

$$\mathfrak{a} + (f_1 - a_1, \dots, f_m - a_m) \neq \mathbb{k}[x_1, \dots, x_n].$$

Equivalently, it is enough to prove that

$$(\bar{f}_1 - a_1, \dots, \bar{f}_m - a_m)B \neq B. \quad (\text{I.1})$$

Let $\mathfrak{m}_a = (y_1 - a_1, \dots, y_m - a_m) \subset A$. Under the inclusion $A \subset B$, the ideal $\mathfrak{m}_a B$ is exactly

$$(\bar{f}_1 - a_1, \dots, \bar{f}_m - a_m)B.$$

Since B is a finite A -module and $B \neq 0$, Nakayama's lemma (see the next exercise) gives $\mathfrak{m}_a B \neq B$. Therefore (I.1) holds, and the proof is complete. $\square$

Exercise I.D.6. Prove the following form of Nakayama's lemma. Let $A \rightarrow B$ be an injective homomorphism of rings, with $B \neq 0$, and assume that B is a finitely generated A -module. If $\mathfrak{a} \subsetneq A$ is a proper ideal, then $\mathfrak{a}B \neq B$.

Solution. Since B is a finitely generated A -module, there are $b_1, \dots, b_n \in B$ such that $B = b_1 A + \dots + b_n A$. Suppose, for a contradiction, that $\mathfrak{a}B = B$. Then there are elements $a_{i,j} \in \mathfrak{a}$, for $1 \leq i, j \leq n$, such that

$$b_i = \sum_{1 \leq j \leq n} a_{i,j} b_j, \quad \text{for } 1 \leq i \leq n.$$

Hence the matrix

$$M = (\delta_{i,j} - a_{i,j})_{1 \leq i, j \leq n}$$

annihilates the column vector $(b_1, \dots, b_n)^t$. By the determinant trick, $d = \det(M)$ annihilates each b_i , and therefore annihilates B . Since $A \rightarrow B$ is injective, the A -module B is faithful: if $cB = 0$, then in particular $c \cdot 1_B = 0$, so $c = 0$ in A . Thus $d = 0$ in A .

On the other hand, expanding the determinant gives

$$d = 1 + \alpha$$

for some $\alpha \in \mathfrak{a}$. Since $d = 0$, we get $1 = -\alpha \in \mathfrak{a}$, contradicting the fact that $\mathfrak{a}$ is proper. Hence $\mathfrak{a}B \neq B$.

Exercise I.D.7. Again in the spirit of Example I.D.3.

$$B = \mathbb{k}[x, y]/(xy - 1) \simeq \mathbb{k}[x, x^{-1}]$$

and set $u = x + y = x + x^{-1}$.

- i) Show that the homomorphism $\mathbb{k}[u] \hookrightarrow B$ is injective.
- ii) Show that x is integral over $\mathbb{k}[u]$ and that B is generated by $1, x$ as a $\mathbb{k}[u]$ -module.
- iii) Describe the fibres of the corresponding map $\mathbf{V}(xy - 1) \rightarrow \mathbb{k}$, $(x, y) \mapsto x + y$.

Solution. The element x^{-1} is not integral over $\mathbb{k}[x]$. Indeed, if there was a monic polynomial $f \in \mathbb{k}[x]$ of degree m vanishing at x^{-1} , then looking at $x^{m-1}f$ we see that x^{-1} would then lie in $\mathbb{k}[x]$, which is not the case. Then, $u = x + x^{-1}$ is also transcendental: if $P(u) = 0$ for a non-zero polynomial P , multiplication by a sufficiently high power of x would give a non-zero polynomial relation for x . Since $y = u - x$ and $xy = 1$, the element x satisfies

$$x^2 - ux + 1 = 0.$$

Thus x is integral over $\mathbb{k}[u]$ and every element of $B = \mathbb{k}[u][x]$ reduces to $a(u) + b(u)x$. Over $u = c$, the possible values of x are the roots of

$$x^2 - cx + 1 = 0,$$

and $y = x^{-1}$. Hence every fibre has at most two points; over an algebraically closed field it has two points counted with multiplicity.

Example I.D.8 (An elliptic curve as a branched double cover). Assume that $\mathbb{k}$ is algebraically closed of characteristic different from 2, and choose $\lambda \in \mathbb{k} \setminus \{0, 1\}$. Consider

$$B = \mathbb{k}[x, y] / (y^2 - x(x-1)(x-\lambda)).$$

The image of x is algebraically independent over $\mathbb{k}$, while the image of y is integral over $\mathbb{k}[x]$, since it satisfies the monic equation

$$T^2 - x(x-1)(x-\lambda) = 0.$$

Hence B is finite over $\mathbb{k}[x]$. More precisely,

$$B = \mathbb{k}[x] \oplus \mathbb{k}[x]y$$

as a $\mathbb{k}[x]$ -module. Geometrically, the corresponding map

$$\mathbf{V}(y^2 - x(x-1)(x-\lambda)) \longrightarrow \mathbb{k}, \quad (x, y) \longmapsto x,$$

is generically two-to-one. Indeed, the fibre over $a \in \mathbb{k}$ is determined by

$$y^2 = a(a-1)(a-\lambda).$$

It consists of two distinct points when

$$a \notin \{0, 1, \lambda\},$$

and of a single point when $a \in \{0, 1, \lambda\}$. Thus the affine double cover is branched over 0, 1, and λ .

II Projective algebraic sets

Further reading. For projective algebraic sets, homogeneous coordinates, and morphisms, see [Har77, Chapter I, §§2–3]; [Hul03] and [Hol12] contain many concrete examples.

Throughout this section, $\mathbb{k}$ is a field. We consider the polynomial ring

$$A = \mathbb{k}[x_0, \dots, x_n],$$

endowed with its standard grading by total degree.

II.A Projective space and projective subspaces

Definition II.A.1. Let $n \geq 0$. The n -dimensional *projective space* over $\mathbb{k}$ is the quotient

$$\mathbf{P}_{\mathbb{k}}^n = (\mathbb{k}^{n+1} \setminus \{0\}) / \mathbb{k}^*,$$

where $\lambda \in \mathbb{k}^*$ acts by

$$\lambda \cdot (a_0, \dots, a_n) = (\lambda a_0, \dots, \lambda a_n).$$

The class of a non-zero vector $(a_0, \dots, a_n)$ is denoted by

$$[a_0 : \dots : a_n].$$

The entries $a_0, \dots, a_n$ are called homogeneous coordinates. By convention, $\mathbf{P}(\{0\}) = \emptyset$.

Thus, a point of $\mathbf{P}_{\mathbb{k}}^n$ is identified with a one-dimensional linear subspace of $\mathbb{k}^{n+1}$. For $0 \leq i \leq n$, set

$$U_i = \{[a_0 : \dots : a_n] \in \mathbf{P}_{\mathbb{k}}^n \mid a_i \neq 0\}.$$

Remark II.A.2. Every point of U_i has a unique representative whose i -th coordinate is equal to 1. We therefore obtain a bijection

$$\varphi_i : U_i \longrightarrow \mathbb{k}^n, \quad [a_0 : \cdots : a_n] \longmapsto \left(\frac{a_0}{a_i}, \dots, \widehat{\frac{a_i}{a_i}}, \dots, \frac{a_n}{a_i} \right).$$

The subsets $U_0, \dots, U_n$ cover $\mathbf{P}_{\mathbb{k}}^n$ and are called the standard affine charts of projective space.

Remark II.A.3. In particular,

$$\mathbf{P}_{\mathbb{k}}^n = U_0 \sqcup H_{\infty}, \quad H_{\infty} = \{[a_0 : \cdots : a_n] \mid a_0 = 0\} \simeq \mathbf{P}_{\mathbb{k}}^{n-1}.$$

Under the identification $U_0 \simeq \mathbb{k}^n$, the subset H_{∞} is called the hyperplane at infinity.

Definition II.A.4. Let $V = \mathbb{k}^{n+1}$ and let $W \subset V$ be a non-zero vector subspace. The associated projective subspace is

$$\mathbf{P}(W) = \{[w] \in \mathbf{P}(V) \mid 0 \neq w \in W\}.$$

If $\dim(W) = r + 1$, then $\mathbf{P}(W)$ is called a projective r -plane. Projective 1-planes and projective $(n - 1)$ -planes are called, respectively, projective lines and projective hyperplanes.

If W is the common kernel of linear forms $\ell_1, \dots, \ell_s \in V^{\vee}$, then

$$\mathbf{P}(W) = \{[a] \in \mathbf{P}(V) \mid \ell_1(a) = \cdots = \ell_s(a) = 0\}.$$

Thus projective subspaces are described by homogeneous linear equations.

Example II.A.5. We have

$$\mathbf{P}_{\mathbb{k}}^1 = U_0 \sqcup \{[0 : 1]\} \simeq \mathbb{k} \sqcup \{\infty\}.$$

The point $[0 : 1]$ is the point at infinity of the affine line $U_0 = \{[1 : t] \mid t \in \mathbb{k}\}$.

Exercise II.A.6. Show that each map $\varphi_i : U_i \rightarrow \mathbb{k}^n$ is a bijection. Describe explicitly the change of coordinates from the chart U_i to the chart U_j on $U_i \cap U_j$.

Solution. The inverse of φ_i inserts the coordinate 1 in the i -th position. For instance, for $i = 0$ it is

$$(u_1, \dots, u_n) \longmapsto [1 : u_1 : \cdots : u_n].$$

On $U_i \cap U_j$, write $u_{\ell} = a_{\ell}/a_i$, with $u_i = 1$. Since $u_j \neq 0$, the coordinates $v_{\ell} = a_{\ell}/a_j$ in the chart U_j are

$$v_i = \frac{1}{u_j}, \quad v_{\ell} = \frac{u_{\ell}}{u_j} \quad \text{for } \ell \neq i, j, \quad v_j = 1.$$

Exercise II.A.7. Let

$$p = [a_0 : \cdots : a_n], \quad q = [b_0 : \cdots : b_n]$$

be distinct points of $\mathbf{P}_{\mathbb{k}}^n$.

- i) Show that there is a unique projective line L containing p and q , namely $L = \mathbf{P}(\langle a, b \rangle)$.
- ii) Show that a point $[x_0 : \cdots : x_n]$ lies on L if and only if the matrix

$$\begin{pmatrix} a_0 & \cdots & a_n \\ b_0 & \cdots & b_n \\ x_0 & \cdots & x_n \end{pmatrix}$$

has rank at most 2. Deduce that L is defined by its 3×3 minors.

iii) When $n = 2$, write the resulting single linear equation explicitly as a determinant.

Solution. Since $p \neq q$, the vectors a and b are linearly independent. Hence $\mathbf{P}(\langle a, b \rangle)$ is a projective line containing p and q . Any projective line containing them comes from a two-dimensional vector subspace containing a and b , so it is the same line.

A vector x belongs to $\langle a, b \rangle$ precisely when the three rows a, b, x are linearly dependent, which is equivalent to the vanishing of all 3×3 minors. For $n = 2$ this gives the equation

$$\det \begin{pmatrix} a_0 & a_1 & a_2 \\ b_0 & b_1 & b_2 \\ x_0 & x_1 & x_2 \end{pmatrix} = 0.$$

II.B Projective equations and homogenisation

We see here how to define algebraic subsets of the projective space. These are associated with homogenous ideals, the graded ring structure of the polynomial ring acquire thus a particular prominence when dealing with subsets of the projective space.

II.B.1 Homogeneous ideals and projective subsets

We start by recalling the definition of homogeneous polynomials and homogeneous ideals. This refers to the usual grading of the polynomial ring, where scalars have degree zero and each variable has degree one.

Definition II.B.1. A polynomial $F \in A$ is *homogeneous of degree d* if every monomial appearing in F has total degree d . Equivalently,

$$F(\lambda x_0, \dots, \lambda x_n) = \lambda^d F(x_0, \dots, x_n)$$

is a polynomial identity. An ideal $\mathfrak{a} \subset A$ is *homogeneous* if it is generated by homogeneous polynomials. Equivalently, whenever $F \in \mathfrak{a}$ and

$$F = F_0 + F_1 + \dots + F_d$$

is its decomposition into homogeneous components, then every F_i belongs to $\mathfrak{a}$.

If F is homogeneous, the condition $F(a_0, \dots, a_n) = 0$ does not change when the vector $(a_0, \dots, a_n)$ is multiplied by a non-zero scalar. It therefore makes sense on projective space.

Definition II.B.2. Let $\mathfrak{a} \subset A$ be a homogeneous ideal. Its *projective zero locus* is

$$\mathbf{V}_+(\mathfrak{a}) = \{[a_0 : \dots : a_n] \in \mathbf{P}_{\mathbb{k}}^n \mid F(a_0, \dots, a_n) = 0 \text{ for every homogeneous } F \in \mathfrak{a}\}.$$

A subset $X \subset \mathbf{P}_{\mathbb{k}}^n$ of the form $X = \mathbf{V}_+(\mathfrak{a})$ is called a projective algebraic set.

For a homogeneous polynomial F , we use the notation

$$D_+(F) = \mathbf{P}_{\mathbb{k}}^n \setminus \mathbf{V}_+(F) = \{p \in \mathbf{P}_{\mathbb{k}}^n \mid F(p) \neq 0\}.$$

At this point, this is only notation for a subset of projective space. Later, after introducing the Zariski topology, we shall see that the subsets $\mathbf{V}_+(\mathfrak{a})$ are closed and that the subsets $D_+(F)$ are open. Notice that

$$U_i = D_+(x_i).$$

If $X \subset \mathbf{P}_{\mathbb{k}}^n$ is fixed, we also write $D_+(F)$ for $X \cap D_+(F)$ when no confusion can arise. Given a subset $X \subset \mathbf{P}_{\mathbb{k}}^n$, its homogeneous vanishing ideal is

$$\mathbf{I}_+(X) = \bigoplus_{d \geq 0} \{F \in A_d \mid F(p) = 0 \text{ for every } p \in X\}.$$

This is a homogeneous ideal of A . The quotient

$$\mathbb{k}[X] = A/\mathbf{I}_+(X)$$

is called the homogeneous coordinate ring of X . It is usually denoted in the same way as the coordinate ring of an affine algebraic subset.

Example II.B.3. A projective subspace is a projective algebraic set defined by homogeneous linear equations. The most familiar example of higher degree equations is provided by the projective plane conic:

$$C = \mathbf{V}_+(x_0x_2 - x_1^2) \subset \mathbf{P}_{\mathbb{k}}^2$$

On the chart U_0 , with affine coordinates $u = x_1/x_0$ and $v = x_2/x_0$, its equation becomes $v = u^2$.

II.B.2 Dehomogenisation, affine charts and projective closure

Let $F \in A$ be homogeneous. Its dehomogenisation with respect to x_i is the polynomial obtained by setting $x_i = 1$. Let us denote it by f . Under the bijection $U_i \simeq \mathbb{k}^n$, the subset

$$\mathbf{V}(f) = \mathbf{V}_+(F) \cap U_i, \quad \text{for } 0 \leq i \leq n.$$

is therefore the affine algebraic set defined by this dehomogenised polynomial. The same statement holds for homogeneous ideals. Here, the equality displayed above in really means that, under the identification $U_i \simeq \mathbb{k}^n$, the subset $\mathbf{V}_+(F) \cap U_i$ corresponds to $\mathbf{V}(f)$.

Let us look at the other direction. Let

$$R = \mathbb{k}[y_1, \dots, y_n]$$

and let $f \in R$ have degree d . The homogenisation of f is

$$f^h(x_0, \dots, x_n) = x_0^d f\left(\frac{x_1}{x_0}, \dots, \frac{x_n}{x_0}\right).$$

This is a homogeneous polynomial of degree d and satisfies

$$f^h(1, y_1, \dots, y_n) = f(y_1, \dots, y_n).$$

Definition II.B.4. For an ideal $\mathfrak{a} \subset R$, define its homogenisation by

$$\mathfrak{a}^h = (f^h \mid f \in \mathfrak{a}) \subset A.$$

It is important here to homogenise every element of $\mathfrak{a}$, and not merely an arbitrarily chosen family of generators.

Let us look at the geometric side of the story. Consider the map

$$\iota : \mathbb{k}^n \longrightarrow \mathbf{P}_{\mathbb{k}}^n, \quad (a_1, \dots, a_n) \longmapsto [1 : a_1 : \dots : a_n].$$

For an affine algebraic set $X \subset \mathbb{k}^n$, define its projective homogenisation by

$$X^h = \mathbf{V}_+(\mathbf{I}(X)^h).$$

Remark II.B.5. For any affine algebraic subset, we have

$$X^h \cap U_0 = \iota(X).$$

Indeed, dehomogenising the equations of $\mathbf{I}(X)^h$ with respect to x_0 gives back the equations of $\mathbf{I}(X)$. When $\mathbb{k}$ is algebraically closed, once the Zariski topology has been introduced, we shall see that X^h is precisely the closure of $\iota(X)$ in $\mathbf{P}_{\mathbb{k}}^n$.

Exercise II.B.6. Compute the projective homogenisations in $\mathbf{P}_{\mathbb{k}}^2$ of the affine curves

$$y = x^2, \quad xy = 1.$$

Determine their points on the line at infinity $\mathbf{V}_+(x_0)$.

Solution. We use homogeneous coordinates $[x_0 : x_1 : x_2]$, with $x = x_1/x_0$ and $y = x_2/x_0$ on U_0 . The equation $y = x^2$ becomes

$$x_0x_2 - x_1^2 = 0.$$

On the line $x_0 = 0$, this gives $x_1 = 0$, hence the unique point at infinity is $[0 : 0 : 1]$.

The equation $xy = 1$ becomes

$$x_1x_2 - x_0^2 = 0.$$

On the line $x_0 = 0$, it gives $x_1x_2 = 0$, hence the two points at infinity are

$$[0 : 1 : 0] \quad \text{and} \quad [0 : 0 : 1].$$

Exercise II.B.7. In $\mathbb{k}[x, y]$, the ideal (x, y) is also generated by x and $y + x^2$. Homogenise these two different systems of generators and compare the resulting projective zero loci in $\mathbf{P}_{\mathbb{k}}^2$.

Solution. Using coordinates $[x_0 : x_1 : x_2]$, the homogenisations of x and y are x_1 and x_2 . Their common zero locus is the single point

$$\mathbf{V}_+(x_1, x_2) = \{[1 : 0 : 0]\}.$$

On the other hand, the homogenisations of x and $y + x^2$ are

$$x_1, \quad x_0x_2 + x_1^2.$$

Their common zero locus is defined by $x_1 = 0$ and $x_0x_2 = 0$, hence it is

$$\{[1 : 0 : 0], [0 : 0 : 1]\}.$$

Thus homogenising only a chosen system of generators may introduce extra points at infinity. Homogenising every element of the ideal removes this ambiguity.

Exercise II.B.8. In $\mathbf{P}_{\mathbb{k}}^2$ with coordinates $[x : y : z]$, let

$$M = \begin{pmatrix} x & 0 \\ y & y \\ 0 & z \end{pmatrix}.$$

Show that the 2×2 minors of M generate the homogeneous ideal

$$(xy, xz, yz),$$

and that its zero locus consists of the three coordinate points.

More generally, let $\ell_1, \dots, \ell_4$ be four lines in $\mathbf{P}_{\mathbb{k}}^2$, no three concurrent. Show that their six pairwise intersection points are defined by the maximal minors of

$$\begin{pmatrix} \ell_1 & 0 & 0 \\ -\ell_2 & \ell_2 & 0 \\ 0 & -\ell_3 & \ell_3 \\ 0 & 0 & -\ell_4 \end{pmatrix}.$$

Solution. The minors of M are xy, xz, yz . Their common zero locus consists of points at which at most one coordinate is non-zero, hence of $[1 : 0 : 0]$, $[0 : 1 : 0]$, and $[0 : 0 : 1]$.

The four maximal minors of the second matrix are, up to signs,

$$\ell_2\ell_3\ell_4, \quad \ell_1\ell_3\ell_4, \quad \ell_1\ell_2\ell_4, \quad \ell_1\ell_2\ell_3.$$

At a common zero at least two of the ℓ_i must vanish. Conversely, every pairwise intersection of the four lines annihilates all four products. Since no three lines are concurrent, this gives exactly six points.

II.C Morphisms of projective algebraic sets

II.C.1 Rational functions on projective algebraic sets

A polynomial function of the homogeneous coordinates does not in general define a function on projective space: replacing $(x_0, \dots, x_n)$ by $(\lambda x_0, \dots, \lambda x_n)$ changes a homogeneous polynomial of degree d by the factor λ^d . Quotients of homogeneous polynomials of the same degree, however, are unchanged.

Definition II.C.1. A *rational function on $\mathbf{P}_{\mathbb{k}}^n$* is an equivalence class of fractions

$$\frac{F}{G},$$

where $F, G \in \mathbb{k}[x_0, \dots, x_n]$ are homogeneous of the same degree and $G \neq 0$. Two such fractions are equivalent when

$$FG' = F'G.$$

The fraction F/G is defined at a point $[a_0 : \dots : a_n]$ whenever $G(a_0, \dots, a_n) \neq 0$.

The equal-degree condition guarantees that

$$\frac{F(\lambda a_0, \dots, \lambda a_n)}{G(\lambda a_0, \dots, \lambda a_n)} = \frac{F(a_0, \dots, a_n)}{G(a_0, \dots, a_n)},$$

so the value does not depend on the chosen homogeneous coordinates.

More generally, let $X \subset \mathbf{P}_{\mathbb{k}}^n$ be a projective algebraic set. Two equal-degree fractions F/G and F'/G' define the same rational function on X when

$$FG' - F'G$$

vanishes identically on X . If X is irreducible, these rational functions form a field, denoted $K(X)$.

Let H be homogeneous of degree $d > 0$. On

$$D_+(H) = \mathbf{P}_{\mathbb{k}}^n \setminus \mathbf{V}_+(H)$$

we may divide by powers of H . Thus every expression

$$\frac{F}{H^m}, \quad \deg(F) = md,$$

defines a geometrically regular function on $D_+(H)$. More generally, on $X \cap D_+(H)$ we use the same expressions, considered modulo the homogeneous equations of X .

For the standard chart $U_i = D_+(x_i)$, these functions are polynomials in the ratios

$$\frac{x_0}{x_i}, \dots, \frac{\widehat{x_i}}{x_i}, \dots, \frac{x_n}{x_i}.$$

Indeed, a monomial of degree m divided by x_i^m becomes a monomial in these ratios. This recovers the usual identification

$$U_i \simeq \mathbb{k}^n.$$

Later, after localisation has been introduced, the ring of these functions will be denoted

$$\mathbb{k}[x_0, \dots, x_n][H^{-1}]_0.$$

The subscript 0 records exactly the equal-degree condition above.

II.C.2 Polynomial maps in local charts

A tuple of homogeneous polynomials of the same degree transforms in the same way under a rescaling of the homogeneous coordinates. This gives the basic form of a map between projective algebraic sets.

Definition II.C.2. Let $X \subset \mathbf{P}_{\mathbb{k}}^n$ and $Y \subset \mathbf{P}_{\mathbb{k}}^m$ be projective algebraic sets, and let $\varphi : X \rightarrow Y$ be a map. We say that φ is regular at a point $p \in X$ if there exist a homogeneous polynomial G and homogeneous polynomials

$$F_0, \dots, F_m \in \mathbb{k}[x_0, \dots, x_n]$$

of the same degree such that:

- (i) $G(p) \neq 0$,
- (ii) the polynomials $F_0, \dots, F_m$ do not vanish simultaneously at any point of $X \cap D_+(G)$;
- (iii) for every $q \in X \cap D_+(G)$, we have

$$\varphi(q) = [F_0(q) : \dots : F_m(q)].$$

The map φ is a morphism if it is regular at every point of X .

The polynomial G is auxiliary: it does not enter the formula for φ , but only specifies a distinguished subset on which one homogeneous formula $[F_0 : \dots : F_m]$ is valid. There is no further relation required between G and the F_i , except that the latter must have no common zero on $X \cap D_+(G)$. The condition $G(p) \neq 0$ is independent of the chosen homogeneous representative of p , because G is homogeneous.

Example II.C.3. Let $F_0, \dots, F_m$ be homogeneous polynomials of the same degree which do not vanish simultaneously at any point of X . If

$$[F_0(p) : \dots : F_m(p)] \in Y$$

for every $p \in X$, then

$$X \longrightarrow Y, \quad p \longmapsto [F_0(p) : \dots : F_m(p)]$$

is a morphism.

Exercise II.C.4. Show that the identity map of a projective algebraic set is a morphism and that the composition of two morphisms of projective algebraic sets is a morphism.

Solution. The identity is represented by the homogeneous coordinates $x_0, \dots, x_n$.

Let $\varphi : X \rightarrow Y$ and $\psi : Y \rightarrow Z$ be morphisms. Near a point $p \in X$, write

$$\varphi = [F_0 : \dots : F_m]$$

with the F_i homogeneous of the same degree. Near $\varphi(p)$, write

$$\psi = [G_0 : \dots : G_r],$$

where the G_j are homogeneous of the same degree. Then, after restricting to a suitable subset containing p , the composition is given by

$$\psi \circ \varphi = [G_0(F_0, \dots, F_m) : \dots : G_r(F_0, \dots, F_m)].$$

These polynomials are homogeneous of the same degree and do not vanish simultaneously on a sufficiently small distinguished subset containing p .

Example II.C.5. Let $u : \mathbb{k}^{n+1} \rightarrow \mathbb{k}^{m+1}$ be a linear map and let $X \subset \mathbf{P}_{\mathbb{k}}^n$ be a projective algebraic set such that

$$X \cap \mathbf{P}(\ker u) = \emptyset.$$

Then

$$X \longrightarrow \mathbf{P}_{\mathbb{k}}^m, \quad [v] \longmapsto [u(v)]$$

is a morphism. In particular, an injective linear map u induces a morphism $\mathbf{P}_{\mathbb{k}}^n \rightarrow \mathbf{P}_{\mathbb{k}}^m$.

Remark II.C.6. The formula

$$[x_0 : x_1 : x_2] \longmapsto [x_1 : x_2]$$

does not define a morphism on all of $\mathbf{P}_{\mathbb{k}}^2$, since both coordinates vanish at $[1 : 0 : 0]$. At every other point, however, it is given by a valid local formula in the sense of the preceding definition. Later, this will be interpreted as a rational map

$$\mathbf{P}_{\mathbb{k}}^2 \dashrightarrow \mathbf{P}_{\mathbb{k}}^1,$$

called the projection from the point $[1 : 0 : 0]$.

Example II.C.7 (The quadratic Veronese morphism). Consider the morphism

$$\nu_2 : \mathbf{P}_{\mathbb{k}}^1 \longrightarrow \mathbf{P}_{\mathbb{k}}^2, \quad [s : t] \longmapsto [s^2 : st : t^2].$$

Its image is contained in the conic

$$C = \mathbf{V}_+(x_0x_2 - x_1^2) \subset \mathbf{P}_{\mathbb{k}}^2,$$

since $s^2t^2 - (st)^2 = 0$. Conversely, let $[x_0 : x_1 : x_2] \in C$. We define a map

$$\psi : C \longrightarrow \mathbf{P}_{\mathbb{k}}^1$$

using two local formulas.

- On the subset $C \cap D_+(x_0)$, set

$$\psi([x_0 : x_1 : x_2]) = [x_0 : x_1].$$

- On the subset $C \cap D_+(x_2)$, set

$$\psi([x_0 : x_1 : x_2]) = [x_1 : x_2].$$

These two subsets cover C . Indeed, if $x_0 = x_2 = 0$, then the equation

$$x_0x_2 = x_1^2$$

implies $x_1 = 0$, which is impossible for a point of projective space.

Moreover, the two formulas agree on their intersection. Indeed, $[x_0 : x_1] = [x_1 : x_2]$ if and only if $x_0x_2 = x_1^2$, which is precisely the equation defining C . The map ψ is inverse to ν_2 . For instance, on $C \cap D_+(x_0)$, the relation $x_0x_2 = x_1^2$ gives

$$\nu_2([x_0 : x_1]) = [x_0^2 : x_0x_1 : x_1^2] = [x_0 : x_1 : x_2].$$

The computation on $C \cap D_+(x_2)$ is analogous. Thus

$$\mathbf{P}_{\mathbb{k}}^1 \simeq C.$$

This example also illustrates the role of the polynomial G in Definition II.C.2. On $C \cap D_+(x_0)$, we take

$$G = x_0, \quad (F_0, F_1) = (x_0, x_1),$$

whereas on $C \cap D_+(x_2)$, we take

$$G = x_2, \quad (F_0, F_1) = (x_1, x_2).$$

Neither pair defines the inverse morphism everywhere on C : the pair (x_0, x_1) vanishes simultaneously at $[0 : 0 : 1]$, while the pair (x_1, x_2) vanishes simultaneously at $[1 : 0 : 0]$. Hence the inverse admits homogeneous coordinate formulas locally, but neither of these formulas works globally.

In general, no single pair of homogeneous polynomials defines ψ on the whole conic. Indeed, assume that $\mathbb{k}$ is algebraically closed. Suppose that homogeneous polynomials F_0, F_1 of degree d define ψ everywhere on C . Composing with ν_2 , we obtain

$$[F_0(s^2, st, t^2) : F_1(s^2, st, t^2)] = [s : t].$$

Hence

$$F_0(s^2, st, t^2) = Hs, \quad F_1(s^2, st, t^2) = Ht$$

for some homogeneous polynomial $H \in \mathbb{k}[s, t]$ of degree $2d - 1$. Since $\mathbb{k}$ is algebraically closed and $\deg(H) > 0$, the polynomial H vanishes at some point of $\mathbf{P}_{\mathbb{k}}^1$. At the corresponding point of C , both F_0 and F_1 vanish, a contradiction.

Example II.C.8. Looking back at Example I.D.8, we see that the its projective homogenisation of the elliptic curve we say as branched cover of the affine line is the cubic curve

$$\mathbf{V}(x_2^2x_0 - x_1(x_1 - x_0)(x_1 - \lambda x_0)) \subset \mathbf{P}_{\mathbb{k}}^2.$$

The projection onto the x -coordinate extends to a degree-two map to $\mathbf{P}_{\mathbb{k}}^1$, branched also at the point at infinity. This projective curve is an elliptic curve.

Exercise II.C.9. Here is one more exercise about elementary equations of Segre and Veronese varieties.

i) Consider the Segre map

$$\sigma : \mathbf{P}_{\mathbb{k}}^1 \times \mathbf{P}_{\mathbb{k}}^1 \longrightarrow \mathbf{P}_{\mathbb{k}}^3, \quad ([s_0 : s_1], [t_0 : t_1]) \longmapsto [s_0t_0 : s_0t_1 : s_1t_0 : s_1t_1].$$

Compute the kernel of

$$\mathbb{k}[z_{00}, z_{01}, z_{10}, z_{11}] \longrightarrow \mathbb{k}[s_0, s_1, t_0, t_1], \quad z_{ij} \longmapsto s_it_j,$$

and deduce that the image is the quadric $z_{00}z_{11} - z_{01}z_{10} = 0$.

ii) Consider the quadratic Veronese map

$$\nu_2 : \mathbf{P}_{\mathbb{k}}^2 \longrightarrow \mathbf{P}_{\mathbb{k}}^5, \quad [x_0 : x_1 : x_2] \longmapsto [x_0^2 : x_0x_1 : x_0x_2 : x_1^2 : x_1x_2 : x_2^2].$$

Arrange the six target coordinates as the symmetric matrix

$$Z = \begin{pmatrix} z_{00} & z_{01} & z_{02} \\ z_{01} & z_{11} & z_{12} \\ z_{02} & z_{12} & z_{22} \end{pmatrix}.$$

Verify that every 2×2 minor of Z vanishes on the image. Show that their common projective zero locus consists precisely of symmetric matrices of rank one, hence is the image of ν_2 . The homogeneous ideal of the Veronese surface is generated by these minors.

Solution. For the Segre map, the determinant $z_{00}z_{11} - z_{01}z_{10}$ lies in the kernel. Conversely, modulo this relation every monomial can be reduced so that it does not contain both z_{00} and z_{11} ; their images are distinct monomials in the s_i, t_j , which proves that the kernel is the principal ideal generated by the determinant. A point of the quadric is a non-zero 2×2 matrix of rank one, hence an outer product $(s_0, s_1)^t(t_0, t_1)$.

For the Veronese map one has $Z = (x_i x_j)_{i,j}$, a matrix of rank one, so all 2×2 minors vanish. Conversely, a non-zero symmetric matrix of rank one is projectively of the form vv^t and therefore comes from $[v] \in \mathbf{P}_{\mathbb{k}}^2$. The standard determinantal calculation shows that the kernel of $z_{ij} \mapsto x_i x_j$ is generated by these minors.

III Localisation

Further reading. For further details on localisation, local rings, and residue fields, see [Liu02]. These constructions are used throughout [Har77, Chapter II].

We define the localisation of a module or ring at a multiplicative subset S .

III.A Localisation and inversion of elements

Definition III.A.1. A subset S of A is a *multiplicative set* if $1 \in S$ and S is closed under multiplication, i.e., given $a, b \in S$, we have $ab \in S$.

Remark III.A.2. If $\mathfrak{p}$ is a prime ideal of A , then $A \setminus \mathfrak{p}$ is multiplicative. Conversely, if $A \setminus \mathfrak{p}$ is multiplicative and $\mathfrak{p}$ is an ideal, then $\mathfrak{p}$ is prime.

Let M be an A -module and S a multiplicative subset of A . We describe the elements of the localisation $S^{-1}M$. We consider the equivalence relation

$$\frac{x}{s} \sim \frac{y}{t} \iff \exists u \in S, u(tx - sy) = 0.$$

Then the localisation is

$$S^{-1}M = \left\{ \frac{x}{s} \mid x \in M, s \in S \right\} / \sim.$$

When $M = A$, the localisation $S^{-1}A$ is a ring, with the usual operations on fractions. If $S = \{1, f, f^2, \dots\}$, we write A_f for $S^{-1}A$; if $\mathfrak{p}$ is a prime ideal and $S = A \setminus \mathfrak{p}$, we write $A_{\mathfrak{p}}$.

III.B Properties of the localisation

III.B.1 Universal property of localisation

We start by the universal property of localisation, with respect to homomorphisms of rings $A \rightarrow B$ sending each element of S to invertible elements of B .

Proposition III.B.1. *Let $\varphi : A \rightarrow B$ be a ring homomorphism such that $\varphi(s)$ is invertible in B for every $s \in S$. Then there is a unique ring homomorphism*

$$\bar{\varphi} : S^{-1}A \longrightarrow B, \quad \frac{a}{s} \longmapsto \varphi(a)\varphi(s)^{-1},$$

such that $\bar{\varphi}(a/1) = \varphi(a)$ for every $a \in A$.

Proof. The displayed formula is forced by the requirement that every $s \in S$ become invertible. It is well defined by the equivalence relation defining $S^{-1}A$, and it plainly gives the required homomorphism. $\square$

Example III.B.2. Let $A = \mathbb{k}[x]$ and let $S = \{1, x, x^2, \dots\}$. Then $S^{-1}A = \mathbb{k}[x, x^{-1}]$, the ring of Laurent polynomials. Geometrically, this corresponds to removing the point 0 from the affine line.

Example III.B.3. Let $A = \mathbb{k}[x]$ and let $\mathfrak{p} = (x)$. Then $A_{\mathfrak{p}} = \mathbb{k}[x]_{(x)}$ consists of fractions f/g with $g(0) \neq 0$. Thus we invert precisely the polynomials which do not vanish at the origin. The unique maximal ideal is generated by x .

Example III.B.4. Let $A = \mathbb{k}[x, y]/(xy)$ and localise at the multiplicative set $S = \{1, x, x^2, \dots\}$. In $S^{-1}A$, the element x becomes invertible. Since $xy = 0$, this forces $y = 0$. Hence

$$S^{-1}A \simeq \mathbb{k}[x, x^{-1}].$$

Thus, after making x invertible, the relation $xy = 0$ forces $y = 0$.

Exercise III.B.5. Let $A = \mathbb{k}[\varepsilon]/(\varepsilon^2)$.

- i) Compute A_{ε} .
- ii) Show that every element $a + b\varepsilon$ with $a \neq 0$ is a unit and deduce that $A_{(\varepsilon)} \simeq A$.
- iii) Compute $A_{1+\varepsilon}$ and explain the contrast between localising at a nilpotent element and at a unit.

Solution. Since $\varepsilon^2 = 0$, the multiplicative set generated by ε contains 0. Hence $A_{\varepsilon} = 0$. If $a \neq 0$, then

$$(a + b\varepsilon)^{-1} = a^{-1} - a^{-2}b\varepsilon,$$

so the complement of (ε) already consists of units and $A_{(\varepsilon)} = A$. Finally $1 + \varepsilon$ is a unit, so localising at it changes nothing: $A_{1+\varepsilon} \simeq A$.

III.B.2 Pull-back properties of localisation

In the following lemmas, we let S be a multiplicative subset of a ring A and put $B = S^{-1}A$. We write $\varphi : A \rightarrow B$ for the localisation map, sending $x \in A$ to $x/1 \in B$. Given an ideal $\mathfrak{a}$ of A , we set

$$S^{-1}\mathfrak{a} = \{x/s \mid x \in \mathfrak{a}, s \in S\}.$$

Lemma III.B.6. *For any ideal $\mathfrak{a}$ of A , put $\mathfrak{b} = S^{-1}\mathfrak{a}$. Then $\mathfrak{b}$ is the ideal of B generated by $\varphi(\mathfrak{a})$. Moreover:*

i) We have $S^{-1}\mathfrak{a} \neq B$ if and only if $\mathfrak{a} \cap S = \emptyset$.

ii) If $\mathfrak{a}$ is prime and $\mathfrak{a} \cap S = \emptyset$, then $\mathfrak{b}$ is prime and $\mathfrak{a} = \varphi^{-1}(\mathfrak{b})$.

Proof. Let us check that $\mathfrak{b}$ is an ideal of B . Already, $0 \in \mathfrak{b}$. Further, $\mathfrak{b}$ is an abelian subgroup of B , as for x/s and y/s in $\mathfrak{b}$, so $x, y \in \mathfrak{a}$ and $s, t \in S$, we have

$$\frac{x}{s} - \frac{y}{t} = \frac{xt - ys}{st} \in \mathfrak{b}.$$

Also, if $z/v \in B$, for some $z \in A$ and $v \in S$, we have $xz \in \mathfrak{a}$ so

$$\frac{x}{s} \cdot \frac{z}{v} = \frac{xz}{sv} \in \mathfrak{b}.$$

The ideal $\mathfrak{b}$ contains $\varphi(\mathfrak{a})$, as any element of $\varphi(\mathfrak{a})$ takes the form $x/1 \in \mathfrak{b}$ for some $x \in \mathfrak{a}$. Also, any ideal of B containing $\varphi(\mathfrak{a})$ contains all elements of the form x/s , with $x \in \mathfrak{a}$ and $s \in S$, because $x/1 \in \varphi(\mathfrak{a})$ and $x/s = x/1 \cdot 1/s$. So $\mathfrak{b}$ is the ideal of B generated by $\varphi(\mathfrak{a})$.

To prove i), assume first that $\mathfrak{a} \cap S \neq \emptyset$. Then, choose $s \in \mathfrak{a} \cap S$ and note that $s/1$ lies in $\mathfrak{b}$. Then $1 = s/1 \cdot 1/s \in \mathfrak{b}$, so $\mathfrak{b} = B$. Conversely, suppose $\mathfrak{a} \cap S = \emptyset$ and assume, by contradiction, that $\mathfrak{b} = B$. Then there are $x \in \mathfrak{a}$ and $s \in S$ such that $x/s = 1$, which means $t(x - s) = 0$ for some $t \in S$. Hence $\mathfrak{a} \ni xt = ts \in S$, which is absurd. So $\mathfrak{b} \neq B$.

Let us prove ii), so assume that $\mathfrak{a}$ is prime. To see that $\mathfrak{b}$ is prime, first recall that $\mathfrak{b} \neq B$ because $\mathfrak{a} \cap S = \emptyset$. Then, take $x, y \in A$ and $s, t \in S$ and assume $x/s \cdot y/t \in \mathfrak{b}$. So there are $z \in \mathfrak{a}$ and $v \in S$ such that $xy/st = z/v$. Then, there is $w \in S$ such that

$$w(vxy - stz) = 0.$$

So we get $wvxy = wstz \in \mathfrak{a}$. Since $wv \in S$ and $S \cap \mathfrak{a} = \emptyset$, we get $xy \in \mathfrak{a}$, because $\mathfrak{a}$ is prime. Thus $x \in \mathfrak{a}$ or $y \in \mathfrak{a}$, which says that $x/s \in \mathfrak{b}$ or $y/t \in \mathfrak{b}$.

Finally, let us prove that $\mathfrak{a} = \varphi^{-1}(\mathfrak{b})$. Given $x \in \mathfrak{a}$, we have $\varphi(x) = x/1 \in \mathfrak{b}$, so $x \in \varphi^{-1}(\mathfrak{b})$. So the inclusion $\mathfrak{a} \subset \varphi^{-1}(\mathfrak{b})$ is proved. Conversely, let $x \in \varphi^{-1}(\mathfrak{b})$, so $x/1 \in \mathfrak{b}$. Then there are $y \in \mathfrak{a}$ and $s \in S$ such that $x/1 = y/s$. Hence, for some $t \in S$, we must have $t(sx - y) = 0$. Therefore $tsx = ty \in \mathfrak{a}$, so that $x \in \mathfrak{a}$ because $ts \in S$, so $ts \notin \mathfrak{a}$, and $\mathfrak{a}$ is prime. $\square$

Lemma III.B.7. For an ideal $\mathfrak{b} \subset B$, $\mathfrak{b} = S^{-1}(\varphi^{-1}(\mathfrak{b}))$. In particular, if $\mathfrak{b} \neq B$, then $\varphi^{-1}(\mathfrak{b}) \cap S = \emptyset$.

Proof. An element of $S^{-1}(\varphi^{-1}(\mathfrak{b}))$ takes the form x/s , with $x \in \varphi^{-1}(\mathfrak{b})$ and $s \in S$. Then, $x/1 \in \mathfrak{b}$, which is an ideal of B , so $x/s = x/1 \cdot 1/s \in \mathfrak{b}$. This shows that $S^{-1}(\varphi^{-1}(\mathfrak{b})) \subset \mathfrak{b}$. Conversely, let $x/s \in \mathfrak{b}$. Then $x/1 = x/s \cdot s/1 \in \mathfrak{b}$, so that $x \in \varphi^{-1}(\mathfrak{b})$. This gives $x/s \in S^{-1}(\varphi^{-1}(\mathfrak{b}))$.

Concerning the last statement, we note that, if $\mathfrak{b} \neq B$, then $S^{-1}(\varphi^{-1}(\mathfrak{b})) = \mathfrak{b} \neq B$, so by Lemma III.B.6 we must have $\varphi^{-1}(\mathfrak{b}) \cap S = \emptyset$. $\square$

III.C Localisation and the fraction field

In the next proposition $\text{Frac}(A)$ denotes the fraction field of an integral domain A .

Exercise III.C.1. Let A be an integral domain and let $S = A \setminus \{0\}$. Show that $S^{-1}A$ is a field.

Solution. Every element of $S^{-1}A$ has the form a/s with $a \in A$ and $s \neq 0$. If $a \neq 0$, then

$$\frac{a}{s} \cdot \frac{s}{a} = 1.$$

Thus every non-zero element is invertible, so $S^{-1}A$ is a field. This field is precisely $\text{Frac}(A)$.

Proposition III.C.2. *If A is an integral domain and $0 \notin S$, then $S^{-1}A$ is an integral domain, namely the subring of $\text{Frac}(A)$ consisting of fractions with denominator in S .*

Proof. First, $0 \neq 1$ in $S^{-1}A$, since $0 \notin S$. We see that $S^{-1}A$ is an integral domain: assume $(a/s) \cdot (b/t) = 0$, then $u(ab) = 0$ for some $u \in S$, so $a = 0$ or $b = 0$. Also, set $T = A \setminus \{0\}$. By construction $T^{-1}A$ is identified with $\text{Frac}(A)$. If a/s is an element of $S^{-1}A$, we can see a/s in $\text{Frac}(A)$ and this defines an injective assignment $S^{-1}A \rightarrow \text{Frac}(A)$ which is a ring homomorphism. Indeed, if b/t represents the same element as a/s then $u(at - bs) = 0$ for some $u \in S$, hence $at = bs$ because A is an integral domain, so $a/s = b/t$ in $\text{Frac}(A)$. $\square$

III.D Local rings

Recall that a *local ring* is a ring having a unique maximal ideal. For instance, any field is a local ring, with (0) as maximal ideal. Another classical example of local ring is the ring of power series in one variable $\mathbb{k}[[T]]$ over a field $\mathbb{k}$. The maximal ideal consists of power series with vanishing constant term.

Proposition III.D.1. *Let $\mathfrak{p}$ be a prime ideal of A . Then an element $x/s \in A_{\mathfrak{p}}$ is invertible if and only if $x \notin \mathfrak{p}$. In addition, $\mathfrak{p}A_{\mathfrak{p}}$ is a maximal ideal of $A_{\mathfrak{p}}$, actually the only maximal ideal of this ring.*

Proof. Let $x \in A$ and $s \in A \setminus \mathfrak{p}$. If $x \notin \mathfrak{p}$, then s/x is an inverse of x/s in $A_{\mathfrak{p}}$.

Conversely, assume that $x \in \mathfrak{p}$ and that x/s is invertible. Then there is $y/t \in A_{\mathfrak{p}}$ such that

$$\frac{x}{s} \frac{y}{t} = 1.$$

Hence there exists $u \in A \setminus \mathfrak{p}$ such that

$$u(xy - st) = 0.$$

It follows that $ust = uxy \in \mathfrak{p}$. Since $u, s, t \notin \mathfrak{p}$ and $\mathfrak{p}$ is prime, this is impossible. Thus x/s is invertible if and only if $x \notin \mathfrak{p}$.

The non-invertible elements of $A_{\mathfrak{p}}$ are therefore exactly the elements of $\mathfrak{p}A_{\mathfrak{p}}$. This set is an ideal, so it is the unique maximal ideal of $A_{\mathfrak{p}}$. $\square$

Lemma III.D.2. *For any prime ideal $\mathfrak{p}$ of a ring A , we have $\text{Frac}(A/\mathfrak{p}) \simeq A_{\mathfrak{p}}/\mathfrak{p}A_{\mathfrak{p}}$.*

Proof. The localisation map $\varphi : A \rightarrow A_{\mathfrak{p}}$, $x \mapsto x/1$, induces a map

$$\bar{\varphi} : A/\mathfrak{p} \rightarrow A_{\mathfrak{p}}/\mathfrak{p}A_{\mathfrak{p}}.$$

This map is injective. Indeed, if $x/1 \in \mathfrak{p}A_{\mathfrak{p}}$, then there are $p \in \mathfrak{p}$ and $s \in A \setminus \mathfrak{p}$ such that $x/1 = p/s$ in $A_{\mathfrak{p}}$. Hence there exists $u \in A \setminus \mathfrak{p}$ such that

$$u(sx - p) = 0.$$

In particular, $usx \in \mathfrak{p}$. Since $u, s \notin \mathfrak{p}$ and $\mathfrak{p}$ is prime, we get $x \in \mathfrak{p}$.

Since $A_{\mathfrak{p}}/\mathfrak{p}A_{\mathfrak{p}}$ is a field and every non-zero element of $A/\mathfrak{p}$ becomes invertible in it, the map $\bar{\varphi}$ extends to an injective field homomorphism

$$\text{Frac}(A/\mathfrak{p}) \rightarrow A_{\mathfrak{p}}/\mathfrak{p}A_{\mathfrak{p}}.$$

It is surjective because every element of $A_{\mathfrak{p}}/\mathfrak{p}A_{\mathfrak{p}}$ can be written as the class of x/s , with $x \in A$ and $s \in A \setminus \mathfrak{p}$, and this is the image of $\bar{x}\bar{s}^{-1} \in \text{Frac}(A/\mathfrak{p})$. $\square$

Exercise III.D.3. Let $A = \mathbb{k}[x]$ and $S = \{1, x, x^2, \dots\}$. Show that

$$S^{-1}A \simeq \mathbb{k}[x, x^{-1}].$$

Solution. The localisation $S^{-1}A$ is obtained by making x invertible. Hence every element of $S^{-1}A$ can be written as $f(x)/x^n$, with $f(x) \in \mathbb{k}[x]$. This is exactly a Laurent polynomial, i.e. an element of $\mathbb{k}[x, x^{-1}]$.

Exercise III.D.4. Let $A = \mathbb{k}[x]$ and $\mathfrak{p} = (x - a)$, for some $a \in \mathbb{k}$. Describe the elements of $A_{\mathfrak{p}}$ and its maximal ideal.

Solution. We have

$$A_{\mathfrak{p}} = \left\{ \frac{f}{g} \mid f, g \in \mathbb{k}[x], g(a) \neq 0 \right\}.$$

Indeed, the elements outside $\mathfrak{p}$ are precisely the polynomials which do not vanish at a . The unique maximal ideal is

$$\mathfrak{p}A_{\mathfrak{p}} = (x - a)A_{\mathfrak{p}}.$$

Exercise III.D.5. Let

$$A = \mathbb{k}[x, y]/(xy)$$

and let $S = \{1, x, x^2, \dots\}$. Compute $S^{-1}A$.

Solution. In $S^{-1}A$, the element x is invertible. Since $xy = 0$ in A , we get $y = x^{-1}xy = 0$ in $S^{-1}A$. Therefore

$$S^{-1}A \simeq \mathbb{k}[x, x^{-1}].$$

IV Sheaves

Further reading. For sheaves, stalks, and sheafification, see [Har77, Chapter II, §1] and [Liu02]. The geometric role of sheaves in scheme theory is illustrated throughout [EH00].

IV.A Sheaves of abelian groups

Let X be a topological space.

Definition IV.A.1. A *presheaf of sets* on X is the data of a set $\mathcal{F}(U)$ for every open subset $U \subset X$, together with restriction maps, for each inclusion of open sets $V \subset U$:

$$\rho_{U,V}: \mathcal{F}(U) \longrightarrow \mathcal{F}(V), \quad s \longmapsto s|_V.$$

The notation $s|_V$ just means $\rho_{U,V}(s)$, for all $s \in \mathcal{F}(U)$. For every inclusion $W \subset V \subset U$, we must have $\rho_{U,U}(s) = s$ and $\rho_{V,W}(\rho_{U,V}(s))$, i.e. $s|_U = s$ and $(s|_V)|_W = s|_W$.

If for every open subset U of X , $\mathcal{F}(U)$ is a set equipped with some algebraic structure (typically an abelian group, a vector space, a ring, or a module over a ring) and for each inclusion of open subsets $V \subset U$, the restriction map $\rho_{U,V}$ is a morphism of that algebraic structure, then $\mathcal{F}$ is a presheaf with that structure. We say that $\mathcal{F}$ is a sheaf of abelian groups, of rings, etc. Note that these structures can be layered. For instance, if $\mathcal{F}$ is a sheaf of rings, then a sheaf of $\mathcal{F}$ -modules is a sheaf $\mathcal{E}$ such that each $\mathcal{E}(U)$ is an $\mathcal{F}(U)$ -module and, for each inclusion $V \subset U$, the restriction map $\mathcal{E}(U) \rightarrow \mathcal{E}(V)$ is compatible with the restriction map $\mathcal{F}(U) \rightarrow \mathcal{F}(V)$.

Definition IV.A.2. A presheaf $\mathcal{F}$ is a *sheaf* if it satisfies the following two conditions. Let $U = \bigcup_{i \in I} U_i$ be an open covering of an open subset U of X .

- i) If $s, t \in \mathcal{F}(U)$ and $s|_{U_i} = t|_{U_i}$ for all $i \in I$, then $s = t$.

ii) If, for all $i \in I$, $s_i \in \mathcal{F}(U_i)$ are sections such that

$$s_i|_{U_i \cap U_j} = s_j|_{U_i \cap U_j}, \quad \text{for all } i, j \in I,$$

then there exists a section $s \in \mathcal{F}(U)$ such that

$$s|_{U_i} = s_i, \quad \text{for all } i.$$

A sheaf is a presheaf whose sections are determined locally and glue from compatible local data.

Exercise IV.A.3. Let X be a topological space. For every open subset $U \subset X$, let

$$\mathcal{C}_X^0(U) = \{\text{continuous functions } U \rightarrow \mathbb{R}\}.$$

Show that $\mathcal{C}_X^0$ is a sheaf of abelian groups. If X is a smooth manifold, define similarly $\mathcal{C}_X^\infty(U)$ as the set of smooth functions $U \rightarrow \mathbb{R}$. Show that $\mathcal{C}_X^\infty$ is a sheaf.

Solution. Addition is pointwise, and restrictions are homomorphisms of abelian groups. If two continuous functions on U agree on every member of an open cover, they agree at every point of U . Conversely, compatible continuous functions $s_i : U_i \rightarrow \mathbb{R}$ glue to a function $s : U \rightarrow \mathbb{R}$ by setting $s(x) = s_i(x)$ for any i such that $x \in U_i$. This is well defined by compatibility, and it is continuous because continuity is local on the source. The same proof works for smooth functions on a smooth manifold, since smoothness is also local on the source.

Exercise IV.A.4. Let X be a topological space and let A be an abelian group. Define a presheaf by

$$\mathcal{F}(U) = A$$

for every non-empty open subset $U \subset X$, put $\mathcal{F}(\emptyset) = 0$, and let all restriction maps between non-empty open subsets be the identity. Is this presheaf always a sheaf? What happens if U is disconnected?

Solution. This presheaf is not always a sheaf. Suppose that $U = U_1 \sqcup U_2$ is a disjoint union of two non-empty open subsets. Choose two distinct elements $a, b \in A$. The sections $a \in \mathcal{F}(U_1)$ and $b \in \mathcal{F}(U_2)$ are automatically compatible on the empty intersection, but they cannot be glued to a single element of A on U , because its restriction to both U_1 and U_2 would have to be the same element.

On the other hand, locally constant functions with values in A do form a sheaf on X , see Example IV.D.8.

Exercise IV.A.5. Let X be a complex manifold. For every open subset $U \subset X$, let

$$\mathcal{O}_X(U) = \{\text{holomorphic functions } U \rightarrow \mathbb{C}\}.$$

Show that $\mathcal{O}_X$ is a sheaf of complex vector spaces.

Solution. The sheaf axioms are proved exactly as for continuous functions, using the fact that holomorphicity is local. Thus compatible holomorphic functions on an open cover glue to a holomorphic function.

Exercise IV.A.6. Let $U \subset \mathbb{C}$ be a connected open subset containing at least two points.

- i) Show that the ring $\mathcal{O}(U)$ of holomorphic functions has no zero divisors.
- ii) Show that the ring $\mathcal{C}^0(U)$ of continuous functions usually has zero divisors: choose two disjoint closed discs contained in U and construct non-zero continuous functions with disjoint supports.
- iii) Explain why connectedness alone does not force a ring of functions to be an integral domain.

Solution. If $fg = 0$ with $f, g \in \mathcal{O}(U)$ and $f \neq 0$, then the set on which f does not vanish is a non-empty open subset. The function g vanishes there, so the identity theorem gives $g = 0$ on the connected set U .

Choose disjoint closed discs $D_1, D_2 \subset U$. There are non-zero continuous functions f_i supported in small neighbourhoods of D_i . Their supports are disjoint, hence $f_1 f_2 = 0$. Thus $\mathcal{C}^0(U)$ has zero divisors even though U is connected. The difference is analytic rigidity, not topology alone.

IV.B Morphisms

The definitions above describe the local sections of a sheaf. We now explain how sheaves are compared and transported. These constructions will later enter the definition of a morphism of ringed spaces and, in particular, of a morphism of schemes.

Definition IV.B.1. Let $\mathcal{F}$ and $\mathcal{G}$ be presheaves on a topological space X . A *morphism of presheaves*

$$\varphi : \mathcal{F} \longrightarrow \mathcal{G}$$

is a collection of maps

$$\varphi_U : \mathcal{F}(U) \longrightarrow \mathcal{G}(U),$$

one for every open subset $U \subset X$, such that for every inclusion $V \subset U$ the diagram

$$\begin{array}{ccc} \mathcal{F}(U) & \xrightarrow{\varphi_U} & \mathcal{G}(U) \\ \downarrow & & \downarrow \\ \mathcal{F}(V) & \xrightarrow{\varphi_V} & \mathcal{G}(V) \end{array}$$

commutes. If $\mathcal{F}$ and $\mathcal{G}$ are sheaves, we call φ a morphism of sheaves.

Example IV.B.2 (The de Rham differential). Let X be a smooth manifold. For each $p \geq 0$, let $\mathcal{A}_X^p$ denote the presheaf of smooth differential p -forms. Since smooth p -forms glue over open coverings, this is a sheaf of real vector spaces. Exterior differentiation defines a morphism of sheaves of real vector spaces

$$d : \mathcal{A}_X^p \longrightarrow \mathcal{A}_X^{p+1}, \quad \omega \longmapsto d\omega,$$

and $d \circ d = 0$. For $p = 0$ this is

$$d : \mathcal{C}_X^\infty \longrightarrow \mathcal{A}_X^1, \quad f \longmapsto df.$$

It is not a morphism of $\mathcal{C}_X^\infty$ -modules, since

$$d(hf) = h df + f dh.$$

Thus, when speaking of a morphism of sheaves, it is important to specify which algebraic structure is being preserved.

Example IV.B.3 (The $\bar{\partial}$ -operator). Let X be a complex manifold and let $\mathcal{A}_X^{0,q}$ be the sheaf of smooth $(0, q)$ -forms. The operator

$$\bar{\partial} : \mathcal{A}_X^{0,q} \longrightarrow \mathcal{A}_X^{0,q+1}$$

is a morphism of sheaves of complex vector spaces and satisfies $\bar{\partial}^2 = 0$. In degree zero,

$$\mathcal{O}_X = \ker(\bar{\partial} : \mathcal{A}_X^{0,0} \rightarrow \mathcal{A}_X^{0,1}).$$

Thus holomorphic functions form the sheaf of solutions of a local differential equation.

IV.C Stalks, direct and inverse images

Here we define stalks and direct images of sheaves. These account for the local behaviour of a sheaf around a point and for the local sections on preimages of open sets with respect to a continuous map.

IV.C.1 Direct limits

Let us define direct limits of abelian groups. We start from a directed set $(I, \leq)$ and a family of abelian groups $(F_i | i \in I)$ together with morphisms $f_{i,j} : F_i \rightarrow F_j$ for all pairs $i, j \in I$ with $i \leq j$, with the properties:

- i) $f_{i,i} = \text{id}_{F_i}$, for all $i \in I$,
- ii) $f_{i,k} = f_{j,k} \circ f_{i,j}$, for all $i, j, k \in I$ with $i \leq j \leq k$.

Definition IV.C.1. We define the direct limit:

$$\varinjlim_{i \in I} F_i = \bigsqcup_{i \in I} F_i / \sim$$

Here, the equivalence relation is defined by declaring that, given $a \in F_i$ and $b \in F_j$ for some $i, j \in I$, then $a \sim b$ if there is $k \in I$ with $k \geq i$ and $k \geq j$ such that $f_{i,k}(a) = f_{j,k}(b)$. It turns out that the direct limit is also an abelian group and that there are morphisms

$$f_i : F_i \rightarrow F = \varinjlim_{i \in I} F_i, \quad \text{for all } i \in I,$$

such that

$$f_j \circ f_{i,j} = f_i, \quad \text{for all } i, j \in I \text{ with } i \leq j.$$

The construction of direct limit works for modules over a ring, or for vector spaces, and many more general categories.

The direct limit has the following universal property. Given an abelian group G , (or module over a ring, etc) and morphisms $g_i : F_i \rightarrow G$ for $i \in I$, such that $g_j \circ f_{i,j} = g_i$ for $i, j \in I$ with $i \leq j$, then there is a unique morphism $h : F \rightarrow G$ such that $h \circ f_i = g_i$ for all $i \in I$.

IV.C.2 The stalk

Let $\mathcal{F}$ be a sheaf of sets on a topological space X . Then, for all $x \in X$, we define the stalk of $\mathcal{F}$ at x as:

$$\mathcal{F}_x = \varinjlim_{U \ni x} \mathcal{F}(U).$$

Formally, this means the following. We consider the set $\mathcal{U}_x$ of open subsets of X containing x . If $U, V \in \mathcal{U}_x$ and $V \subset U$, then we have a map $\rho_{U,V} : \mathcal{F}(U) \rightarrow \mathcal{F}(V)$. This forms a directed set and the restriction maps satisfy the conditions i) and ii) So we have:

$$\mathcal{F}_x = \bigsqcup_{U \in \mathcal{U}_x} \mathcal{F}(U) / \sim,$$

where, for all $U, U' \in \mathcal{U}_x$ and for $s \in \mathcal{F}(U)$ and $t \in \mathcal{F}(U')$, we put $s \sim t$ if and only if there is $W \in \mathcal{U}_x$, with $W \subset U \cap U'$ and $\rho_{U,W}(s) = \rho_{U',W}(t)$.

Equivalently, an element of $\mathcal{F}_x$ is represented by a pair (U, s) , where U is an open neighbourhood of x and $s \in \mathcal{F}(U)$. Two pairs (U, s) and (V, t) define the same element of $\mathcal{F}_x$ if there exists an open neighbourhood $W \subset U \cap V$ of x such that

$$s|_W = t|_W.$$

The class of (U, s) in $\mathcal{F}_x$ is called the *germ* of s at x and is denoted by s_x .

Remark IV.C.2. A morphism $\varphi : \mathcal{F} \rightarrow \mathcal{G}$ induces, for every $x \in X$, a map on stalks

$$\varphi_x : \mathcal{F}_x \longrightarrow \mathcal{G}_x, \quad s_x \longmapsto \varphi_U(s)_x,$$

where s is any representative of the germ s_x on a neighbourhood U of x . Compatibility with restriction makes this map well defined.

Exercise IV.C.3. Prove that this is well-defined and that for all open subsets $U \subset X$ containing x we get an induced map $\mathcal{F}(U) \rightarrow \mathcal{F}_x$.

Solution. The relation is reflexive and symmetric by definition. For transitivity, assume that (U, s) is equivalent to (V, t) and that (V, t) is equivalent to (W, r) . Then there are neighbourhoods $U_1 \subset U \cap V$ and $U_2 \subset V \cap W$ of x such that $s|_{U_1} = t|_{U_1}$ and $t|_{U_2} = r|_{U_2}$. On $U_1 \cap U_2$, all three restrictions agree, so (U, s) is equivalent to (W, r) .

If U is an open neighbourhood of x , the map $\mathcal{F}(U) \rightarrow \mathcal{F}_x$ sends s to the class of (U, s) . It is compatible with restriction: if $V \subset U$ is another neighbourhood of x , then (U, s) and $(V, s|_V)$ define the same germ.

Example IV.C.4. Every germ of a holomorphic function at the origin has a convergent power series expansion, and two germs are equal if and only if their power series are equal. Thus

$$\mathcal{O}_{\mathbb{C},0} \simeq \mathbb{C}\{z\},$$

the ring of convergent power series. This ring is local, and its unique maximal ideal is

$$\mathfrak{m}_0 = \{f_0 \mid f(0) = 0\} = (z),$$

and a germ is invertible precisely when its constant term is non-zero.

Note that the holomorphic functions $f(z) = 0$ and $g(z) = z$ have the same value at the origin, but they do not define the same germ there. Indeed, they do not agree on any neighbourhood of 0. The stalk $\mathcal{O}_{\mathbb{C},0}$ therefore remembers the behaviour of a function near 0, not merely its value at 0.

Example IV.C.5 (Different representatives of the same germ). At the origin of $\mathbb{C}$, consider

$$f(z) = \frac{1}{1-z}$$

on $\mathbb{C} \setminus \{1\}$ and

$$g(z) = \sum_{n \geq 0} z^n$$

on the unit disc. The two functions agree on a neighbourhood of 0, hence they define the same element of the stalk $\mathcal{O}_{\mathbb{C},0}$, although their domains and their descriptions are different.

Remark IV.C.6 (A contrast with smooth germs). The smooth function

$$h(x) = \begin{cases} e^{-1/x^2}, & x \neq 0, \\ 0, & x = 0, \end{cases}$$

has zero Taylor series at the origin, but its germ is not zero. Holomorphic germs are determined by their Taylor series; smooth germs are not.

Exercise IV.C.7. In $\mathcal{O}_{\mathbb{C},0}$, determine which of the following germs are invertible:

$$z, \quad 1+z, \quad e^z - 1, \quad \cos z.$$

For each non-invertible germ, determine its order of vanishing.

Solution. A holomorphic germ is invertible exactly when its value at 0 is non-zero. Thus $1+z$ and $\cos z$ are invertible. The germs z and $e^z - 1$ are not; both have order of vanishing one, since

$$e^z - 1 = z + \frac{z^2}{2!} + \cdots$$

IV.D Defining a sheaf from compatible data

IV.D.1 The sheafification

There is a way to describe the sheaf associated with a presheaf, which goes through looking at the stalks of the presheaf at each point.

Lemma IV.D.1. *Let $\mathcal{F}$ be a presheaf of abelian groups on a topological space X . Then there is a sheaf $\mathcal{F}^+$ and a morphism $\mathcal{F} \rightarrow \mathcal{F}^+$ which is universal with respect to morphisms from $\mathcal{F}$ to sheaves.*

Proof. The definition of $\mathcal{F}^+$ goes as follows. Let U be an open subset of X . Then we consider the set of functions

$$s : U \rightarrow \prod_{x \in U} \mathcal{F}_x$$

such that $s(x) \in \mathcal{F}_x$, for all $x \in U$. Among these functions, we consider those functions s such that, for all $x \in U$, there is an open subset $V \subset U$, with $x \in V$, and there is $t \in \mathcal{F}(V)$ such that, for any point $y \in V$, the canonical germ map $\mathcal{F}(V) \rightarrow \mathcal{F}_y$ takes t to $s(y)$.

This defines the set $\mathcal{F}^+(U)$ and we get a morphism $\mathcal{F} \rightarrow \mathcal{F}^+$ by sending $s \in \mathcal{F}(U)$ to its local values $(s_x)_{x \in U}$, where s_x is the image of s in $\mathcal{F}_x$. From the very definition of $\mathcal{F}^+(U)$, the image of s is in $\mathcal{F}^+(U)$. The resulting map is universal with respect to morphisms $\psi : \mathcal{F} \rightarrow \mathcal{G}$ of presheaves of abelian groups, where $\mathcal{G}$ is a sheaf. Indeed, we first observe that $\mathcal{F}_x^+ \simeq \mathcal{F}_x$, for all $x \in X$. Then, one checks that, given ψ and an open subset $U \subset X$, we get maps $\psi : \mathcal{F}_x \rightarrow \mathcal{G}_x$ and an induced map $\mathcal{F}_x^+ \rightarrow \mathcal{G}_x$, which gives a map $\mathcal{F}^+(U) \rightarrow \mathcal{G}(U)$, which is uniquely defined and commutes with our morphism $\mathcal{F} \rightarrow \mathcal{F}^+$. $\square$

Exercise IV.D.2. Fill in the details of the above proof.

Solution. The restrictions on $\mathcal{F}^+$ are obtained by restricting functions $U \rightarrow \prod_{x \in U} \mathcal{F}_x$. The locality condition is local by definition, so these restrictions are well defined. The sheaf axioms are also immediate from the description by functions: if two sections have the same restrictions to an open cover, then they have the same value in each stalk; and compatible local sections glue by defining the value at each point using any member of the cover containing that point.

The morphism $\mathcal{F} \rightarrow \mathcal{F}^+$ sends a section to its germs. If $\mathcal{G}$ is a sheaf and $\psi : \mathcal{F} \rightarrow \mathcal{G}$ is a morphism of presheaves, then ψ induces maps on stalks $\psi_x : \mathcal{F}_x \rightarrow \mathcal{G}_x$. Given $s \in \mathcal{F}^+(U)$, define a section of $\mathcal{G}$ locally as follows: near every $x \in U$, the section s is represented by some $t \in \mathcal{F}(V)$, and we take $\psi(t) \in \mathcal{G}(V)$. These local sections agree on overlaps because they have the same germs and $\mathcal{G}$ is a sheaf. Hence they glue to a unique section of $\mathcal{G}(U)$. This gives the unique morphism $\mathcal{F}^+ \rightarrow \mathcal{G}$ extending ψ .

IV.D.2 The étalé space of a presheaf

Let $\mathcal{F}$ be a presheaf of sets on a topological space X . The étalé space of $\mathcal{F}$ is the disjoint union of stalks

$$E(\mathcal{F}) := \coprod_{x \in X} \mathcal{F}_x.$$

The étalé space is equipped with the natural projection $\pi : E(\mathcal{F}) \rightarrow X$, defined by $\xi \in \mathcal{F}_x \mapsto x$. Every section $s \in \mathcal{F}(U)$ defines a map

$$\tilde{s} : U \rightarrow E(\mathcal{F}), \quad x \mapsto s_x.$$

Lemma IV.D.3. *The subsets of the form $\tilde{s}(U) = \{s_x \mid x \in U\}$, with $U \subset X$ open and $s \in \mathcal{F}(U)$, for a basis of a topology on $E(\mathcal{F})$.*

Proof. To see this, consider U, V open subsets of X and let $s \in \mathcal{F}(U)$ and $t \in \mathcal{F}(V)$. Take $\xi \in \tilde{s}(U) \cap \tilde{t}(V)$. Writing $x = \pi(\xi)$, we have $s_x = t_x$. By the definition of equality of germs, there is an open neighbourhood W of x , with $W \subset U \cap V$ together with $v \in \mathcal{F}(W)$, such that $s|_W = v = t|_W$. It follows that

$$\xi \in \tilde{v}(W) \subset \tilde{s}(U) \cap \tilde{t}(V).$$

Therefore ξ has a neighborhood in $E(\mathcal{F})$ consisting of an open subset from our basis, contained in $\tilde{s}(U) \cap \tilde{t}(V)$. Hence the subsets above do form a basis. $\square$

Definition IV.D.4. The étalé space of $\mathcal{F}$ is the disjoint union of stalks $E(\mathcal{F}) := \coprod_{x \in X} \mathcal{F}_x$, equipped with the topology defined by Lemma IV.D.3.

Remark IV.D.5. For every such basic open subset, the restriction

$$\pi|_{\tilde{s}(U)} : \tilde{s}(U) \longrightarrow U$$

is a homeomorphism, with inverse $\tilde{s}$. Hence π is a local homeomorphism.

Remark IV.D.6. The space $E(\mathcal{F})$ should be thought of as the space of all possible local values of sections of $\mathcal{F}$. A section of π over an open subset $U \subset X$ is a map

$$\sigma : U \longrightarrow E(\mathcal{F})$$

such that

$$\pi \circ \sigma = \text{id}_U.$$

Thus σ chooses, for every $x \in U$, a germ $\sigma(x) \in \mathcal{F}_x$. Continuity means precisely that this choice is locally represented by a section of the original presheaf.

Remark IV.D.7. This resembles the usual notion of a section of a vector bundle $p : E \rightarrow X$: in both cases, a section chooses a point in the fibre over every point of the base.

There is nevertheless an important difference. For a vector bundle, the fibre E_x is a finite-dimensional vector space. For the étalé space $E(\mathcal{F})$, the fibre over x is the stalk $\mathcal{F}_x$, whose elements are germs of local sections. Moreover, $\pi : E(\mathcal{F}) \rightarrow X$ is a local homeomorphism, whereas the projection of a vector bundle of positive rank is not. In particular, if $\mathcal{E}$ is the sheaf of sections of a vector bundle $E \rightarrow X$, the étalé space $E(\mathcal{E})$ is not the total space E : its fibre at x is the space $\mathcal{E}_x$ of germs of sections near x , rather than the vector-space fibre E_x .

Example IV.D.8. Let X be a topological space and let A be an abelian group. Consider the constant presheaf $\mathcal{F}$ defined by

$$\mathcal{F}(U) = A$$

for every non-empty open subset $U \subset X$, with identity restriction maps, and by $\mathcal{F}(\emptyset) = 0$. For every $x \in X$, the stalk is naturally isomorphic to A , $\mathcal{F}_x \simeq A$. Therefore, as a set over X , the étalé space is

$$E(\mathcal{F}) = \coprod_{x \in X} A \simeq X \times A.$$

Here A is equipped with the discrete topology. Indeed, the basic open subset associated with an element $a \in \mathcal{F}(U) = A$ is

$$\tilde{a}(U) = U \times \{a\}.$$

A section of the projection $\pi : X \times A \rightarrow X$ over an open subset $U \subset X$ is of the form

$$x \longmapsto (x, f(x))$$

for a function $f: U \rightarrow A$. This section is continuous if and only if f is locally constant. Consequently,

$$\mathcal{F}^+(U) = \{\text{locally constant functions } U \rightarrow A\}.$$

Thus the sheafification of the constant presheaf is the sheaf of locally constant A -valued functions. In particular, sheafification allows a section to take different constant values on different connected components. This is precisely the gluing which was missing from the constant presheaf.

IV.D.3 Sheaves on a basis of the topology

Let X be a topological space and let $\mathcal{B}$ be a basis of its topology. Let us assume that the basis $\mathcal{B}$ is stable under finite intersections. In several situations, it is possible to prescribe sections only on the open sets belonging to $\mathcal{B}$, in a compatible way, and this is enough to define a sheaf on all open sets. More precisely, we say that $\mathcal{F}$ *satisfies sheaf condition on $\mathcal{B}$* , if for every $U \in \mathcal{B}$, we are given a set $\mathcal{F}(U)$ and, for every inclusion $V \subset U$ of elements of $\mathcal{B}$, a restriction map

$$\mathcal{F}(U) \longrightarrow \mathcal{F}(V), \quad s \longmapsto s|_V.$$

The first assumption we need is that, for all $W \subset V \subset U$ in $\mathcal{B}$, these maps satisfy

$$s|_U = s, \quad (s|_V)|_W = s|_W$$

The *sheaf condition on $\mathcal{B}$* requires that the sheaf axioms hold for coverings of elements of $\mathcal{B}$ by elements of $\mathcal{B}$, namely for all covering

$$U = \bigcup_{i \in I} U_i, \quad U, U_i \in \mathcal{B},$$

the following properties hold.

- i) If $s, t \in \mathcal{F}(U)$ satisfy $s|_{U_i} = t|_{U_i}$ for every $i \in I$, then $s = t$.
- ii) If $s_i \in \mathcal{F}(U_i)$ satisfy $s_i|_{U_i \cap U_j} = s_j|_{U_i \cap U_j}$ for all $i, j \in I$, then there exists a unique $s \in \mathcal{F}(U)$ such that $s|_{U_i} = s_i$ for every $i \in I$.

Proposition IV.D.9 (Defining a sheaf on a basis). *Assume $\mathcal{F}$ satisfies the sheaf condition on $\mathcal{B}$. Then there is a sheaf on X , unique up to unique isomorphism, whose sections and restriction maps on the elements of $\mathcal{B}$ are the ones given above.*

Remark IV.D.10. It is enough to check compatibility on basis elements contained in intersections: every point of $B \cap B'$ has such a neighbourhood. This proposition will later be applied to the principal opens $D(f)$ of $\text{Spec}(A)$, with sections given by localisations A_f or F_f .

Proof. We first prove uniqueness. Let $\mathcal{G}$ be a sheaf on X such that, for every $V \in \mathcal{B}$,

$$\mathcal{G}(V) \simeq \mathcal{F}(V)$$

the isomorphisms being compatible with the restriction maps induced by open subsets of $\mathcal{B}$.

Let $U \subset X$ be an arbitrary open subset. Since $\mathcal{B}$ is a basis, the elements of $\mathcal{B}$ contained in U form an open covering of U . Having this in mind, for an arbitrary open subset $U \subset X$, we define

$$\widehat{\mathcal{F}}(U) = \left\{ (s_V)_{\substack{V \in \mathcal{B} \\ V \subset U}} \mid \begin{array}{l} s_V \in \mathcal{F}(V), \\ s_V|_{V \cap W} = s_W|_{V \cap W} \text{ for all } V, W \in \mathcal{B} \text{ contained in } U \end{array} \right\}.$$

The compatibility condition makes sense because $\mathcal{B}$ is stable under finite intersections. If $U' \subset U$ are open subsets, define the restriction map

$$\widehat{\mathcal{F}}(U) \longrightarrow \widehat{\mathcal{F}}(U'), \quad (s_V)_{\substack{V \in \mathcal{B} \\ V \subset U}} \longmapsto (s_V)_{\substack{V \in \mathcal{B} \\ V \subset U'}}.$$

In other words, the restriction map one simply forgets the components indexed by basis elements which are not contained in U' . These maps make $\widehat{\mathcal{F}}$ into a presheaf on X .

If $\mathcal{G}$ is a sheaf as above, we have, for any open subset U of X , an identification

$$\mathcal{G}(U) \simeq \widehat{\mathcal{F}}(U),$$

which sends $s \in \mathcal{G}(U)$ to $(s|_V)_V$, where $V \in \mathcal{B}$ and $V \subset U$. This is a bijection since, by the sheaf axiom, a section of $\mathcal{G}(U)$ is uniquely determined by its restrictions to these basis elements, and every compatible family of such sections glues uniquely. Thus, if an extension exists, it is uniquely determined (up to a unique isomorphism).

We check that $\widehat{\mathcal{F}}$ agrees with the given presheaf on $\mathcal{B}$. Let $U \in \mathcal{B}$. There is a natural map

$$\alpha_U: \mathcal{F}(U) \longrightarrow \widehat{\mathcal{F}}(U), \quad s \longmapsto (s|_V)_{\substack{V \in \mathcal{B} \\ V \subset U}}.$$

Conversely, define

$$\beta_U: \widehat{\mathcal{F}}(U) \longrightarrow \mathcal{F}(U), \quad (s_V)_{\substack{V \in \mathcal{B} \\ V \subset U}} \longmapsto s_U.$$

Clearly,

$$\beta_U \circ \alpha_U = \text{id}_{\mathcal{F}(U)}.$$

Moreover, if $(s_V) \in \widehat{\mathcal{F}}(U)$ and $V \in \mathcal{B}$ is contained in U , then compatibility applied to U and V gives $s_U|_V = s_V$, because $U \cap V = V$. Hence

$$\alpha_U \circ \beta_U = \text{id}_{\widehat{\mathcal{F}}(U)}.$$

Thus α_U is a bijection, compatible with restriction maps.

We now prove that $\widehat{\mathcal{F}}$ is a sheaf. Let $(U_i \mid i \in I)$ be an arbitrary open covering of an open subset U of X . We first prove locality. Let $s, t \in \widehat{\mathcal{F}}(U)$ and suppose that $s|_{U_i} = t|_{U_i}$ for every $i \in I$. Write

$$s = (s_V)_{\substack{V \in \mathcal{B} \\ V \subset U}}, \quad t = (t_V)_{\substack{V \in \mathcal{B} \\ V \subset U}}.$$

Fix $V \in \mathcal{B}$ contained in U . The basis elements $W \in \mathcal{B}$ such that

$$W \subset V \cap U_i$$

for some $i \in I$ form a covering of V . On every such W , we have

$$s_V|_W = s_W = t_W = t_V|_W.$$

By the locality axiom for coverings of basis elements, we obtain

$$s_V = t_V.$$

This holds for every $V \in \mathcal{B}$ contained in U , and therefore $s = t$.

We now prove gluing. Suppose that, for every $i \in I$, we are given $s_i \in \widehat{\mathcal{F}}(U_i)$, and that $s_i|_{U_i \cap U_j} = s_j|_{U_i \cap U_j}$ for all $i, j \in I$. Fix $V \in \mathcal{B}$ contained in U . Consider all basis elements $W \in \mathcal{B}$ such that

$$W \subset V \cap U_i$$

for some $i \in I$. These basis elements cover V . For such a W , choose an index $i \in I$ such that $W \subset U_i$, and let

$$t_W \in \mathcal{F}(W)$$

be the W -component of s_i . This does not depend on the choice of $i \in I$. Indeed, if also $W \subset U_j$, for some $j \in I$ then the equality of s_i and s_j on $U_i \cap U_j$ implies that their W -components coincide.

The sections t_W are compatible on pairwise intersections. Hence, by the gluing axiom for coverings by basis elements, there exists a unique section $s_V \in \mathcal{F}(V)$ such that $s_V|_W = t_W$ for every such W . We claim that the family

$$(s_V)_{\substack{V \in \mathcal{B} \\ V \subset U}}$$

is compatible. Let $V, V' \in \mathcal{B}$ be contained in U . The basis elements $W \in \mathcal{B}$ satisfying

$$W \subset V \cap V' \cap U_i$$

for some $i \in I$ cover $V \cap V'$. On every such W , the sections $s_V|_{V \cap V'}$ and $s_{V'}|_{V \cap V'}$ have the same restriction. By locality on the basis,

$$s_V|_{V \cap V'} = s_{V'}|_{V \cap V'}.$$

In conclusion, the family (s_V) defines a section

$$s \in \widehat{\mathcal{F}}(U).$$

By construction, $s|_{U_i} = s_i$ for every $i \in I$. The uniqueness of the section we just constructed is a consequence of locality of $\mathcal{F}$ on the basis $\mathcal{B}$. Therefore $\widehat{\mathcal{F}}$ is a sheaf. $\square$

Proposition IV.D.11 (Defining a morphism on a basis). *Let X be a topological space and let $\mathcal{B}$ be a basis of its topology, stable under finite intersections. Let $\mathcal{F}$ and $\mathcal{G}$ be sheaves on X . Suppose that, for every $V \in \mathcal{B}$, a map $\varphi_V : \mathcal{F}(V) \rightarrow \mathcal{G}(V)$ is given, and that these maps are compatible with restrictions. Then there exists a unique morphism of sheaves*

$$\varphi : \mathcal{F} \rightarrow \mathcal{G}$$

whose map on sections over every $V \in \mathcal{B}$ is φ_V . If $\mathcal{F}$ and $\mathcal{G}$ are sheaves of abelian groups, rings, or modules, and the maps φ_V preserve the corresponding algebraic structure, then so does φ .

Proof. Let $U \subset X$ be an arbitrary open subset and let $s \in \mathcal{F}(U)$. For every basis element $V \in \mathcal{B}$ contained in U , as in the previous proof, we define

$$t_V = \varphi_V(s|_V) \in \mathcal{G}(V).$$

We claim that these sections are compatible. Indeed, let $V, W \in \mathcal{B}$ be contained in U . Since $\mathcal{B}$ is stable under finite intersections, $V \cap W$ belongs to $\mathcal{B}$. Compatibility with restrictions gives

$$t_V|_{V \cap W} = \varphi_{V \cap W}(s|_{V \cap W}) = t_W|_{V \cap W}.$$

Since the basis elements contained in U cover U , the sheaf axiom for $\mathcal{G}$ gives a unique section $t \in \mathcal{G}(U)$ such that $t|_V = t_V$ for every $V \in \mathcal{B}$ contained in U . We define $\varphi_U(s) = t$.

We check that these maps commute with restrictions. Let $U' \subset U$. The two sections

$$\varphi_U(s)|_{U'} \quad \text{and} \quad \varphi_{U'}(s|_{U'})$$

have the same restriction to every basis element $V \in \mathcal{B}$ contained in U' , namely $\varphi_V(s|_V)$. By the locality axiom for $\mathcal{G}$, they are equal. Hence the maps φ_U define a morphism of sheaves

$$\varphi : \mathcal{F} \longrightarrow \mathcal{G}.$$

If the maps φ_V preserve an algebraic structure, then so do the maps φ_U . Indeed, the relevant identities may be checked after restriction to the basis elements contained in U , where they follow from the corresponding identities for φ_V .

Finally, the morphism is unique. If $\psi : \mathcal{F} \longrightarrow \mathcal{G}$ is another morphism whose maps on the elements of $\mathcal{B}$ are the φ_V , then, for every $s \in \mathcal{F}(U)$, the sections $\varphi_U(s)$ and $\psi_U(s)$ have the same restrictions to all basis elements contained in U . The locality axiom gives $\varphi_U(s) = \psi_U(s)$. Thus $\varphi = \psi$. $\square$

Exercise IV.D.12 (Transition functions of a vector bundle). Let X be a smooth manifold, respectively a complex manifold, and let $p : E \rightarrow X$ be a smooth, respectively holomorphic, vector bundle of rank r . Let $(U_i \mid i \in I)$ be a trivialising open covering of X for E , and choose trivialisations

$$\tau_i : E|_{U_i} \xrightarrow{\cong} U_i \times \mathbb{K}^r,$$

where $\mathbb{K} = \mathbb{R}$ in the smooth case and $\mathbb{K} = \mathbb{C}$ in the holomorphic case.

i) Show that, on U_{ij} , the map

$$\tau_i \circ \tau_j^{-1} : U_{ij} \times \mathbb{K}^r \longrightarrow U_{ij} \times \mathbb{K}^r$$

has the form

$$(x, v) \longmapsto (x, g_{ij}(x)v),$$

where

$$g_{ij} : U_{ij} \longrightarrow \mathrm{GL}_r(\mathbb{K})$$

is smooth, respectively holomorphic.

ii) Show that the transition functions satisfy

$$g_{ii} = \mathbf{1}_r, \quad g_{ji} = g_{ij}^{-1}, \quad g_{ij}g_{jk} = g_{ik}$$

on every triple intersection U_{ijk} .

iii) Let $\mathcal{E}_i$ be the sheaf of smooth, respectively holomorphic, maps $U \longrightarrow \mathbb{K}^r$ on open subsets $U \subset U_i$. Show that multiplication by g_{ij} defines an isomorphism of sheaves

$$\varphi_{ij} : \mathcal{E}_j|_{U_{ij}} \xrightarrow{\cong} \mathcal{E}_i|_{U_{ij}}, \quad s \longmapsto g_{ij}s,$$

and that the cocycle condition for the φ_{ij} is precisely the identity

$$g_{ij}g_{jk} = g_{ik}.$$

iv) Conversely, suppose that smooth, respectively holomorphic, maps $g_{ij} : U_{ij} \longrightarrow \mathrm{GL}_r(\mathbb{K})$ are given and satisfy the cocycle condition. Show that the trivial bundles $U_i \times \mathbb{K}^r$ can be glued to a smooth, respectively holomorphic, vector bundle of rank r on X .

Solution. On U_{ij} , the map $\tau_i \circ \tau_j^{-1}$ preserves the projection to U_{ij} and is linear on each fibre. Hence it is of the form

$$(x, v) \mapsto (x, g_{ij}(x)v)$$

for a map $g_{ij} : U_{ij} \rightarrow \mathrm{GL}_r(\mathbb{K})$. Its regularity follows from that of the trivialisations. We have

$$g_{ii} = \mathbf{1}_r$$

because $\tau_i \circ \tau_i^{-1}$ is the identity, and

$$g_{ji} = g_{ij}^{-1}$$

because $\tau_j \circ \tau_i^{-1}$ is the inverse of $\tau_i \circ \tau_j^{-1}$. On a triple intersection,

$$(\tau_i \circ \tau_j^{-1}) \circ (\tau_j \circ \tau_k^{-1}) = \tau_i \circ \tau_k^{-1},$$

and therefore

$$g_{ij}g_{jk} = g_{ik}.$$

If $s : U \rightarrow \mathbb{K}^r$ is a local section on $U \subset U_{ij}$, define

$$\varphi_{ij}(s)(x) = g_{ij}(x)s(x).$$

This gives an isomorphism of sheaves, with inverse given by multiplication by g_{ji} . Moreover,

$$\varphi_{ij} \circ \varphi_{jk}(s) = g_{ij}g_{jk}s,$$

so the cocycle condition for the sheaves is equivalent to

$$g_{ij}g_{jk} = g_{ik}.$$

Conversely, define an equivalence relation on the disjoint union

$$\coprod_{i \in I} U_i \times \mathbb{K}^r$$

by declaring

$$(x, v)_j \sim (x, g_{ij}(x)v)_i$$

whenever $x \in U_{ij}$. The cocycle condition shows that this is an equivalence relation. The quotient carries natural local trivialisations over the U_i , and these endow it with the structure of a smooth, respectively holomorphic, vector bundle of rank r .

Exercise IV.D.13. Let $\mathbf{P}_{\mathbb{C}}^1 = \mathbb{C} \cup \{\infty\}$, and let z denote the standard affine coordinate on $U_0 = \mathbf{P}_{\mathbb{C}}^1 \setminus \{\infty\} \simeq \mathbb{C}$. For $d \in \mathbb{Z}$, define a presheaf $\mathcal{O}_{\mathbf{P}_{\mathbb{C}}^1}(d[\infty])$ over an open subset $U \subset \mathbf{P}_{\mathbb{C}}^1$ as the set of meromorphic functions f on U , whose divisor of poles is bounded by $d[\infty]$. In other words, $\mathcal{O}_{\mathbf{P}_{\mathbb{C}}^1}(d[\infty])(U)$ is the set of meromorphic functions f on U such that

$$\mathrm{ord}_p(f) \geq 0 \quad \text{for every } p \in U \setminus \{\infty\},$$

and, if $\infty \in U$,

$$\mathrm{ord}_{\infty}(f) \geq -d.$$

- i) Check directly that $U \mapsto \mathcal{O}_{\mathbf{P}_{\mathbb{C}}^1}(d[\infty])(U)$ is a sheaf of $\mathcal{O}_{\mathbf{P}_{\mathbb{C}}^1}$ -modules.

ii) Prove that, for $d \geq 0$,

$$\Gamma(\mathbf{P}_{\mathbb{C}}^1, \mathcal{O}_{\mathbf{P}_{\mathbb{C}}^1}(d[\infty])) = \{a_0 + a_1z + \cdots + a_dz^d \mid a_i \in \mathbb{C}\}.$$

Deduce that $\dim_{\mathbb{C}} \Gamma(\mathbf{P}_{\mathbb{C}}^1, \mathcal{O}_{\mathbf{P}_{\mathbb{C}}^1}(d[\infty])) = d + 1$.

iii) Show that $\Gamma(\mathbf{P}_{\mathbb{C}}^1, \mathcal{O}_{\mathbf{P}_{\mathbb{C}}^1}(d[\infty])) = 0$ when $d < 0$.

iv) Let $U_{\infty} = \mathbf{P}_{\mathbb{C}}^1 \setminus \{0\}$, and let $w = \frac{1}{z}$ be the standard coordinate near ∞ . Show that the restrictions of $\mathcal{O}_{\mathbf{P}_{\mathbb{C}}^1}(d[\infty])$ to U_0 and U_{∞} are trivial. More precisely, show that $e_0 = 1$ is a generator on U_0 , while $e_{\infty} = w^{-d}$ is a generator on U_{∞} . Check that, on $U_0 \cap U_{\infty}$, $e_{\infty} = z^d e_0$. Conclude that $\mathcal{O}_{\mathbf{P}_{\mathbb{C}}^1}(d[\infty])$ is obtained by gluing two trivial line bundles with transition function z^d .

Solution. i) The condition on the orders of zeros and poles is local. Indeed, a meromorphic function belongs to $\mathcal{O}_{\mathbf{P}_{\mathbb{C}}^1}(d[\infty])(U)$ if and only if it satisfies the required inequalities in a neighbourhood of every point of U .

Suppose that $U = \bigcup_i U_i$ and that

$$f_i \in \mathcal{O}_{\mathbf{P}_{\mathbb{C}}^1}(d[\infty])(U_i)$$

are sections agreeing on all intersections $U_i \cap U_j$. Since meromorphic functions form a sheaf, the f_i glue uniquely to a meromorphic function f on U . The inequalities on the orders of f hold locally, because they hold for every f_i . Hence

$$f \in \mathcal{O}_{\mathbf{P}_{\mathbb{C}}^1}(d[\infty])(U).$$

Thus the assignment is a sheaf.

It is also a sheaf of $\mathcal{O}_{\mathbf{P}_{\mathbb{C}}^1}$ -modules. Indeed, if f is a local section and g is holomorphic, then

$$\text{ord}_p(gf) = \text{ord}_p(g) + \text{ord}_p(f) \geq \text{ord}_p(f),$$

since $\text{ord}_p(g) \geq 0$. Hence multiplication by holomorphic functions preserves the prescribed pole bound.

ii) Let

$$f \in \Gamma(\mathbf{P}_{\mathbb{C}}^1, \mathcal{O}_{\mathbf{P}_{\mathbb{C}}^1}(d[\infty])).$$

The function f has no poles on the affine chart

$$U_0 \simeq \mathbb{C}.$$

Therefore $f|_{U_0}$ is an entire function. Moreover, f is meromorphic at ∞ , so $f|_{U_0}$ is a rational function. A rational function with no finite poles is a polynomial. Thus

$$f(z) = a_0 + a_1z + \cdots + a_mz^m$$

for some $m \geq 0$.

Near ∞ , write $w = 1/z$. Then

$$f = a_0 + a_1w^{-1} + \cdots + a_mw^{-m}.$$

If $a_m \neq 0$, the pole order of f at ∞ is m . Equivalently,

$$\text{ord}_{\infty}(f) = -m.$$

The condition

$$\text{ord}_\infty(f) \geq -d$$

is therefore equivalent to $m \leq d$. Consequently,

$$\Gamma(\mathbf{P}_{\mathbb{C}}^1, \mathcal{O}_{\mathbf{P}_{\mathbb{C}}^1}(d[\infty])) = \{a_0 + a_1 z + \cdots + a_d z^d \mid a_i \in \mathbb{C}\}.$$

This vector space has basis

$$1, z, \dots, z^d,$$

and hence dimension $d + 1$.

iii) Suppose that $d < 0$, and let f be a global section. Since

$$\text{ord}_\infty(f) \geq -d > 0,$$

the function f is holomorphic at ∞ and vanishes there. It is also holomorphic at every finite point. Therefore f is a global holomorphic function on $\mathbf{P}_{\mathbb{C}}^1$.

Every global holomorphic function on $\mathbf{P}_{\mathbb{C}}^1$ is constant. Since $f(\infty) = 0$, this constant is zero. Hence

$$\Gamma(\mathbf{P}_{\mathbb{C}}^1, \mathcal{O}_{\mathbf{P}_{\mathbb{C}}^1}(d[\infty])) = 0.$$

iv) Since $\infty \notin U_0$, sections of $\mathcal{O}_{\mathbf{P}_{\mathbb{C}}^1}(d[\infty])$ on an open subset of U_0 are simply holomorphic functions. Thus

$$\mathcal{O}_{\mathbf{P}_{\mathbb{C}}^1}(d[\infty])|_{U_0} \simeq \mathcal{O}_{U_0},$$

with generator

$$e_0 = 1.$$

On U_∞ , set

$$e_\infty = w^{-d}.$$

Let $V \subseteq U_\infty$ be open and let

$$f \in \mathcal{O}_{\mathbf{P}_{\mathbb{C}}^1}(d[\infty])(V).$$

The condition at ∞ is

$$\text{ord}_\infty(f) \geq -d.$$

Since

$$\text{ord}_\infty(w^d) = d,$$

we obtain

$$\text{ord}_\infty(w^d f) \geq 0.$$

Hence $w^d f$ is holomorphic at ∞ . It is also holomorphic away from ∞ , so there exists a unique holomorphic function $g \in \mathcal{O}_{U_\infty}(V)$ such that

$$f = g w^{-d} = g e_\infty.$$

Therefore e_∞ generates the restriction to U_∞ , and

$$\mathcal{O}_{\mathbf{P}_{\mathbb{C}}^1}(d[\infty])|_{U_\infty} \simeq \mathcal{O}_{U_\infty}.$$

On the overlap $U_0 \cap U_\infty$, we have $w = 1/z$. Therefore

$$e_\infty = w^{-d} = z^d.$$

Since $e_0 = 1$, this gives

$$e_\infty = z^d e_0.$$

Thus $\mathcal{O}_{\mathbf{P}_{\mathbb{C}}^1}(d[\infty])$ is obtained by gluing the two trivial line bundles on U_0 and U_∞ using the invertible function

$$z^d \in \mathcal{O}^\times(U_0 \cap U_\infty).$$

IV.E Properties of sheaves

In this subsection, $\mathcal{F}$ and $\mathcal{G}$ are sheaves of abelian groups on a topological space X .

IV.E.1 Kernels, images, cokernels stalks

Let us define the kernel, cokernel and image of a morphism of sheaves $\varphi : \mathcal{F} \longrightarrow \mathcal{G}$

Definition IV.E.1. The *kernel* of φ is the sheaf $\ker(\varphi)$ defined by

$$\ker(\varphi)(U) = \ker(\varphi_U : \mathcal{F}(U) \longrightarrow \mathcal{G}(U))$$

for every open subset $U \subset X$. The *presheaf image* of φ is the presheaf $\mathrm{Im}^{\mathrm{pre}}(\varphi)$ defined by

$$\mathrm{Im}^{\mathrm{pre}}(\varphi)(U) = \mathrm{Im}(\mathcal{F}(U) \longrightarrow \mathcal{G}(U)).$$

The *image sheaf* of φ is its sheafification $\mathrm{Im}(\varphi) = (\mathrm{Im}^{\mathrm{pre}}(\varphi))^+$. The *presheaf cokernel* of φ is the presheaf $\mathrm{Coker}^{\mathrm{pre}}(\varphi)$ defined by

$$\mathrm{Coker}^{\mathrm{pre}}(\varphi)(U) = \mathcal{G}(U) / \varphi_U(\mathcal{F}(U)).$$

The *cokernel sheaf* of φ is $\mathrm{Coker}(\varphi) = (\mathrm{Coker}^{\mathrm{pre}}(\varphi))^+$. We say that φ is *injective* if

$$\ker(\varphi) = 0,$$

and *surjective* if

$$\mathrm{Im}(\varphi) = \mathcal{G}.$$

Equivalently, φ is surjective if

$$\mathrm{Coker}(\varphi) = 0.$$

Remark IV.E.2. As we will see, the kernel is already a sheaf. Consequently

$$\varphi \text{ is injective} \iff \varphi_U : \mathcal{F}(U) \longrightarrow \mathcal{G}(U) \text{ is injective for every open } U \subset X.$$

The analogous assertion for surjectivity is false, as the next example indicates. Certainly, if every map

$$\varphi_U : \mathcal{F}(U) \longrightarrow \mathcal{G}(U)$$

is surjective, then φ is surjective as a morphism of sheaves. But the converse need not hold. Surjectivity as a morphism of sheaves means instead that sections lift *locally*: for every open subset $U \subset X$, every $s \in \mathcal{G}(U)$ and every $x \in U$, there exist an open neighbourhood $V \subset U$ of x and a section $t \in \mathcal{F}(V)$ such that

$$\varphi_V(t) = s|_V.$$

Example IV.E.3 (A surjective sheaf morphism which is not surjective on sections). Let

$$X = S^1 = \{z \in \mathbb{C} \mid |z| = 1\}.$$

Let $\mathcal{C}_X^0(\mathbb{R})$ be the sheaf of continuous real-valued functions and let $\mathcal{C}_X^0(S^1)$ be the sheaf of continuous S^1 -valued functions, where the latter is regarded as a sheaf of abelian groups under multiplication. Consider the morphism

$$\exp : \mathcal{C}_X^0(\mathbb{R}) \longrightarrow \mathcal{C}_X^0(S^1), \quad f \longmapsto e^{2\pi i f}.$$

This morphism is surjective on every stalk. Indeed, let

$$g : U \longrightarrow S^1$$

be continuous and let $x \in U$. After shrinking U around x , the image of g is contained in an open arc of S^1 on which the exponential map admits a continuous inverse. Thus g has a continuous logarithm locally around x .

However, the map on global sections

$$\mathcal{C}_X^0(\mathbb{R})(X) \longrightarrow \mathcal{C}_X^0(S^1)(X)$$

is not surjective. Indeed, the identity map $\iota : S^1 \longrightarrow S^1$, defined just as $z \longmapsto z$, has no continuous logarithm. To see this, suppose that there were a continuous map

$$h : S^1 \longrightarrow \mathbb{R}$$

such that

$$e^{2\pi i h(z)} = z.$$

For $t \in [0, 1]$, the function

$$h(e^{2\pi i t}) - t$$

takes values in $\mathbb{Z}$. Since it is continuous, it is constant. But its values at $t = 0$ and $t = 1$ differ by 1, a contradiction.

Thus $\exp$ is surjective as a morphism of sheaves, but it is not surjective on global sections. This example also shows why the presheaf cokernel must be sheafified. The class of ι is nonzero in

$$\mathcal{C}_X^0(S^1)(X) / \exp(\mathcal{C}_X^0(\mathbb{R})(X)),$$

but its restriction to every sufficiently small open arc is zero. Therefore the presheaf cokernel does not satisfy the locality axiom.

Exercise IV.E.4 (The kernel of a morphism of sheaves). Let $\varphi : \mathcal{F} \longrightarrow \mathcal{G}$ be a morphism of sheaves of abelian groups on X . For every open subset $U \subset X$, put

$$\mathcal{K}(U) = \ker(\varphi_U : \mathcal{F}(U) \longrightarrow \mathcal{G}(U)).$$

- i) Show that the restriction maps of $\mathcal{F}$ induce restriction maps on the groups $\mathcal{K}(U)$.
- ii) Prove that $\mathcal{K}$ is a sheaf.
- iii) Show that, for every $x \in X$, the natural map

$$\mathcal{K}_x \longrightarrow \ker(\varphi_x)$$

is an isomorphism.

- iv) Deduce that the following conditions are equivalent:

- (a) φ is injective;
- (b) φ_U is injective for every open subset $U \subset X$;
- (c) φ_x is injective for every $x \in X$.

Solution. Let $V \subset U$ be open and let $s \in \mathcal{K}(U)$. Since φ commutes with restrictions,

$$\varphi_V(s|_V) = \varphi_U(s)|_V = 0.$$

Thus $s|_V \in \mathcal{K}(V)$, proving i).

For ii), let $U = \bigcup_i U_i$ and suppose that sections

$$s_i \in \mathcal{K}(U_i)$$

agree on all intersections. Since $\mathcal{F}$ is a sheaf, they glue uniquely to a section

$$s \in \mathcal{F}(U).$$

The sections $\varphi_{U_i}(s_i)$ are all zero. Hence

$$\varphi_U(s)|_{U_i} = 0$$

for every i . By the locality axiom for $\mathcal{G}$,

$$\varphi_U(s) = 0.$$

Thus $s \in \mathcal{K}(U)$, and $\mathcal{K}$ is a sheaf.

For iii), the natural map

$$\mathcal{K}_x \longrightarrow \mathcal{F}_x$$

has image contained in $\ker(\varphi_x)$. It is injective because $\mathcal{K}$ is a subsheaf of $\mathcal{F}$.

Let now $\sigma \in \ker(\varphi_x)$. Choose an open neighbourhood U of x and a section $s \in \mathcal{F}(U)$ representing σ . The equality

$$\varphi_x(\sigma) = 0$$

means that, after shrinking U around x , one has

$$\varphi_U(s) = 0.$$

Thus s is locally a section of $\mathcal{K}$, and its germ maps to σ . Therefore

$$\mathcal{K}_x \simeq \ker(\varphi_x).$$

Finally,

$$\varphi \text{ is injective} \iff \ker(\varphi) = 0.$$

A sheaf is zero if and only if all its groups of sections are zero, and also if and only if all its stalks are zero. Using iii), this proves iv).

IV.E.2 Injectivity and surjectivity on stalks

Let $\varphi : \mathcal{F} \rightarrow \mathcal{G}$ be a morphism of sheaves on X .

Lemma IV.E.5. *Let $\varphi : \mathcal{F} \rightarrow \mathcal{G}$ be a morphism of sheaves of abelian groups on X . Then:*

- i) *the morphism φ is injective if and only if $\varphi_x : \mathcal{F}_x \rightarrow \mathcal{G}_x$ is injective for every $x \in X$;*
- ii) *the morphism φ is surjective if and only if φ_x is surjective for every $x \in X$;*
- iii) *the morphism φ is an isomorphism if and only if φ_x is an isomorphism for every $x \in X$.*

Proof. Assertion i) follows from Exercise IV.E.4. For the image sheaf, there is a natural identification

$$\mathrm{Im}(\varphi)_x \simeq \mathrm{Im}(\varphi_x).$$

Indeed, sheafification does not change stalks, and a germ in the presheaf image is represented by a section of the form $\varphi(s)$ on some neighbourhood of x . Similarly,

$$\mathrm{Coker}(\varphi)_x \simeq \mathrm{Coker}(\varphi_x).$$

To see this directly, let a germ represented by $t \in \mathcal{G}(U)$ map to zero in $\text{Coker}(\varphi_x)$. Then $t_x = \varphi_x(s_x)$ for some $s_x \in \mathcal{F}_x$. Choose a representative $s \in \mathcal{F}(V)$ of s_x . After shrinking around x , one has $t|_V = \varphi_V(s)$, so the germ of the class of t in the presheaf cokernel is zero. Consequently, we get ii), as

$$\text{Coker}(\varphi) = 0 \iff \text{Coker}(\varphi_x) = 0 \text{ for every } x \in X.$$

Suppose now that every φ_x is an isomorphism. By i), φ is injective. Let $u \in \mathcal{G}(U)$. By the surjectivity of the maps on stalks, every point $x \in U$ has an open neighbourhood $U_x \subset U$ and a section $s_x \in \mathcal{F}(U_x)$ such that $\varphi_{U_x}(s_x) = u|_{U_x}$. On an intersection $U_x \cap U_y$, the images of s_x and s_y agree. Since φ is injective, one has

$$s_x|_{U_x \cap U_y} = s_y|_{U_x \cap U_y}.$$

The sections s_x therefore glue to a section $s \in \mathcal{F}(U)$ satisfying $\varphi_U(s) = u$. Thus every φ_U is bijective, and φ is an isomorphism. The converse is immediate. $\square$

IV.F Functorial construction with sheaves

IV.F.1 Direct image

Let $f: X \rightarrow Y$ be a continuous map and let $\mathcal{F}$ be a sheaf of sets on X . The *direct image* of $\mathcal{F}$ is the sheaf $f_*\mathcal{F}$ on Y defined by

$$(f_*\mathcal{F})(V) = \mathcal{F}(f^{-1}(V))$$

for every open subset $V \subset Y$. The restriction maps are induced by the restriction maps of $\mathcal{F}$.

Exercise IV.F.1. Let $f: X \rightarrow Y$ be a continuous map and let $\mathcal{F}$ be a sheaf on X . Show directly from the definition that $f_*\mathcal{F}$ is a sheaf on Y .

Solution. Let $V = \bigcup_i V_i$ be an open cover in Y . Then

$$f^{-1}(V) = \bigcup_i f^{-1}(V_i)$$

is an open cover in X . Sections of $f_*\mathcal{F}$ on V are sections of $\mathcal{F}$ on $f^{-1}(V)$. The uniqueness and gluing axioms for $f_*\mathcal{F}$ therefore follow immediately from the corresponding axioms for $\mathcal{F}$ applied to the cover $(f^{-1}(V_i))$.

Example IV.F.2 (A skyscraper sheaf). Let x be a point of a topological space X and let $i: \{x\} \hookrightarrow X$ be the inclusion. Let W be a vector space, regarded as a sheaf on the one-point space. Then

$$(i_*W)(U) = \begin{cases} W, & x \in U, \\ 0, & x \notin U. \end{cases}$$

This sheaf is called the *skyscraper sheaf* supported at x with fibre W . Its stalk is W at x and zero at every other point.

Example IV.F.3 (The map to a point). Let $p: X \rightarrow \{*\}$ be the unique map. Then

$$(p_*\mathcal{F})(\{*\}) = \mathcal{F}(X).$$

Thus direct image to a point is the operation of taking global sections.

Example IV.F.4 (An open inclusion). Let $j : \mathbb{C}^* \hookrightarrow \mathbb{C}$ be the inclusion. Then

$$(j_* \mathcal{O}_{\mathbb{C}^*})(U) = \mathcal{O}(U \cap \mathbb{C}^*).$$

The stalk $(j_* \mathcal{O}_{\mathbb{C}^*})_0$ consists of germs of holomorphic functions on punctured neighbourhoods of the origin. It contains, for instance, the germs of $1/z$ and $e^{1/z}$. Therefore direct image by an open inclusion is not the same as extension by zero.

Example IV.F.5 (A double cover). Consider locally the map

$$\pi : \mathbb{C} \longrightarrow \mathbb{C}, \quad w \longmapsto z = w^2.$$

At the branch point, the map of local rings is

$$\mathbb{C}\{z\} \longrightarrow \mathbb{C}\{w\}, \quad z \longmapsto w^2.$$

Every convergent power series in w has a unique decomposition into its even and odd parts,

$$h(w) = a(w^2) + w b(w^2).$$

Consequently

$$\mathbb{C}\{w\} = \mathbb{C}\{z\} \oplus w \mathbb{C}\{z\}$$

as a $\mathbb{C}\{z\}$ -module. Sheaf-theoretically, $\pi_* \mathcal{O}_{\mathbb{C}}$ is locally a free $\mathcal{O}_{\mathbb{C}}$ -module of rank two. This is the local analytic form of the algebraic decomposition

$$\mathbb{k}[x, y]/(y^2 - f(x)) = \mathbb{k}[x] \oplus \mathbb{k}[x]y$$

which appeared in the elliptic-curve example.

Exercise IV.F.6. Let $i : Z \hookrightarrow X$ be the inclusion of a closed subset and let $\mathcal{F}$ be a sheaf on Z . Show that

$$(i_* \mathcal{F})_x \simeq \begin{cases} \mathcal{F}_x, & x \in Z, \\ 0, & x \notin Z, \end{cases}$$

when $\mathcal{F}$ is a sheaf of abelian groups.

Solution. If $x \notin Z$, there is an open neighbourhood U of x disjoint from Z , so $(i_* \mathcal{F})(U) = \mathcal{F}(\emptyset) = 0$. Hence the stalk is zero. If $x \in Z$, the open neighbourhoods $U \cap Z$ form a neighbourhood basis of x in Z , and the definition of the two stalks gives $(i_* \mathcal{F})_x \simeq \mathcal{F}_x$.

IV.F.2 Inverse image

Let $f : X \rightarrow Y$ be continuous and let $\mathcal{G}$ be a sheaf on Y . The assignment

$$U \longmapsto \varinjlim_{U \subset f^{-1}(V)} \mathcal{G}(V),$$

where V runs through the open subsets of Y , is in general only a presheaf. Its sheafification is called the *inverse-image sheaf* and is denoted by $f^{-1}\mathcal{G}$. Its stalks have the particularly simple description

$$(f^{-1}\mathcal{G})_x \simeq \mathcal{G}_{f(x)}.$$

Thus inverse image transports germs near $f(x)$ to germs near x .

If X and Y are smooth manifolds and f is smooth, or if they are complex manifolds and f is holomorphic, composition of functions gives a natural morphism

$$f^{-1}\mathcal{O}_Y \longrightarrow \mathcal{O}_X, \quad g \longmapsto g \circ f.$$

Chapter 2

Affine schemes

This chapter contains a review of the basic concepts and constructions concerning affine schemes. Throughout this chapter, rings are commutative and unital, and we allow the zero ring, whose unit is 0. For every ring A , we denote by

$$\gamma_A : \mathbb{Z} \longrightarrow A, \quad n \longmapsto n1_A,$$

the unique unital ring homomorphism from $\mathbb{Z}$ to A .

I Prime spectrum of a ring

Further reading. For affine spectra and the Zariski topology, see [Har77, Chapter II, §2], [EH00], and [Liu02, Chapter 2].

We introduce the set of prime ideals of a ring and basic properties of the spectrum.

I.A Prime ideals

We define prime ideals and the prime spectrum. An ideal $\mathfrak{p}$ of A is *prime* if $\mathfrak{p} \neq A$ and, for all $x, y \in A$, whenever $xy \in \mathfrak{p}$, we must have $x \in \mathfrak{p}$ or $y \in \mathfrak{p}$. This is equivalent to the fact that $A/\mathfrak{p}$ is an integral domain.

Definition I.A.1. Let A be a ring. We denote by $\text{Spec}(A)$ the set of prime ideals of A , called the *prime spectrum* of A . When we speak of a prime ideal, we exclude the ideal A , sometimes called the *unit ideal*. In particular, if A is the zero ring, then $\text{Spec}(A) = \emptyset$.

Example I.A.2. Let us describe the prime spectrum of some well-known rings.

1. The prime spectrum of $\mathbb{Z}$ consists of the ideal (0) , together with all ideals of the form (p) , where p is a prime number.
2. More generally, the prime spectrum of a principal ideal domain A consists of the ideal (0) , as well as ideals of the form (f) , where f is irreducible in A .
3. The prime spectrum of a field $\mathbb{k}$ is given by the single ideal (0) .

Example I.A.3. Let $A = \mathbb{k}[x]/(x^2)$ be the ring of dual numbers. The prime ideals of A correspond to the prime ideals of $\mathbb{k}[x]$ containing (x^2) . Since every such prime ideal contains x , there is only one prime ideal:

$$(\bar{x}) \subset A.$$

Thus $\text{Spec}(\mathbb{k}[x]/(x^2))$ has a single point. Note that (0) is not a prime ideal.

I.B Functoriality of the prime spectrum

We explain how a ring homomorphism induces a map of spectra. Let $\varphi : A \rightarrow B$ be a ring homomorphism. Then we define:

$$\text{Spec}(B) \rightarrow \text{Spec}(A), \quad \mathfrak{q} \mapsto \varphi^{-1}(\mathfrak{q}).$$

This is well defined. Indeed, if $\mathfrak{q}$ is a prime ideal of B , then $\varphi^{-1}(\mathfrak{q})$ is also a prime ideal. To see this, note that for any ideal $\mathfrak{q}$ of B , $\varphi^{-1}(\mathfrak{q})$ is the kernel of the composition $A \xrightarrow{\varphi} B \xrightarrow{\pi} B/\mathfrak{q}$. Since $B/\mathfrak{q}$ is an integral domain, this kernel is prime.

Exercise I.B.1. Let $\varphi : A \rightarrow B$ be a morphism of rings. Prove that if φ and ψ are composable ring homomorphisms, then

$$f_{\psi \circ \varphi} = f_{\varphi} \circ f_{\psi}.$$

Next, show that if φ is surjective, then

$$f_{\varphi} : \text{Spec}(B) \rightarrow \text{Spec}(A)$$

is injective.

Solution. Let $\varphi : A \rightarrow B$ and $\psi : B \rightarrow C$, and let $\mathfrak{r} \in \text{Spec}(C)$. Then

$$f_{\psi \circ \varphi}(\mathfrak{r}) = (\psi \circ \varphi)^{-1}(\mathfrak{r}) = \varphi^{-1}(\psi^{-1}(\mathfrak{r})) = f_{\varphi}(f_{\psi}(\mathfrak{r})).$$

Hence $f_{\psi \circ \varphi} = f_{\varphi} \circ f_{\psi}$.

If φ is surjective and $\mathfrak{q}_1, \mathfrak{q}_2 \in \text{Spec}(B)$ have the same inverse image in A , then every element of $\mathfrak{q}_1$ is of the form $\varphi(a)$ with $a \in A$. It lies in $\mathfrak{q}_1$ if and only if $a \in \varphi^{-1}(\mathfrak{q}_1) = \varphi^{-1}(\mathfrak{q}_2)$, hence if and only if $\varphi(a) \in \mathfrak{q}_2$. Thus $\mathfrak{q}_1 = \mathfrak{q}_2$.

II The Zariski topology

The prime spectrum of a ring A is naturally equipped with the Zariski topology. The closed sets of this topology are the subsets of the form $\mathbb{V}(\mathfrak{a})$, where $\mathfrak{a}$ is an ideal of A .

II.A Closed subsets of the affine spectrum

We define closed subsets of the spectrum and study basic set-theoretic properties.

II.A.1 Closed sets and ideals

Let C be a subset of A .

Definition II.A.1. Let C be any subset of A . Define

$$\mathbb{V}(C) := \{\mathfrak{p} \in \text{Spec}(A) \mid C \subset \mathfrak{p}\}.$$

Let $\mathfrak{a}$ be the ideal of A generated by C . Then, given a prime ideal $\mathfrak{p}$ of A , we have $\mathfrak{a} \subset \mathfrak{p}$ if and only if $C \subset \mathfrak{p}$. For this reason we usually consider subsets $\mathbb{W}(\mathfrak{a})$ where $\mathfrak{a}$ is an ideal of A . We have

$$\mathbb{W}(\{0\}) = \text{Spec}(A), \quad \mathbb{W}(A) = \emptyset.$$

Remark II.A.2. Let $\mathfrak{a}$ be an ideal of A and consider the projection $\pi : A \rightarrow A/\mathfrak{a}$. Then we have

$$\mathbb{W}(\mathfrak{a}) := \text{Im}(\text{Spec}(A/\mathfrak{a}) \xrightarrow{f_\pi} \text{Spec}(A)).$$

This establishes a bijection:

$$\{\mathfrak{p} \in \text{Spec}(A) \mid \mathfrak{p} \supset \mathfrak{a}\} \leftrightarrow \{\mathfrak{q} \in \text{Spec}(A/\mathfrak{a})\}.$$

Indeed, by definition, a prime ideal $\mathfrak{p}$, we have $\mathfrak{p} \in \mathbb{W}(\mathfrak{a})$ if and only if $\mathfrak{p} \supset \mathfrak{a}$. Now, if $\mathfrak{p} \supset \mathfrak{a}$, then we see that $\pi^{-1}(\pi(\mathfrak{p})) = \mathfrak{p}$, because for any element $f \in \pi^{-1}(\pi(\mathfrak{p}))$, we have $\pi(f) \in \pi(\mathfrak{p})$ so $\pi(f) = \pi(g)$ for some $g \in \mathfrak{p}$, so $\pi(f - g) \in \mathfrak{a} \subset \mathfrak{p}$ and thus $f \in \mathfrak{p}$. Also, $\pi(\mathfrak{p})$ is a prime ideal of $A/\mathfrak{a}$ in this case.

So given $\mathfrak{p} \in \text{Spec}(A)$, the condition that $\mathfrak{p} \supset \mathfrak{a}$ is equivalent to $\mathfrak{p} = \pi^{-1}(\mathfrak{q})$, where $\mathfrak{q}$ is a prime ideal of $A/\mathfrak{a}$.

Remark II.A.3. The notion of Zariski closed subset of $\text{Spec}(A)$ parallels the definition of the affine algebraic subsets of $\mathbb{k}^n$. Indeed, let $A = \mathbb{k}[x_1, \dots, x_n]$. Then, among the points of $\text{Spec}(A)$, it is natural to consider the maximal ideals of the form $\mathfrak{m}_a = (x_1 - a_1, \dots, x_n - a_n)$, where $a = (a_1, \dots, a_n)$ lies in $\mathbb{k}^n$. The scheme $\text{Spec}(A)$ is called the n -th affine space over $\mathbb{k}$, denoted by $\mathbb{A}_{\mathbb{k}}^n$.

For a given ideal $\mathfrak{a}$ of A , a point $\mathfrak{m}_a$ lies in $\mathbb{W}(\mathfrak{a})$ if and only if $\mathfrak{a} \subset \mathfrak{m}_a$, which means that every $f \in \mathfrak{a}$ lies in $\mathfrak{m}_a$, in other words, that $f(a) = 0$. So we have

$$\mathfrak{m}_a \in \mathbb{W}(\mathfrak{a}) \Leftrightarrow a \in \mathbb{V}(\mathfrak{a}).$$

Example II.A.4. If $A = \mathbb{k}[x]$, then

$$\mathbb{W}(x) = \{(x)\}, \quad \mathbb{W}(0) = \text{Spec}(\mathbb{k}[x]).$$

If $\mathbb{k}$ is algebraically closed, the closed points of $\text{Spec}(\mathbb{k}[x])$ are the ideals $(x - a)$, for $a \in \mathbb{k}$. On the other hand, (0) is a point whose closure is the whole space, since every prime ideal contains 0.

II.A.2 First properties of closed sets

Associating $\mathbb{W}(\mathfrak{a})$ to $\mathfrak{a}$ is an inclusion-reversing operation, namely, given ideals $\mathfrak{a}$ and $\mathfrak{b}$ of A , we have $\mathbb{W}(\mathfrak{b}) \subset \mathbb{W}(\mathfrak{a})$ if $\mathfrak{a} \subset \mathfrak{b}$. The next properties are crucial to define the subsets of the form $\mathbb{W}(\mathfrak{a})$, where $\mathfrak{a}$ runs in the set of all ideals of A , as the closed subsets of a topology.

Lemma II.A.5. *We have the following equalities of closed subsets.*

1. For all ideals $\mathfrak{a}, \mathfrak{b}$ of A : $\mathbb{W}(\mathfrak{a}) \cup \mathbb{W}(\mathfrak{b}) = \mathbb{W}(\mathfrak{ab})$, where $\mathfrak{ab} = (xy \mid x \in \mathfrak{a}, y \in \mathfrak{b})$.
2. If $(\mathfrak{a}_i)_{i \in I}$ is a family of ideals and $\mathfrak{a} = \sum_{i \in I} \mathfrak{a}_i$, then $\mathbb{W}(\mathfrak{a}) = \bigcap_{i \in I} \mathbb{W}(\mathfrak{a}_i)$.

Proof. For the first item, of course $\mathbb{W}(\mathfrak{ab}) \supset \mathbb{W}(\mathfrak{a}) \cup \mathbb{W}(\mathfrak{b})$. For the reverse inclusion, let $\mathfrak{p} \supset \mathfrak{ab}$ be a prime ideal. If $\mathfrak{p}$ contains neither $\mathfrak{a}$ nor $\mathfrak{b}$, then we may pick $x \in \mathfrak{a} \setminus \mathfrak{p}$, $y \in \mathfrak{b} \setminus \mathfrak{p}$. Then $xy \in \mathfrak{ab} \subset \mathfrak{p}$, a contradiction. About the second item, note that a prime ideal contains $\sum_{i \in I} \mathfrak{a}_i$ if and only if it contains each $\mathfrak{a}_i$, for all $i \in I$. $\square$

Proposition II.A.6. *Let $\mathfrak{p} \in \text{Spec}(A)$. Then $\overline{\{\mathfrak{p}\}} = \mathbb{W}(\mathfrak{p})$. The ideal $\mathfrak{p}$ is maximal if and only if $\{\mathfrak{p}\}$ is closed in $\text{Spec}(A)$, which is to say, if and only if $\mathbb{W}(\mathfrak{p}) = \{\mathfrak{p}\}$.*

Proof. A point $\mathfrak{q}$ lies in $\overline{\{\mathfrak{p}\}}$ if and only if, for all ideals $\mathfrak{a}$ of A with $\mathfrak{p} \in \mathbb{W}(\mathfrak{a})$, one has $\mathfrak{q} \in \mathbb{W}(\mathfrak{a})$. In other words, $\mathfrak{q}$ lies in $\overline{\{\mathfrak{p}\}}$ if and only if, for all ideals $\mathfrak{a}$ of A with $\mathfrak{a} \subset \mathfrak{p}$, one has $\mathfrak{a} \subset \mathfrak{q}$. Therefore, any point $\mathfrak{q} \in \mathbb{W}(\mathfrak{p})$ lies in $\overline{\{\mathfrak{p}\}}$, as $\mathfrak{q}$ satisfies $\mathfrak{p} \subset \mathfrak{q}$, so $\mathfrak{a} \subset \mathfrak{p}$ implies $\mathfrak{a} \subset \mathfrak{q}$. Conversely, given $\mathfrak{q} \in \overline{\{\mathfrak{p}\}}$, choosing $\mathfrak{a} = \mathfrak{p}$, we have $\mathfrak{a} \subset \mathfrak{a} = \mathfrak{p}$, so $\mathfrak{p} = \mathfrak{a} \subset \mathfrak{q}$. Therefore $\mathfrak{q} \in \mathbb{W}(\mathfrak{p})$.

For the second statement, if $\mathfrak{p}$ is maximal, then the only point of $\overline{\{\mathfrak{p}\}}$ is $\mathfrak{p}$. Indeed, if a prime ideal $\mathfrak{q}$ lies in $\overline{\{\mathfrak{p}\}}$, using the above characterisation and choosing $\mathfrak{a} = \mathfrak{p}$ we have $\mathfrak{a} = \mathfrak{p} \subset \mathfrak{q}$, so $\mathfrak{q} = \mathfrak{p}$ since $\mathfrak{p}$ is maximal. Conversely, if $\mathfrak{p}$ is not maximal, then there is a maximal ideal $\mathfrak{q}$ strictly containing $\mathfrak{p}$, so that $\mathfrak{q} \in \mathbb{W}(\mathfrak{p}) \setminus \{\mathfrak{p}\}$. $\square$

Lemma II.A.7. *Let $\varphi : A \rightarrow B$ be a ring morphism. Then $f_\varphi : \text{Spec}(B) \rightarrow \text{Spec}(A)$ is continuous.*

Proof. Given an ideal $\mathfrak{a}$ of A , we have

$$f_\varphi^{-1}(\mathbb{W}(\mathfrak{a})) = \mathbb{W}(\varphi(\mathfrak{a})).$$

Indeed, on one hand, by definition a prime ideal $\mathfrak{q}$ of B contains $\varphi(\mathfrak{a})$ if and only if $\mathfrak{q} \in \mathbb{W}(\varphi(\mathfrak{a}))$. On the other hand $\mathfrak{q}$ lies in $f_\varphi^{-1}(\mathbb{W}(\mathfrak{a}))$ if and only if $f_\varphi(\mathfrak{q}) \in \mathbb{W}(\mathfrak{a})$, which happens if and only if $\varphi^{-1}(\mathfrak{q})$ contains $\mathfrak{a}$ and applying φ we see that this amounts precisely to $\mathfrak{q} \supset \varphi(\mathfrak{a})$. $\square$

II.B Intersection of primes and the radical

It turns out that the radical of an ideal is the intersection of prime ideals containing it.

Lemma II.B.1. *Let $\mathfrak{a}$ be an ideal of a ring A . Then:*

$$\sqrt{\mathfrak{a}} = \bigcap_{\mathfrak{p} \in \mathbb{W}(\mathfrak{a})} \mathfrak{p}.$$

In particular, the nilradical $\text{Rad}(A) = \sqrt{0}$ is the intersection of all prime ideals of A .

Proof. Let us check that the left-hand side is contained in the right-hand side. Given a prime ideal $\mathfrak{p} \in \mathbb{W}(\mathfrak{a})$, by definition $\mathfrak{a} \subset \mathfrak{p}$, so $\sqrt{\mathfrak{a}} \subset \sqrt{\mathfrak{p}} = \mathfrak{p}$.

To check the opposite inclusion, we first consider the ring $B = A/\mathfrak{a}$ and the projection $\varphi : A \rightarrow B$. Then $\varphi^{-1}(\text{Rad}(B)) = \sqrt{\mathfrak{a}}$ and sending $\mathfrak{q} \in \mathbb{W}(0)$ to $\varphi^{-1}(\mathfrak{q})$ gives a bijection of $\mathbb{W}(0)$ and $\mathbb{W}(\mathfrak{a})$. Therefore, up to replacing A by B and $\mathfrak{a}$ by 0 , it suffices to prove the statement for $\mathfrak{a} = 0$.

So let us show that an element $a \in A$ lying in all prime ideals $\mathfrak{p}$ of A is nilpotent. Suppose the contrary, so that the localisation A_a is a non-zero ring. Then this ring has a prime ideal $\mathfrak{q}$. Pulling back $\mathfrak{q}$ to A , we get a prime ideal $\mathfrak{p}$, which does not contain a . A contradiction. $\square$

Exercise II.B.2. Show the equality of subsets $\mathbb{W}(\mathfrak{a}) = \mathbb{W}(\sqrt{\mathfrak{a}})$ for every ideal $\mathfrak{a} \subset A$.

Solution. A prime ideal $\mathfrak{p}$ contains $\mathfrak{a}$ if and only if it contains $\sqrt{\mathfrak{a}}$. One implication is immediate from $\mathfrak{a} \subset \sqrt{\mathfrak{a}}$. Conversely, if $\mathfrak{a} \subset \mathfrak{p}$, then $\sqrt{\mathfrak{a}} \subset \sqrt{\mathfrak{p}} = \mathfrak{p}$ because $\mathfrak{p}$ is radical.

Lemma II.B.3. *Given ideals $\mathfrak{a}$ and $\mathfrak{b}$ of A , we have $\mathbb{W}(\mathfrak{a}) \subset \mathbb{W}(\mathfrak{b})$ if and only if $\mathfrak{b} \subset \sqrt{\mathfrak{a}}$. Moreover:*

$$\mathbb{W}(\mathfrak{a}) = \emptyset \Leftrightarrow \mathfrak{a} = A, \quad \mathbb{W}(\mathfrak{a}) = \text{Spec}(A) \Leftrightarrow \mathfrak{a} \subset \text{Rad}(A).$$

Proof. For the statement about inclusions, assume $\mathbb{W}(\mathfrak{a}) \subset \mathbb{W}(\mathfrak{b})$. Then $\mathfrak{b} \subset \sqrt{\mathfrak{b}} \subset \sqrt{\mathfrak{a}}$. Conversely, suppose $\mathfrak{b} \subset \sqrt{\mathfrak{a}}$. Then, given a prime ideal $\mathfrak{p} \in \mathbb{W}(\mathfrak{a})$, we have $\mathfrak{b} \subset \sqrt{\mathfrak{a}} \subset \sqrt{\mathfrak{p}} = \mathfrak{p}$, so $\mathfrak{p} \in \mathbb{W}(\mathfrak{b})$.

Let us show the statement appearing in the last display. If $\mathbb{W}(\mathfrak{a}) = \emptyset$, then there is no prime ideal containing $\mathfrak{a}$, so $\mathfrak{a} = A$, because any ideal different from A is contained in some maximal (hence prime) ideal. Conversely, if $\mathfrak{a} = A$, then there is no prime ideal containing $\mathfrak{a}$, so $\mathbb{W}(\mathfrak{a}) = \emptyset$. Also, if $\mathbb{W}(\mathfrak{a}) = \text{Spec}(A)$ then any prime ideal of A contains $\mathfrak{a}$, so $\mathfrak{a}$ is contained in the intersection of all prime ideals of A , i.e., in $\text{Rad}(A)$. Conversely, if $\mathfrak{a} \subset \text{Rad}(A)$, then $\mathfrak{a}$ is contained in every prime ideal of A so $\mathbb{W}(\mathfrak{a}) = \text{Spec}(A)$. $\square$

Exercise II.B.4. Compute the image of the map on spectra in each of the following cases, and compare it with $\mathbb{W}(\ker \varphi)$.

- (i) The quotient map $A \rightarrow A/\mathfrak{a}$.
- (ii) The localisation map $A \rightarrow S^{-1}A$.
- (iii) The homomorphism $\gamma_{\mathbb{Q}} : \mathbb{Z} \hookrightarrow \mathbb{Q}$.

Solution. For a quotient, contraction gives a bijection between the primes of $A/\mathfrak{a}$ and the primes of A containing $\mathfrak{a}$. Thus the image is exactly $\mathbb{W}(\mathfrak{a}) = \mathbb{W}(\ker \varphi)$.

The image of $\text{Spec}(S^{-1}A) \rightarrow \text{Spec}(A)$ is

$$\{\mathfrak{p} \in \text{Spec}(A) \mid \mathfrak{p} \cap S = \emptyset\}.$$

It is usually not closed, although its closure is $\mathbb{W}(\ker(A \rightarrow S^{-1}A))$.

Finally, $\text{Spec}(\mathbb{Q})$ has one point, whose image in $\text{Spec}(\mathbb{Z})$ is (0) . However, $\ker(\gamma_{\mathbb{Q}}) = 0$. Since all (prime) ideals of $\mathbb{Z}$ contain 0, we get

$$\overline{\{(0)\}} = \mathbb{W}(0) = \text{Spec}(\mathbb{Z}).$$

Thus the image consists of a single point, but this point is dense.

II.C Dominant morphisms

A *dominant morphism* refers to a morphism of topological spaces such that the image is dense in the target. We use the notion introduced previously, namely, the *radical* of a ring.

Example II.C.1. The inclusion $\mathbb{k}[x] \subset \mathbb{k}[x, y]$ induces the projection

$$\text{Spec}(\mathbb{k}[x, y]) \rightarrow \text{Spec}(\mathbb{k}[x]).$$

Its image is dense. Geometrically, this says that the projection from the affine plane to the affine line is dominant.

Proposition II.C.2. Let $\varphi : A \rightarrow B$ be an injective morphism of rings. Then the induced map $f_{\varphi} : \text{Spec}(B) \rightarrow \text{Spec}(A)$ is dominant.

Proof. We identify A with its image in B . Since the nilradical of a ring is the set of nilpotent elements, we have $\text{Rad}(B) \cap A \subset \text{Rad}(A)$. Therefore, given an ideal $\mathfrak{a}$ of A , we have $\mathfrak{a} \subset \text{Rad}(A)$, as soon as $\mathfrak{a} \subset \text{Rad}(B) \cap A$.

Now, take a closed subset $\mathbb{W}(\mathfrak{a}) \subset \text{Spec}(A)$, for some ideal $\mathfrak{a}$ of A , and assume that $\mathbb{W}(\mathfrak{a})$ contains the image of f_{φ} , so that $\mathfrak{a}$ is contained in $\mathfrak{q} \cap A$, for all prime ideals $\mathfrak{q}$ of B . Then, by Lemma II.B.1, we have

$$\mathfrak{a} \subset \bigcap_{\mathfrak{q} \in \text{Spec}(B)} (\mathfrak{q} \cap A) = \text{Rad}(B) \cap A \subset \text{Rad}(A).$$

Then $\mathbb{W}(\mathfrak{a}) = \text{Spec}(A)$, by Lemma II.B.3. So there is no closed subset of $\text{Spec}(A)$, strictly contained in $\text{Spec}(A)$, containing the image of f_{φ} , which says that the closure of the image of f_{φ} is $\text{Spec}(A)$. $\square$

Exercise II.C.3. Let $\varphi : A \rightarrow B$ be a morphism of rings and let $f : \text{Spec}(B) \rightarrow \text{Spec}(A)$ be the induced morphism. Generalise the previous proposition by showing that f is dominant if and only if

$$\ker(\varphi) \subset \text{Rad}(A).$$

Deduce that, if A is reduced, then f is dominant if and only if φ is injective.

Solution. Take a closed subset $\mathbb{V}(\mathfrak{a}) \subset \text{Spec}(A)$, for some ideal $\mathfrak{a}$ of A , and assume that $\mathbb{V}(\mathfrak{a})$ contains the image of f_φ . So $\mathfrak{a}$ is contained in $\varphi^{-1}(\mathfrak{q})$, for all prime ideals $\mathfrak{q}$ of B . Then, by Lemma II.B.1

$$\mathfrak{a} \subset \bigcap_{\mathfrak{q} \in \text{Spec}(B)} \varphi^{-1}(\mathfrak{q}) = \varphi^{-1} \left(\bigcap_{\mathfrak{q} \in \text{Spec}(B)} \mathfrak{q} \right) = \varphi^{-1}(\text{Rad}(B)).$$

In other words, $\mathbb{V}(\mathfrak{a})$ contains the image of f if and only if

$$\varphi(\mathfrak{a}) \subset \bigcap_{\mathfrak{q} \in \text{Spec}(B)} \mathfrak{q} = \text{Rad}(B).$$

Now, assume that f_φ is dominant. Then, for any ideal $\mathfrak{a}$ of A such that $\mathbb{V}(\mathfrak{a})$ contains the image of φ , (i.e., $\varphi(\mathfrak{a}) \subset \text{Rad}(B)$), we must have $\mathbb{V}(\mathfrak{a}) = \text{Spec}(A)$, i.e. $\mathfrak{a} \subset \text{Rad}(A)$. In particular, $\mathfrak{a} = \ker(\varphi)$ gives $\varphi(\mathfrak{a}) = (0) \subset \text{Rad}(B)$ so $\ker(\varphi) \subset \text{Rad}(A)$.

Conversely, assume $\ker(\varphi) \subset \text{Rad}(A)$. Let $\mathfrak{a}$ be an ideal of A such that $\mathbb{V}(\mathfrak{a})$ contains the image of φ , i.e., $\varphi(\mathfrak{a}) \subset \text{Rad}(B)$. We want to show that $\mathfrak{a} \subset \text{Rad}(A)$, so that $\mathbb{V}(\mathfrak{a}) = \text{Spec}(A)$. Pick $a \in \mathfrak{a}$. Then $\varphi(a) \in \text{Rad}(B)$, so there is $m \in \mathbb{N}$ such that $\varphi(a)^m = 0$. Hence $\varphi(a^m) = 0$, so that $a^m \in \ker(\varphi) \subset \text{Rad}(A)$. Thus, there must be an integer p such that $(a^m)^p = 0$. Hence $a^{pm} = 0$, so that $a \in \text{Rad}(A)$. In conclusion, $\mathfrak{a} \subset \text{Rad}(A)$, which finishes the proof.

Remark II.C.4. Dominance should not be checked only on maximal ideals in arbitrary rings. For example, the homomorphism

$$\gamma_{\mathbb{Q}} : \mathbb{Z} \hookrightarrow \mathbb{Q}$$

induces

$$\text{Spec}(\mathbb{Q}) \rightarrow \text{Spec}(\mathbb{Z}).$$

The image consists only of the generic point (0) , but this point is dense in $\text{Spec}(\mathbb{Z})$. Hence the morphism is dominant, although no closed point $(p) \in \text{Spec}(\mathbb{Z})$ lies in the image.

Remark II.C.5. If $\mathbb{k}$ is algebraically closed and A, B are finitely generated reduced $\mathbb{k}$ -algebras, then maximal ideals correspond to $\mathbb{k}$ -rational points by the Nullstellensatz. In this setting, a morphism

$$f : \text{Spec}(B) \rightarrow \text{Spec}(A)$$

is dominant if and only if the image on closed points is Zariski dense. Equivalently, for affine varieties over $\mathbb{k}$, a morphism is dominant if its image contains a dense subset of classical points.

Exercise II.C.6. Let $\mathbb{k}$ be a field and consider the inclusion

$$\varphi : \mathbb{k}[x] \hookrightarrow \mathbb{k}[x, x^{-1}].$$

Describe the image of the induced morphism $\text{Spec}(\mathbb{k}[x, x^{-1}]) \rightarrow \text{Spec}(\mathbb{k}[x])$. What is its image on closed points when $\mathbb{k}$ is algebraically closed?

Solution. The ring $\mathbb{k}[x, x^{-1}]$ is the localisation of $\mathbb{k}[x]$ at the element x . Therefore the morphism

$$\text{Spec}(\mathbb{k}[x, x^{-1}]) \rightarrow \text{Spec}(\mathbb{k}[x])$$

identifies $\text{Spec}(\mathbb{k}[x, x^{-1}])$ with the principal open subset

$$D(x) = \{\mathfrak{p} \in \text{Spec}(\mathbb{k}[x]) \mid x \notin \mathfrak{p}\}.$$

Thus the image is precisely $D(x)$. The morphism is dominant because the ring morphism $\mathbb{k}[x] \hookrightarrow \mathbb{k}[x, x^{-1}]$ is injective.

Assume now that $\mathbb{k}$ is algebraically closed. The closed points of $\text{Spec}(\mathbb{k}[x])$ are the maximal ideals

$$(x - a), \quad a \in \mathbb{k}.$$

The point $(x - a)$ belongs to $D(x)$ if and only if

$$x \notin (x - a).$$

This happens if and only if $a \neq 0$. Hence the image on closed points is

$$\{(x - a) \mid a \in \mathbb{k}, a \neq 0\}.$$

So the closed point (x) is missing, but the image is still dense in $\text{Spec}(\mathbb{k}[x])$.

Remark II.C.7. This example shows that a dominant morphism need not be surjective, even on closed points. Dominance only means that the image is dense.

II.D Open and principal open subsets

We define the Zariski topology on $\text{Spec}(A)$ using the sets $\mathbb{W}(\mathfrak{a})$, where $\mathfrak{a}$ is an ideal of A .

Definition II.D.1. We consider the topology on $\text{Spec}(A)$ whose open subsets are $D(\mathfrak{a}) := \text{Spec}(A) \setminus \mathbb{W}(\mathfrak{a})$, where $\mathfrak{a}$ is an ideal of A . For $f \in A$, we have the set $D(f) = \text{Spec}(A) \setminus \mathbb{W}(f)$. This is called a *principal open subset*. Let us spell out:

$$D(f) = \{\mathfrak{p} \in \text{Spec}(A) \mid f \notin \mathfrak{p}\}.$$

When f runs in A , the principal open subsets $D(f)$ form a base of the Zariski topology. Indeed, for any ideal $\mathfrak{a}$ of A we have

$$D(\mathfrak{a}) = \{\mathfrak{p} \in \text{Spec}(A) \mid \exists f \in \mathfrak{a} \setminus \mathfrak{p}\} = \bigcup_{f \in \mathfrak{a}} \{\mathfrak{p} \in \text{Spec}(A) \mid f \notin \mathfrak{p}\} = \bigcup_{f \in \mathfrak{a}} D(f).$$

Lemma II.D.2. Let $X = \text{Spec}(A)$ and consider a collection $(f_j \mid j \in J)$ of elements of A . The open subsets $(D(f_j) \mid j \in J)$ cover X if and only if $(f_j \mid j \in J)$ generate the unit ideal. Therefore, X is quasi-compact.

Proof. The subsets $(D(f_j) \mid j \in J)$ cover X if and only if, for any prime ideal $\mathfrak{p}$ of A , there is $j \in J$ such that f_j does not belong to $\mathfrak{p}$, i.e., no prime ideal $\mathfrak{p}$ of A contains all the f_j 's. Let $\mathfrak{a}$ be the ideal generated by $(f_j \mid j \in J)$. If $\mathfrak{a}$ is the unit ideal, then it is contained in no prime ideal, so no prime ideal contains all the f_j 's, and conversely, if $\mathfrak{a}$ is not the unit ideal then it is contained in some prime ideal $\mathfrak{p}$, which therefore contains all the f_j 's. So the first statement is proved.

About quasi-compactness, let $(U_i \mid i \in I)$ be an open cover of X and write $U_i = X \setminus \mathbb{W}(\mathfrak{a}_i)$ for some ideal $\mathfrak{a}_i$ of A , for all $i \in I$. Then we write $D(\mathfrak{a}_i) = \bigcup_{f \in \mathfrak{a}_i} D(f)$, so

$$X = \bigcup_{i \in I} \bigcup_{f \in \mathfrak{a}_i} D(f).$$

Therefore the ideal generated by all elements belonging to the ideals $\mathfrak{a}_i$ is the unit ideal. By the definition of an ideal generated by a set, there are finitely many elements $f_1, \dots, f_r$, with each f_j belonging to some $\mathfrak{a}_{i_j}$, and coefficients $u_1, \dots, u_r \in A$ such that

$$1 = u_1 f_1 + \dots + u_r f_r.$$

Hence $D(f_1), \dots, D(f_r)$ cover X , and therefore $U_{i_1}, \dots, U_{i_r}$ already cover X . Thus X is quasi-compact. $\square$

Exercise II.D.3. Let A be a Noetherian ring. Then every open subset of $\text{Spec}(A)$ is quasi-compact.

Solution. Since A is Noetherian, every ideal is finitely generated. Therefore every closed subset of $\text{Spec}(A)$ is of the form $\mathbb{V}(\mathfrak{a})$ with $\mathfrak{a}$ finitely generated. This implies that every descending chain of closed subsets stabilizes, so $\text{Spec}(A)$ is a Noetherian topological space.

In a Noetherian topological space, every open subset is quasi-compact. Indeed, if an open subset U had an open cover with no finite subcover, choose such a counterexample maximal among open subsets, or equivalently use the descending chain condition on closed complements. The standard Noetherian induction argument then gives a contradiction.

II.D.1 Localisation and principal open subsets

Let $f \in A$ and consider $D(f) = \{\mathfrak{p} \in \text{Spec}(A) \mid f \notin \mathfrak{p}\}$. Recall that the localised ring A_f is $A[f^{-1}] = S^{-1}A$ with $S = \{f^n \mid n \in \mathbb{N}\}$.

Exercise II.D.4. Show that, for all $f, g \in A$, we have $D(f) \cap D(g) = D(fg)$.

Solution. A prime ideal $\mathfrak{p}$ lies in $D(f) \cap D(g)$ if and only if $f \notin \mathfrak{p}$ and $g \notin \mathfrak{p}$. Since $\mathfrak{p}$ is prime, whenever $f \notin \mathfrak{p}$ and $g \notin \mathfrak{p}$, we have $fg \notin \mathfrak{p}$. Also, if $f \in \mathfrak{p}$ or $g \in \mathfrak{p}$, we have $fg \in \mathfrak{p}$.

Lemma II.D.5. Let $f, g \in A$. Then $D(f) \subset D(g)$ if and only if g divides a power of f , i.e. if and only if $f^m \in (g)$ for some $m \geq 1$. In this case there is a morphism $A_g \rightarrow A_f$, compatible with the localisation maps from A .

Proof. We have $D(f) \subset D(g)$ if and only if $\mathbb{V}(g) \subset \mathbb{V}(f)$. By Lemma II.B.1, this is equivalent to

$$\sqrt{(g)} \supset \sqrt{(f)}.$$

In turn, this means that $f^m \in (g)$ for some $m \geq 1$, i.e. that g divides a power of f .

If this is the case, write $f^m = gh$ for some $h \in A$. Then g becomes invertible in A_f , with inverse h/f^m . By the universal property of localisation, see Proposition III.B.1, the localisation map $A \rightarrow A_f$ therefore factors through a ring morphism

$$A_g \longrightarrow A_f.$$

By the uniqueness of the factorisation, the resulting composition $A \rightarrow A_g \rightarrow A_f$ agrees with the localisation $A \rightarrow A_f$. $\square$

Example II.D.6 (The double point and localisation). Consider the ring of dual numbers

$$A = \mathbb{k}[x]/(x^2)$$

and write $\bar{x}$ for the class of x in A . As we said, the affine scheme $X = \text{Spec}(A)$ has a single point, namely the prime ideal $\mathfrak{p} = (\bar{x})$. Indeed, prime ideals of A correspond to prime ideals of $\mathbb{k}[x]$ containing (x^2) , and any such prime ideal contains x .

Let us look at the principal open subsets of X . First,

$$D(0) = \{\mathfrak{p} \in \text{Spec}(A) \mid 0 \notin \mathfrak{p}\} = \emptyset,$$

because every ideal contains 0. Correspondingly,

$$A_0 = 0,$$

since localising at 0 forces 0 to become invertible, hence forces $1 = 0$. On the other hand,

$$D(\bar{x}) = \{\mathfrak{p} \in \operatorname{Spec}(A) \mid \bar{x} \notin \mathfrak{p}\} = \emptyset,$$

because the unique prime ideal $\mathfrak{p} = (\bar{x})$ contains $\bar{x}$. In terms of localisation, we have

$$A_{\bar{x}} = 0.$$

Indeed, in $A_{\bar{x}}$, the element $\bar{x}$ becomes invertible. But $\bar{x}^2 = 0$. Multiplying by $\bar{x}^{-2}$, we get $1 = 0$. Thus the localisation is the zero ring. Finally,

$$D(1) = X,$$

and

$$A_1 = A.$$

More generally, if $a \in A$, then $D(a) = X$ precisely when $a \notin (\bar{x})$, i.e. when a has non-zero constant term. In that case a is already invertible in A , so

$$A_a \simeq A.$$

If $a \in (\bar{x})$, then a is nilpotent, hence $A_a = 0$, and $D(a) = \emptyset$.

II.D.2 Residue field

Let A be a ring. We introduce a basic object attached to a point of $\operatorname{Spec}(A)$ given by a prime ideal $\mathfrak{p}$ of A , namely the fraction field of the integral domain $A/\mathfrak{p}$.

Definition II.D.7. Let $\mathfrak{p}$ be a prime ideal of A . Then the *residue field* $\kappa(\mathfrak{p})$ is defined as

$$\kappa(\mathfrak{p}) = \operatorname{Frac}(A/\mathfrak{p}).$$

Example II.D.8. For $A = \mathbb{k}[x]$, the residue field at the prime ideal (0) is

$$\kappa((0)) = \mathbb{k}(x),$$

while the residue field at $(x - a)$, for $a \in \mathbb{k}$, is

$$\kappa((x - a)) \simeq \mathbb{k}.$$

If $f \in \mathbb{k}[x]$ is irreducible, then the residue field at (f) is the field obtained by adjoining one root of f to $\mathbb{k}$. Indeed, since (f) is maximal,

$$\kappa((f)) = \mathbb{k}[x]/(f).$$

Lemma II.D.9. Let S be a multiplicative subset of A and $\mathfrak{a}$ be an ideal of A . Set T for the image of S in $B = A/\mathfrak{a}$. Then there is an isomorphism of rings

$$S^{-1}A/S^{-1}\mathfrak{a} \simeq T^{-1}B.$$

Proof. One first checks that T is a multiplicative subset of B . Then, we define a map $S^{-1}A \rightarrow T^{-1}B$ by sending a/s to $\bar{a}/\bar{s}$, where the canonical projection $A \rightarrow A/\mathfrak{a}$, a surjective morphism of rings, is denoted by $a \mapsto \bar{a}$. One verifies that this map is well-defined, for $a/s = b/t$ implies $u(at - bs) = 0$ for some $u \in S$ so that $\bar{u}(\bar{a}\bar{t} - \bar{b}\bar{s}) = \bar{0}$ with $\bar{u} \in T$ so that $\bar{a}/\bar{s} = \bar{b}/\bar{t}$.

The map $S^{-1}A \rightarrow T^{-1}B$ is surjective. An element a/s lies in its kernel if and only if $\bar{a}/\bar{s} = 0$ in $T^{-1}B$, namely if and only if there exists $u \in S$ such that $ua \in \mathfrak{a}$. This is exactly the condition that a/s belong to $S^{-1}\mathfrak{a}$. Hence the kernel is $S^{-1}\mathfrak{a}$. □

The previous lemma recovers Lemma III.D.2. Indeed, when $S = A \setminus \mathfrak{p}$, with $\mathfrak{p}$ a prime ideal of A , by definition $S^{-1}A = A_{\mathfrak{p}}$ and $S^{-1}\mathfrak{p} = \mathfrak{p}A_{\mathfrak{p}}$ so we get that $A_{\mathfrak{p}}/\mathfrak{p}A_{\mathfrak{p}}$ is isomorphic to the localisation of $A/\mathfrak{p}$ at its zero prime ideal, which is the fraction field of $A/\mathfrak{p}$, i.e. the residue field $\kappa(\mathfrak{p})$. This says, in particular, that $\mathfrak{p}A_{\mathfrak{p}}$ is a maximal ideal of $A_{\mathfrak{p}}$. But we know that more is true about $A_{\mathfrak{p}}$. Namely, $A_{\mathfrak{p}}$ is a local ring, with maximal ideal $\mathfrak{p}A_{\mathfrak{p}}$, as we saw in Proposition III.D.1.

Exercise II.D.10. Compute the residue fields of the following points.

- i) The points (0) and (p) of $\text{Spec}(\mathbb{Z})$, where p is a prime number.
- ii) The point (f) of $\text{Spec}(\mathbb{F}_p[t])$, where $f \in \mathbb{F}_p[t]$ is irreducible of degree d . Are all these residue fields isomorphic?
- iii) The points (p) and $(p, \bar{f})$ of $\text{Spec}(\mathbb{Z}[t])$, where p is a prime number and $\bar{f} \in \mathbb{F}_p[t]$ is irreducible.

Solution. Recall that, for a prime ideal $\mathfrak{p} \subset A$, the residue field is

$$\kappa(\mathfrak{p}) = \text{Frac}(A/\mathfrak{p}).$$

- i) In $\text{Spec}(\mathbb{Z})$, the ideal (0) is prime and

$$\kappa((0)) = \text{Frac}(\mathbb{Z}) = \mathbb{Q}.$$

If p is a prime number, then (p) is maximal and

$$\kappa((p)) = \text{Frac}(\mathbb{Z}/(p)) = \mathbb{F}_p.$$

- ii) In $\text{Spec}(\mathbb{F}_p[t])$, the ideal (0) is prime and

$$\kappa((0)) = \text{Frac}(\mathbb{F}_p[t]) = \mathbb{F}_p(t).$$

If $f \in \mathbb{F}_p[t]$ is irreducible, then (f) is maximal and

$$\kappa((f)) = \text{Frac}(\mathbb{F}_p[t]/(f)) = \mathbb{F}_p[t]/(f).$$

This is a finite extension of $\mathbb{F}_p$ of degree $\deg(f)$.

- iii) In $\text{Spec}(\mathbb{Z}[t])$, the ideal (p) is prime because

$$\mathbb{Z}[t]/(p) \simeq \mathbb{F}_p[t],$$

which is an integral domain. Therefore

$$\kappa((p)) = \text{Frac}(\mathbb{Z}[t]/(p)) = \text{Frac}(\mathbb{F}_p[t]) = \mathbb{F}_p(t).$$

Now let $\bar{f} \in \mathbb{F}_p[t]$ be irreducible and consider the ideal

$$\mathfrak{p} = (p, \bar{f}) \subset \mathbb{Z}[t],$$

where, more precisely, we choose a lift $f \in \mathbb{Z}[t]$ of $\bar{f}$ and write

$$\mathfrak{p} = (p, f).$$

Then

$$\mathbb{Z}[t]/(p, f) \simeq \mathbb{F}_p[t]/(\bar{f}),$$

which is a field. Hence $\mathfrak{p}$ is maximal and

$$\kappa(\mathfrak{p}) = \mathbb{F}_p[t]/(\bar{f}).$$

II.D.3 Localisation and homeomorphism

We study how prime ideals and localisation interact, proving that the map on spectra is a homeomorphism onto its image. Let S be a multiplicative subset of a ring A and let $\varphi : A \rightarrow B$ be the localisation map, with $B = S^{-1}A$. Recall that the map $f_\varphi : \text{Spec}(B) \rightarrow \text{Spec}(A)$ sends a prime ideal $\mathfrak{q}$ of B to $\varphi^{-1}(\mathfrak{q})$, a prime ideal of A . Put

$$D(S) = \{\mathfrak{p} \in \text{Spec}(A) \mid \mathfrak{p} \cap S = \emptyset\}.$$

Proposition II.D.11. *The map $f_\varphi : \text{Spec}(B) \rightarrow \text{Spec}(A)$ is injective and gives a homeomorphism $\text{Spec}(B) \simeq D(S)$. In particular, $\text{Spec}(A_f)$ is homeomorphic to $D(f)$.*

Proof. Let $\mathfrak{b}$ be a prime ideal of $S^{-1}A$ and set $\mathfrak{a} = \varphi^{-1}(\mathfrak{b})$. By Lemma III.B.7 in Chapter 1, we have $\mathfrak{b} = S^{-1}\mathfrak{a}$, so $\mathfrak{b}$ is determined by its inverse image. This proves injectivity.

Now we apply Lemma III.B.6. Since $\mathfrak{b}$ is prime, $\varphi^{-1}(\mathfrak{b})$ is a prime ideal of A that does not intersect S , and conversely, if $\mathfrak{a} \in \text{Spec}(A)$ and $\mathfrak{a} \cap S = \emptyset$, then $S^{-1}\mathfrak{a}$ is a prime ideal of $S^{-1}A$. These constructions are inverse to each other, so the image of f_φ is $D(S)$.

It remains to check the topology. The map f_φ is continuous, so we only need to check that it also closed. Consider an ideal $\mathfrak{b} \subset B$ and look at $\varphi^{-1}(\mathfrak{b}) \subset A$. Then we claim

$$f_\varphi(\mathbb{W}(\mathfrak{b})) = \mathbb{W}(\varphi^{-1}(\mathfrak{b})) \cap D(S).$$

Indeed, a prime ideal $\mathfrak{p}$ is written as $\mathfrak{p} = \varphi^{-1}(\mathfrak{q})$, for some prime ideal $\mathfrak{q}$ of B , if and only if $\mathfrak{p} \cap S = \emptyset$ and $\mathfrak{q} = S^{-1}(\mathfrak{p})$. Also, if $\mathfrak{q} = S^{-1}(\mathfrak{p})$ contains $\mathfrak{b}$, then applying φ^{-1} we get, again by Lemma III.B.6 in Chapter 1:

$$\mathfrak{p} = \varphi^{-1}(S^{-1}(\mathfrak{p})) \supset \varphi^{-1}(\mathfrak{b}).$$

Conversely, if $\mathfrak{p}$ contains $\varphi^{-1}(\mathfrak{b})$, then by Lemma III.B.7 in Chapter 1, we get

$$\mathfrak{q} = S^{-1}(\mathfrak{p}) \supset S^{-1}(\varphi^{-1}(\mathfrak{b})) = \mathfrak{b}.$$

In conclusion we proved:

$$\begin{aligned} f_\varphi(\mathbb{W}(\mathfrak{b})) &= \{\varphi^{-1}(\mathfrak{q}) \mid \mathfrak{q} \in \text{Spec}(B), \mathfrak{q} \supset \mathfrak{b}\} = \\ &= \{\mathfrak{p} \in \text{Spec}(A) \mid \mathfrak{p} \cap S = \emptyset, S^{-1}(\mathfrak{p}) \supset \mathfrak{b}\} = \mathbb{W}(\varphi^{-1}(\mathfrak{b})) \cap D(S). \end{aligned}$$

This shows that f_φ is a homeomorphism onto its image. Taking $S = \{f^m \mid m \in \mathbb{N}\}$ gives the homeomorphism $\text{Spec}(A_f) \simeq D(f)$. $\square$

Exercise II.D.12. Let $A = \mathbb{k}[x]$. Describe the points of $\text{Spec}(A_f)$ for

$$f = x, \quad f = x(x-1), \quad f = (x-a)^m,$$

where $a \in \mathbb{k}$ and $m \geq 1$.

Solution. By the localisation theorem for spectra, the prime ideals of A_f are in bijection with the prime ideals $\mathfrak{p} \subset A$ such that

$$\mathfrak{p} \cap \{1, f, f^2, \dots\} = \emptyset.$$

This is equivalent to saying that $f \notin \mathfrak{p}$. Hence

$$\text{Spec}(A_f) \simeq D(f) = \{\mathfrak{p} \in \text{Spec}(A) \mid f \notin \mathfrak{p}\}.$$

Now take $A = \mathbb{k}[x]$. The prime ideals of $\mathbb{k}[x]$ are (0) and the ideals (p) , where $p \in \mathbb{k}[x]$ is irreducible.

If $f = x$, then $\text{Spec}(A_x)$ corresponds to the points

$$(0), \quad (p) \text{ with } p \neq x.$$

Thus we remove the closed point (x) .

If $f = x(x-1)$, then $\text{Spec}(A_f)$ corresponds to the points

$$(0), \quad (p) \text{ with } p \neq x, x-1.$$

Thus we remove the two closed points (x) and $(x-1)$.

Finally, if $f = (x-a)^m$, then

$$D((x-a)^m) = D(x-a).$$

Therefore $\text{Spec}(A_f)$ corresponds to the points

$$(0), \quad (p) \text{ with } p \neq x-a.$$

So we remove only the closed point $(x-a)$.

Definition II.D.13. A *discrete valuation ring*, or DVR, is a principal ideal domain A which has a unique non-zero prime ideal. Equivalently, it is a local principal ideal domain which is not a field. If A is a DVR, its unique maximal ideal is principal. A generator π of this maximal ideal is called a *uniformiser*.

Exercise II.D.14. Let A be a discrete valuation ring, with uniformiser π and fraction field K . Show

$$\text{Spec}(A) = \{(0), (\pi)\}.$$

Which point is closed? What is the closure of the point (0) ?

Solution. Since A is a principal ideal domain, every non-zero prime ideal is generated by an irreducible element. Since A is a DVR, it has a unique non-zero prime ideal, namely its maximal ideal (π) . Therefore the only prime ideals of A are

$$(0), \quad (\pi).$$

Hence

$$\text{Spec}(A) = \{(0), (\pi)\}.$$

The point (π) is closed because (π) is maximal. The point (0) is the generic point. Its closure is

$$\overline{\{(0)\}} = \mathbb{V}(0) = \text{Spec}(A).$$

Thus $\text{Spec}(A)$ has two points: a generic point and a closed point.

III The structure sheaf

Further reading. The structure sheaf of an affine scheme is treated in [Har77, Chapter II, §2] and [Liu02, Chapter 2]; [EH00] emphasises the geometric meaning through examples.

We construct the structure sheaf on $X = \text{Spec}(A)$ and show that affine spectra are naturally locally ringed spaces.

III.A The principal-open description and associated modules

One way to define the structure sheaf on $X = \operatorname{Spec}(A)$ is by prescribing its values on the basis of principal open subsets. This goes by setting, for all $f \in A$:

$$\mathcal{O}_X(D(f)) \simeq A_f.$$

We now verify the sheaf condition directly on this basis. This is needed to justify that the local sections we just gave glue to define a sheaf on X . Also, this is needed to extend the construction from rings to modules.

The principal open subsets $(D(f) \mid f \in A)$ form a basis of the Zariski topology. An inclusion $D(f) \subset D(g)$ induces the compatible localisation map $A_g \rightarrow A_f$. Having this in mind, we state the following result.

Lemma III.A.1. *Let F be an A -module. For all $f \in A$, put $\mathcal{F}(D(f)) = F_f$. Then $\mathcal{F}$ satisfies the sheaf condition on open coverings of $D(f)$ by principal open subsets.*

Since the principal open sets of the form $D(f)$, with $f \in A$, form a basis of the Zariski topology on $X = \operatorname{Spec}(A)$, according to the previous lemma, for every module F , we get a sheaf $\mathcal{F}$ on X such that for all $f \in A$, $\mathcal{F}(D(f)) = F_f$.

Definition III.A.2. The sheaf $\mathcal{F}$ associated with $F = A$ according to the previous lemma is a sheaf of rings on X , called the *structure sheaf* and denoted by $\mathcal{O}_X$. For any A -module F , the sheaf $\mathcal{F}$ associated with F is a sheaf of $\mathcal{O}_X$ -modules, and it is denoted by $\tilde{F}$.

Proof of Lemma III.A.1. Replacing A by A_f , it is enough to check the sheaf condition for a covering of $X = \operatorname{Spec}(A)$ by principal open subsets. By quasi-compactness, we may assume that the covering is finite, say

$$X = \bigcup_{1 \leq i \leq \ell} D(f_i).$$

Equivalently, the elements $f_1, \dots, f_\ell$ generate the unit ideal.

First, let $v \in F$ be a section whose image in every F_{f_i} is zero. Then, for each i , there is an integer $n_i \geq 0$ such that $f_i^{n_i} v = 0$ in F . Since the opens $D(f_i^{n_i}) = D(f_i)$ still cover X , the elements $f_i^{n_i}$ generate the unit ideal. We can write

$$1 = \sum_{1 \leq i \leq \ell} a_i f_i^{n_i}$$

for suitable $a_i \in A$, $1 \leq i \leq \ell$. Multiplying by v , we get $v = 0$. This proves the locality condition.

Now let $v_i \in F_{f_i}$ be sections which agree on the overlaps, i.e. whose images in $F_{f_i f_j}$ coincide for all i, j . For $1 \leq i \leq \ell$, there are $y_i \in F$ and $r_i \in \mathbb{N}$ such that

$$v_i = \frac{y_i}{f_i^{r_i}}$$

Let $r = \max\{r_1, \dots, r_\ell\}$. Then, for $1 \leq i \leq \ell$ we let $x_i = y_i f_i^{r-r_i}$ and rewrite v_i as

$$v_i = \frac{y_i f_i^{r-r_i}}{f_i^r} = \frac{x_i}{f_i^r}.$$

Since v_i and v_j agree in $F_{f_i f_j}$, there is an integer $n_{ij} \geq 0$ such that

$$(f_i f_j)^{n_{ij}} (f_j^r x_i - f_i^r x_j) = 0. \quad (\text{III.1})$$

Choose an integer N such that $N \geq r + n_{i,j}$ for all i, j . Since the opens $D(f_i^N)$ still cover X , we can write

$$1 = \sum_{1 \leq i \leq \ell} c_i f_i^N$$

for some $c_i \in A$. Define

$$v = \sum_{1 \leq i \leq \ell} c_i f_i^{N-r} x_i \in F.$$

Now, fix an integer k , with $1 \leq k \leq \ell$. Then we compute

$$f_k^T v - x_k = \sum_{1 \leq i \leq \ell} c_i f_i^{N-r} (f_k^T x_i - f_i^T x_k).$$

Multiplying by f_k^{N-r} gives:

$$f_k^{N-r} (f_k^T v - x_k) = \sum_{1 \leq i \leq \ell} c_i (f_i f_k)^{N-r} (f_k^T x_i - f_i^T x_k) = 0,$$

by virtue of (III.1), because $N - r \geq n_{i,k}$ for $1 \leq i, k \leq \ell$. Hence v maps to $x_k / f_k^r = v_k$ in F_{f_k} . This proves the existence of a gluing section, and hence the sheaf condition on the basis. $\square$

Example III.A.3. If $\mathfrak{a} \subset A$ is an ideal, then $\widetilde{\mathfrak{a}}$ is a sheaf of ideals of $\mathcal{O}_X$. The quotient map $A \rightarrow A/\mathfrak{a}$ sheafifies to an exact sequence

$$0 \rightarrow \widetilde{\mathfrak{a}} \rightarrow \mathcal{O}_X \rightarrow \widetilde{A/\mathfrak{a}} \rightarrow 0.$$

III.B Equivalent definition by sheafification

Let A be a ring and put $X = \text{Spec}(A)$. Let us check that our definition of the sheaf of regular functions is the sheafification of a presheaf, easily described by localisation. For an open subset $U \subset X$, define

$$S_U := \bigcap_{\mathfrak{p} \in U} (A \setminus \mathfrak{p}) = A \setminus \bigcup_{\mathfrak{p} \in U} \mathfrak{p}.$$

Then S_U is a multiplicative subset of A . If $U \subset U'$, then $S_{U'} \subset S_U$. This induces a ring homomorphism

$$S_{U'}^{-1} A \rightarrow S_U^{-1} A.$$

Thus $U \mapsto S_U^{-1} A$ defines a presheaf of rings on the topological space X , denoted by ${}^{\text{pre}}\mathcal{O}_X$. The sheaf of rings associated with ${}^{\text{pre}}\mathcal{O}_X$ is precisely $\mathcal{O}_X$. This is the content of the next result, whose upshot is that the definition by sheafification agrees with the construction on the basis of principal open subsets.

Proposition III.B.1. *For every $f \in A$, there are natural isomorphisms*

$$S_{D(f)}^{-1} A \simeq A_f, \quad \mathcal{O}_X(D(f)) \simeq A_f.$$

Proof. Since f belongs to $S_{D(f)}$, there is a natural map $A_f \rightarrow S_{D(f)}^{-1} A$. Conversely, if $g \in S_{D(f)}$, then $D(f) \subset D(g)$. By Lemma II.D.5, some power of f lies in (g) , so g is invertible in A_f . The universal property of localisation gives a map $S_{D(f)}^{-1} A \rightarrow A_f$. The two maps are mutually inverse. For the compatibility with sheafification, use Lemma III.A.1: the assignment $D(f) \mapsto A_f$ satisfies the sheaf condition on the principal-open basis. The isomorphisms just constructed identify the restriction of ${}^{\text{pre}}\mathcal{O}_X$ to that basis with this assignment. Hence its sheafification has $\mathcal{O}_X(D(f)) \simeq A_f$. $\square$

Exercise III.B.2. For an open subset $U \subset X$, describe the elements of $\mathcal{O}_X(U)$ using Lemma IV.D.1.

Solution. A section of $\mathcal{O}_X(U)$ is a function

$$s : U \rightarrow \prod_{\mathfrak{p} \in U} A_{\mathfrak{p}}$$

such that $s(\mathfrak{p}) \in A_{\mathfrak{p}}$ for every $\mathfrak{p} \in U$, and which is locally given by a fraction. More precisely, for every $\mathfrak{p} \in U$, there is an open neighbourhood $V \subset U$ of $\mathfrak{p}$ and elements $a, t \in A$, with $t \notin \mathfrak{q}$ for all $\mathfrak{q} \in V$, such that

$$s(\mathfrak{q}) = \frac{a}{t} \in A_{\mathfrak{q}} \quad \text{for all } \mathfrak{q} \in V.$$

In other words, regular functions on U are functions on prime ideals which are locally represented by fractions with denominators not vanishing on the neighbourhood under consideration.

Example III.B.3. Let $A = \mathbb{k}[x]$ and $X = \text{Spec}(A)$. For the principal open subset

$$D(x) = X \setminus \mathbb{W}(x),$$

we have

$$\mathcal{O}_X(D(x)) \simeq A_x = \mathbb{k}[x, x^{-1}].$$

Thus the function x^{-1} is a regular function on $D(x)$, although it is not a regular function on all of X .

Exercise III.B.4. Let $A = \mathbb{k}[x]$ and $X = \text{Spec}(A)$. Compute

$$\mathcal{O}_X(D(x)), \quad \mathcal{O}_X(D(x-a)), \quad \mathcal{O}_X(D(x(x-1))).$$

Solution. For every $f \in A$, we have

$$\mathcal{O}_X(D(f)) \simeq A_f.$$

Therefore

$$\mathcal{O}_X(D(x)) \simeq \mathbb{k}[x, x^{-1}],$$

and

$$\mathcal{O}_X(D(x-a)) \simeq \mathbb{k}[x, (x-a)^{-1}].$$

Finally,

$$\mathcal{O}_X(D(x(x-1))) \simeq \mathbb{k}[x, (x(x-1))^{-1}].$$

Equivalently, this is the subring of $\mathbb{k}(x)$ consisting of rational functions whose poles are allowed only at 0 and 1.

Exercise III.B.5. Let

$$A = \mathbb{k}[x]/(x^2)$$

and let $X = \text{Spec}(A)$.

- i) Describe the points of X .
- ii) Describe the open subsets of X .
- iii) Compute the residue field $\kappa(\mathfrak{p})$ at each point.

Solution. The prime ideals of $A = \mathbb{k}[x]/(x^2)$ correspond to the prime ideals of $\mathbb{k}[x]$ containing (x^2) . Such a prime ideal must contain x , hence the only prime ideal is

$$\mathfrak{p} = (\bar{x}).$$

Therefore X has a single point. Since X has only one point, its open subsets are

$$\emptyset \quad \text{and} \quad X.$$

Finally, the residue field is

$$\kappa(\mathfrak{p}) = \text{Frac}(A/\mathfrak{p}).$$

Since

$$A/\mathfrak{p} \simeq \mathbb{k},$$

we get

$$\kappa(\mathfrak{p}) \simeq \mathbb{k}.$$

Remark III.B.6. The schemes

$$\text{Spec}(\mathbb{k}) \quad \text{and} \quad \text{Spec}(\mathbb{k}[x]/(x^2))$$

have the same underlying topological space: a single point. However, they are not isomorphic as schemes, because their local rings are different. In $\mathbb{k}[x]/(x^2)$, the class of x is a non-zero nilpotent element.

Exercise III.B.7. Let $A = \mathbb{k}[x, y]$ and $X = \text{Spec}(A)$. Compute

$$\mathcal{O}_X(D(x)), \quad \mathcal{O}_X(D(xy)).$$

Assume that $\mathbb{k}$ is algebraically closed. What do these open subsets look like on closed points?

Solution. We have

$$\mathcal{O}_X(D(x)) \simeq A_x = \mathbb{k}[x, y, x^{-1}].$$

Similarly,

$$\mathcal{O}_X(D(xy)) \simeq A_{xy} = \mathbb{k}[x, y, (xy)^{-1}].$$

Since inverting xy is the same as inverting both x and y , we also have

$$A_{xy} \simeq \mathbb{k}[x, y, x^{-1}, y^{-1}].$$

If $\mathbb{k}$ is algebraically closed, the closed points of X are the points $(a, b) \in \mathbb{k}^2$. The open subset $D(x)$ consists of the closed points with $a \neq 0$. The open subset $D(xy) = D(x) \cap D(y)$ consists of the closed points with

$$a \neq 0, \quad b \neq 0.$$

Thus $D(x)$ is the affine plane with the line $x = 0$ removed, and $D(xy)$ is the affine plane with both coordinate axes removed.

Example III.B.8 (Sheafification is needed to define the structure sheaf). Assume that $\mathbb{k}$ is algebraically closed. Let

$$A = \mathbb{k}[x, y, z, w]/(xw - yz), \quad X = \text{Spec}(A).$$

We want to see that ${}^{\text{pre}}\mathcal{O}_X$ is not the same as $\mathcal{O}_X$.

We still denote by x, y, z, w the images of the variables in A . Consider the prime ideal $\mathfrak{p} = (x, y) \subset A$ and the open subset $U = X \setminus V(\mathfrak{p})$. Since $V(\mathfrak{p}) = V(x, y)$, we have

$$U = D(x) \cup D(y).$$

We are brought to consider

$$S_U = \{f \in A \mid U \subset D(f)\}.$$

Let us show that

$$S_U = \mathbb{k} \setminus \{0\}.$$

By definition, if $f \in S_U$, then $U \subset D(f)$, or equivalently

$$\mathbb{W}(f) \subset X \setminus U = \mathbb{W}(x, y).$$

We claim that this is impossible for a non-unit f . To see this, use a well-known parametrisation of the quadric cone:

$$A \simeq \mathbb{k}[ac, ad, bc, bd] \subset \mathbb{k}[a, b, c, d],$$

where

$$x = ac, \quad y = ad, \quad z = bc, \quad w = bd.$$

Suppose that $f \in A$ satisfies

$$\mathbb{W}(f) \subset \mathbb{W}(x, y).$$

Let F be its image in $\mathbb{k}[a, b, c, d]$. Pulling back the inclusion above to $\text{Spec } \mathbb{k}[a, b, c, d]$, we get

$$\mathbb{W}(F) \subset \mathbb{W}(ac, ad).$$

Thus

$$ac, ad \in \sqrt{(F)}.$$

Let h be an irreducible factor of F . Since $ac, ad \in \sqrt{(F)}$, the element h divides both ac and ad . As c and d are relatively prime, this forces

$$h \mid a.$$

Therefore every irreducible factor of F is associated with a , and hence

$$F = \lambda a^n$$

for some $\lambda \in \mathbb{k}^\times$ and some $n \geq 0$. But every element of

$$\mathbb{k}[ac, ad, bc, bd]$$

is a sum of monomials having the same total degree in a, b as in c, d . The monomial a^n has bidegree $(n, 0)$, so it belongs to $\mathbb{k}[ac, ad, bc, bd]$ only if $n = 0$. Thus F is constant, and so f is a unit. This proves

$$S_U = \mathbb{k} \setminus \{0\}.$$

Consequently,

$${}^{\text{pre}}\mathcal{O}_X(U) = S_U^{-1}A = A.$$

Now we are in position to show that ${}^{\text{pre}}\mathcal{O}_X(U) = S_U^{-1}A$ is not the same as $\mathcal{O}_X(U)$. On $D(x)$ and $D(y)$, respectively, we have the sections

$$\frac{z}{x} \in A_x, \quad \frac{w}{y} \in A_y.$$

These two sections agree on the intersection

$$D(x) \cap D(y) = D(xy).$$

Indeed, in A we have the relation $xw = yz$, and therefore, in A_{xy} , we obtain

$$\frac{z}{x} = \frac{w}{y}$$

Hence the sheaf axiom for $\mathcal{O}_X$ gives a section $s \in \mathcal{O}_X(U)$ such that

$$s|_{D(x)} = \frac{z}{x}, \quad s|_{D(y)} = \frac{w}{y}.$$

We now show that this section does not come from ${}^{\text{pre}}\mathcal{O}_X(U) = A$. Indeed, if $s \in A$, then $z = xs$ in A . Under the embedding $A \subset \mathbb{k}[a, b, c, d]$, this would give $bc = ac \cdot s$. Since $\mathbb{k}[a, b, c, d]$ is a domain, we could cancel c and obtain

$$b = a \cdot s,$$

which is impossible in the polynomial ring $\mathbb{k}[a, b, c, d]$. In conclusion, we proved that

$$s \in \mathcal{O}_X(U) \quad \text{but} \quad s \notin {}^{\text{pre}}\mathcal{O}_X(U).$$

Hence the presheaf ${}^{\text{pre}}\mathcal{O}_X$ is not a sheaf.

Remark III.B.9. The point of the example is that $\mathcal{O}_X$ allows denominators locally. On $D(x)$ the section is written as z/x , while on $D(y)$ it is written as w/y . These two local expressions agree on the overlap and therefore glue. The presheaf ${}^{\text{pre}}\mathcal{O}_X$ asks instead for a single denominator which is invertible on all of U . In this example no non-unit of A is invertible on all of U , so the presheaf only gives A on U and misses the glued section.

IV Affine schemes as locally ringed spaces

We define a category of spaces, called locally ringed spaces, where schemes live. A locally ringed space is a topological space equipped with a sheaf of rings such that the stalk at each point is a local ring. Morphisms in this category are defined in a subtle way, in that they should combine continuous maps of topological spaces and a ring homomorphism on local sections of the structure sheaves, which respects the locality condition about maximal ideals on stalks. Schemes are locally ringed spaces which are locally isomorphic to the spectrum of a ring.

IV.A The affine spectrum as a locally ringed space

Here we see how to identify the category of rings with a full subcategory of locally ringed spaces.

IV.A.1 The stalk of the structure sheaf and the local ring

Let A be a ring and let $\mathfrak{p} \in X = \text{Spec}(A)$. Let us identify the stalk of the structure sheaf $\mathcal{O}_X$ at a point $\mathfrak{p}$ with the local ring $A_{\mathfrak{p}}$.

Proposition IV.A.1. *Let A be a ring and let $\mathfrak{p} \in \text{Spec}(A)$. For every $f \notin \mathfrak{p}$, consider the localisation A_f . The restriction homomorphisms associated with inclusions of principal neighbourhoods of $\mathfrak{p}$ form a filtered direct system, and there is a natural isomorphism*

$$\varinjlim_{f \notin \mathfrak{p}} A_f \simeq A_{\mathfrak{p}}.$$

Proof. For every $f \notin \mathfrak{p}$, the image of f is invertible in $A_{\mathfrak{p}}$. By the universal property of localisation, there is therefore a unique homomorphism $A_f \rightarrow A_{\mathfrak{p}}$ compatible with the homomorphism $A \rightarrow A_{\mathfrak{p}}$. These homomorphisms are compatible with the restriction maps. Indeed, if $D(g) \subset D(f)$ and $\mathfrak{p} \in D(g)$, the diagram

$$\begin{array}{ccc} A_f & \xrightarrow{\quad} & A_g \\ & \searrow & \swarrow \\ & A_{\mathfrak{p}} & \end{array}$$

commutes by the uniqueness part of the universal property of localisation. Hence, these morphisms induce a homomorphism

$$\Phi : \varinjlim_{f \notin \mathfrak{p}} A_f \rightarrow A_{\mathfrak{p}}.$$

We first show that Φ is surjective. Every element of $A_{\mathfrak{p}}$ may be written as

$$\frac{a}{s}, \quad a \in A, \quad s \notin \mathfrak{p}.$$

But a/s is already an element of A_s , and its image under $A_s \rightarrow A_{\mathfrak{p}}$ is the given element. Hence Φ is surjective. We now prove injectivity. Let an element of the direct limit be represented by

$$\frac{a}{f^n} \in A_f, \quad f \notin \mathfrak{p},$$

and suppose that its image in $A_{\mathfrak{p}}$ is zero. By the definition of localisation, there exists $t \notin \mathfrak{p}$ such that $ta = 0$ in A . Consider the smaller principal neighbourhood $D(ft) = D(f) \cap D(t)$ of $\mathfrak{p}$. Under the restriction homomorphism $A_f \rightarrow A_{ft}$, the element t becomes invertible. Consequently,

$$\frac{a}{f^n} = \frac{ta}{tf^n} = 0$$

in A_{ft} . Thus the original element becomes zero at a later stage of the direct system, and hence represents zero in the direct limit. Therefore Φ is injective. $\square$

Corollary IV.A.2. *Let $X = \operatorname{Spec}(A)$ and let $\mathfrak{p} \in X$. Then there is a natural isomorphism*

$$\mathcal{O}_{X,\mathfrak{p}} \simeq A_{\mathfrak{p}}.$$

Proof. The principal open subsets $D(f)$ with $f \notin \mathfrak{p}$ form a basis of neighbourhoods of $\mathfrak{p}$. Since

$$\mathcal{O}_X(D(f)) = A_f,$$

the definition of the stalk gives

$$\mathcal{O}_{X,\mathfrak{p}} = \varinjlim_{\mathfrak{p} \in U} \mathcal{O}_X(U) \simeq \varinjlim_{f \notin \mathfrak{p}} A_f.$$

The conclusion now follows from Proposition IV.A.1. $\square$

Exercise IV.A.3. Let $A = \mathbb{k}[x]$ and $X = \operatorname{Spec}(A)$. For $a \in \mathbb{k}$, compute the stalks

$$\mathcal{O}_{X,(0)}, \quad \mathcal{O}_{X,(x-a)}$$

Solution. For every prime ideal $\mathfrak{p} \subset A$, we have

$$\mathcal{O}_{X,\mathfrak{p}} \simeq A_{\mathfrak{p}}.$$

Therefore

$$\mathcal{O}_{X,(0)} \simeq A_{(0)} = \mathbb{k}(x),$$

the rational function field of the affine line.

On the other hand,

$$\mathcal{O}_{X,(x-a)} \simeq A_{(x-a)} = \left\{ \frac{f}{g} \in \mathbb{k}(x) \mid g(a) \neq 0 \right\}.$$

This is the local ring of rational functions which are regular at the point a .

IV.A.2 Ringed spaces and locally ringed spaces

Definition IV.A.4. A *ringed space* is a topological space equipped with a sheaf of rings. A *locally ringed space* is a ringed space such that all stalks of its sheaf of rings are local rings.

Given a ringed space $(X, \mathcal{O}_X)$, we often write just X , meaning that X is silently equipped with some sheaf of rings, usually denoted by $\mathcal{O}_X$. In case we want to refer to X as the topological space only, without the datum of the structure sheaf, sometimes we write $|X|$. Given a point x of the ringed space X , we write $\mathcal{O}_{X,x}$ for the stalk at x of the sheaf of rings on X , so by definition,

$$\mathcal{O}_{X,x} = \varinjlim_{U \ni x} \mathcal{O}_X(U).$$

Lemma IV.A.5. For any ring A , the pair $X = (\text{Spec}(A), \mathcal{O}_X)$ is a locally ringed space.

Proof. By Corollary IV.A.2, the stalk of $\mathcal{O}_X$ at a prime ideal $\mathfrak{p}$ is naturally isomorphic to $A_{\mathfrak{p}}$. This ring is local, with maximal ideal $\mathfrak{p}A_{\mathfrak{p}}$, by Proposition III.D.1. Hence all stalks of $\mathcal{O}_X$ are local rings. $\square$

Exercise IV.A.6. Let X be either a smooth manifold, with sheaf $\mathcal{C}_X^\infty$, or a complex manifold, with sheaf $\mathcal{O}_X$. Show that the stalk at a point $x \in X$ is a local ring. Describe its maximal ideal and its residue field.

Solution. In the smooth case, the stalk $\mathcal{C}_{X,x}^\infty$ is the ring of germs of smooth real-valued functions at x . Evaluation at x gives a surjective morphism

$$\mathcal{C}_{X,x}^\infty \rightarrow \mathbb{R}.$$

Its kernel is the ideal of germs vanishing at x :

$$\mathfrak{m}_x = \{f \in \mathcal{C}_{X,x}^\infty \mid f(x) = 0\}.$$

A germ f is invertible if and only if $f(x) \neq 0$, because then $1/f$ is defined and smooth near x . Hence $\mathfrak{m}_x$ is the unique maximal ideal, and

$$\mathcal{C}_{X,x}^\infty / \mathfrak{m}_x \simeq \mathbb{R}.$$

In the complex case, the same argument applies with holomorphic functions. The maximal ideal of $\mathcal{O}_{X,x}$ is

$$\mathfrak{m}_x = \{f \in \mathcal{O}_{X,x} \mid f(x) = 0\},$$

and the residue field is

$$\mathcal{O}_{X,x} / \mathfrak{m}_x \simeq \mathbb{C}.$$

IV.B Morphisms of locally ringed spaces

Given a continuous map $f : X \rightarrow Y$, where $(X, \mathcal{O}_X)$ and $(Y, \mathcal{O}_Y)$ are ringed spaces, recall that we have the direct image sheaf on Y , whose sections on an open subset U of Y are given by:

$$f_*(\mathcal{O}_X) : U \mapsto \mathcal{O}_X(f^{-1}(U)).$$

Let $x \in X$ and set $y = f(x)$. Then, there is a map

$$f_*(\mathcal{O}_X)_y \rightarrow \mathcal{O}_{X,x}. \quad (\text{IV.1})$$

Indeed, by definition, we have

$$f_*(\mathcal{O}_X)_y = \varinjlim_{V \ni y} f_*(\mathcal{O}_X(V)) = \varinjlim_{V \ni y} \mathcal{O}_X(f^{-1}(V)). \quad (\text{IV.2})$$

Then, for any open subset $V \subset Y$ with $y \in V$, we have that $U = f^{-1}(V)$ is an open subset of X containing x . Thus, $\mathcal{O}_X(f^{-1}(V))$ maps to $\mathcal{O}_{X,x}$. Hence we get maps $\mathcal{O}_X(f^{-1}(V)) \rightarrow \mathcal{O}_{X,x}$, which are compatible with the maps induced by inclusions of the form $V \subset V' \ni y$. This gives (IV.1).

To relate the structure sheaves of X and Y , we use a morphism $f^\# : \mathcal{O}_Y \rightarrow f_*(\mathcal{O}_X)$. In other words, for each open subset V of Y , we need a morphism of rings:

$$f_V^\# : \mathcal{O}_Y(V) \rightarrow \mathcal{O}_X(f^{-1}(V)).$$

We need also that these ring homomorphisms are compatible with restrictions, namely, for $V \subset U$ open subsets of Y , we need the next diagram to commute

$$\begin{array}{ccc} \mathcal{O}_Y(U) & \xrightarrow{f_U^\#} & \mathcal{O}_X(f^{-1}(U)) \\ \rho_{U,V} \downarrow & & \downarrow \rho_{f^{-1}(U), f^{-1}(V)} \\ \mathcal{O}_Y(V) & \xrightarrow{f_V^\#} & \mathcal{O}_X(f^{-1}(V)) \end{array}$$

Definition IV.B.1. A *morphism of ringed spaces* $(X, \mathcal{O}_X) \rightarrow (Y, \mathcal{O}_Y)$ is a pair $(f, f^\#)$ where $f : X \rightarrow Y$ is a continuous map of topological spaces and $f^\# : \mathcal{O}_Y \rightarrow f_*(\mathcal{O}_X)$ is a morphism of sheaves of rings. A *morphism of locally ringed spaces* $(X, \mathcal{O}_X) \rightarrow (Y, \mathcal{O}_Y)$ is a morphism of ringed spaces $(f, f^\#)$ such that, for all $x \in X$, the induced local map $\mathcal{O}_{Y,f(x)} \rightarrow \mathcal{O}_{X,x}$ sends the maximal ideal of $\mathcal{O}_{Y,f(x)}$ into the maximal ideal of $\mathcal{O}_{X,x}$.

Example IV.B.2 (A morphism of ringed spaces which is not local). Let $X = Y = \{*\}$ be the one-point topological space. Put

$$\mathcal{O}_Y = \mathbb{k}[t]_{(t)}, \quad \mathcal{O}_X = \mathbb{k}(t),$$

viewed as sheaves of rings on the one-point space. The inclusion

$$\mathbb{k}[t]_{(t)} \hookrightarrow \mathbb{k}(t)$$

defines a morphism of ringed spaces

$$(X, \mathcal{O}_X) \rightarrow (Y, \mathcal{O}_Y).$$

However, this morphism is not a morphism of locally ringed spaces.

Indeed, the ring $\mathbb{k}[t]_{(t)}$ is local, with maximal ideal

$$(t)\mathbb{k}[t]_{(t)}.$$

The ring $\mathbb{k}(t)$ is a field, hence its maximal ideal is (0) . The induced map on stalks is

$$\mathbb{k}[t]_{(t)} \hookrightarrow \mathbb{k}(t).$$

But the element t belongs to the maximal ideal of $\mathbb{k}[t]_{(t)}$, while its image in $\mathbb{k}(t)$ is non-zero, hence invertible. Therefore

$$t \notin (0).$$

So the maximal ideal of $\mathbb{k}[t]_{(t)}$ is not sent into the maximal ideal of $\mathbb{k}(t)$, and the morphism is not local.

Exercise IV.B.3. Let $X = Y = \{*\}$ be the one-point space. Let R and S be local rings, and view them as sheaves of rings on Y and X respectively. Show that a ring homomorphism

$$\varphi : R \rightarrow S$$

defines a morphism of ringed spaces

$$(X, S) \rightarrow (Y, R).$$

Show that this morphism is a morphism of locally ringed spaces if and only if the ring homomorphism $R \rightarrow S$ is local, i.e. if and only if

$$\varphi(\mathfrak{m}_R) \subset \mathfrak{m}_S.$$

Solution. Since X and Y each have only one open subset different from the empty set, a sheaf of rings on such a space is just a ring. A morphism of ringed spaces

$$(X, S) \rightarrow (Y, R)$$

therefore amounts to a ring homomorphism

$$R \rightarrow S.$$

The condition for this morphism to be a morphism of locally ringed spaces is that the induced map on stalks sends the maximal ideal of R into the maximal ideal of S :

$$\varphi(\mathfrak{m}_R) \subset \mathfrak{m}_S.$$

This is precisely the usual definition of a local homomorphism of local rings.

Now take

$$R = \mathbb{k}[t]_{(t)}, \quad S = \mathbb{k}(t).$$

Then R is local with maximal ideal (t) , while S is a field, with maximal ideal (0) . The inclusion sends t to a non-zero element of $\mathbb{k}(t)$, hence to an invertible element. Thus

$$\varphi((t)) \not\subset (0),$$

so the corresponding morphism of ringed spaces is not a morphism of locally ringed spaces.

Proposition IV.B.4. Every ring homomorphism $\varphi : A \rightarrow B$ induces a morphism of locally ringed spaces

$$\mathrm{Spec}(B) \longrightarrow \mathrm{Spec}(A).$$

This construction is contravariantly functorial in the ring.

Proof. Let A be a ring and let X be the ringed space $\text{Spec}(A)$. So X is a topological space equipped with the sheaf of rings $\mathcal{O}_X$. We said that X is a locally ringed space as for all $\mathfrak{p} \in X$, the stalk $\mathcal{O}_{X,\mathfrak{p}} \simeq A_{\mathfrak{p}}$ is a local ring, with maximal ideal $\mathfrak{p}A_{\mathfrak{p}}$.

Let $\varphi : A \rightarrow B$ be a ring homomorphism and set $X = \text{Spec}(B)$, $Y = \text{Spec}(A)$. Then we proved that the map $f_{\varphi} : X \rightarrow Y$ sending a prime ideal $\mathfrak{q} \subset B$ to $\varphi^{-1}(\mathfrak{q})$ is continuous for the Zariski topology.

We need to define a morphism of sheaves of rings $f_{\varphi}^{\#} : \mathcal{O}_Y \rightarrow f_{\varphi*}(\mathcal{O}_X)$. By Proposition IV.D.11, it is enough to define $f_{\varphi}^{\#}$ on the principal open subsets of Y and to check that these maps are compatible with restrictions. So let $a \in A$ and consider the principal open subsets $D(a) \subset Y$ and $D(b) \subset X$, where we put $b = \varphi(a)$. We note that $f_{\varphi}^{-1}(D(a)) = D(b)$. Indeed:

$$\begin{aligned} f_{\varphi}^{-1}(D(a)) &= \{\mathfrak{p} \in X \mid f_{\varphi}(\mathfrak{p}) \in D(a)\} = \\ &= \{\mathfrak{p} \in X \mid a \notin \varphi^{-1}(\mathfrak{p})\} = \\ &= \{\mathfrak{p} \in X \mid \varphi(a) \notin \mathfrak{p}\} = D(b). \end{aligned}$$

Then we should define:

$$f_{D(a)}^{\#} : A_a \simeq \mathcal{O}_Y(D(a)) \rightarrow \mathcal{O}_X(f_{\varphi}^{-1}(D(a))) \simeq \mathcal{O}_X(D(b)) \simeq B_b$$

This map is nothing but the localisation of φ at a . Since these maps are compatible with restriction to further principal open subsets, we get the desired map of sheaves $f_{\varphi}^{\#}$. For any prime ideal $\mathfrak{p} \subset B$ and $\mathfrak{q} = \varphi^{-1}(\mathfrak{p}) \subset A$, by construction, the induced map

$$A_{\mathfrak{q}} \simeq \mathcal{O}_{Y,\mathfrak{q}} \rightarrow B_{\mathfrak{p}} \simeq \mathcal{O}_{X,\mathfrak{p}}$$

is given by localising φ at $\mathfrak{q}$. Since $\mathfrak{q} = \varphi^{-1}(\mathfrak{p})$, we also have $\mathfrak{q}A_{\mathfrak{q}} = \varphi^{-1}(\mathfrak{p}B_{\mathfrak{p}})$. In other words, the maximal ideals correspond to each other.

So $(f_{\varphi}, f_{\varphi}^{\#})$ defines a morphism of locally ringed spaces. The compatibility with composition of ring homomorphisms follows directly from the construction on principal open subsets. $\square$

Exercise IV.B.5. Let $f : X \rightarrow Y$ be a continuous map.

- i) Assume that X and Y are smooth manifolds and that f is a smooth map. Show that pullback of smooth functions defines a morphism of locally ringed spaces

$$(X, \mathcal{C}_X^{\infty}) \longrightarrow (Y, \mathcal{C}_Y^{\infty}).$$

- ii) Assume that X and Y are complex manifolds and that f is a holomorphic map. Show that pullback of holomorphic functions defines a morphism of locally ringed spaces

$$(X, \mathcal{O}_X) \longrightarrow (Y, \mathcal{O}_Y).$$

Solution. We treat the smooth case first. For every open subset $V \subset Y$, define

$$f_V^{\#} : \mathcal{C}_Y^{\infty}(V) \longrightarrow \mathcal{C}_X^{\infty}(f^{-1}(V))$$

by

$$f_V^{\#}(g) = g \circ f.$$

Since f is smooth, the pullback $g \circ f$ is a smooth function on $f^{-1}(V)$. These maps are compatible with restrictions: if $V' \subset V$, then

$$(g|_{V'}) \circ f = (g \circ f)|_{f^{-1}(V')}.$$

Therefore the maps $f_V^\#$ define a morphism of sheaves of rings

$$f^\# : \mathcal{C}_Y^\infty \longrightarrow f_* \mathcal{C}_X^\infty.$$

Together with the continuous map $f : X \rightarrow Y$, this gives a morphism of ringed spaces.

We now check that this morphism is local. Let $x \in X$. The induced map on stalks is

$$f_x^\# : \mathcal{C}_{Y, f(x)}^\infty \longrightarrow \mathcal{C}_{X, x}^\infty, \quad [g]_{f(x)} \longmapsto [g \circ f]_x.$$

The maximal ideal of $\mathcal{C}_{Y, f(x)}^\infty$ is

$$\mathfrak{m}_{f(x)} = \{[g]_{f(x)} \mid g(f(x)) = 0\}.$$

Similarly, the maximal ideal of $\mathcal{C}_{X, x}^\infty$ is

$$\mathfrak{m}_x = \{[h]_x \mid h(x) = 0\}.$$

If $[g]_{f(x)} \in \mathfrak{m}_{f(x)}$, then $g(f(x)) = 0$. Hence

$$(g \circ f)(x) = g(f(x)) = 0,$$

so

$$[g \circ f]_x \in \mathfrak{m}_x.$$

Thus

$$f_x^\#(\mathfrak{m}_{f(x)}) \subset \mathfrak{m}_x,$$

which is precisely the condition for a morphism of locally ringed spaces.

The holomorphic case is identical. If $f : X \rightarrow Y$ is holomorphic and $g \in \mathcal{O}_Y(V)$, then $g \circ f$ is holomorphic on $f^{-1}(V)$. Thus we obtain maps

$$f_V^\# : \mathcal{O}_Y(V) \longrightarrow \mathcal{O}_X(f^{-1}(V)), \quad g \longmapsto g \circ f,$$

compatible with restrictions, hence a morphism of sheaves

$$f^\# : \mathcal{O}_Y \longrightarrow f_* \mathcal{O}_X.$$

On stalks,

$$f_x^\# : \mathcal{O}_{Y, f(x)} \longrightarrow \mathcal{O}_{X, x}$$

sends the germ of a holomorphic function g at $f(x)$ to the germ of $g \circ f$ at x . Since

$$g(f(x)) = 0 \implies (g \circ f)(x) = 0,$$

the maximal ideal of germs vanishing at $f(x)$ is sent into the maximal ideal of germs vanishing at x . Hence this is again a morphism of locally ringed spaces.

Theorem IV.B.6. *Let A and B be rings. Taking global sections gives a natural bijection*

$$\mathrm{Hom}_{(\mathrm{LRS})}(\mathrm{Spec}(B), \mathrm{Spec}(A)) \simeq \mathrm{Hom}_{(\mathrm{Ring})}(A, B).$$

Under this bijection, a ring homomorphism $\varphi : A \rightarrow B$ corresponds to the morphism $\mathrm{Spec}(B) \rightarrow \mathrm{Spec}(A)$ constructed in Proposition IV.B.4.

Proof. Put $X = \operatorname{Spec}(B)$, and $Y = \operatorname{Spec}(A)$. Recall that

$$\mathcal{O}_Y(Y) = A, \quad \mathcal{O}_X(X) = B.$$

Thus, taking global sections defines a ring homomorphism

$$\operatorname{Hom}_{(\operatorname{LRS})}(X, Y) \longrightarrow \operatorname{Hom}_{(\operatorname{Ring})}(A, B), \quad (f, f^\#) \longmapsto f_Y^\#.$$

Conversely, Proposition IV.B.4 associates with every ring homomorphism

$$\varphi : A \longrightarrow B$$

a morphism of locally ringed spaces

$$(f_\varphi, f_\varphi^\#) : X \longrightarrow Y.$$

By construction, the map induced on global sections is φ . Indeed, the map on global sections described in the proof of Proposition IV.B.4 corresponds to localising φ at the unit of A and B , which just gives back φ . Thus the next composition is the identity:

$$\operatorname{Hom}_{(\operatorname{Ring})}(A, B) \longrightarrow \operatorname{Hom}_{(\operatorname{LRS})}(X, Y) \longrightarrow \operatorname{Hom}_{(\operatorname{Ring})}(A, B)$$

It remains to prove the converse. Let $(f, f^\#) : X \longrightarrow Y$ be a morphism of locally ringed spaces, and let $\varphi = f_Y^\# : A \longrightarrow B$ be the induced map on global sections. We shall prove that

$$(f, f^\#) = (f_\varphi, f_\varphi^\#).$$

We first determine the underlying map of topological spaces. Let $\mathfrak{q} \in \operatorname{Spec}(B)$ and put $\mathfrak{p} = f(\mathfrak{q}) \in \operatorname{Spec}(A)$. The morphism of locally ringed spaces induces a local homomorphism

$$f_\mathfrak{q}^\# : \mathcal{O}_{Y, \mathfrak{p}} \longrightarrow \mathcal{O}_{X, \mathfrak{q}}.$$

Using the descriptions of the stalks of affine schemes, this is a local homomorphism

$$f_\mathfrak{q}^\# : A_\mathfrak{p} \longrightarrow B_\mathfrak{q}.$$

Moreover, the diagram

$$\begin{array}{ccc} A & \xrightarrow{\varphi} & B \\ \downarrow & & \downarrow \\ A_\mathfrak{p} & \xrightarrow{f_\mathfrak{q}^\#} & B_\mathfrak{q} \end{array}$$

commutes, where the vertical arrows are the localisation homomorphisms. Let $a \in A$. Suppose first that $a \in \mathfrak{p}$. Then

$$\frac{a}{1} \in \mathfrak{p}A_\mathfrak{p},$$

the maximal ideal of $A_\mathfrak{p}$. Since $f_\mathfrak{q}^\#$ is local, it sends this element into the maximal ideal $\mathfrak{q}B_\mathfrak{q}$ of $B_\mathfrak{q}$. By commutativity of the diagram,

$$\frac{\varphi(a)}{1} \in \mathfrak{q}B_\mathfrak{q}.$$

This implies that $\varphi(a) \in \mathfrak{q}$. Indeed, if $\varphi(a)/1 \in \mathfrak{q}B_\mathfrak{q}$, there exists $s \notin \mathfrak{q}$ such that $s\varphi(a) \in \mathfrak{q}$. Since $\mathfrak{q}$ is prime and $s \notin \mathfrak{q}$, we obtain $\varphi(a) \in \mathfrak{q}$.

Conversely, suppose that $a \notin \mathfrak{p}$. Then $a/1$ is invertible in $A_{\mathfrak{p}}$. Ring homomorphisms send invertible elements to invertible elements, so $\frac{\varphi(a)}{1} = f_{\mathfrak{q}}^{\#}\left(\frac{a}{1}\right)$ is invertible in $B_{\mathfrak{q}}$. Hence it does not belong to the maximal ideal $\mathfrak{q}B_{\mathfrak{q}}$, and therefore $\varphi(a) \notin \mathfrak{q}$. We have proved that

$$a \in \mathfrak{p} \iff \varphi(a) \in \mathfrak{q}.$$

Consequently,

$$f(\mathfrak{q}) = \mathfrak{p} = \varphi^{-1}(\mathfrak{q}) = f_{\varphi}(\mathfrak{q}).$$

This holds for every $\mathfrak{q} \in \text{Spec}(B)$, so

$$f = f_{\varphi}.$$

It remains to identify the morphism of structure sheaves. Let $a \in A$. Since the underlying continuous map is f_{φ} , already we have

$$f^{-1}(D(a)) = D(\varphi(a)).$$

Therefore the component of $f^{\#}$ on the principal open subset $D(a)$ is a ring homomorphism

$$f_{D(a)}^{\#} : \mathcal{O}_Y(D(a)) \longrightarrow \mathcal{O}_X(D(\varphi(a))),$$

or, using the description of the structure sheaf on principal open subsets,

$$f_{D(a)}^{\#} : A_a \longrightarrow B_{\varphi(a)}.$$

Compatibility of $f^{\#}$ with restriction from Y to $D(a)$ gives a commutative diagram

$$\begin{array}{ccc} A & \xrightarrow{\varphi} & B \\ \downarrow & & \downarrow \\ A_a & \xrightarrow{f_{D(a)}^{\#}} & B_{\varphi(a)}. \end{array}$$

The element $\varphi(a)$ is invertible in $B_{\varphi(a)}$. By the universal property of localisation, there is a unique ring homomorphism $A_a \longrightarrow B_{\varphi(a)}$ making this diagram commute. It is the localisation of φ , explicitly:

$$\frac{b}{a^n} \mapsto \frac{\varphi(b)}{\varphi(a)^n}.$$

Thus

$$f_{D(a)}^{\#} = (f_{\varphi}^{\#})_{D(a)}$$

for every principal open subset $D(a) \subset Y$.

The principal open subsets form a basis of the topology of Y . By Proposition IV.D.11, a morphism of sheaves is uniquely determined by its maps on the principal open subsets. Hence

$$f^{\#} = f_{\varphi}^{\#}.$$

Therefore $(f, f^{\#}) = (f_{\varphi}, f_{\varphi}^{\#})$. The two constructions are inverse to each other, proving the desired bijection. Its naturality follows from the contravariant functoriality proved in Proposition IV.B.4. $\square$

Definition IV.B.7. An *affine scheme* is a locally ringed space isomorphic to $\text{Spec}(A)$ for some ring A .

V Affine subschemes

Further reading. For open and closed subschemes and their scheme structures, see [Har77, Chapter II, §§2–3], [Liu02, Chapter 2], and the examples in [EH00].

V.A Open subsets

Let us review how to define subschemes of an affine scheme. Let A be a ring and let $X = \operatorname{Spec}(A)$. For a principal open subset $D(a) \subset X$, the localisation map $A \rightarrow A_a$ induces an isomorphism of schemes

$$\operatorname{Spec}(A_a) \xrightarrow{\sim} D(a).$$

Thus principal open subsets of affine schemes are affine schemes.

More generally, an arbitrary open subset $U \subset X$ inherits a scheme structure by restriction:

$$(U, \mathcal{O}_X|_U).$$

Such an open subscheme need not be affine. The inclusion $j : U \rightarrow X$ is a morphism of schemes. It is moreover a monomorphism: if $f_1, f_2 : Y \rightarrow U$ are morphisms of schemes such that $j \circ f_1 = j \circ f_2$, then $f_1 = f_2$ set-theoretically, and the maps on structure sheaves agree because they agree after composition with the inclusion.

Example V.A.1. In $X = \operatorname{Spec}(\mathbb{k}[x])$, the principal open subset $D(x)$ is the complement of the closed point (x) . As a scheme,

$$D(x) \simeq \operatorname{Spec}(\mathbb{k}[x, x^{-1}]).$$

Example V.A.2. Let $A = \mathbb{k}[x, y]$ and $X = \operatorname{Spec}(A) = \mathbb{A}_{\mathbb{k}}^2$. The principal open subset

$$D(x) \subset X$$

is affine and isomorphic to

$$\operatorname{Spec}(\mathbb{k}[x, y, x^{-1}]).$$

On closed points, assuming that $\mathbb{k}$ is algebraically closed, this is the affine plane with the line $x = 0$ removed.

V.B Closed affine subschemes

Let A be a ring and set $X = \operatorname{Spec}(A)$. Then, given an ideal $\mathfrak{a} \subset A$, we have a ring $B = A/\mathfrak{a}$ and a surjective morphism of rings $\varphi : A \rightarrow B$. Put $Y = \operatorname{Spec}(B)$. The morphism φ induces a homeomorphism $i : Y \rightarrow \mathbb{V}(\mathfrak{a}) \subset X$. This also gives a morphism of sheaves:

$$i^\# : \mathcal{O}_X \rightarrow i_*(\mathcal{O}_Y).$$

On global sections, this morphism is the quotient map

$$\varphi : A = \mathcal{O}_X(X) \rightarrow i_*(\mathcal{O}_Y)(i^{-1}(X)) = \mathcal{O}_Y(Y) = B.$$

The morphism of sheaves $i^\#$ is obtained by localising this quotient map on principal open subsets $D(a) \subset X$ for $a \in A$. Indeed, we have

$$\begin{aligned} i^{-1}(D(a)) &= \{\mathfrak{q} \in \operatorname{Spec}(B) \mid a \notin i(\mathfrak{q})\} = \\ &= \{\mathfrak{q} \in \operatorname{Spec}(B) \mid a \notin \varphi^{-1}(\mathfrak{q})\} = \\ &= \{\mathfrak{q} \in \operatorname{Spec}(B) \mid \varphi(a) \notin \mathfrak{q}\} = D(\varphi(a)). \end{aligned}$$

Then for each $a \in A$, we just consider $i^\#$ to be defined at $D(a)$ by the localisation $A_a \rightarrow B_{\varphi(a)}$ of φ at a .

A closed affine subscheme Y of an affine scheme $X = \operatorname{Spec}(A)$ is, by definition, a closed subscheme arising in the above form, namely $Y = \operatorname{Spec}(B)$, for some surjective morphism of rings $\varphi : A \rightarrow B$. The ideal $\mathfrak{a} = \ker(\varphi)$ is the *ideal defining Y as a subscheme of X* .

Example V.B.1. Let $X = \operatorname{Spec}(\mathbb{k}[x, y]) = \mathbb{A}_{\mathbb{k}}^2$. The ideal (x, y) defines the origin:

$$\operatorname{Spec}(\mathbb{k}[x, y]/(x, y)) \simeq \operatorname{Spec}(\mathbb{k}).$$

The ideal (x^2, y) defines a non-reduced closed subscheme supported at the origin:

$$\operatorname{Spec}(\mathbb{k}[x, y]/(x^2, y)).$$

Its underlying closed subset is the same point, but the scheme remembers an infinitesimal direction along the x -axis.

Exercise V.B.2. If A and B are rings, then there is a homeomorphism

$$\operatorname{Spec}(A \times B) \simeq \operatorname{Spec}(A) \sqcup \operatorname{Spec}(B).$$

The two subsets $\operatorname{Spec}(A)$ and $\operatorname{Spec}(B)$ are open and closed in $\operatorname{Spec}(A \times B)$.

Solution. Let $e_A = (1, 0)$, and $e_B = (0, 1)$ be the two standard idempotents of $A \times B$. They satisfy

$$e_A e_B = 0, \quad e_A + e_B = 1.$$

If $\mathfrak{p} \subset A \times B$ is a prime ideal, then from $e_A e_B = 0 \in \mathfrak{p}$ we get $e_A \in \mathfrak{p}$ or $e_B \in \mathfrak{p}$. Since $e_A + e_B = 1$, they cannot both belong to $\mathfrak{p}$.

If $e_B \in \mathfrak{p}$, then $\mathfrak{p}$ is of the form

$$\mathfrak{q} \times B$$

for a unique prime ideal $\mathfrak{q} \subset A$. Similarly, if $e_A \in \mathfrak{p}$, then $\mathfrak{p}$ is of the form

$$A \times \mathfrak{r}$$

for a unique prime ideal $\mathfrak{r} \subset B$. Thus, as sets,

$$\operatorname{Spec}(A \times B) = \operatorname{Spec}(A) \sqcup \operatorname{Spec}(B).$$

Moreover, these two pieces are open and closed. Indeed,

$$D(e_A) \simeq \operatorname{Spec}(A), \quad D(e_B) \simeq \operatorname{Spec}(B),$$

and

$$D(e_A) \cap D(e_B) = \emptyset, \quad D(e_A) \cup D(e_B) = \operatorname{Spec}(A \times B).$$

Finally, the structure sheaf restricts correctly, because

$$(A \times B)_{e_A} \simeq A, \quad (A \times B)_{e_B} \simeq B.$$

Therefore the above bijection is an isomorphism of schemes.

More generally, for finitely many rings $A_1, \dots, A_s$, one has

$$\operatorname{Spec} \left(\prod_{i=1}^s A_i \right) \simeq \bigsqcup_{i=1}^s \operatorname{Spec}(A_i).$$

Exercise V.B.3. Let $X = \operatorname{Spec}(A)$, and let $Y, Z \subset X$ be closed subschemes defined by ideals $\mathfrak{a}, \mathfrak{b} \subset A$. Show that the closed subscheme defined by $\mathfrak{a} \cap \mathfrak{b}$ is the smallest closed subscheme of X containing both Y and Z . This closed subscheme is called the *scheme-theoretic union* of Y and Z .

Show also that $\mathfrak{a} \cap \mathfrak{b}$ and $\mathfrak{a}\mathfrak{b}$ define the same underlying closed subset of X . Give an example where the corresponding closed subschemes are not isomorphic.

Solution. Let $W \subset X$ be a closed subscheme defined by an ideal $\mathfrak{c} \subset A$. Then W contains Y if and only if the closed immersion

$$Y = \operatorname{Spec}(A/\mathfrak{a}) \longrightarrow X$$

factors through $W = \operatorname{Spec}(A/\mathfrak{c})$. Algebraically, this is equivalent to the existence of a surjective morphism

$$A/\mathfrak{c} \longrightarrow A/\mathfrak{a},$$

or equivalently to

$$\mathfrak{c} \subset \mathfrak{a}.$$

Similarly, W contains Z if and only if $\mathfrak{c} \subset \mathfrak{b}$. Therefore W contains both Y and Z if and only if

$$\mathfrak{c} \subset \mathfrak{a} \cap \mathfrak{b}.$$

Thus the largest ideal defining a closed subscheme containing both Y and Z is $\mathfrak{a} \cap \mathfrak{b}$. Since larger ideals give smaller closed subschemes, the closed subscheme defined by $\mathfrak{a} \cap \mathfrak{b}$ is the smallest closed subscheme of X containing both Y and Z .

Now compare $\mathfrak{a} \cap \mathfrak{b}$ and $\mathfrak{a}\mathfrak{b}$. Since

$$\mathfrak{a}\mathfrak{b} \subset \mathfrak{a} \cap \mathfrak{b},$$

we have

$$\mathbb{V}(\mathfrak{a} \cap \mathfrak{b}) \subset \mathbb{V}(\mathfrak{a}\mathfrak{b}).$$

Conversely, let $\mathfrak{p} \in \mathbb{V}(\mathfrak{a}\mathfrak{b})$. Then $\mathfrak{a}\mathfrak{b} \subset \mathfrak{p}$. Since $\mathfrak{p}$ is prime, we must have

$$\mathfrak{a} \subset \mathfrak{p} \quad \text{or} \quad \mathfrak{b} \subset \mathfrak{p}.$$

In either case,

$$\mathfrak{a} \cap \mathfrak{b} \subset \mathfrak{p}.$$

Thus

$$\mathbb{V}(\mathfrak{a}\mathfrak{b}) \subset \mathbb{V}(\mathfrak{a} \cap \mathfrak{b}).$$

Hence

$$\mathbb{V}(\mathfrak{a}\mathfrak{b}) = \mathbb{V}(\mathfrak{a} \cap \mathfrak{b}).$$

So the two ideals define the same underlying closed subset of X .

However, they need not define the same closed subscheme. For example, take

$$A = \mathbb{k}[x], \quad \mathfrak{a} = \mathfrak{b} = (x).$$

Then

$$\mathfrak{a} \cap \mathfrak{b} = (x), \quad \mathfrak{a}\mathfrak{b} = (x^2).$$

The closed subscheme defined by $\mathfrak{a} \cap \mathfrak{b}$ is

$$\operatorname{Spec}(\mathbb{k}[x]/(x)) \simeq \operatorname{Spec}(\mathbb{k}),$$

whereas the closed subscheme defined by $\mathfrak{a}\mathfrak{b}$ is

$$\operatorname{Spec}(\mathbb{k}[x]/(x^2)).$$

These schemes are not isomorphic, because $\mathbb{k}[x]/(x) \simeq \mathbb{k}$ is reduced, whereas $\mathbb{k}[x]/(x^2)$ has a non-zero nilpotent element, namely the class of x .

VI Subschemes of the affine space

Let B be a commutative ring and $A = B[x_1, \dots, x_n]$. Then $\mathbb{A}_B^n = \text{Spec}(A)$ is the *affine n -space over B* . The inclusion $B \subset A$ induces a surjective morphism $\mathbb{A}_B^n \rightarrow \text{Spec}(B)$. We will mostly consider the case when B is a field.

VI.A Subschemes of the affine line over a field

Definition VI.A.1. An *affine scheme of finite type over $\mathbb{k}$* is the spectrum of a finitely generated $\mathbb{k}$ -algebra, namely a scheme of the form

$$\text{Spec}(\mathbb{k}[x_1, \dots, x_n]/\mathfrak{a})$$

for some ideal $\mathfrak{a} \subset \mathbb{k}[x_1, \dots, x_n]$. Equivalently, it is a closed subscheme of an affine space over $\mathbb{k}$.

We call such a scheme an *affine variety* when its coordinate ring is an integral domain. Some authors use the word “variety” for more general reduced schemes; the convention should always be stated explicitly.

Proposition VI.A.2. A closed subscheme $X \subset \mathbb{A}_{\mathbb{k}}^1$ is either $\emptyset$, or $\mathbb{A}_{\mathbb{k}}^1$, or the disjoint union of schemes of the form $\text{Spec}(\mathbb{k}[x]/(f^m))$, where $f \in \mathbb{k}[x]$ is an irreducible polynomial in $\mathbb{k}[x]$ and $m \geq 1$.

Proof. By definition, a closed subscheme of $\mathbb{A}_{\mathbb{k}}^1$ has the form $\text{Spec}(\mathbb{k}[x]/\mathfrak{a})$ for some ideal $\mathfrak{a}$ of $\mathbb{k}[x]$, so that $\mathfrak{a} = (g)$, for some $g \in \mathbb{k}[x]$, since $\mathbb{k}[x]$ is a principal ideal domain. If $g = 0$ then $X = \mathbb{A}_{\mathbb{k}}^1$ and, if g is a non-zero constant, then $X = \emptyset$. If g is non-constant, then we can write $g = f_1^{m_1} \cdots f_s^{m_s}$, in an essentially unique way, where $m_i \geq 1$ are integers and f_i are pairwise coprime irreducible polynomials in $\mathbb{k}[x]$, for $1 \leq i \leq s$. Then $\mathbb{V}(g)$ is the disjoint union of the subschemes $\mathbb{V}(f_i^{m_i})$, for $1 \leq i \leq s$. Indeed, by the Chinese remainder theorem, we have

$$\mathbb{k}[x]/(g) \simeq \prod_{1 \leq i \leq s} \mathbb{k}[x]/(f_i^{m_i}).$$

So taking spectra (see the next exercise), we get

$$X = \text{Spec}(\mathbb{k}[x]/(g)) \simeq \bigsqcup_{1 \leq i \leq s} \text{Spec}(\mathbb{k}[x]/(f_i^{m_i})).$$

□

Exercise VI.A.3. Let $A = \mathbb{k}[x]$ and let $\mathfrak{a} = (x^2(x-1)^3)$. Show that $\mathfrak{a} = (x^2) \cap ((x-1)^3)$. Describe the corresponding closed subscheme of $\mathbb{A}_{\mathbb{k}}^1$.

Solution. The ideals (x^2) and $((x-1)^3)$ are coprime, because x and $x-1$ are coprime. Hence

$$(x^2) \cap ((x-1)^3) = (x^2)((x-1)^3) = (x^2(x-1)^3).$$

Thus

$$\text{Spec}(\mathbb{k}[x]/(x^2(x-1)^3))$$

is the scheme-theoretic union of the double point at 0 and the triple point at 1.

VI.B Points of the affine plane over a field

Let $\mathbb{k}$ be an algebraically closed field and consider $\mathbb{A}_{\mathbb{k}}^2 = \text{Spec}(A)$, with $A = \mathbb{k}[x, y]$. Then, a subscheme of $\mathbb{A}_{\mathbb{k}}^2$ is given by an ideal $\mathfrak{a}$ of A . By the Nullstellensatz, we have $\mathbf{V}(\mathfrak{a}) = \emptyset$ if and only if $\sqrt{\mathfrak{a}} = A$, i.e. if there is $n \in \mathbb{N}$ such that $1^n \in \mathfrak{a}$, which of course means that $1 \in \mathfrak{a}$, so that $\mathfrak{a} = A$. Therefore, for any proper ideal $\mathfrak{a} \subsetneq A$, we have $\mathbf{V}(\mathfrak{a}) \neq \emptyset$ and any given point $a \in \mathbf{V}(\mathfrak{a})$ gives a maximal ideal $\mathfrak{m}_a$ which contains $\mathfrak{a}$.

The point about the For the affine plane, we have a classification of prime ideals. There are principal prime ideals, for example the ideal $(y - x^2)$ is prime because $\mathbb{k}[x, y]/(y - x^2) \simeq \mathbb{k}[x]$ is an integral domain. The point is that we can classify all other prime ideals and they are all maximal. This relies on the fact that $A = R[x_2]$ where $R = \mathbb{k}[x_1]$ is a principal ideal domain. More generally, we have the following result.

Proposition VI.B.1. *Let R be a principal ideal domain, let y be an indeterminate, and put $B = R[y]$. Let $\mathfrak{p}$ be a prime ideal of B .*

- i) *If $\mathfrak{p} \cap R = (0)$, then $\mathfrak{p}$ is principal.*
- ii) *If $\mathfrak{p} \cap R = (r) \neq (0)$, then r is irreducible and either $\mathfrak{p} = (r)$, or $\mathfrak{p} = (r, g)$, where, setting $\kappa = R/(r)$, the image $\bar{g} \in \kappa[y]$ is irreducible. In the latter case, $\mathfrak{p}$ is maximal.*

Proof. Set $K = \text{Frac}(R)$. Assume first that $\mathfrak{p} \cap R = (0)$. If $\mathfrak{p} = (0)$ there is nothing to prove. Otherwise, $\mathfrak{p} \cap (R \setminus \{0\}) = \emptyset$ so localising at $R \setminus \{0\}$ gives a non-zero prime ideal $\mathfrak{p}K[y]$ of the principal ideal domain $K[y]$. Hence

$$\mathfrak{p}K[y] = (F)$$

for some irreducible polynomial $F \in K[y]$. After multiplying by a non-zero scalar, we may choose a primitive polynomial $f \in R[y]$ associated with F . Since $f \in \mathfrak{p}K[y]$, there exists $s \in R \setminus \{0\}$ such that $sf \in \mathfrak{p}$. As $s \notin \mathfrak{p}$ and $\mathfrak{p}$ is prime, we get $f \in \mathfrak{p}$. Thus $(f) \subset \mathfrak{p}$.

Conversely, if $g \in \mathfrak{p}$, then $g \in (f)$ in $K[y]$. Since f is primitive, Gauss's lemma implies that f divides g already in $R[y]$. Therefore $\mathfrak{p} \subset (f)$, and hence $\mathfrak{p} = (f)$. Recall how this works: write $g = fh$ for some $h \in K[y]$. Then, for some $c = a/b \in K$, with $a, b \in R$ coprime, we have $h = ch_0$ with $h_0 \in R[y]$ primitive. Then $g = cfh_0$ gives that b must divide all coefficients of fh_0 , because $g \in R[y]$. But fh_0 is primitive, so b must be a unit. Hence c lies in R and $g = ch_0f$ happens in $R[y]$, so f divides g in $R[y]$.

Now assume that $\mathfrak{p} \cap R = (r) \neq (0)$. Since this is a non-zero prime ideal of the principal ideal domain R , the element r is irreducible. Also, $\kappa = R/(r)$ is a field. Modulo (r) , the ideal

$$\bar{\mathfrak{p}} = \mathfrak{p}/(r)$$

is a prime ideal of the principal ideal domain

$$\kappa[y].$$

Thus either $\bar{\mathfrak{p}} = (0)$, in which case $\mathfrak{p} = (r)$, or $\bar{\mathfrak{p}} = (\bar{g})$ for an irreducible polynomial $\bar{g}$. Choosing a lift $g \in R[y]$, we obtain $\mathfrak{p} = (r, g)$. In this case

$$B/\mathfrak{p} \simeq \kappa[y]/(\bar{g})$$

is a field, so $\mathfrak{p}$ is maximal. □

Corollary VI.B.2. *Let $\mathbb{k}$ be an algebraically closed field. Then any prime ideal $\mathfrak{p}$ of $A = \mathbb{k}[x_1, x_2]$ is (0) , or (f) with $f \in A$ irreducible, or $\mathfrak{p} = (x_1 - a_1, x_2 - a_2)$ with $a = (a_1, a_2) \in \mathbb{k}^2$.*

About residue fields, we have the following observations.

- If $\mathfrak{p}$ is maximal, then we have $\mathfrak{p} = \mathfrak{m}_a$ for some $a \in \mathbb{k}^2$. We have $A_{\mathfrak{m}_a} = \{f/g \in \mathbb{k}(x_1, x_2) \mid g(a) \neq 0\}$. Also, we have $\kappa(\mathfrak{m}_a) \simeq \mathbb{k}$, for all $a \in \mathbb{k}^2$.
- Principal prime ideals are precisely the ideals of the form $\mathfrak{p} = (p)$ where $p \in A$ is an irreducible polynomial. We have $A_{(p)} = \{f/g \in \mathbb{k}(x_1, x_2) \mid p \nmid g\}$. The residue field $\kappa((p)) \simeq \text{Frac}(A/(p))$ depends strongly on p itself.
- The ideal $(0) = \{0\}$ is also prime and $A_{(0)} = \mathbb{k}(x_1, x_2) = \kappa((0))$.

VI.C Closed subschemes of affine space over a field

Let $\mathbb{k}$ be a field and let

$$X = \text{Spec}(\mathbb{k}[x_1, \dots, x_n]/\mathfrak{a})$$

be a closed subscheme of $\mathbb{A}_{\mathbb{k}}^n$. If the ideal $\mathfrak{a}$ is complicated, one way to understand X is to write $\mathfrak{a}$ as an intersection of simpler ideals.

Definition VI.C.1. Let A be a ring and let

$$Y_1, \dots, Y_r \subset \text{Spec}(A)$$

be closed subschemes defined by ideals

$$\mathfrak{q}_1, \dots, \mathfrak{q}_r \subset A.$$

The *scheme-theoretic union* of $Y_1, \dots, Y_r$ is the closed subscheme defined by the ideal

$$\mathfrak{q}_1 \cap \dots \cap \mathfrak{q}_r.$$

Proposition VI.C.2. Let A be a ring and let $\mathfrak{q}_1, \dots, \mathfrak{q}_r \subset A$ be ideals. Put

$$\mathfrak{a} = \mathfrak{q}_1 \cap \dots \cap \mathfrak{q}_r.$$

Then the closed subscheme

$$X = \text{Spec}(A/\mathfrak{a})$$

is the scheme-theoretic union of the closed subschemes

$$X_i = \text{Spec}(A/\mathfrak{q}_i).$$

Moreover, the underlying closed subset of X is the union of the underlying closed subsets of the X_i :

$$\mathbb{W}(\mathfrak{a}) = \mathbb{W}(\mathfrak{q}_1) \cup \dots \cup \mathbb{W}(\mathfrak{q}_r).$$

Proof. The first statement is just the definition of the scheme-theoretic union.

Let us prove the statement about the underlying closed subsets. It is enough to treat the case of two ideals. Let $\mathfrak{q}_1, \mathfrak{q}_2 \subset A$. Since

$$\mathfrak{q}_1 \mathfrak{q}_2 \subset \mathfrak{q}_1 \cap \mathfrak{q}_2,$$

we have

$$\mathbb{W}(\mathfrak{q}_1 \cap \mathfrak{q}_2) \subset \mathbb{W}(\mathfrak{q}_1 \mathfrak{q}_2).$$

Conversely, let $\mathfrak{p} \in \mathbb{W}(\mathfrak{q}_1 \mathfrak{q}_2)$. Then

$$\mathfrak{q}_1 \mathfrak{q}_2 \subset \mathfrak{p}.$$

Since $\mathfrak{p}$ is prime, we have

$$\mathfrak{q}_1 \subset \mathfrak{p} \quad \text{or} \quad \mathfrak{q}_2 \subset \mathfrak{p}.$$

Hence $\mathfrak{p} \in \mathbb{V}(\mathfrak{q}_1) \cup \mathbb{V}(\mathfrak{q}_2)$. Since

$$\mathbb{V}(\mathfrak{q}_1) \cup \mathbb{V}(\mathfrak{q}_2) \subset \mathbb{V}(\mathfrak{q}_1 \cap \mathfrak{q}_2)$$

is immediate, we get

$$\mathbb{V}(\mathfrak{q}_1 \cap \mathfrak{q}_2) = \mathbb{V}(\mathfrak{q}_1) \cup \mathbb{V}(\mathfrak{q}_2).$$

The general case follows by induction. □

Remark VI.C.3. The scheme-theoretic union depends on the ideals themselves, not only on their radicals. Thus it remembers possible nilpotent structure. For instance, the ideals

$$(x) \quad \text{and} \quad (x^2)$$

in $\mathbb{k}[x]$ define the same underlying closed subset of $\mathbb{A}_{\mathbb{k}}^1$, namely the origin, but the corresponding closed subschemes are different.

Example VI.C.4. Let $A = \mathbb{k}[x, y]$, and $\mathfrak{a} = (xy)$. Then $(xy) = (x) \cap (y)$. Therefore $\text{Spec}(A/(xy))$ is the scheme-theoretic union of the two coordinate axes: $\text{Spec}(A/(x))$ and $\text{Spec}(A/(y))$.

Remark VI.C.5. In a Noetherian ring, every proper ideal admits a primary decomposition

$$\mathfrak{a} = \mathfrak{q}_1 \cap \cdots \cap \mathfrak{q}_r.$$

This gives a way to view a closed subscheme

$$\text{Spec}(A/\mathfrak{a})$$

as a scheme-theoretic union of closed subschemes defined by the ideals $\mathfrak{q}_i$. Later, after introducing irreducible components and generic points, we will see how primary decomposition also detects the components of the underlying topological space and possible embedded components.

Exercise VI.C.6. Let $A = \mathbb{k}[x]$ and compare the subschemes of $\mathbb{A}_{\mathbb{k}}^1$ defined by $(x) \cap (x)$ and $(x) \cdot (x)$.

Solution. We have

$$(x) \cap (x) = (x),$$

whereas

$$(x)(x) = (x^2).$$

Thus $(x) \cap (x)$ defines the reduced origin

$$\text{Spec}(\mathbb{k}[x]/(x)) \simeq \text{Spec}(\mathbb{k}),$$

while $(x)(x)$ defines the double point

$$\text{Spec}(\mathbb{k}[x]/(x^2)).$$

The two closed subschemes have the same underlying closed subset, but they are not isomorphic as schemes: $\mathbb{k}[x]/(x)$ is reduced, whereas $\mathbb{k}[x]/(x^2)$ has a non-zero nilpotent element.

VI.D Arithmetic affine schemes

Further reading. For schemes over $\text{Spec}(\mathbb{Z})$, reduction modulo primes, and the arithmetic viewpoint, see [Liu02]; [EH00] offers complementary geometric examples.

Every (unital) ring A has a unique morphism $\mathbb{Z} \rightarrow A$. We denote this morphism by γ_A . Thus every affine scheme has a canonical morphism $\pi_A = \text{Spec}(\gamma_A) : \text{Spec}(A) \rightarrow \text{Spec}(\mathbb{Z})$. This single map records the characteristics in which the scheme has points.

Proposition VI.D.1. *Let A be an integral domain and let $\pi_A = \text{Spec}(\gamma_A) : \text{Spec}(A) \rightarrow \text{Spec}(\mathbb{Z})$ be its structure morphism.*

(i) *If $\text{char}(A) = p > 0$, then the image of π_A is $\{(p)\}$.*

(ii) *If $\text{char}(A) = 0$, then*

$$\text{Im}(\pi_A) = \{(0)\} \cup \{(p) \mid pA \neq A\}.$$

In particular, π_A is dominant.

Proof. The image of a prime $\mathfrak{q} \subset A$ is its contraction $\gamma_A^{-1}(\mathfrak{q})$. If A has characteristic p , every such contraction is (p) . Suppose that A has characteristic zero. The prime $(0) \subset A$ maps to $(0) \subset \mathbb{Z}$. Moreover, a prime of A maps to (p) precisely when it contains $\gamma_A(p) = p1_A$, or equivalently when it contains pA . Such a prime exists precisely when pA is proper. Finally, (0) is dense in $\text{Spec}(\mathbb{Z})$. $\square$

For a general $\mathbb{Z}$ -algebra, reduction modulo a prime packages all points of a fixed residue characteristic.

Proposition VI.D.2. *Let A be a ring and let p be a prime number. The closed immersion*

$$\text{Spec}(A/pA) \rightarrow \text{Spec}(A)$$

has image

$$\mathbb{W}(pA) = \{\mathfrak{q} \in \text{Spec}(A) \mid \text{char } \kappa(\mathfrak{q}) = p\}.$$

Equivalently, these are exactly the points mapping to (p) in $\text{Spec}(\mathbb{Z})$.

Proof. Prime ideals of A/pA correspond to prime ideals of A containing pA . For $\mathfrak{q} \in \text{Spec}(A)$, the contraction $\gamma_A^{-1}(\mathfrak{q})$ is the kernel of the composite

$$\mathbb{Z} \xrightarrow{\gamma_A} A \rightarrow \kappa(\mathfrak{q}),$$

which is precisely $\gamma_{\kappa(\mathfrak{q})}$. It is (p) exactly when $\kappa(\mathfrak{q})$ has characteristic p . $\square$

Example VI.D.3. Let $X = \text{Spec}(\mathbb{Z}[x]/(x^2 - 1))$. For an odd prime p ,

$$X_p = \text{Spec}(\mathbb{F}_p[x]/(x^2 - 1)) \simeq \text{Spec}(\mathbb{F}_p) \coprod \text{Spec}(\mathbb{F}_p),$$

because $x^2 - 1 = (x - 1)(x + 1)$ has two distinct factors. At $p = 2$ the two factors coincide:

$$X_2 = \text{Spec}(\mathbb{F}_2[x]/((x - 1)^2)).$$

Its underlying space has one point, but its ring contains a nonzero nilpotent. Thus reduction may change both the number of points and the scheme structure.

Example VI.D.4 (An arithmetic subscheme of the plane). Let

$$Y = \operatorname{Spec}(\mathbb{Z}[x, y]/(xy - 2)) \subset \mathbb{A}_{\mathbb{Z}}^2.$$

For $p \neq 2$, reduction gives

$$Y_p = \operatorname{Spec}(\mathbb{F}_p[x, y]/(xy - 2)) \simeq \operatorname{Spec}(\mathbb{F}_p[x, x^{-1}]),$$

a hyperbola. At $p = 2$ the equation becomes $xy = 0$, so

$$Y_2 = \operatorname{Spec}(\mathbb{F}_2[x, y]/(xy))$$

is the union of the two coordinate axes. This example exhibits, in one equation, why a family over $\operatorname{Spec}(\mathbb{Z})$ must be studied prime by prime.

VI.E Rings of integers in number fields

VI.E.1 Number fields

Definition VI.E.1. A *number field* is a finite field extension $K/\mathbb{Q}$.

The basic examples are $\mathbb{Q}$, quadratic fields $\mathbb{Q}(\sqrt{d})$ for squarefree $d \in \mathbb{Z}$, and cyclotomic fields $\mathbb{Q}(\zeta_n)$. By contrast, $\mathbb{R}$, $\mathbb{C}$ and $\mathbb{Q}(t)$ are not number fields: none is finite over $\mathbb{Q}$.

VI.E.2 Rings of integers of number fields

Definition VI.E.2. Let K be a number field. An element $\alpha \in K$ is an *algebraic integer* if it satisfies a monic polynomial with coefficients in $\mathbb{Z}$. The algebraic integers in K form a ring, denoted by $\mathcal{O}_K$ and called the *ring of integers* of K .

For example,

$$\mathcal{O}_{\mathbb{Q}} = \mathbb{Z}, \quad \mathcal{O}_{\mathbb{Q}(i)} = \mathbb{Z}[i].$$

More generally, if d is squarefree, then

$$\mathcal{O}_{\mathbb{Q}(\sqrt{d})} = \begin{cases} \mathbb{Z}\left[\frac{1 + \sqrt{d}}{2}\right], & d \equiv 1 \pmod{4}, \\ \mathbb{Z}[\sqrt{d}], & d \not\equiv 1 \pmod{4}. \end{cases}$$

The homomorphism $\gamma_{\mathcal{O}_K} : \mathbb{Z} \rightarrow \mathcal{O}_K$ induces

$$\operatorname{Spec}(\gamma_{\mathcal{O}_K}) : \operatorname{Spec}(\mathcal{O}_K) \longrightarrow \operatorname{Spec}(\mathbb{Z}).$$

Its generic point has residue field K , while the closed points over (p) are the prime ideals of

$$\mathcal{O}_K/p\mathcal{O}_K.$$

Thus factorisation modulo p translates directly into the geometry over the prime (p) .

Example VI.E.3 (The Gaussian integers). For $K = \mathbb{Q}(i)$, write

$$\mathcal{O}_K = \mathbb{Z}[i] \simeq \mathbb{Z}[x]/(x^2 + 1).$$

Three reductions display the basic possibilities:

p	$x^2 + 1$ in $\mathbb{F}_p[x]$	$\text{Spec}(\mathcal{O}_K/p\mathcal{O}_K)$
2	$(x+1)^2$	one non-reduced point with residue field $\mathbb{F}_2$
3	irreducible	one point with residue field $\mathbb{F}_9$
5	$(x-2)(x+2)$	two points with residue field $\mathbb{F}_5$.

Repeated, irreducible and split factorisations therefore produce three different kinds of reductions.

Exercise VI.E.4. Let $K = \mathbb{Q}(\sqrt{2})$, so that $\mathcal{O}_K = \mathbb{Z}[\sqrt{2}]$. Describe

$$\text{Spec}(\mathcal{O}_K/p\mathcal{O}_K)$$

for $p = 2, 3, 7$, including the residue fields and any non-reduced structure.

Solution. We have

$$\mathcal{O}_K/p\mathcal{O}_K \simeq \mathbb{F}_p[x]/(x^2 - 2).$$

For $p = 2$, this is $\mathbb{F}_2[x]/(x^2)$, so there is one non-reduced point with residue field $\mathbb{F}_2$. Over $\mathbb{F}_3$, the polynomial $x^2 - 2$ is irreducible, so there is one point with residue field $\mathbb{F}_9$. Finally,

$$x^2 - 2 = (x - 3)(x + 3) \quad \text{in } \mathbb{F}_7[x],$$

so the reduction at 7 is the disjoint union of two reduced $\mathbb{F}_7$ -points.

Chapter 3

Schemes

I Schemes

Further reading. For the basic language of schemes, local rings, closed subschemes and morphisms, see [Liu02, Chapter 2] and [Har77, Chapter II, §2]; [EH00, Chapters I–II] gives a particularly geometric introduction.

I.A Schemes as locally ringed spaces

I.A.1 Locally ringed spaces

We defined a *locally ringed space* is a topological space X equipped with a sheaf of rings $\mathcal{O}_X$ such that, for every point $x \in X$, the stalk $\mathcal{O}_{X,x}$ is a local ring. We also defined morphisms of locally ringed spaces.

Definition I.A.1. A *scheme* is a locally ringed space $(X, \mathcal{O}_X)$ admitting an open covering $X = \bigcup_{i \in I} U_i$ such that, for every $i \in I$, the locally ringed space $(U_i, \mathcal{O}_X|_{U_i})$ is isomorphic to $\text{Spec}(A_i)$ for some ring A_i . The maximal ideal of the local ring $\mathcal{O}_{X,x}$ is denoted by $\mathfrak{m}_{X,x}$, and the residue field at x is

$$\kappa(x) = \mathcal{O}_{X,x} / \mathfrak{m}_{X,x}.$$

Remark I.A.2. Let $U \simeq \text{Spec}(A)$ be an affine open subset of a scheme X , and let $x \in U$ correspond to a prime ideal $\mathfrak{p} \subset A$. Then

$$\mathcal{O}_{X,x} \simeq A_{\mathfrak{p}}, \quad \mathfrak{m}_{X,x} \simeq \mathfrak{p}A_{\mathfrak{p}}, \quad \kappa(x) \simeq \text{Frac}(A/\mathfrak{p}).$$

Thus the local ring and the residue field may be computed in any affine neighbourhood of the point.

Exercise I.A.3. Let $U \subset X$ be an open subset of a scheme. Prove that $(U, \mathcal{O}_X|_U)$ is a scheme.

Solution. Let $x \in U$. Since X is a scheme, there is an affine open neighbourhood $V = \text{Spec}(A)$ of x in X . The intersection $U \cap V$ is an open neighbourhood of x in V . Since the principal open subsets form a basis of the topology of $\text{Spec}(A)$, there is an element $f \in A$ such that

$$x \in D(f) \subset U \cap V.$$

Moreover,

$$(D(f), \mathcal{O}_X|_{D(f)}) \simeq \text{Spec}(A_f),$$

so $D(f)$ is an affine open neighbourhood of x contained in U . Therefore U is covered by affine open subschemes and is a scheme.

I.A.2 Morphisms of schemes

The category (Sch) of schemes is the full subcategory of the category of locally ringed spaces whose objects are schemes. In other words, given two schemes X and Y , morphisms of schemes $X \rightarrow Y$ are just morphisms $X \rightarrow Y$ as locally ringed spaces.

Definition I.A.4. Let $(X, \mathcal{O}_X)$ and $(Y, \mathcal{O}_Y)$ be schemes. A *morphism of schemes*

$$f: X \longrightarrow Y$$

is a morphism of locally ringed spaces. Thus it consists of a continuous map $f: X \rightarrow Y$ and a morphism of sheaves of rings

$$f^\#: \mathcal{O}_Y \longrightarrow f_* \mathcal{O}_X$$

such that, for every $x \in X$, the induced homomorphism

$$f_x^\#: \mathcal{O}_{Y, f(x)} \longrightarrow \mathcal{O}_{X, x}$$

is local, that is,

$$(f_x^\#)^{-1}(\mathfrak{m}_{X, x}) = \mathfrak{m}_{Y, f(x)}.$$

Exercise I.A.5 (Morphisms are local on the source). Let $X = \bigcup_i U_i$ be an open covering and let Y be a scheme. Suppose that morphisms $f_i: U_i \rightarrow Y$ are given and that

$$f_i|_{U_i \cap U_j} = f_j|_{U_i \cap U_j}$$

for all i, j . Prove that there is a unique morphism $f: X \rightarrow Y$ whose restriction to U_i is f_i .

Solution. The continuous maps underlying the f_i glue uniquely to a continuous map $f: X \rightarrow Y$. For every open subset $V \subset Y$, the maps on sections

$$\mathcal{O}_Y(V) \longrightarrow \mathcal{O}_X(f^{-1}(V) \cap U_i)$$

agree on pairwise overlaps. The sheaf axiom for $\mathcal{O}_X$ glues them uniquely to a homomorphism

$$\mathcal{O}_Y(V) \longrightarrow \mathcal{O}_X(f^{-1}(V)).$$

These homomorphisms commute with restriction and define $f^\#: \mathcal{O}_Y \rightarrow f_* \mathcal{O}_X$. Locality on stalks can be checked on the open cover, so $(f, f^\#)$ is a morphism of schemes. Every part of the construction is forced by the restrictions to the U_i , which proves uniqueness.

I.B Quasi-coherent sheaves

Let $(X, \mathcal{O}_X)$ be a scheme. A *sheaf of $\mathcal{O}_X$ -modules* is a sheaf $\mathcal{F}$ such that $\mathcal{F}(U)$ is an $\mathcal{O}_X(U)$ -module for every open subset $U \subset X$, compatibly with restriction maps.

Definition I.B.1. A sheaf of $\mathcal{O}_X$ -modules $\mathcal{F}$ is *quasi-coherent* if every point of X has an affine open neighbourhood $U = \text{Spec}(A)$ for which there is an A -module M and an isomorphism

$$\mathcal{F}|_U \simeq \widetilde{M}.$$

Here $\widetilde{M}$ is the associated sheaf of Definition III.A.2. Equivalently, on such an affine open,

$$\mathcal{F}(D(f)) \simeq M_f \quad (f \in A),$$

compatibly with restrictions.

Thus quasi-coherence is a local condition: it says that the sheaf is obtained locally from modules by localisation. The structure sheaf $\mathcal{O}_X$ and every finite direct sum $\mathcal{O}_X^{\oplus r}$ are quasi-coherent.

I.B.1 Generating sections

Let $\mathcal{F}$ be an $\mathcal{O}_X$ -module and let $s_1, \dots, s_r \in \Gamma(\mathcal{F})$. Their *evaluation morphism* is

$$\mathrm{ev}_{s_1, \dots, s_r} : \mathcal{O}_X^{\oplus r} \longrightarrow \mathcal{F}, \quad (a_1, \dots, a_r) \longmapsto \sum_{i=1}^r a_i s_i.$$

The sections $s_1, \dots, s_r$ *generate* $\mathcal{F}$ if this morphism is surjective.

Lemma I.B.2 (Generation may be checked on stalks). *The sections $s_1, \dots, s_r$ generate $\mathcal{F}$ if and only if, for every $x \in X$, their germs $(s_1)_x, \dots, (s_r)_x$ generate the $\mathcal{O}_{X,x}$ -module $\mathcal{F}_x$.*

Proof. The stalk at x of the evaluation morphism is

$$\mathcal{O}_{X,x}^{\oplus r} \longrightarrow \mathcal{F}_x, \quad (a_1, \dots, a_r) \longmapsto \sum_i a_i (s_i)_x.$$

A morphism of sheaves of modules is surjective if and only if it is surjective on every stalk, which gives the assertion. $\square$

Example I.B.3. Let $X = \mathrm{Spec}(A)$ and let M be generated by $m_1, \dots, m_r$. The elements $m_i \in M = \widetilde{M}(X)$ define global sections of $\widetilde{M}$, and their germs generate

$$\widetilde{M}_{\mathfrak{p}} \simeq M_{\mathfrak{p}}$$

for every $\mathfrak{p} \in X$. Hence they generate $\widetilde{M}$. For instance, for $A = \mathbb{k}[x, y]$ and $p = (a, b) \in \mathbb{k}^2$, the sections $x - a$ and $y - b$ generate the quasi-coherent ideal sheaf $\mathcal{I}_p$ which is the sheafification of $\mathfrak{m}_p = (x - a, y - b)$. We have a presentation in terms of free sheaves:

$$0 \rightarrow \mathcal{O}_{\mathbb{A}_{\mathbb{k}}^2} \xrightarrow{\begin{pmatrix} b - y \\ x - a \end{pmatrix}} \mathcal{O}_{\mathbb{A}_{\mathbb{k}}^2}^{\oplus 2} \xrightarrow{(x-a, y-b)} \mathfrak{m}_p \rightarrow 0.$$

Exercise I.B.4 (Generating the structure sheaf). Let $f_1, \dots, f_r \in \Gamma(\mathcal{O}_X)$. Show that they generate $\mathcal{O}_X$ if and only if the open subsets

$$X_{f_i} = \{x \in X \mid (f_i)_x \text{ is invertible in } \mathcal{O}_{X,x}\}$$

cover X . If $X = \mathrm{Spec}(A)$ and the $f_i \in A$, show that this is equivalent to $(f_1, \dots, f_r) = A$.

Solution. The germs $(f_i)_x$ generate the local ring $\mathcal{O}_{X,x}$ precisely when at least one of them is a unit: a proper ideal of a local ring is contained in its maximal ideal. Lemma I.B.2 proves the first assertion. On $\mathrm{Spec}(A)$ one has $X_{f_i} = D(f_i)$, and these principal opens cover X if and only if

$$\mathbb{W}(f_1, \dots, f_r) = \emptyset,$$

equivalently $(f_1, \dots, f_r) = A$.

I.B.2 Finite type and finite presentation

An $\mathcal{O}_X$ -module $\mathcal{F}$ is *of finite type* if every point has an open neighbourhood U on which there is a surjection

$$\mathcal{O}_U^{\oplus r} \twoheadrightarrow \mathcal{F}|_U$$

for some finite r . It is *of finite presentation* if locally there is an exact sequence

$$\mathcal{O}_U^{\oplus m} \longrightarrow \mathcal{O}_U^{\oplus r} \longrightarrow \mathcal{F}|_U \longrightarrow 0$$

with m and r finite. Thus finite presentation asks for finitely many local generators and finitely many relations among them.

Example I.B.5. If M is a finitely presented A -module with presentation

$$A^{\oplus m} \xrightarrow{P} A^{\oplus r} \longrightarrow M \longrightarrow 0,$$

then localisation gives the corresponding presentation of M_f over A_f for every $f \in A$. The associated sheaf therefore has a presentation

$$\mathcal{O}_{\mathrm{Spec}(A)}^{\oplus m} \xrightarrow{\tilde{P}} \mathcal{O}_{\mathrm{Spec}(A)}^{\oplus r} \longrightarrow \widetilde{M} \longrightarrow 0.$$

For $M = A/(f_1, \dots, f_m)$, the first map is the row $(f_1, \dots, f_m)$.

I.B.3 Quasi-coherent sheaves on an affine scheme

Write (Mod_A) for the category of A -modules and $(\mathrm{QCoh}(X))$ for the category of quasi-coherent $\mathcal{O}_X$ -modules. We say that a functor is *exact* if it sends exact sequences to exact sequences.

Theorem I.B.6 (The sheaves-modules equivalence for affine schemes). *Let A be a ring and set $X = \mathrm{Spec}(A)$. The functors*

$$(\mathrm{Mod}_A) \xrightarrow{\widetilde{(-)}} (\mathrm{QCoh}(X)), \quad (\mathrm{QCoh}(X)) \xrightarrow{\Gamma} (\mathrm{Mod}_A)$$

are exact quasi-inverse equivalences. Under this equivalence, finitely generated modules correspond to quasi-coherent sheaves of finite type, and finitely presented modules correspond to quasi-coherent sheaves of finite presentation.

Proof. Let M be an A -module. By construction,

$$\Gamma(\widetilde{M}) = \widetilde{M}(D(1)) = M.$$

The key statement is that, for a quasi-coherent sheaf $\mathcal{F}$ and $f \in A$, the natural homomorphism

$$\Gamma(\mathcal{F})_f \longrightarrow \mathcal{F}(D(f))$$

is an isomorphism. To see this, choose a finite principal-open covering on which $\mathcal{F}$ is associated with modules. The sheaf axiom expresses global sections as the equaliser of the sections on this covering and on its pairwise intersections. Localising that finite equaliser at f gives the corresponding equaliser over $D(f)$, because localisation is exact. This is precisely the displayed homomorphism.

Taking $M = \Gamma(\mathcal{F})$, these isomorphisms on all principal opens assemble to a natural isomorphism $\widetilde{M} \simeq \mathcal{F}$. Thus the two functors are quasi-inverse. The functor $M \mapsto \widetilde{M}$ is exact because its maps on stalks are localisations, and localisation is exact. Its quasi-inverse Γ is therefore exact on quasi-coherent sheaves over X .

Finally, finite generators and finite relations sheafify to local finite presentations. Conversely, a finite principal-open cover and the usual clearing-of-denominators argument turn local finite generators and relations into a finite presentation of $\Gamma(\mathcal{F})$. This proves the last assertion. $\square$

Exercise I.B.7 (Modules and sheaves on an affine scheme). Let $X = \operatorname{Spec}(A)$ and let M, N be A -modules.

- (i) Prove that the natural map

$$\operatorname{Hom}_A(M, N) \longrightarrow \operatorname{Hom}_{\mathcal{O}_X}(\widetilde{M}, \widetilde{N})$$

is a bijection.

- (ii) If $M = A/(f_1, \dots, f_r)$, compute the stalk $\widetilde{M}_{\mathfrak{p}}$ and show that it is zero precisely when some f_i does not belong to $\mathfrak{p}$.
- (iii) Explain why kernels, cokernels and short exact sequences of quasi-coherent sheaves on X may be computed on the corresponding A -modules.

Solution. For a sheaf morphism $\widetilde{M} \rightarrow \widetilde{N}$, take global sections and use Theorem I.B.6; this is inverse to sheafification and proves (i). For (ii),

$$\widetilde{M}_{\mathfrak{p}} \simeq M_{\mathfrak{p}} \simeq A_{\mathfrak{p}}/(f_1, \dots, f_r)A_{\mathfrak{p}}.$$

This module is zero exactly when the ideal generated by the f_i becomes the unit ideal in $A_{\mathfrak{p}}$, equivalently when at least one f_i is not in $\mathfrak{p}$. Part (iii) follows from the exactness of the two quasi-inverse functors.

Exercise I.B.8. Let $\mathcal{F}$ be a quasi-coherent $\mathcal{O}_X$ -module.

- (i) Show that $\mathcal{F}|_U$ is quasi-coherent for every open subset $U \subset X$.
- (ii) Let $X = \bigcup_i U_i$ be an open covering. Suppose that $\mathcal{F}|_{U_i}$ is quasi-coherent for every i . Show that $\mathcal{F}$ is quasi-coherent.

Solution. Both assertions follow from the fact that quasi-coherence may be checked on an affine open covering. For the first, if $x \in U$, choose an affine open neighbourhood V of x on which $\mathcal{F}$ is associated with a module. A principal open neighbourhood $D(f)$ of x contained in $U \cap V$ is affine, and

$$\mathcal{F}|_{D(f)} \simeq \widetilde{M_f}.$$

For the second, cover every U_i by affine opens on which $\mathcal{F}|_{U_i}$ is associated with a module. Together these affine opens cover X , proving that $\mathcal{F}$ is quasi-coherent.

I.C Closed subschemes

Definition I.C.1. A morphism $i: Z \rightarrow X$ is a *closed immersion* if X admits an affine open covering $U_{\alpha} = \operatorname{Spec}(A_{\alpha})$ such that $i^{-1}(U_{\alpha})$ is affine and the homomorphism

$$A_{\alpha} \longrightarrow \Gamma(i^{-1}(U_{\alpha}), \mathcal{O}_Z)$$

induced by i is surjective. Equivalently, on every such chart the restriction of i is isomorphic over U_{α} to

$$\operatorname{Spec}(A_{\alpha}/\mathfrak{a}_{\alpha}) \longrightarrow \operatorname{Spec}(A_{\alpha})$$

for some ideal $\mathfrak{a}_{\alpha} \subset A_{\alpha}$. A *closed subscheme* of X is a scheme Z equipped with a closed immersion $i: Z \hookrightarrow X$.

Let X be a scheme. A *sheaf of ideals* of $\mathcal{O}_X$ is a subsheaf $\mathcal{I} \subset \mathcal{O}_X$ such that $\mathcal{I}(U)$ is an ideal of $\mathcal{O}_X(U)$ for every open subset $U \subset X$. It is *quasi-coherent* if it is quasi-coherent as an $\mathcal{O}_X$ -module. Thus, on an affine open $U = \operatorname{Spec}(A)$,

$$\mathcal{I}|_U \simeq \widetilde{\mathfrak{a}}$$

for an ideal $\mathfrak{a} \subset A$, and $\mathcal{I}(D(f)) = \mathfrak{a}A_f$.

Proposition I.C.2. *Closed subschemes of X , up to unique isomorphism over X , are in bijection with quasi-coherent sheaves of ideals of $\mathcal{O}_X$. The correspondence sends a closed immersion $i: Z \hookrightarrow X$ to*

$$\mathcal{I}_{Z/X} = \ker(\mathcal{O}_X \longrightarrow i_*\mathcal{O}_Z).$$

Conversely, if $\mathcal{I}|_U \simeq \tilde{\mathfrak{a}}$ on $U = \operatorname{Spec}(A)$, the corresponding closed subscheme restricts to $\operatorname{Spec}(A/\mathfrak{a}) \hookrightarrow \operatorname{Spec}(A)$.

Proof. Let $i: Z \hookrightarrow X$ be a closed immersion. On an affine chart $U = \operatorname{Spec}(A)$ as in the definition, write

$$i^{-1}(U) = \operatorname{Spec}(A/\mathfrak{a}).$$

The morphism $\mathcal{O}_X \rightarrow i_*\mathcal{O}_Z$ restricts to the sheafification of $A \rightarrow A/\mathfrak{a}$. It is therefore surjective, and its kernel on U is $\tilde{\mathfrak{a}}$. Hence $\mathcal{I}_{Z/X}$ is a quasi-coherent ideal sheaf and

$$i_*\mathcal{O}_Z \simeq \mathcal{O}_X/\mathcal{I}_{Z/X}.$$

Conversely, let $\mathcal{I} \subset \mathcal{O}_X$ be quasi-coherent. Define a closed subset

$$|Z| = \{x \in X \mid \mathcal{I}_x \subset \mathfrak{m}_{X,x}\}.$$

On $U = \operatorname{Spec}(A)$ with $\mathcal{I}|_U \simeq \tilde{\mathfrak{a}}$, this subset is $\mathbb{V}(\mathfrak{a})$. If $j: |Z| \hookrightarrow X$ is the inclusion, put

$$\mathcal{O}_Z = j^{-1}(\mathcal{O}_X/\mathcal{I}).$$

Locally, $(|Z|, \mathcal{O}_Z)$ is $\operatorname{Spec}(A/\mathfrak{a})$, so it is a scheme and $j: Z \hookrightarrow X$ is a closed immersion. The two constructions are inverse on every affine open chart, and therefore globally. $\square$

Example I.C.3 (Restricting a closed subscheme). Let $i: Z \hookrightarrow X$ be the closed subscheme defined by a quasi-coherent ideal sheaf $\mathcal{I}$, and let $U \subset X$ be open. Then $Z \cap U = i^{-1}(U)$ is a closed subscheme of U , defined by $\mathcal{I}|_U$. Hence closed subschemes restrict naturally to open subsets; on an affine chart $V = \operatorname{Spec}(A)$ this is just the familiar passage from an ideal $\mathfrak{a} \subset A$ to its localisations $\mathfrak{a}_f$.

Exercise I.C.4 (Gluing closed subschemes). Let $X = \bigcup_i U_i$ be an open covering. For every i , let $Z_i \hookrightarrow U_i$ be a closed subscheme, and suppose that the restrictions of Z_i and Z_j to $U_i \cap U_j$ agree. Show that there is a unique closed subscheme $Z \hookrightarrow X$ whose restriction to U_i is Z_i .

Solution. Let $\mathcal{I}_i \subset \mathcal{O}_{U_i}$ be the quasi-coherent ideal sheaf defining Z_i . The hypothesis identifies $\mathcal{I}_i|_{U_i \cap U_j}$ with $\mathcal{I}_j|_{U_i \cap U_j}$. The sheaf gluing axiom therefore gives a unique sheaf of ideals $\mathcal{I} \subset \mathcal{O}_X$ such that $\mathcal{I}|_{U_i} = \mathcal{I}_i$. It is quasi-coherent because this property is local. The ideal sheaf $\mathcal{I}$ defines the required closed subscheme. Uniqueness follows from the uniqueness of the glued ideal sheaf and the correspondence between quasi-coherent ideal sheaves and closed subschemes.

Definition I.C.5. A morphism $j: Z \rightarrow X$ is an *immersion* (or *locally closed immersion*) if it factors as

$$Z \xrightarrow{i} U \xrightarrow{u} X,$$

where i is a closed immersion and u is an open immersion. A *subscheme* of X is a scheme Z equipped with an immersion $Z \rightarrow X$. Thus open and closed subschemes are the two special cases.

Remark I.C.6. The factorisation is local on X : the image of an immersion is a locally closed subset, and Z is a closed subscheme of a suitable open subscheme $U \subset X$.

Exercise I.C.7 (A closed subscheme and its first thickening). Let $\mathcal{I} \subset \mathcal{O}_X$ be a quasi-coherent ideal sheaf. Denote by $Z = \mathbb{V}(\mathcal{I})$ and $Z^{(2)} = \mathbb{V}(\mathcal{I}^2)$ the corresponding closed subschemes.

- (i) Show that $\mathcal{I}^2$ is quasi-coherent.
- (ii) Prove that Z and $Z^{(2)}$ have the same underlying closed subset.
- (iii) Find an example where $\mathcal{I} \neq 0$ and $Z = Z^{(2)}$.

Solution. On an affine open $U = \text{Spec}(A)$, write $\mathcal{I}|_U \simeq \tilde{\mathfrak{a}}$. Then

$$\mathcal{I}^2|_U \simeq \widetilde{(\mathfrak{a}^2)},$$

so $\mathcal{I}^2$ is quasi-coherent. Since $\sqrt{\mathfrak{a}^2} = \sqrt{\mathfrak{a}}$, the closed subsets $\mathbb{W}(\mathfrak{a})$ and $\mathbb{W}(\mathfrak{a}^2)$ agree on every affine chart. Thus $|Z| = |Z^{(2)}|$. Closed subschemes correspond to their ideal sheaves, so $\mathcal{I}^2 \neq \mathcal{I}$ implies that Z and $Z^{(2)}$ are different as closed subschemes of X . For $X = \text{Spec}(\mathbb{k}[t])$ and $\mathcal{I} = (t)$, they are

$$\text{Spec}(\mathbb{k}[t]/(t)) \quad \text{and} \quad \text{Spec}(\mathbb{k}[t]/(t^2)).$$

The first is a reduced point and the second is a non-reduced double point with the same support.

An example as requested is the following. Take $X = \text{Spec}(\mathbb{k} \times \mathbb{k})$ and $\mathcal{I} = \mathbb{k} \times \{0\}$. Then $\mathcal{I}^2 = \mathcal{I}$.

Exercise I.C.8 (The local form of a closed immersion). Let $i: Z \hookrightarrow X$ be the closed subscheme defined by a quasi-coherent ideal sheaf $\mathcal{I}$.

- (i) Show that $|i|: |Z| \rightarrow |X|$ is a homeomorphism onto the closed subset

$$\{x \in X \mid \mathcal{I}_x \subset \mathfrak{m}_{X,x}\}.$$

- (ii) If $U \subset X$ is open, show that $i^{-1}(U)$ is the closed subscheme of U defined by $\mathcal{I}|_U$.
- (iii) For $z \in Z$, prove that

$$\mathcal{O}_{Z,z} \simeq \mathcal{O}_{X,i(z)}/\mathcal{I}_{i(z)} \quad \text{and} \quad \kappa(z) \simeq \kappa(i(z)).$$

Solution. All three assertions are local on X . Take an affine neighbourhood $U = \text{Spec}(A)$ of $i(z)$ on which $\mathcal{I}$ corresponds to an ideal $\mathfrak{a} \subset A$. Then

$$i^{-1}(U) = \text{Spec}(A/\mathfrak{a}) \longrightarrow \text{Spec}(A).$$

Prime ideals of $A/\mathfrak{a}$ correspond to the primes of A containing $\mathfrak{a}$, and this identifies $\text{Spec}(A/\mathfrak{a})$ homeomorphically with $\mathbb{W}(\mathfrak{a})$. Restricting to an open subset localises both A and $\mathfrak{a}$, which proves the second assertion. Finally, if z corresponds to $\bar{\mathfrak{p}} = \mathfrak{p}/\mathfrak{a} \subset A/\mathfrak{a}$, then

$$(A/\mathfrak{a})_{\bar{\mathfrak{p}}} \simeq A_{\mathfrak{p}}/\mathfrak{a}A_{\mathfrak{p}},$$

where $\bar{\mathfrak{p}}$ denotes the image of $\mathfrak{p}$ in $A/\mathfrak{a}$. Taking the quotient by the maximal ideal gives the same residue field $\text{Frac}(A/\mathfrak{p})$ on both sides.

Exercise I.C.9 (Unions and intersections of closed subschemes). Let $Z_1, Z_2 \hookrightarrow X$ be closed subschemes with quasi-coherent ideal sheaves $\mathcal{I}_1, \mathcal{I}_2$.

- (i) Show that $\mathcal{I}_1 + \mathcal{I}_2$ and $\mathcal{I}_1 \cap \mathcal{I}_2$ are quasi-coherent.
- (ii) Define the scheme-theoretic intersection and union by

$$Z_1 \cap Z_2 = \mathbb{W}(\mathcal{I}_1 + \mathcal{I}_2), \quad Z_1 \cup Z_2 = \mathbb{W}(\mathcal{I}_1 \cap \mathcal{I}_2),$$

and show that their underlying subsets are respectively $|Z_1| \cap |Z_2|$ and $|Z_1| \cup |Z_2|$.

- (iii) For the two coordinate axes in $\mathbb{A}_{\mathbb{k}}^2$, compute both closed subschemes and compare their local rings at the origin.

Solution. On an affine open $U = \operatorname{Spec}(A)$, let $\mathcal{I}_1$ and $\mathcal{I}_2$ correspond to ideals $\mathfrak{a}_1, \mathfrak{a}_2 \subset A$. Localisation commutes with finite sums and intersections, so the two sheaves restrict to

$$\widetilde{\mathfrak{a}_1 + \mathfrak{a}_2} \quad \text{and} \quad \widetilde{\mathfrak{a}_1 \cap \mathfrak{a}_2}.$$

They are therefore quasi-coherent. The identities

$$\mathbb{V}(\mathfrak{a}_1 + \mathfrak{a}_2) = \mathbb{V}(\mathfrak{a}_1) \cap \mathbb{V}(\mathfrak{a}_2), \quad \mathbb{V}(\mathfrak{a}_1 \cap \mathfrak{a}_2) = \mathbb{V}(\mathfrak{a}_1) \cup \mathbb{V}(\mathfrak{a}_2)$$

prove the assertion about underlying subsets.

For the axes, $\mathfrak{a}_1 = (x)$ and $\mathfrak{a}_2 = (y)$. Hence their intersection as closed subschemes is defined by

$$(x) + (y) = (x, y),$$

whereas their union is defined by

$$(x) \cap (y) = (xy).$$

At the origin the union has local ring

$$\mathbb{k}[x, y]_{(x, y)} / (xy),$$

while the two axes have local rings $\mathbb{k}[y]_{(y)}$ and $\mathbb{k}[x]_{(x)}$. The first ring has two branches and zero divisors; the other two are domains.

Example I.C.10 (A sheaf of ideals which is not quasi-coherent). Let

$$A = \mathbb{k}[x]_{(x)}, \quad X = \operatorname{Spec}(A).$$

The scheme X has two points: the generic point $\eta = (0)$ and the closed point $s = (x)$. Its nonempty open subsets are

$$U_0 = D(x) = \{\eta\}, \quad X,$$

with

$$\mathcal{O}_X(U_0) = \mathbb{k}(x), \quad \mathcal{O}_X(X) = A.$$

Define a sheaf of ideals $\mathcal{I}$ by

$$\mathcal{I}(X) = 0, \quad \mathcal{I}(U_0) = \mathbb{k}(x),$$

with the only possible restriction map. This is a sheaf of ideals, but it is not quasi-coherent: if $\mathcal{I} \simeq \widetilde{\mathfrak{a}}$ for an ideal $\mathfrak{a} \subset A$, then $\mathfrak{a} = \mathcal{I}(X) = 0$, hence $\mathcal{I}(U_0) = \mathfrak{a}A_x = 0$, a contradiction.

I.D Gluing schemes

Further reading. For gluing and the basic non-affine examples, see [EH00, Chapter I] and [Liu02, Chapter 2]. The construction of projective space by affine charts is also discussed in [Har77, Chapter II, §2].

I.D.1 The gluing lemma

Many schemes are constructed by gluing simpler schemes along open subschemes. The tool to do this is the following lemma.

Lemma I.D.1 (Gluing lemma). *Let $(X_i)_{i \in I}$ be a family of schemes. For every $i, j \in I$, let $X_{ij} \subset X_i$ be an open subscheme and let*

$$\varphi_{ij}: X_{ij} \xrightarrow{\sim} X_{ji}$$

be an isomorphism. Assume that

$$(i) \quad X_{ii} = X_i \text{ and } \varphi_{ii} = \text{id}_{X_i};$$

$$(ii) \quad \varphi_{ji} = \varphi_{ij}^{-1};$$

(iii) for all i, j, k , one has

$$\varphi_{ij}(X_{ij} \cap X_{ik}) = X_{ji} \cap X_{jk}$$

and, on $X_{ij} \cap X_{ik}$:

$$\varphi_{jk} \circ \varphi_{ij} = \varphi_{ik}$$

Then there exists a scheme X with an open covering $X = \bigcup_{i \in I} U_i$ and isomorphisms $X_i \simeq U_i$ which identify the transition maps on $U_i \cap U_j$ with the maps φ_{ij} . The scheme X is unique up to unique isomorphism compatible with these identifications.

Proof. We consider the disjoint union of the underlying topological spaces:

$$\tilde{X} = \coprod_{i \in I} X_i$$

If $x \in X_i$ and $y \in X_j$, define $x \sim y$ if $x \in X_{ij}$ and $y = \varphi_{ij}(x)$. We first check that this is an equivalence relation. It is reflexive because $X_{ii} = X_i$, and $\varphi_{ii} = \text{id}_{X_i}$. It is symmetric because $\varphi_{ji} = \varphi_{ij}^{-1}$. For transitivity, suppose that

$$x \in X_i, \quad y \in X_j, \quad z \in X_k$$

satisfy $y = \varphi_{ij}(x)$, and $z = \varphi_{jk}(y)$. Then

$$y \in X_{ji} \cap X_{jk}.$$

By the hypothesis

$$\varphi_{ij}(X_{ij} \cap X_{ik}) = X_{ji} \cap X_{jk},$$

it follows that

$$x \in X_{ij} \cap X_{ik}.$$

The cocycle condition therefore gives

$$z = \varphi_{jk}(\varphi_{ij}(x)) = \varphi_{ik}(x),$$

so $x \sim z$. Thus $\sim$ is an equivalence relation.

As next step, we consider the quotient space

$$X = \tilde{X} / \sim$$

and denote by $q : \tilde{X} \rightarrow X$ the quotient map. We equip X with the quotient topology. For each $i \in I$, let $q_i : X_i \rightarrow X$ be the restriction of q , and put $U_i = q_i(X_i)$. We claim that U_i is open and that

$$q_i : X_i \rightarrow U_i$$

is a homeomorphism. Indeed, for every $j \in I$, $q^{-1}(U_i) \cap X_j = X_{ji}$. The right-hand side is open in X_j , and hence $q^{-1}(U_i)$ is open in the disjoint union. By definition of the quotient topology, U_i is open in X .

The map q_i is clearly surjective. It is injective because if $x, x' \in X_i$ and $q_i(x) = q_i(x')$, then

$$x' = \varphi_{ii}(x) = x.$$

Finally, let $W \subset X_i$ be open. For every $j \in I$, one has

$$q^{-1}(q_i(W)) \cap X_j = \varphi_{ij}(W \cap X_{ij}),$$

which is open in X_j . Thus $q_i(W)$ is open in X . Hence q_i is an open continuous bijection, and therefore a homeomorphism onto U_i . Under these identifications,

$$U_i \cap U_j = q_i(X_{ij}) = q_j(X_{ji}),$$

and the transition homeomorphism

$$q_j^{-1} \circ q_i : X_{ij} \rightarrow X_{ji}$$

is precisely φ_{ij} .

We now construct the structure sheaf on X . Let $V \subset X$ be open, and set

$$V_i = q_i^{-1}(V \cap U_i) \subset X_i.$$

Then $\varphi_{ij}(V_i \cap X_{ij}) = V_j \cap X_{ji}$. Define $\mathcal{O}_X(V)$ to be the set of families

$$(s_i)_{i \in I}, \quad s_i \in \mathcal{O}_{X_i}(V_i),$$

such that, for every $i, j \in I$,

$$\varphi_{ij}^\#(s_j|_{V_j \cap X_{ji}}) = s_i|_{V_i \cap X_{ij}}.$$

Here φ_{ij} is the morphism induced by the isomorphism $\varphi_{ij}^\# : \mathcal{O}_{X_j}(V_j \cap X_{ji}) \rightarrow \mathcal{O}_{X_i}(V_i \cap X_{ij})$. If $V' \subset V$, restriction is defined componentwise:

$$(s_i)_{i \in I} \mapsto (s_i|_{V'_i})_{i \in I}.$$

The compatibility condition is preserved by restriction, so this defines a presheaf of rings on X .

Let us check the sheaf axioms. We start by the locality axiom. Suppose first that $V = \bigcup_{\alpha \in L} V^\alpha$ is an open covering of V and that two elements of $\mathcal{O}_X(V)$ have the same restriction to V^α , for all $\alpha \in L$. For each $i \in I$, their i -th components agree on the open cover

$$V_i = \bigcup_{\alpha \in L} q_i^{-1}(V^\alpha \cap U_i).$$

Since $\mathcal{O}_{X_i}$ is a sheaf, their i -th components are equal. Hence the two families are equal.

Concerning the gluing axiom, suppose that we are given sections

$$s^\alpha \in \mathcal{O}_X(V^\alpha), \quad \alpha \in L,$$

with the condition that on intersections $V^\alpha \cap V^\beta$ these sections agree, for all $\alpha, \beta \in L$. For each $i \in I$, their i -th components glue, by the sheaf axiom for $\mathcal{O}_{X_i}$, to a section $s_i \in \mathcal{O}_{X_i}(V_i)$. The transition compatibility for the s_i may be checked on the open cover induced by covering $(V^\alpha \mid \alpha \in L)$ of V , where it holds by hypothesis. Therefore the family $(s_i)_{i \in I}$ belongs to $\mathcal{O}_X(V)$, and it is the unique section restricting to all the s^α . Thus $\mathcal{O}_X$ is a sheaf of rings.

Next, we show that the restriction of $\mathcal{O}_X$ to U_i is naturally isomorphic to $\mathcal{O}_{X_i}$. Let $W \subset U_i$ be an open subset and, for $i \in I$, write $W_i = q_i^{-1}(W)$. Then projection onto the i -th component gives a homomorphism

$$\mathcal{O}_X(W) \longrightarrow \mathcal{O}_{X_i}(W_i).$$

This homomorphism is injective: if the i -th component is zero, then the compatibility condition forces every other component to be zero. It is also surjective. Indeed, given $s_i \in \mathcal{O}_{X_i}(W_i)$, define, for every $j \in I$, a section on

$$W_j = q_j^{-1}(W \cap U_j) = \varphi_{ij}(W_i \cap X_{ij})$$

by transporting $s_i|_{W_i \cap X_{ij}}$ through the isomorphism φ_{ij} . The cocycle condition implies that these transported sections are compatible on triple overlaps. Thus they define an element of $\mathcal{O}_X(W)$. Consequently,

$$(U_i, \mathcal{O}_X|_{U_i}) \simeq (X_i, \mathcal{O}_{X_i})$$

as locally ringed spaces. Since the U_i form an open covering of X and each U_i is isomorphic to the scheme X_i , every point of X has an affine open neighbourhood. Hence $(X, \mathcal{O}_X)$ is a scheme.

It remains to prove uniqueness of the scheme we constructed this way. Suppose that Y is another scheme with an open covering

$$Y = \bigcup_{i \in I} V_i$$

and isomorphisms $\alpha_i : X_i \xrightarrow{\sim} V_i$ having transition maps φ_{ij} . The morphisms $\alpha_i \circ q_i^{-1} : U_i \longrightarrow Y$ agree on every intersection $U_i \cap U_j$, precisely because the transition maps are the φ_{ij} . By Exercise I.A.5, they glue uniquely to a morphism

$$F : X \longrightarrow Y.$$

Applying the same construction in the other direction gives a morphism

$$G : Y \longrightarrow X.$$

The compositions $G \circ F$ and $F \circ G$ restrict to the identity on every member of the respective open coverings. Hence

$$G \circ F = \text{id}_X, \quad F \circ G = \text{id}_Y.$$

Thus F is an isomorphism. Its uniqueness follows again because its restriction to every U_i is prescribed. $\square$

Remark I.D.2 (Gluing morphisms). Let X and Y be schemes and let $X = \bigcup_{i \in I} U_i$ be an open covering. Giving a morphism $f : X \rightarrow Y$ is equivalent to giving morphisms $f_i : U_i \rightarrow Y$ such that

$$f_i|_{U_i \cap U_j} = f_j|_{U_i \cap U_j}$$

for every $i, j \in I$. This is the morphism-gluing statement proved in Exercise I.A.5.

I.D.2 The line with two origins

Let $\mathbb{k}$ be a field and let

$$X_1 = \operatorname{Spec}(\mathbb{k}[x_1]), \quad X_2 = \operatorname{Spec}(\mathbb{k}[x_2]).$$

The principal opens $D(x_1)$ and $D(x_2)$ have coordinate rings

$$\mathbb{k}[x_1, x_1^{-1}], \quad \mathbb{k}[x_2, x_2^{-1}].$$

We identify $D(x_1)$ with $D(x_2)$ by $x_1 \mapsto x_2$. The resulting scheme X is called the *affine line with two origins*. Let $o_i \in X_i$ be the origin. The points o_1 and o_2 remain distinct in X , while the complements $X_i \setminus \{o_i\}$ are identified.

A global function on X is a pair

$$(f_1, f_2) \in \mathbb{k}[x_1] \times \mathbb{k}[x_2]$$

whose restrictions agree on the common punctured line. Hence

$$\Gamma(\mathcal{O}_X) = \{(f_1, f_2) \mid f_1 = f_2 \text{ in } \mathbb{k}[x, x^{-1}]\} \simeq \mathbb{k}[x].$$

The two origins cannot be distinguished by global functions. We shall interpret this observation through the affinisation morphism below and use it to prove that X is not affine.

Exercise I.D.3. Let X be the affine line with two origins.

- (i) Show that deleting either origin gives a scheme isomorphic to $\mathbb{A}_{\mathbb{k}}^1$.
- (ii) Show that any open neighbourhood of o_1 meets every open neighbourhood of o_2 .
- (iii) Compute the local rings $\mathcal{O}_{X, o_1}$ and $\mathcal{O}_{X, o_2}$.

Solution. (i) Let X_1 and X_2 be the two affine charts, with origins o_1 and o_2 . After deleting o_1 , the remaining part of X_1 is the common punctured line, which is already contained in the image of X_2 . Hence

$$X \setminus \{o_1\} = X_2 \simeq \mathbb{A}_{\mathbb{k}}^1.$$

Similarly,

$$X \setminus \{o_2\} = X_1 \simeq \mathbb{A}_{\mathbb{k}}^1.$$

- (ii) Let V_1 and V_2 be open neighbourhoods of o_1 and o_2 in X . Then $V_i \cap X_i$ is a nonempty open subset of the irreducible scheme $X_i \simeq \operatorname{Spec}(\mathbb{k}[x_i])$. Every nonempty open subset of X_i contains its generic point. The generic points of X_1 and X_2 are identified in the common punctured line. Therefore this common generic point belongs to $V_1 \cap V_2$, and

$$V_1 \cap V_2 \neq \emptyset.$$

- (iii) Since X_i is an open neighbourhood of o_i , the local ring in X is the same as the local ring in that chart:

$$\mathcal{O}_{X, o_1} \simeq \mathbb{k}[x_1]_{(x_1)}, \quad \mathcal{O}_{X, o_2} \simeq \mathbb{k}[x_2]_{(x_2)}.$$

I.D.3 Projective line

We start with the same ingredients as for the line with two origins, so we let $\mathbb{k}$ be a field and let

$$X_1 = \operatorname{Spec}(\mathbb{k}[x_1]), \quad X_2 = \operatorname{Spec}(\mathbb{k}[x_2]).$$

The principal opens $D(x_1)$ and $D(x_2)$ have coordinate rings

$$\mathbb{k}[x_1, x_1^{-1}], \quad \mathbb{k}[x_2, x_2^{-1}].$$

Now identify $D(x_1)$ with $D(x_2)$ by $x_1 \mapsto x_2^{-1}$. The resulting scheme is the projective line $\mathbb{P}_{\mathbb{k}}^1$.

Its global functions will be computed together with those of $\mathbb{P}_B^n$ in Proposition I.D.6.

Exercise I.D.4. Let U_0 and U_1 be the two standard affine charts of $\mathbb{P}_{\mathbb{k}}^1$.

- (i) Show that $U_0 \cap U_1 \simeq \mathbb{G}_{m, \mathbb{k}}$.
- (ii) Let ∞ be the origin in the second chart. Show that $\mathbb{P}_{\mathbb{k}}^1 \setminus \{\infty\} \simeq \mathbb{A}_{\mathbb{k}}^1$ and compute $\kappa(\infty)$.
- (iii) Describe all closed points of $\mathbb{P}_{\mathbb{k}}^1$ and their residue fields. Which of them are $\mathbb{k}$ -rational?

Solution. Write

$$U_0 = \operatorname{Spec}(\mathbb{k}[x]), \quad U_1 = \operatorname{Spec}(\mathbb{k}[y]),$$

with $x = y^{-1}$ on the overlap.

- (i) The overlap is

$$D(x) = \operatorname{Spec}(\mathbb{k}[x, x^{-1}]) = \mathbb{G}_{m, \mathbb{k}}.$$

- (ii) Every point of U_1 other than its origin belongs to $D(y)$ and hence to U_0 . Therefore

$$\mathbb{P}_{\mathbb{k}}^1 \setminus \{\infty\} = U_0 \simeq \mathbb{A}_{\mathbb{k}}^1.$$

Moreover,

$$\mathcal{O}_{\mathbb{P}_{\mathbb{k}}^1, \infty} \simeq \mathbb{k}[y]_{(y)}, \quad \kappa(\infty) \simeq \mathbb{k}.$$

- (iii) Besides ∞ , the closed points are the points of U_0 defined by monic irreducible polynomials $f \in \mathbb{k}[x]$. Their residue fields are

$$\kappa(x_f) = \mathbb{k}[x]/(f).$$

Consequently, a closed point is $\mathbb{k}$ -rational precisely when it is ∞ or is defined by a linear polynomial $x - a$.

I.D.4 Higher-dimensional projective space

More generally, consider a ring B and the polynomial algebra

$$A = B[x_0, \dots, x_n].$$

For $0 \leq i \leq n$, set

$$R_i = (A[x_i^{-1}])_0 = B \left[\frac{x_0}{x_i}, \dots, \frac{\widehat{x_i}}{x_i}, \dots, \frac{x_n}{x_i} \right] \simeq B[t_1, \dots, t_n]$$

and

$$U_i = \operatorname{Spec}(R_i) \simeq \mathbb{A}_B^n.$$

For $i \neq j$, the principal open subset

$$D\left(\frac{x_j}{x_i}\right) \subset U_i$$

has coordinate ring

$$(A[(x_i x_j)^{-1}])_0,$$

and the same ring is obtained from the j -th chart. These identifications satisfy the cocycle condition, so the U_i glue to a scheme denoted by $\mathbb{P}_B^n$. For the moment, the subscript records the coefficient ring used in the construction. In Subsection II.A.2 we shall glue the compatible maps from the charts to $\text{Spec}(B)$ and thereby make precise in what sense $\mathbb{P}_B^n$ is a scheme over B .

Example I.D.5 (Transition functions on $\mathbb{P}_B^2$). On the chart U_0 , use coordinates

$$u = \frac{x_1}{x_0}, \quad v = \frac{x_2}{x_0}.$$

On U_1 , use

$$s = \frac{x_0}{x_1}, \quad t = \frac{x_2}{x_1}.$$

On the overlap $u \neq 0$, the transition functions are

$$s = u^{-1}, \quad t = vu^{-1}.$$

The subscript 0 means the degree-zero part of the graded ring obtained after inverting the indicated homogeneous element. Explicitly,

$$(A[x_i^{-1}])_0 = \left\{ \frac{a}{x_i^m} \mid a \in A_m \right\}.$$

This notation will be used systematically below; in particular, it is different from the local ring $A_{(f)}$ at the prime ideal (f) .

Proposition I.D.6 (Global functions on projective space). Assume $n \geq 1$. Then

$$\Gamma(\mathcal{O}_{\mathbb{P}_B^n}) = B.$$

In particular, $\Gamma(\mathcal{O}_{\mathbb{P}_{\mathbb{k}}^1}) = \mathbb{k}$.

Proof. Regard all the rings R_i as subrings of

$$B[x_0^{\pm 1}, \dots, x_n^{\pm 1}]_0.$$

Let

$$L = B[x_0^{\pm 1}, \dots, x_n^{\pm 1}]_0.$$

The Laurent monomials

$$x_0^{a_0} \cdots x_n^{a_n}, \quad a_0 + \cdots + a_n = 0,$$

form a B -basis of L . The subring R_i is generated by the monomials x_j/x_i , with $j \neq i$. Thus a Laurent monomial belongs to R_i only if

$$a_j \geq 0 \quad \text{for every } j \neq i.$$

If a Laurent monomial belongs to every R_i , then, since $n \geq 1$, each exponent a_j is nonnegative: choose some $i \neq j$. Since their sum is zero, all the exponents are zero. Consequently, the only Laurent monomial which can occur in an element of every R_i is 1, and therefore

$$\bigcap_{i=0}^n R_i = B.$$

A global section on $\mathbb{P}_B^n$ is a compatible family

$$(s_i), \quad s_i \in R_i.$$

After restricting to the common open subset on which all the x_i are invertible, all the s_i define the same element of L . Conversely, an element of $\bigcap_i R_i$ gives such a compatible family. Hence

$$\Gamma(\mathcal{O}_{\mathbb{P}_B^n}) \simeq \bigcap_{i=0}^n R_i = B.$$

□

I.E Affine targets and affinisation

The following adjunction explains why global functions control every morphism from an arbitrary scheme to an affine scheme.

I.E.1 The affine-target adjunction

Theorem I.E.1 (Morphisms to an affine scheme). *Let X be a scheme and let A be a ring. There is a natural bijection*

$$\mathrm{Hom}_{(\mathrm{Sch})}(X, \mathrm{Spec}(A)) \simeq \mathrm{Hom}_{(\mathrm{Ring})}(A, \Gamma(\mathcal{O}_X)).$$

Proof. A morphism $f: X \rightarrow \mathrm{Spec}(A)$ gives, by taking global sections, a ring homomorphism

$$\varphi_f: A = \Gamma(\mathcal{O}_{\mathrm{Spec}(A)}) \longrightarrow \Gamma(\mathcal{O}_X).$$

Conversely, let

$$\varphi: A \longrightarrow B = \Gamma(\mathcal{O}_X)$$

be a ring homomorphism. For $x \in X$, let

$$\rho_x: B \longrightarrow \mathcal{O}_{X,x}$$

be the canonical map. We define

$$f_\varphi(x) = (\rho_x \circ \varphi)^{-1}(\mathfrak{m}_{X,x}) \in \mathrm{Spec}(A).$$

Let $U = \mathrm{Spec}(R)$ be an affine open subset of X , and let $\rho_U: B \rightarrow R$ be the restriction map. For $a \in A$, set $b = \rho_U(\varphi(a))$. Then

$$f_\varphi^{-1}(D(a)) \cap U = D(b).$$

Indeed, a prime $\mathfrak{p} \subset R$ belongs to the left-hand side if and only if a does not belong to

$$(A \xrightarrow{\varphi} B \xrightarrow{\rho_U} R \longrightarrow R_{\mathfrak{p}})^{-1}(\mathfrak{p}R_{\mathfrak{p}}),$$

which is equivalent to $b \notin \mathfrak{p}$. Hence f_φ is continuous.

On the open subset $D(b) = f_\varphi^{-1}(D(a)) \cap U$, the homomorphism $\rho_U \circ \varphi$ localizes to

$$A_a \longrightarrow R_b = \Gamma(D(b), \mathcal{O}_X).$$

These maps are compatible on intersections of affine open subsets and therefore glue to a morphism of sheaves

$$f_\varphi^\# : \mathcal{O}_{\text{Spec}(A)} \longrightarrow (f_\varphi)_* \mathcal{O}_X.$$

The induced homomorphisms on stalks are local by the definition of $f_\varphi(x)$. Thus $(f_\varphi, f_\varphi^\#)$ is a morphism of schemes. The two constructions are inverse to one another. $\square$

Example I.E.2 (A polynomial map). The homomorphism

$$\mathbb{k}[u, v] \longrightarrow \mathbb{k}[t], \quad u \longmapsto t, \quad v \longmapsto t^2,$$

defines a morphism

$$\mathbb{A}_{\mathbb{k}}^1 \longrightarrow \mathbb{A}_{\mathbb{k}}^2.$$

On closed points, it is the familiar map

$$a \longmapsto (a, a^2).$$

Its image is contained in the closed subscheme defined by $v - u^2$.

Exercise I.E.3. Let $\varphi: A \rightarrow B$ be a ring homomorphism and let $f: \text{Spec}(B) \rightarrow \text{Spec}(A)$ be the associated morphism.

- (i) Prove that $f(\mathfrak{q}) = \varphi^{-1}(\mathfrak{q})$ for every prime ideal $\mathfrak{q} \subset B$.
- (ii) Show that $f^{-1}(D(a)) = D(\varphi(a))$.
- (iii) Verify directly that the induced map $A_a \rightarrow B_{\varphi(a)}$ is the map on sections over these principal open subsets.

Solution. (i) By construction of the morphism associated with a ring homomorphism, its underlying map is contraction of prime ideals:

$$f(\mathfrak{q}) = \varphi^{-1}(\mathfrak{q}).$$

The inverse image is prime because $A/\varphi^{-1}(\mathfrak{q})$ embeds into the integral domain $B/\mathfrak{q}$.

- (ii) Let $a \in A$. Then

$$\begin{aligned} \mathfrak{q} \in f^{-1}(D(a)) &\iff a \notin f(\mathfrak{q}) \\ &\iff a \notin \varphi^{-1}(\mathfrak{q}) \\ &\iff \varphi(a) \notin \mathfrak{q} \\ &\iff \mathfrak{q} \in D(\varphi(a)). \end{aligned}$$

Therefore

$$f^{-1}(D(a)) = D(\varphi(a)).$$

- (iii) We have

$$\mathcal{O}_{\text{Spec}(A)}(D(a)) = A_a, \quad \mathcal{O}_{\text{Spec}(B)}(D(\varphi(a))) = B_{\varphi(a)}.$$

The map on sections sends

$$\frac{c}{a^n} \longmapsto \frac{\varphi(c)}{\varphi(a)^n}.$$

It is exactly the localisation of $\varphi: A \rightarrow B$ at the multiplicative set generated by a .

Exercise I.E.4. Fill in the remaining details in the proof of Theorem I.E.1: verify compatibility on overlaps and check explicitly that the two constructions are inverse.

Solution. Let

$$\varphi : A \longrightarrow \Gamma(\mathcal{O}_X)$$

be a ring homomorphism. For an affine open subset $U = \text{Spec}(R) \subset X$, let $\rho_U : \Gamma(\mathcal{O}_X) \rightarrow R$ be restriction. If $a \in A$ and $b_U = \rho_U(\varphi(a))$, the construction gives a homomorphism

$$A_a \longrightarrow R_{b_U} = \Gamma(D(b_U), \mathcal{O}_X).$$

We first verify compatibility. Let $U = \text{Spec}(R)$ and $V = \text{Spec}(S)$ be affine open subsets, and let W be an affine open subset of

$$D(b_U) \cap D(b_V) \subset U \cap V.$$

Both maps, after restriction to W , send

$$\frac{c}{a^n} \longmapsto \frac{\varphi(c)|_W}{\varphi(a)|_W^n}.$$

They therefore agree on W . Since such affine open subsets cover the overlap, the local morphisms of sheaves agree on overlaps and glue to a morphism

$$f_\varphi^\# : \mathcal{O}_{\text{Spec}(A)} \longrightarrow (f_\varphi)_* \mathcal{O}_X.$$

We now check that the two constructions are inverse. Start with a morphism

$$f : X \longrightarrow \text{Spec}(A)$$

and let

$$\varphi = f_{\text{Spec}(A)}^\# : A \longrightarrow \Gamma(\mathcal{O}_X)$$

be the induced map on global sections. For $x \in X$, put $\mathfrak{p} = f(x)$. The map

$$A \longrightarrow \mathcal{O}_{X,x}$$

factors through the local homomorphism

$$A_{\mathfrak{p}} = \mathcal{O}_{\text{Spec}(A), \mathfrak{p}} \longrightarrow \mathcal{O}_{X,x}.$$

Since this homomorphism is local, the inverse image of $\mathfrak{m}_{X,x}$ in $A_{\mathfrak{p}}$ is $\mathfrak{p}A_{\mathfrak{p}}$. Its inverse image in A is $\mathfrak{p}$. Hence

$$f_\varphi(x) = \varphi_x^{-1}(\mathfrak{m}_{X,x}) = \mathfrak{p} = f(x).$$

Thus the two maps agree on points. On each principal open subset $D(a)$, both maps on sections are obtained by localising the same global homomorphism φ . Therefore the morphisms of sheaves also agree, and $f_\varphi = f$.

Conversely, start with φ . In the construction of f_φ , take the principal open subset $D(1) = \text{Spec}(A)$. The induced map on sections is precisely

$$A = A_1 \longrightarrow \Gamma(\mathcal{O}_X),$$

namely φ . Thus taking global sections recovers the original ring homomorphism.

Exercise I.E.5. Prove that, for every scheme X ,

$$\mathrm{Hom}_{(\mathrm{Sch})}(X, \mathbb{A}_{\mathbb{Z}}^1) \simeq \Gamma(\mathcal{O}_X).$$

Under this bijection, which morphism corresponds to a global section $s \in \Gamma(\mathcal{O}_X)$?

Solution. Since

$$\mathbb{A}_{\mathbb{Z}}^1 = \mathrm{Spec}(\mathbb{Z}[t]),$$

Theorem I.E.1 gives

$$\mathrm{Hom}_{(\mathrm{Sch})}(X, \mathbb{A}_{\mathbb{Z}}^1) \simeq \mathrm{Hom}_{(\mathrm{Ring})}(\mathbb{Z}[t], \Gamma(\mathcal{O}_X)).$$

A ring homomorphism from $\mathbb{Z}[t]$ to $\Gamma(\mathcal{O}_X)$ restricts to $\gamma_{\Gamma(\mathcal{O}_X)}$ on $\mathbb{Z}$ and is uniquely determined by the image of t , which may be any element of $\Gamma(\mathcal{O}_X)$. Hence

$$\mathrm{Hom}_{(\mathrm{Sch})}(X, \mathbb{A}_{\mathbb{Z}}^1) \simeq \Gamma(\mathcal{O}_X).$$

The global section s corresponds to the unique morphism

$$f_s : X \longrightarrow \mathbb{A}_{\mathbb{Z}}^1$$

whose pullback of the coordinate function t is s :

$$f_s^\#(t) = s.$$

Corollary I.E.6. *The scheme $\mathrm{Spec}(\mathbb{Z})$ is a final object of the category (Sch) .*

Proof. For every scheme X , there is the unique unital ring homomorphism

$$\gamma_{\Gamma(\mathcal{O}_X)} : \mathbb{Z} \longrightarrow \Gamma(\mathcal{O}_X).$$

The result follows from Theorem I.E.1. □

I.E.2 Global sections as left adjoint of affine spectrum

Here is a categorical version of the result concerning morphisms towards an affine scheme.

Proposition I.E.7 (Categorical form of the affine-target adjunction). *Taking global sections and taking spectra define functors*

$$\Gamma : (\mathrm{Sch}) \longrightarrow (\mathrm{Ring}^{\mathrm{op}}), \quad \mathrm{Spec} : (\mathrm{Ring}^{\mathrm{op}}) \longrightarrow (\mathrm{Sch}).$$

The functor Γ is left adjoint to Spec , and Spec is fully faithful.

Proof. The adjunction is Theorem I.E.1, written as

$$\mathrm{Hom}_{(\mathrm{Sch})}(X, \mathrm{Spec}(A)) \simeq \mathrm{Hom}_{(\mathrm{Ring}^{\mathrm{op}})}(\Gamma(\mathcal{O}_X), A).$$

Taking $X = \mathrm{Spec}(B)$ and using $\Gamma(\mathcal{O}_{\mathrm{Spec}(B)}) \simeq B$ gives

$$\mathrm{Hom}_{(\mathrm{Sch})}(\mathrm{Spec}(B), \mathrm{Spec}(A)) \simeq \mathrm{Hom}_{(\mathrm{Ring}^{\mathrm{op}})}(B, A),$$

which is the full faithfulness of Spec . □

I.E.3 The affinisation morphism

For every scheme X , the identity of $\Gamma(\mathcal{O}_X)$ corresponds under the affine-target adjunction to a canonical morphism

$$\eta_X : X \longrightarrow \operatorname{Spec}(\Gamma(\mathcal{O}_X)).$$

Remark I.E.8. The morphism η_X is called the *affinisation morphism* of X ; it is the canonical morphism to the affine scheme of global functions. By the construction in Theorem I.E.1, it sends a point $x \in X$ to

$$\eta_X(x) = \ker(\Gamma(\mathcal{O}_X) \longrightarrow \kappa(x)).$$

Thus x is sent to the ideal of global functions vanishing at x .

Example I.E.9. If $X = \operatorname{Spec}(B)$ is affine, then $\Gamma(\mathcal{O}_X) = B$, and η_X is the identity of $\operatorname{Spec}(B)$ under this identification.

Proposition I.E.10 (Affinisation). *The morphisms η_X are natural in X . Moreover, the following conditions are equivalent:*

- i) X is affine;
- ii) η_X is an isomorphism;
- iii) X belongs to the essential image of Spec .

Proof. If $f : X \rightarrow Y$ is a morphism, both composites in

$$\begin{array}{ccc} X & \xrightarrow{\eta_X} & \operatorname{Spec}(\Gamma(\mathcal{O}_X)) \\ f \downarrow & & \downarrow \operatorname{Spec}(\Gamma(f)) \\ Y & \xrightarrow{\eta_Y} & \operatorname{Spec}(\Gamma(\mathcal{O}_Y)) \end{array}$$

correspond to the same homomorphism $\Gamma(\mathcal{O}_Y) \rightarrow \Gamma(\mathcal{O}_X)$, proving naturality. If η_X is an isomorphism, then X is affine. Conversely, for $X = \operatorname{Spec}(B)$ both η_X and the identity correspond to id_B . The last condition is simply the definition of the essential image. $\square$

Exercise I.E.11. Let X be a scheme. Prove directly, using Theorem I.E.1, that every morphism from X to an affine scheme factors uniquely through η_X . More precisely, for every ring A and every morphism

$$f : X \longrightarrow \operatorname{Spec}(A),$$

show that there is a unique morphism

$$\tilde{f} : \operatorname{Spec}(\Gamma(\mathcal{O}_X)) \longrightarrow \operatorname{Spec}(A)$$

such that

$$f = \tilde{f} \circ \eta_X.$$

Solution. Let

$$B = \Gamma(\mathcal{O}_X).$$

By Theorem I.E.1, the morphism

$$f : X \longrightarrow \operatorname{Spec}(A)$$

corresponds to a ring homomorphism

$$\alpha : A \longrightarrow B.$$

Let

$$\tilde{f} = \text{Spec}(\alpha) : \text{Spec}(B) \longrightarrow \text{Spec}(A).$$

The morphism $\eta_X : X \rightarrow \text{Spec}(B)$ corresponds to id_B . Consequently, the composite

$$\tilde{f} \circ \eta_X : X \longrightarrow \text{Spec}(A)$$

corresponds, on global sections, to

$$A \xrightarrow{\alpha} B \xrightarrow{\text{id}_B} B,$$

which is α . Hence

$$f = \tilde{f} \circ \eta_X.$$

For uniqueness, a morphism

$$g : \text{Spec}(B) \longrightarrow \text{Spec}(A)$$

is uniquely determined by the corresponding ring homomorphism $A \rightarrow B$. If $g \circ \eta_X = f$, then the induced homomorphism on global sections is necessarily α . Therefore $g = \text{Spec}(\alpha) = \tilde{f}$.

I.E.4 What affinisation forgets

The examples constructed by gluing now give immediate tests for non-affineness.

Example I.E.12 (The doubled origin and projective space). Let L be the line with two origins. We computed $\Gamma(\mathcal{O}_L) \simeq \mathbb{k}[x]$. The morphism

$$\eta_L : L \longrightarrow \mathbb{A}_{\mathbb{k}}^1$$

restricts to the usual coordinate morphism on both affine charts and hence identifies the two origins. It is not an isomorphism, so L is not affine.

For $n \geq 1$, we also computed

$$\Gamma(\mathcal{O}_{\mathbb{P}_{\mathbb{k}}^n}) = \mathbb{k}.$$

Thus $\eta_{\mathbb{P}_{\mathbb{k}}^n}$ is the unique morphism to $\text{Spec}(\mathbb{k})$. Since projective space has more than one point, this morphism is not an isomorphism, and $\mathbb{P}_{\mathbb{k}}^n$ is not affine.

II Schemes over a base

II.A The category of schemes over a base

Further reading. For schemes over a base, rational points and the functorial viewpoint, see [Liu02, Chapter 2] and [Har77, Chapter II, §3]; the examples in [EH00, Chapter I] are a useful complement.

Let S be a scheme. The category $(\text{Sch})_S$ of *schemes over S* has as objects morphisms $X \rightarrow S$. A morphism

$$(X \rightarrow S) \longrightarrow (Y \rightarrow S)$$

is a morphism $X \rightarrow Y$ commuting with the structure morphisms to S . Since $\text{Spec}(\mathbb{Z})$ is final in (Sch) , the category $(\text{Sch})_{\text{Spec}(\mathbb{Z})}$ is naturally equivalent to (Sch) .

II.A.1 Morphisms of finite type

Definition II.A.1. A morphism $f : X \rightarrow Y$ is *quasi-compact* if $f^{-1}(V)$ is quasi-compact for every affine open subset $V \subset Y$.

Definition II.A.2. A morphism $f : X \rightarrow Y$ is *locally of finite type* if, for every affine open subset $V = \text{Spec}(A) \subset Y$, the inverse image $f^{-1}(V)$ can be covered by affine open subsets $U_i = \text{Spec}(B_i)$ such that every B_i is a finitely generated A -algebra. It is *of finite type* if it is locally of finite type and quasi-compact.

Equivalently, f is of finite type if, for every affine open subset $V = \text{Spec}(A) \subset Y$, the inverse image $f^{-1}(V)$ has a finite affine covering by schemes $\text{Spec}(B_i)$ with B_i finitely generated over A .

Example II.A.3. A morphism of affine schemes

$$\text{Spec}(B) \longrightarrow \text{Spec}(A)$$

is of finite type if and only if B is a finitely generated A -algebra. Thus the projection $\mathbb{A}_A^n \rightarrow \text{Spec}(A)$ is of finite type.

Every closed immersion is of finite type, since a quotient $A \rightarrow A/\mathfrak{a}$ is generated as an A -algebra by 1. Every principal open immersion $D(f) = \text{Spec}(A_f) \rightarrow \text{Spec}(A)$ is also of finite type because

$$A_f \simeq A[t]/(ft - 1).$$

Exercise II.A.4. i) Show that the composition of two morphisms of finite type is of finite type.

ii) Let A be a ring and $f \in A$. Show directly that the open immersion $D(f) \rightarrow \text{Spec}(A)$ is of finite type.

iii) Show that

$$\text{Spec}(A[x_1, x_2]/(x_1x_2 - f)) \longrightarrow \text{Spec}(A)$$

is of finite type.

iv) Give an example of an A -algebra which is not of finite type over A .

Solution. Finite generation of algebras is transitive: if B is generated over A by $b_1, \dots, b_r$ and C is generated over B by $c_1, \dots, c_s$, then C is generated over A by the b_i and the c_j . Applying this on affine charts, and using quasi-compactness, proves i).

For ii), use

$$A_f \simeq A[t]/(ft - 1).$$

The algebra in iii) is generated over A by the classes of x_1, x_2 . Finally, $A[x_1, x_2, x_3, \dots]$ is not finitely generated as an A -algebra.

II.A.2 Schemes over a ring

Let R be a ring. An R -scheme is a scheme over $\text{Spec}(R)$. By Theorem I.E.1, giving an R -scheme structure on X is equivalent to giving a ring homomorphism

$$R \longrightarrow \Gamma(\mathcal{O}_X).$$

If $U = \text{Spec}(B)$ is an affine open subset of X , restriction gives a ring homomorphism $R \rightarrow B$, so B is naturally an R -algebra.

Example II.A.5. The affine line over R is

$$\mathbb{A}_R^1 = \operatorname{Spec}(R[t]),$$

with structure morphism induced by the inclusion $R \rightarrow R[t]$. More generally,

$$\mathbb{A}_R^n = \operatorname{Spec}(R[t_1, \dots, t_n]).$$

The projective space constructed by gluing also has a canonical structure morphism. Indeed, for

$$R_i = R \left[\frac{x_0}{x_i}, \dots, \frac{\widehat{x_i}}{x_i}, \dots, \frac{x_n}{x_i} \right],$$

the inclusions $R \rightarrow R_i$ define morphisms $U_i = \operatorname{Spec}(R_i) \rightarrow \operatorname{Spec}(R)$. They agree on the overlaps used in the construction of $\mathbb{P}_R^n$, and hence glue to a morphism

$$\mathbb{P}_R^n \longrightarrow \operatorname{Spec}(R).$$

This makes $\mathbb{P}_R^n$ an R -scheme and justifies the notation introduced earlier.

Remark II.A.6 (The affine line over a base). For the structure morphism

$$\pi : \mathbb{A}_R^1 = \operatorname{Spec}(R[t]) \longrightarrow \operatorname{Spec}(R),$$

a point $\mathfrak{q}$ maps to $\mathfrak{q} \cap R$. For $\mathfrak{p} \in \operatorname{Spec}(R)$, the points mapping to $\mathfrak{p}$ are governed by

$$\operatorname{Spec}(\kappa(\mathfrak{p})[t]).$$

Thus they consist of its generic point and the closed points defined by irreducible polynomials over $\kappa(\mathfrak{p})$. This is precisely the polynomial-ring calculation of Chapter 2, now interpreted relative to the base; we shall use it without repeating the affine classification.

II.A.3 The relative adjunction

Let $S = \operatorname{Spec}(R)$. The adjunction restricts to an adjunction

$$\operatorname{Spec} : (R\text{-Alg})^{\text{op}} \longrightarrow (\operatorname{Sch})_S, \quad \Gamma : (\operatorname{Sch})_S \longrightarrow (R\text{-Alg})^{\text{op}}.$$

Thus, if X is an R -scheme and A is an R -algebra, then

$$\operatorname{Hom}_{(\operatorname{Sch})_S}(X, \operatorname{Spec}(A)) \simeq \operatorname{Hom}_{(R\text{-Alg})}(A, \Gamma(\mathcal{O}_X)).$$

Exercise II.A.7. Let A and B be R -algebras. Prove that

$$\operatorname{Hom}_{(\operatorname{Sch})_R}(\operatorname{Spec}(B), \operatorname{Spec}(A)) \simeq \operatorname{Hom}_{(R\text{-Alg})}(A, B).$$

Explain precisely where compatibility with the maps from R enters.

Solution. Let

$$\alpha : R \longrightarrow A, \quad \beta : R \longrightarrow B$$

be the given R -algebra structures. By the affine correspondence, a morphism of schemes

$$\operatorname{Spec}(B) \longrightarrow \operatorname{Spec}(A)$$

is equivalent to a ring homomorphism

$$\varphi : A \longrightarrow B.$$

It is a morphism over $\text{Spec}(R)$ precisely when the diagram

$$\begin{array}{ccc} \text{Spec}(B) & \longrightarrow & \text{Spec}(A) \\ & \searrow & \downarrow \\ & & \text{Spec}(R) \end{array}$$

commutes. Passing to rings reverses the arrows, and this condition becomes

$$\varphi \circ \alpha = \beta.$$

This is exactly the condition that φ be a homomorphism of R -algebras. Hence

$$\text{Hom}_{(\text{Sch})_R}(\text{Spec}(B), \text{Spec}(A)) \simeq \text{Hom}_{(R\text{-Alg})}(A, B).$$

II.A.4 Schemes over a field

When $R = \mathbb{k}$ is a field, a $\mathbb{k}$ -scheme is simply a scheme equipped with a morphism to $\text{Spec}(\mathbb{k})$. The principal examples in this course are

$$\mathbb{A}_{\mathbb{k}}^n = \text{Spec}(\mathbb{k}[t_1, \dots, t_n]) \quad \text{and} \quad \mathbb{P}_{\mathbb{k}}^n \longrightarrow \text{Spec}(\mathbb{k}).$$

Their affine charts and all their transition morphisms are morphisms of $\mathbb{k}$ -schemes. The expression $\mathbb{k}$ -scheme by itself imposes no finiteness condition.

Definition II.A.8. An *algebraic $\mathbb{k}$ -scheme* is a scheme of finite type over a field $\mathbb{k}$.

Proposition II.A.9. Let X be an algebraic $\mathbb{k}$ -scheme and let $U = \text{Spec}(A)$ be an affine open subset of X . Then there exist $n, s \geq 0$ and polynomials

$$f_1, \dots, f_s \in \mathbb{k}[x_1, \dots, x_n]$$

such that

$$U \simeq \mathbb{k}[x_1, \dots, x_n]/(f_1, \dots, f_s).$$

Equivalently, every affine open subset of an algebraic $\mathbb{k}$ -scheme is isomorphic to a closed subscheme

$$\mathbb{V}(f_1, \dots, f_s) \subset \mathbb{A}_{\mathbb{k}}^n.$$

Lemma II.A.10. Let $B \rightarrow A$ be a morphism of rings, and let $f_1, \dots, f_r \in A$ generate the unit ideal. Assume that A_{f_i} is a finitely generated B -algebra for every $1 \leq i \leq r$. Then A is a finitely generated B -algebra.

The proof relies on the next lemma.

Proof. Since the elements $f_1, \dots, f_r$ generate the unit ideal, there are elements $c_1, \dots, c_r \in A$ such that

$$1 = c_1 f_1 + \dots + c_r f_r.$$

For $1 \leq i \leq r$, choose finitely many generators of A_{f_i} as a B -algebra. After writing these generators as fractions, we may write

$$A_{f_i} = B \left[\frac{a_{i,1}}{f_i^{n_{i,1}}}, \dots, \frac{a_{i,m_i}}{f_i^{n_{i,m_i}}} \right]$$

for suitable elements $a_{i,j} \in A$ and integers $n_{i,j} \geq 0$. Let $A' \subset A$ be the B -subalgebra generated by the finitely many elements

$$f_i, \quad c_i, \quad a_{i,j},$$

for $1 \leq i \leq r$ and $1 \leq j \leq m_i$. Then A' is a finitely generated B -algebra. We claim that $A' = A$.

For $1 \leq i \leq r$, the inclusion $A' \subset A$ induces an equality $A'_{f_i} = A_{f_i}$. Indeed, the inclusion $A'_{f_i} \subset A_{f_i}$ is clear, and A'_{f_i} contains all the chosen generators $\frac{a_{i,j}}{f_i^{n_{i,j}}}$ of A_{f_i} as a B -algebra.

Consider now the A' -module $M = A/A'$. The equalities $A'_{f_i} = A_{f_i}$ imply that $M_{f_i} = 0$ for $1 \leq i \leq r$. Let $m \in M$. Since $M_{f_i} = 0$, for $1 \leq i \leq r$, there is an integer $N_i \geq 0$ such that

$$f_i^{N_i} m = 0.$$

The elements $f_1, \dots, f_r$ generate the unit ideal of A' as well, because the elements c_i belong to A' and $c_1 f_1 + \dots + c_r f_r = 1$. Therefore the ideal

$$(f_1^{N_1}, \dots, f_r^{N_r}) \subset A'$$

is also the unit ideal: indeed, its radical contains $(f_1, \dots, f_r) = A'$. Hence there are elements $d_1, \dots, d_r \in A'$ such that $1 = d_1 f_1^{N_1} + \dots + d_r f_r^{N_r}$. Multiplying this equality by m , we obtain

$$m = d_1 f_1^{N_1} m + \dots + d_r f_r^{N_r} m = 0.$$

Thus $M = 0$, so $A = A'$. Since A' is a finitely generated B -algebra, the same holds for A . $\square$

Proof of Proposition II.A.9. Since X is of finite type over $\mathbb{k}$ and open immersions are of finite type, the open subscheme U is of finite type over $\mathbb{k}$. Thus U has an affine open covering by schemes $\text{Spec}(B_i)$ with each B_i a finitely generated $\mathbb{k}$ -algebra. Refining this covering by principal open subsets of $U = \text{Spec}(A)$ and using quasi-compactness, we obtain $f_1, \dots, f_r \in A$ such that

$$U = D(f_1) \cup \dots \cup D(f_r)$$

and every

$$A_{f_i} = \mathcal{O}_U(D(f_i))$$

is a finitely generated $\mathbb{k}$ -algebra. The elements $f_1, \dots, f_r$ generate the unit ideal. Lemma II.A.10 therefore shows that A is a finitely generated $\mathbb{k}$ -algebra.

Choose algebra generators $a_1, \dots, a_n$ of A . They define a surjective $\mathbb{k}$ -algebra homomorphism

$$\mathbb{k}[x_1, \dots, x_n] \twoheadrightarrow A, \quad x_i \mapsto a_i.$$

Its kernel is finitely generated, since $\mathbb{k}[x_1, \dots, x_n]$ is Noetherian by Hilbert's basis theorem. Writing the kernel as $(g_1, \dots, g_s)$ gives

$$A \simeq \mathbb{k}[x_1, \dots, x_n]/(g_1, \dots, g_s),$$

and hence the asserted description as a closed subscheme of affine space. $\square$

II.B The functor of points

Let $p : X \rightarrow S$ be an S -scheme.

Definition II.B.1. An S -valued point of X is a morphism of S -schemes $s : S \rightarrow X$. Equivalently, it is a section of the structure morphism p , that is, $p \circ s = \text{id}_S$. The set of S -valued points is denoted by

$$X_S(S) = \text{Hom}_{(\text{Sch})_S}(S, X).$$

More generally, if $T \rightarrow S$ is an S -scheme, a T -valued point of X is a morphism $T \rightarrow X$ over S . We write $X_S(T) = \text{Hom}_{(\text{Sch})_S}(T, X)$. When the base scheme S is understood, we abbreviate $X_S(T)$ to $X(T)$.

Let X be an S -scheme. The valued-point sets introduced above are functorial in the test scheme. They form the *functor of points*

$$h_X : (\text{Sch})_S^{\text{op}} \longrightarrow (\text{Set}), \quad T \longmapsto X(T).$$

If $g : T' \rightarrow T$ is a morphism of S -schemes, then

$$h_X(g) : X(T) \longrightarrow X(T'), \quad u \longmapsto u \circ g.$$

A morphism $f : X \rightarrow Y$ induces a natural transformation

$$h_f : h_X \longrightarrow h_Y, \quad u \longmapsto f \circ u.$$

We mention along the way that Yoneda's lemma says that every natural transformation $h_X \rightarrow h_Y$ arises in this way from a unique morphism $X \rightarrow Y$. Thus a scheme is completely encoded by all its sets of T -valued points, together with their functoriality in T .

If X is a $\mathbb{k}$ -scheme, restricting to affine test schemes gives a covariant functor

$$(\mathbb{k}\text{-Alg}) \longrightarrow (\text{Set}), \quad R \longmapsto X(R).$$

It is important to allow arbitrary $\mathbb{k}$ -algebras, not only fields: points with values in non-reduced rings detect infinitesimal information which is invisible on ordinary rational points.

Example II.B.2 (Affine valued points). For an affine $\mathbb{k}$ -scheme $X = \text{Spec}(A)$ and a $\mathbb{k}$ -algebra R ,

$$X(R) \simeq \text{Hom}_{(\mathbb{k}\text{-Alg})}(A, R).$$

Thus equations defining A may be evaluated in R ; in particular, $\mathbb{A}_{\mathbb{k}}^n(R) = R^n$. This is the affine dictionary developed in Chapter 2, and we now turn to examples which are not affine.

Definition II.B.3 (The multiplicative group). For a ring R , set

$$\mathbb{G}_{m,R} = \text{Spec}(R[t, t^{-1}]).$$

For every R -algebra A ,

$$\mathbb{G}_{m,R}(A) \simeq A^{\times}.$$

This is a useful affine example of the functorial viewpoint: multiplication and inversion of units come from morphisms of schemes.

Exercise II.B.4. Let $D = \text{Spec}(\mathbb{k}[\varepsilon]/(\varepsilon^2))$. Show that $D(\mathbb{k})$ consists of a single point. If

$$R = \mathbb{k}[\delta]/(\delta^2),$$

show that

$$D(R) = \{b\delta \mid b \in \mathbb{k}\}.$$

Conclude that the functor of points distinguishes D from $\text{Spec}(\mathbb{k})$, even though the two schemes have the same underlying topological space and the same set of $\mathbb{k}$ -rational points.

Solution. An element of $D(R)$ is a $\mathbb{k}$ -algebra homomorphism

$$\mathbb{k}[\varepsilon]/(\varepsilon^2) \longrightarrow R,$$

and is therefore determined by an element $r \in R$ satisfying $r^2 = 0$. For $R = \mathbb{k}$, the only such element is 0. For $R = \mathbb{k}[\delta]/(\delta^2)$, an element has the form $a + b\delta$, and its square is zero only if $a = 0$. Hence the square-zero elements are precisely $b\delta$, with $b \in \mathbb{k}$.

II.C Field-valued and rational points

Definition II.C.1. Let X be a scheme and let K be a field. A K -rational point of X is a K -valued point, namely an element of

$$X(\mathbb{k}) = \text{Hom}_{(\text{Sch})}(\text{Spec}(K), X).$$

If X is a $\mathbb{k}$ -scheme, we talk about K -points of fields K which are extensions of $\mathbb{k}$. More generally, for a $\mathbb{k}$ -algebra R , we put

$$X(R) = \text{Hom}_{(\text{Sch})_{\mathbb{k}}}(\text{Spec}(R), X).$$

Let X be a $\mathbb{k}$ -scheme. A K -rational point determines a point $x \in X$ together with a $\mathbb{k}$ -algebra homomorphism

$$\kappa(x) \longrightarrow K.$$

Indeed, given a morphism $f : \text{Spec}(K) \rightarrow X$, we consider its image, which is a point $x \in X$, and the morphism of stalks $f_x^\# : \mathcal{O}_{X,x} \rightarrow \mathcal{O}_{\text{Spec}(K), (0)} \simeq K$. Then the pull-back of the maximal ideal $(0) \subset K$ is the maximal ideal of $\mathcal{O}_{X,x}$, so $f_x^\#$ factors as $\mathcal{O}_{X,x} \rightarrow \kappa(x) \rightarrow K$, where $\mathcal{O}_{X,x} \rightarrow \kappa(x)$ is the obvious quotient map and $\kappa(x) \rightarrow K$ is a field homomorphism, which is necessarily injective. Thus the $\mathbb{k}$ -rational points of X correspond to the points $x \in X$ whose residue field is contained in K , together with an injection $\kappa(x) \hookrightarrow K$. Conversely, given a point $x \in X$, an injection $\kappa(x) \hookrightarrow K$ gives a map $\mathcal{O}_{X,x} \rightarrow K$ by composition and thus a morphism of schemes $\text{Spec}(K) \rightarrow X$.

If $K = \mathbb{k}$, then $\mathbb{k}$ -points are just points $x \in X$ whose residue field $\kappa(x)$ is $\mathbb{k}$. In particular, a point of the underlying topological space of X need not be $\mathbb{k}$ -rational.

Proposition II.C.2 (Closed points). *Let X be an algebraic $\mathbb{k}$ -scheme. For every closed point $x \in X$, the residue field $\kappa(x)$ is a finite extension of $\mathbb{k}$. The canonical morphism*

$$\text{Spec}(\kappa(x)) \longrightarrow X$$

is therefore a $\kappa(x)$ -valued point of X . Conversely, if $K/\mathbb{k}$ is finite, the image in X of every K -valued point is a closed point.

Proof. Choose an affine open neighbourhood $U = \text{Spec}(A)$ of a closed point x , and let $\mathfrak{m} \subset A$ be the corresponding maximal ideal. The field

$$\kappa(x) = A/\mathfrak{m}$$

is a finitely generated $\mathbb{k}$ -algebra, so Zariski's lemma implies that it is finite over $\mathbb{k}$. The quotient homomorphism $A \rightarrow \kappa(x)$ gives the displayed canonical morphism.

Conversely, let $u : \text{Spec}(K) \rightarrow X$ be a K -valued point and let x be its image. On an affine neighbourhood $\text{Spec}(A)$ of x , the morphism is induced by a homomorphism $A \rightarrow K$. If $K/\mathbb{k}$ is finite, its image is a finite-dimensional domain over $\mathbb{k}$, hence a field. The kernel is therefore maximal, so x is closed. $\square$

Remark II.C.3. The finiteness hypothesis on the value field matters. The inclusion

$$\mathbb{k}[t] \longrightarrow \mathbb{k}(t)$$

defines a $\mathbb{k}(t)$ -valued point of $\mathbb{A}_{\mathbb{k}}^1$ whose image is the generic point (0) , which is not closed.

Example II.C.4 (Points of projective space over a field). The projective scheme $\mathbb{P}_{\mathbb{k}}^n$, constructed below from its standard affine charts, satisfies

$$\mathbb{P}_{\mathbb{k}}^n(K) = (K^{n+1} \setminus \{0\})/K^\times$$

for every field extension $K/\mathbb{k}$. We write its elements as $[a_0 : \cdots : a_n]$.

Projective schemes also have closed points which are not rational. In the standard affine chart $\mathbb{A}_{\mathbb{Q}}^1 \subset \mathbb{P}_{\mathbb{Q}}^1$, the irreducible polynomial $t^2 - 2$ defines a closed point with residue field $\mathbb{Q}(\sqrt{2})$, not a $\mathbb{Q}$ -rational point. Similarly, the non-real closed points of $\mathbb{P}_{\mathbb{R}}^1$ defined by irreducible quadratic polynomials have residue field $\mathbb{C}$.

Exercise II.C.5 (A projective conic). Let

$$C = \{[x : y : z] \in \mathbb{P}_{\mathbb{Q}}^2 \mid x^2 + y^2 = z^2\}.$$

For every field extension $K/\mathbb{Q}$, interpret $C(K)$ as projective solutions of this equation. Show that

$$\mathbb{P}_{\mathbb{Q}}^1 \longrightarrow C, \quad [s : t] \longmapsto [s^2 - t^2 : 2st : s^2 + t^2]$$

parametrises the rational points of C .

Solution. A K -valued point of C is a nonzero triple $(a, b, c) \in K^3$ satisfying $a^2 + b^2 = c^2$, modulo simultaneous multiplication by an element of $K^\times$. The displayed formula gives such a triple.

Conversely, let $[a : b : c] \in C(K)$. If $c + a \neq 0$, take

$$[s : t] = [c + a : b].$$

Using $b^2 = c^2 - a^2$, the image of this point is

$$[2a(c + a) : 2b(c + a) : 2c(c + a)] = [a : b : c].$$

If $c + a = 0$, then $b = 0$ and the point is $[-1 : 0 : 1]$, which is the image of $[0 : 1]$. Hence every rational point occurs.

III Projective schemes

Further reading. For the Proj construction and projective schemes, see [Har77, Chapter II, §2], [Liu02, Chapter 2], and the concrete treatment in [EH00, Chapter II].

The projective spectrum construction gives an intrinsic description of the projective spaces introduced above and, more generally, of closed projective subschemes.

III.A Graded rings and the homogeneous spectrum

Let B be a ring and let

$$A = \bigoplus_{d \geq 0} A_d$$

be a positively graded B -algebra, so that the structure homomorphism $B \rightarrow A$ has image in A_0 , and assume that $A_p A_q \subset A_{p+q}$.

Definition III.A.1. An element of A_d is called *homogeneous of degree d* . An ideal $\mathfrak{a} \subset A$ is *homogeneous* if it is generated by homogeneous elements. Equivalently, whenever

$$a = a_0 + a_1 + \cdots + a_m \in \mathfrak{a}, \quad a_d \in A_d,$$

then every homogeneous component a_d belongs to $\mathfrak{a}$. The ideal

$$A_+ = \bigoplus_{d > 0} A_d$$

is called the *irrelevant ideal*. A homogeneous prime ideal is *relevant* if it does not contain A_+ .

Definition III.A.2. The *projective spectrum* of A is

$$\mathrm{Proj}(A) = \{\mathfrak{p} \subset A \mid \mathfrak{p} \text{ is homogeneous and prime, and } A_+ \not\subset \mathfrak{p}\}.$$

Example III.A.3. Let $A = \mathbb{k}[x_0, x_1]$ with the standard grading. The ideals

$$(0), \quad (x_0), \quad (x_1)$$

are homogeneous and relevant, while (x_0, x_1) is irrelevant and $(x_0 - 1)$ is not homogeneous. If $\mathbb{k}$ is algebraically closed, the closed points of $\mathrm{Proj}(A)$ are the ideals

$$(bx_0 - ax_1), \quad [a : b] \in \mathbf{P}_{\mathbb{k}}^1.$$

The ideal (0) is the generic point.

Exercise III.A.4. Prove that an ideal $\mathfrak{a} \subset A$ is homogeneous if and only if it contains every homogeneous component of each of its elements.

Solution. Assume first that $\mathfrak{a}$ is generated by homogeneous elements. Let $a \in \mathfrak{a}$. We may write

$$a = \sum_{j=1}^r b_j h_j,$$

where the h_j are homogeneous elements of $\mathfrak{a}$. Decompose every b_j into its homogeneous components and collect terms of the same degree. Each homogeneous component of a is then a sum of multiples of the h_j , and therefore belongs to $\mathfrak{a}$.

Conversely, suppose that $\mathfrak{a}$ contains every homogeneous component of each of its elements. Let H be the set of homogeneous elements of $\mathfrak{a}$. If $a \in \mathfrak{a}$ and

$$a = a_0 + \cdots + a_m$$

is its homogeneous decomposition, then every a_d belongs to H and

$$a = a_0 + \cdots + a_m.$$

Thus every element of $\mathfrak{a}$ belongs to the ideal generated by H . Hence $\mathfrak{a}$ is generated by homogeneous elements and is homogeneous.

Exercise III.A.5. Determine the projective spectra of the following graded rings.

- (i) $\mathbb{k}[x]$ with $\deg(x) = 1$;
- (ii) $\mathbb{k}[x_0, x_1]/(x_0 x_1)$ with the standard grading;
- (iii) $\mathbb{k}[x_0, x_1]/(x_0^2)$ with the standard grading.

Compare the underlying topological spaces in (ii) and (iii) with their scheme-theoretic structures.

Solution. (i) Let $A = \mathbb{k}[x]$ with $\deg(x) = 1$. Its homogeneous prime ideals are (0) and (x) , but (x) contains the irrelevant ideal $A_+ = (x)$ and is therefore not relevant. Thus $\mathrm{Proj}(A)$ has one point. Moreover,

$$D_+(x) = \mathrm{Proj}(A), \quad (A[x^{-1}])_0 = \mathbb{k}.$$

Hence

$$\mathrm{Proj}(\mathbb{k}[x]) \simeq \mathrm{Spec}(\mathbb{k}).$$

(ii) Put

$$A = \mathbb{k}[x_0, x_1]/(x_0x_1).$$

The standard opens satisfy

$$(A[x_0^{-1}])_0 \simeq \mathbb{k}, \quad (A[x_1^{-1}])_0 \simeq \mathbb{k}.$$

Indeed, after inverting x_0 , the relation forces $x_1 = 0$, and similarly with the variables exchanged. Their intersection is empty because $x_0x_1 = 0$. Therefore

$$\mathrm{Proj}(A) \simeq \mathrm{Spec}(\mathbb{k}) \coprod \mathrm{Spec}(\mathbb{k}).$$

Its two points correspond to the homogeneous prime ideals (x_0) and (x_1) , and both are reduced.

(iii) Put

$$A = \mathbb{k}[x_0, x_1]/(x_0^2).$$

Every prime ideal contains the nilpotent element x_0 . A relevant prime must not contain x_1 , so the only relevant homogeneous prime is (x_0) . Thus the underlying topological space consists of one point. The open $D_+(x_1)$ is the whole projective spectrum, and

$$(A[x_1^{-1}])_0 \simeq \mathbb{k} \left[\frac{x_0}{x_1} \right] / \left(\left(\frac{x_0}{x_1} \right)^2 \right).$$

Writing $u = x_0/x_1$, we obtain

$$\mathrm{Proj}(A) \simeq \mathrm{Spec}(\mathbb{k}[u]/(u^2)).$$

Hence (ii) is the disjoint union of two reduced points, whereas (iii) is a single non-reduced double point. In particular, the nilpotent structure in (iii) is invisible in its one-point underlying topological space.

III.B The Zariski topology and the structure sheaf in the graded setting

For a homogeneous ideal $\mathfrak{a} \subset A$, define

$$\mathbb{W}_+(\mathfrak{a}) = \{\mathfrak{p} \in \mathrm{Proj}(A) \mid \mathfrak{a} \subset \mathfrak{p}\}.$$

These subsets are the closed subsets of the Zariski topology on $\mathrm{Proj}(A)$. If $f \in A$ is homogeneous of positive degree, set

$$D_+(f) = \mathrm{Proj}(A) \setminus \mathbb{W}_+(f)$$

and write

$$(A[f^{-1}])_0$$

for the degree-zero part of the graded localisation $A[f^{-1}]$. If f has degree d , its elements are represented by fractions a/f^m with $a \in A_{md}$.

Exercise III.B.1. Prove the identities

$$\mathbb{W}_+(\mathfrak{a}) \cup \mathbb{W}_+(\mathfrak{b}) = \mathbb{W}_+(\mathfrak{a}\mathfrak{b}), \quad \bigcap_{\lambda} \mathbb{W}_+(\mathfrak{a}_{\lambda}) = \mathbb{W}_+\left(\sum_{\lambda} \mathfrak{a}_{\lambda}\right).$$

Deduce that the sets $D_+(f)$ form a basis of the Zariski topology.

Solution. Let $\mathfrak{p} \in \text{Proj}(A)$. Since $\mathfrak{p}$ is prime,

$$\mathfrak{a}\mathfrak{b} \subset \mathfrak{p} \iff \mathfrak{a} \subset \mathfrak{p} \text{ or } \mathfrak{b} \subset \mathfrak{p}.$$

Therefore

$$\mathbb{W}_+(\mathfrak{a}\mathfrak{b}) = \mathbb{W}_+(\mathfrak{a}) \cup \mathbb{W}_+(\mathfrak{b}).$$

Similarly,

$$\sum_{\lambda} \mathfrak{a}_{\lambda} \subset \mathfrak{p} \iff \mathfrak{a}_{\lambda} \subset \mathfrak{p} \text{ for every } \lambda,$$

so

$$\mathbb{W}_+\left(\sum_{\lambda} \mathfrak{a}_{\lambda}\right) = \bigcap_{\lambda} \mathbb{W}_+(\mathfrak{a}_{\lambda}).$$

If $\mathfrak{a}$ is homogeneous, it is generated by homogeneous elements. Hence

$$\text{Proj}(A) \setminus \mathbb{W}_+(\mathfrak{a}) = \{\mathfrak{p} \mid \mathfrak{a} \not\subset \mathfrak{p}\} = \bigcup_{\substack{f \in \mathfrak{a} \\ f \text{ homogeneous}}} D_+(f).$$

Thus every open subset is a union of subsets of the form $D_+(f)$. Moreover,

$$D_+(f) \cap D_+(g) = D_+(fg),$$

so these subsets form a basis of the Zariski topology.

Lemma III.B.2 (Affine charts of Proj). *For every homogeneous element $f \in A$ of positive degree, there is a natural homeomorphism*

$$D_+(f) \simeq \text{Spec}((A[f^{-1}])_0).$$

Under this homeomorphism, the principal open subsets on both sides correspond.

Proof. Put $d = \deg(f)$, and $R_f = (A[f^{-1}])_0$. We construct mutually inverse correspondences between the points of $D_+(f)$ and the prime ideals of R_f . Let $\mathfrak{p} \in D_+(f)$. Thus $\mathfrak{p} \subset A$ is a homogeneous prime ideal and $f \notin \mathfrak{p}$. Its extension

$$\mathfrak{p}A[f^{-1}] \subset A[f^{-1}]$$

is a homogeneous prime ideal. Taking its degree-zero part gives a prime ideal

$$\Phi_f(\mathfrak{p}) = (\mathfrak{p}A[f^{-1}])_0 \subset R_f.$$

Indeed, if $u, v \in R_f$ and $uv \in \Phi_f(\mathfrak{p})$, then $uv \in \mathfrak{p}A[f^{-1}]$. Since $\mathfrak{p}A[f^{-1}]$ is prime, either $u \in \mathfrak{p}A[f^{-1}]$ or $v \in \mathfrak{p}A[f^{-1}]$, and hence either $u \in \Phi_f(\mathfrak{p})$ or $v \in \Phi_f(\mathfrak{p})$.

Let us construct the inverse map. Let $\mathfrak{q} \subset R_f$ be a prime ideal. For every homogeneous element $a \in A$ of degree n , the fraction

$$\frac{a^d}{f^n}$$

has degree zero, since $\deg(a^d) = nd = \deg(f^n)$. Define $\Psi_f(\mathfrak{q}) \subset A$ to be the homogeneous ideal whose homogeneous elements are characterised by

$$a \in \Psi_f(\mathfrak{q}) \iff \frac{a^d}{f^n} \in \mathfrak{q}, \quad \deg(a) = n.$$

Let us check that this does define a homogeneous ideal. Suppose that $a, b \in A$ are homogeneous of degree n and that $a, b \in \Psi_f(\mathfrak{q})$. We must prove that $a + b \in \Psi_f(\mathfrak{q})$. Expanding, we obtain

$$\frac{(a+b)^d}{f^n} = \sum_{i=0}^d \binom{d}{i} \frac{a^i b^{d-i}}{f^n}.$$

For $0 \leq i \leq d$, the fraction

$$\frac{a^i b^{d-i}}{f^n}$$

has degree zero, and its d -th power is

$$\left(\frac{a^i b^{d-i}}{f^n} \right)^d = \left(\frac{a^d}{f^n} \right)^i \left(\frac{b^d}{f^n} \right)^{d-i}.$$

This belongs to $\mathfrak{q}$. Since $\mathfrak{q}$ is prime, and hence radical, it follows that

$$\frac{a^i b^{d-i}}{f^n} \in \mathfrak{q}.$$

Consequently

$$\frac{(a+b)^d}{f^n} \in \mathfrak{q},$$

so $a + b \in \Psi_f(\mathfrak{q})$.

If $a \in A_n$ belongs to $\Psi_f(\mathfrak{q})$ and $c \in A_m$ is homogeneous, then

$$\frac{(ac)^d}{f^{n+m}} = \frac{a^d}{f^n} \frac{c^d}{f^m} \in \mathfrak{q}.$$

Thus $ac \in \Psi_f(\mathfrak{q})$. This proves that $\Psi_f(\mathfrak{q})$ is a homogeneous ideal.

It is proper, because

$$\frac{1^d}{f^0} = 1 \notin \mathfrak{q}.$$

Moreover, it is prime. Indeed, if $a \in A_n$ and $b \in A_m$ are homogeneous and $ab \in \Psi_f(\mathfrak{q})$, then

$$\frac{a^d}{f^n} \frac{b^d}{f^m} = \frac{(ab)^d}{f^{n+m}} \in \mathfrak{q}.$$

Since $\mathfrak{q}$ is prime, either

$$\frac{a^d}{f^n} \in \mathfrak{q} \quad \text{or} \quad \frac{b^d}{f^m} \in \mathfrak{q}.$$

Therefore either $a \in \Psi_f(\mathfrak{q})$ or $b \in \Psi_f(\mathfrak{q})$. Since $\Psi_f(\mathfrak{q})$ is homogeneous, this implies that it is a prime ideal.

Finally,

$$f \notin \Psi_f(\mathfrak{q}),$$

because

$$\frac{f^d}{f^d} = 1 \notin \mathfrak{q}.$$

Hence $\Psi_f(\mathfrak{q})$ is a point of $D_+(f)$.

We now check that the two constructions are inverse.

Let $\mathfrak{p} \in D_+(f)$ and let $a \in A_n$ be homogeneous. Then

$$\begin{aligned} a \in \Psi_f(\Phi_f(\mathfrak{p})) &\iff \frac{a^d}{f^n} \in (\mathfrak{p}A[f^{-1}])_0 \\ &\iff a^d \in \mathfrak{p} \\ &\iff a \in \mathfrak{p}. \end{aligned}$$

Here we use that $f \notin \mathfrak{p}$ and that $\mathfrak{p}$ is prime. Thus

$$\Psi_f(\Phi_f(\mathfrak{p})) = \mathfrak{p}.$$

Conversely, let $\mathfrak{q} \in \text{Spec}(R_f)$ and put $\mathfrak{p} = \Psi_f(\mathfrak{q})$. Every element of R_f may be written as

$$u = \frac{a}{f^m}, \quad a \in A_{md}.$$

Then

$$\begin{aligned} u \in (\mathfrak{p}A[f^{-1}])_0 &\iff a \in \mathfrak{p} \\ &\iff \frac{a^d}{f^{md}} \in \mathfrak{q} \\ &\iff u^d \in \mathfrak{q} \\ &\iff u \in \mathfrak{q}. \end{aligned}$$

The last equivalence follows because $\mathfrak{q}$ is radical. Hence

$$\Phi_f(\Psi_f(\mathfrak{q})) = \mathfrak{q}.$$

We have therefore obtained a bijection

$$D_+(f) \longrightarrow \text{Spec}(R_f).$$

It remains to check the topology. Let $g \in A$ be homogeneous of positive degree, say

$$e = \deg(g).$$

The element

$$h_{f,g} = \frac{g^d}{f^e}$$

belongs to R_f . If $\mathfrak{p} \in D_+(f)$ and $\mathfrak{q} = \Phi_f(\mathfrak{p})$, then

$$\begin{aligned} \mathfrak{p} \in D_+(g) &\iff g \notin \mathfrak{p} \\ &\iff g^d \notin \mathfrak{p} \\ &\iff \frac{g^d}{f^e} \notin \mathfrak{q} \\ &\iff \mathfrak{q} \in D(h_{f,g}). \end{aligned}$$

Thus

$$D_+(f) \cap D_+(g) \longleftrightarrow D(h_{f,g}) \subset \text{Spec}(R_f).$$

The standard opens form a basis on the left, and the principal opens form a basis on the right. Hence the bijection is a homeomorphism, and it identifies the indicated basic open subsets. $\square$

Lemma III.B.3 (The standard-open sheaf on Proj). *On the basis of standard open subsets of $\text{Proj}(A)$, the prescription*

$$D_+(f) \longmapsto (A[f^{-1}])_0$$

satisfies the sheaf condition and therefore extends uniquely to a sheaf of rings on $\text{Proj}(A)$. On every $D_+(f)$, the resulting ringed space is the affine scheme

$$\text{Spec}((A[f^{-1}])_0).$$

Proof. For a homogeneous element $f \in A$ of positive degree, put

$$R_f = (A[f^{-1}])_0.$$

By Lemma III.B.2, there is a homeomorphism

$$D_+(f) \simeq \text{Spec}(R_f).$$

We shall show that, under this homeomorphism, the rings assigned to the standard open subsets of $D_+(f)$ are exactly the rings of sections of the usual affine structure sheaf on $\text{Spec}(R_f)$.

Let $g \in A$ be homogeneous of positive degree, and write

$$d = \deg(f), \quad e = \deg(g).$$

As in the proof of Lemma III.B.2, put

$$h_{f,g} = \frac{g^d}{f^e} \in R_f.$$

The intersection

$$D_+(f) \cap D_+(g) = D_+(fg)$$

corresponds under the homeomorphism $D_+(f) \simeq \text{Spec}(R_f)$ to the principal open subset

$$D(h_{f,g}) \subset \text{Spec}(R_f).$$

We claim that there is a natural isomorphism

$$(R_f)_{h_{f,g}} \simeq (A[(fg)^{-1}])_0.$$

The inclusion

$$R_f \subset A[(fg)^{-1}]$$

sends $h_{f,g}$ to a unit, because both f and g are invertible in $A[(fg)^{-1}]$. It therefore induces a homomorphism

$$(R_f)_{h_{f,g}} \longrightarrow (A[(fg)^{-1}])_0.$$

To see that this map is surjective, consider a degree-zero element of $A[(fg)^{-1}]$. It may be written as

$$\frac{a}{f^m g^n},$$

where a is homogeneous and

$$\deg(a) = md + ne.$$

Choose N such that $dN \geq n$. Then

$$\frac{a}{f^m g^n} = \frac{ag^{dN-n}}{f^{m+eN}} = \frac{a}{\left(\frac{g^d}{f^e}\right)^N}.$$

The numerator on the right belongs to R_f , because

$$\begin{aligned}\deg(ag^{dN-n}) &= md + ne + (dN - n)e \\ &= d(m + eN).\end{aligned}$$

Thus every degree-zero element of $A[(fg)^{-1}]$ comes from $(R_f)_{h_{f,g}}$.

The map is also injective. Indeed, both rings are obtained from the same subring of $A[(fg)^{-1}]$ by allowing powers of $h_{f,g}$ in the denominator, and the preceding formula gives an inverse to the displayed homomorphism. Hence

$$(R_f)_{h_{f,g}} \simeq (A[(fg)^{-1}])_0.$$

It follows that, on the chart $D_+(f)$, the assignment

$$D_+(f) \cap D_+(g) \longmapsto (A[(fg)^{-1}])_0$$

is identified with the usual assignment

$$D(h_{f,g}) \longmapsto (R_f)_{h_{f,g}}$$

defining the affine structure sheaf on $\text{Spec}(R_f)$.

The subsets

$$D_+(f) \cap D_+(g),$$

as g ranges over the homogeneous elements of positive degree, form a basis of the topology of $D_+(f)$. Therefore the rings above, with the localisation restriction maps, define on $D_+(f)$ precisely the affine structure sheaf

$$\mathcal{O}_{\text{Spec}(R_f)}.$$

We must check that these affine sheaves agree on the overlaps of two charts. On

$$D_+(f) \cap D_+(g) = D_+(fg),$$

the restriction of the affine sheaf from the chart $D_+(f)$ is determined on standard open subsets by rings of the form

$$(A[(fgh)^{-1}])_0.$$

The restriction from the chart $D_+(g)$ is described by exactly the same rings and the same localisation homomorphisms. Thus the two restricted sheaves are canonically isomorphic.

On a triple intersection

$$D_+(f) \cap D_+(g) \cap D_+(h),$$

these canonical identifications satisfy the cocycle condition, because all of them are induced by the natural localisation maps

$$A \longrightarrow A[(fg)^{-1}] \longrightarrow A[(fgh)^{-1}].$$

The affine structure sheaves on the opens $D_+(f)$ therefore glue to a unique sheaf of rings $\mathcal{O}_{\text{Proj}(A)}$ on $\text{Proj}(A)$.

By construction, its restriction to every standard open $D_+(f)$ is the usual affine structure sheaf of

$$\text{Spec}(R_f) = \text{Spec}((A[f^{-1}])_0).$$

In particular,

$$\mathcal{O}_{\text{Proj}(A)}(D_+(f)) = R_f = (A[f^{-1}])_0.$$

This proves the assertion. $\square$

Example III.B.4 (A non-standard grading on a polynomial ring). Let

$$A = \mathbb{k}[x, y], \quad \deg(x) = 1, \quad \deg(y) = 2.$$

The irrelevant ideal is

$$A_+ = (x, y),$$

so the standard open subsets $D_+(x)$ and $D_+(y)$ cover $\text{Proj}(A)$.

After localizing at x , a monomial $x^i y^j$ has degree zero precisely when

$$i + 2j = 0.$$

It follows that

$$(A[x^{-1}])_0 = \mathbb{k} \left[\frac{y}{x^2} \right].$$

Thus

$$D_+(x) \simeq \text{Spec} \left(\mathbb{k} \left[\frac{y}{x^2} \right] \right) \simeq \mathbb{A}_{\mathbb{k}}^1.$$

Similarly,

$$(A[y^{-1}])_0 = \mathbb{k} \left[\frac{x^2}{y} \right],$$

and hence

$$D_+(y) \simeq \mathbb{A}_{\mathbb{k}}^1.$$

On the intersection $D_+(xy)$, the two coordinates

$$u = \frac{y}{x^2}, \quad v = \frac{x^2}{y}$$

satisfy

$$uv = 1.$$

Therefore the two affine lines are glued by the usual transition $v = u^{-1}$, and we obtain

$$\text{Proj}(A) \simeq \mathbb{P}_{\mathbb{k}}^1.$$

Notice that the grading is essential in the computation: for instance, y/x^2 has degree zero, whereas y/x does not.

Example III.B.5 (Weighted projective lines). Let $A = \mathbb{k}[x, y]$, $\deg(x) = a$, and $\deg(y) = b$, where a, b are positive integers. Put $d = \gcd(a, b)$, $a' = a/d$, and $b' = b/d$. Then

$$\text{Proj}(A) \simeq \mathbb{P}_{\mathbb{k}}^1.$$

Indeed, $D_+(x)$ and $D_+(y)$ cover $\text{Proj}(A)$. A monomial $x^i y^j$ in A_x has degree zero if and only if $ai + bj = 0$. Since $\gcd(a', b') = 1$, this implies that $(i, j) = (-b'n, a'n)$ for some integer $n \geq 0$. Consequently,

$$(A[x^{-1}])_0 = \mathbb{k} \left[\frac{y^{a'}}{x^{b'}} \right], \quad \text{and} \quad (A[y^{-1}])_0 = \mathbb{k} \left[\frac{x^{b'}}{y^{a'}} \right].$$

Hence both $D_+(x)$ and $D_+(y)$ are isomorphic to $\mathbb{A}_{\mathbb{k}}^1$. Writing

$$u = \frac{y^{a'}}{x^{b'}}, \quad v = \frac{x^{b'}}{y^{a'}},$$

we have $uv = 1$ on $D_+(xy)$. The two affine charts are therefore glued by $v = u^{-1}$, exactly as in the standard construction of $\mathbb{P}_{\mathbb{k}}^1$.

Exercise III.B.6 (The weighted projective plane $\mathbb{P}(1, 1, 2)$). Let $A = \mathbb{k}[x, y, z]$, with $\deg(x) = \deg(y) = 1$, $\deg(z) = 2$, and set $X = \text{Proj}(A)$. This scheme is usually denoted by $\mathbb{P}(1, 1, 2)$.

(i) Compute the rings $(A[x^{-1}])_0$ and $(A[y^{-1}])_0$ and show that $D_+(x) \simeq \mathbb{A}_{\mathbb{k}}^2$, and $D_+(y) \simeq \mathbb{A}_{\mathbb{k}}^2$.

(ii) Show that

$$(A[z^{-1}])_0 = \mathbb{k}\left[\frac{x^2}{z}, \frac{xy}{z}, \frac{y^2}{z}\right], \quad D_+(z) \simeq \text{Spec}\left(\frac{\mathbb{k}[u, v, w]}{(uw - v^2)}\right).$$

In this exercise, although the two charts $D_+(x)$ and $D_+(y)$ are affine planes, the third chart is an affine quadric cone.

Solution. (i) In A_x , the elements

$$\frac{y}{x} \quad \text{and} \quad \frac{z}{x^2}$$

have degree zero. Every degree-zero monomial in A_x is a product of powers of these two elements. Therefore

$$A_{(x)} = \mathbb{k}\left[\frac{y}{x}, \frac{z}{x^2}\right] \simeq \mathbb{k}[s, t].$$

Thus

$$D_+(x) \simeq \mathbb{A}_{\mathbb{k}}^2.$$

Similarly,

$$A_{(y)} = \mathbb{k}\left[\frac{x}{y}, \frac{z}{y^2}\right] \simeq \mathbb{k}[s, t],$$

and hence

$$D_+(y) \simeq \mathbb{A}_{\mathbb{k}}^2.$$

(ii) A monomial

$$\frac{x^i y^j}{z^n}$$

has degree zero precisely when

$$i + j = 2n.$$

Every monomial $x^i y^j$ of even total degree is a product of the monomials x^2 , xy and y^2 . It follows that

$$A_{(z)} = \mathbb{k}\left[\frac{x^2}{z}, \frac{xy}{z}, \frac{y^2}{z}\right].$$

(iii) Set

$$U = \frac{x^2}{z}, \quad V = \frac{xy}{z}, \quad W = \frac{y^2}{z}.$$

Since

$$UW = V^2,$$

there is a surjective homomorphism

$$\varphi: \frac{\mathbb{k}[u, v, w]}{(uw - v^2)} \longrightarrow A_{(z)}$$

defined by

$$u \longmapsto U, \quad v \longmapsto V, \quad w \longmapsto W.$$

We show that φ is injective. Using the relation $uw = v^2$, every monomial in u, v, w can be written in one of the forms

$$u^a v^b \quad \text{or} \quad v^b w^c,$$

where in the second family we may assume $c \geq 1$.

Their images are respectively

$$U^a V^b = \frac{x^{2a+b} y^b}{z^{a+b}}$$

and

$$V^b W^c = \frac{x^b y^{b+2c}}{z^{b+c}}.$$

These are distinct Laurent monomials in x, y, z . Hence they are linearly independent over $\mathbb{k}$. It follows that the kernel of φ is zero.

Therefore

$$A_{(z)} \simeq \frac{\mathbb{k}[u, v, w]}{(uw - v^2)},$$

and consequently

$$D_+(z) \simeq \text{Spec} \left(\frac{\mathbb{k}[u, v, w]}{(uw - v^2)} \right).$$

The sheaf furnished by the lemma is denoted $\mathcal{O}_X$, where $X = \text{Proj}(A)$. It equips the projective spectrum with a scheme structure.

III.C Projective schemes and closed subschemes

Definition III.C.1. A *projective B -scheme* is a B -scheme isomorphic to $\text{Proj}(A)$, where A is a positively graded B -algebra generated over B by finitely many homogeneous elements of positive degree and the structure homomorphism $B \rightarrow A_0$ is surjective. Equivalently, it is a closed B -subscheme of some $\mathbb{P}_B^n$.

Remark III.C.2 (Equations for projective schemes). Let $X = \text{Proj}(A)$ be a projective $\mathbb{k}$ -scheme. Choose homogeneous generators $a_0, \dots, a_n$ of A as a $\mathbb{k}$ -algebra, and put $d_i = \deg(a_i) > 0$. Giving $\mathbb{k}[x_0, \dots, x_n]$ the grading $\deg(x_i) = d_i$, the homomorphism

$$\mathbb{k}[x_0, \dots, x_n] \longrightarrow A, \quad x_i \longmapsto a_i,$$

is graded and surjective. Its kernel is a homogeneous ideal. Since $\mathbb{k}[x_0, \dots, x_n]$ is Noetherian, this kernel is generated by finitely many homogeneous polynomials $f_1, \dots, f_r$. Hence

$$X \simeq \text{Proj} \left(\frac{\mathbb{k}[x_0, \dots, x_n]}{(f_1, \dots, f_r)} \right).$$

Thus every projective $\mathbb{k}$ -scheme is described by finitely many homogeneous equations in variables of positive degrees. If all the degrees d_i are equal to one, this exhibits X as a closed subscheme of $\mathbb{P}_{\mathbb{k}}^n$.

Example III.C.3 (The standard charts of projective space). Let $A = B[x_0, \dots, x_n]$ with the standard grading. Then

$$(A[x_i^{-1}])_0 = B \left[\frac{x_0}{x_i}, \dots, \frac{\widehat{x_i}}{x_i}, \dots, \frac{x_n}{x_i} \right].$$

Thus

$$D_+(x_i) \simeq \mathbb{A}_B^n,$$

and the gluing maps are exactly those used earlier to construct $\mathbb{P}_B^n$. Hence

$$\text{Proj}(B[x_0, \dots, x_n]) \simeq \mathbb{P}_B^n.$$

Example III.C.4 (A projective conic). Let

$$A = \mathbb{k}[x_0, x_1, x_2]/(x_0x_2 - x_1^2)$$

with the standard grading. Then $C = \text{Proj}(A)$ is a closed subscheme of $\mathbb{P}_{\mathbb{k}}^2$. On $D_+(x_0)$, putting

$$u = \frac{x_1}{x_0}, \quad v = \frac{x_2}{x_0},$$

the relation becomes $v = u^2$, so this chart is isomorphic to $\mathbb{A}_{\mathbb{k}}^1$. There is a morphism

$$\pi : C \longrightarrow \mathbb{P}_{\mathbb{k}}^1$$

obtained geometrically by projection from the point $p = [1 : 0 : 0] \in C$. The naive formula $[x_0 : x_1 : x_2] \mapsto [x_1 : x_2]$ is undefined at p , but it extends across p : on the two standard charts of C put

$$\pi|_{D_+(x_0)} = [x_0 : x_1], \quad \pi|_{D_+(x_2)} = [x_1 : x_2].$$

On the overlap these expressions agree because $x_0x_2 = x_1^2$. The gluing statement for morphisms therefore gives a well-defined morphism to the non-affine scheme $\mathbb{P}_{\mathbb{k}}^1$.

Its inverse is the quadratic Veronese morphism

$$\nu : \mathbb{P}_{\mathbb{k}}^1 \longrightarrow C, \quad [s : t] \longmapsto [s^2 : st : t^2].$$

The formulas on the standard charts show that $\pi \circ \nu = \text{id}_{\mathbb{P}_{\mathbb{k}}^1}$ and $\nu \circ \pi = \text{id}_C$. Hence $C \simeq \mathbb{P}_{\mathbb{k}}^1$.

Exercise III.C.5. Let $\mathfrak{a} \subset A$ be a homogeneous ideal. Show that

$$\text{Proj}(A/\mathfrak{a}) \longrightarrow \text{Proj}(A)$$

is a closed immersion with image $\mathbb{W}_+(\mathfrak{a})$.

Solution. The quotient map $A \rightarrow A/\mathfrak{a}$ sends a homogeneous prime ideal of $A/\mathfrak{a}$ to its inverse image in A . This gives a bijection between the relevant homogeneous prime ideals of $A/\mathfrak{a}$ and the relevant homogeneous prime ideals of A containing $\mathfrak{a}$. Hence the image on underlying topological spaces is

$$\mathbb{W}_+(\mathfrak{a}).$$

To check the scheme structure, let $f \in A$ be homogeneous of positive degree and write $\bar{f}$ for its image in $A/\mathfrak{a}$. On the standard affine charts, the morphism is

$$\text{Spec}(((A/\mathfrak{a})[\bar{f}^{-1}])_0) \longrightarrow \text{Spec}((A[f^{-1}])_0).$$

There is a natural isomorphism

$$((A/\mathfrak{a})[\bar{f}^{-1}])_0 \simeq (A[f^{-1}])_0 / (\mathfrak{a}A[f^{-1}])_0.$$

Thus the map on every standard affine chart is induced by a surjective ring homomorphism, and is a closed immersion. These local closed immersions agree on overlaps and glue. Therefore

$$\text{Proj}(A/\mathfrak{a}) \longrightarrow \text{Proj}(A)$$

is a closed immersion with image $\mathbb{W}_+(\mathfrak{a})$.

Exercise III.C.6. Continue the computation for the conic

$$C = \text{Proj}(\mathbb{k}[x_0, x_1, x_2]/(x_0x_2 - x_1^2)).$$

Compute the chart $D_+(x_2)$ and its transition map with $D_+(x_0)$. Verify directly that the two formulas defining the projection $\pi : C \rightarrow \mathbb{P}_{\mathbb{k}}^1$ agree on the overlap, and that

$$\nu : \mathbb{P}_{\mathbb{k}}^1 \longrightarrow C, \quad [s : t] \longmapsto [s^2 : st : t^2]$$

is an isomorphism. Deduce a description of $C(K)$ for every field extension $K/\mathbb{k}$, and describe the residue fields of its closed points.

Solution. Let

$$A = \mathbb{k}[x_0, x_1, x_2]/(x_0x_2 - x_1^2).$$

On $D_+(x_2)$, put

$$v = \frac{x_1}{x_2}, \quad w = \frac{x_0}{x_2}.$$

The relation becomes $w = v^2$, so

$$(A[x_2^{-1}])_0 \simeq \mathbb{k}[v].$$

On $D_+(x_0)$ use $u = x_1/x_0$, so that $x_2/x_0 = u^2$. On the overlap,

$$v = \frac{x_1/x_0}{x_2/x_0} = u^{-1}.$$

The two charts are therefore affine lines glued by inversion, exactly as in the construction of $\mathbb{P}_{\mathbb{k}}^1$. On their overlap,

$$[x_0 : x_1] = [x_1 : x_2] \iff x_0x_2 = x_1^2,$$

so the two formulas for π agree. The inverse morphism is induced by

$$[s : t] \longmapsto [s^2 : st : t^2].$$

For every field extension $K/\mathbb{k}$,

$$C(K) = \{[a : b : c] \in \mathbb{P}_{\mathbb{k}}^2(K) \mid ac = b^2\},$$

and the isomorphism identifies this set with $\mathbb{P}_{\mathbb{k}}^1(K)$. It also preserves local rings and residue fields. Hence the closed points of C correspond to ∞ and to monic irreducible polynomials $f \in \mathbb{k}[t]$; their residue fields are respectively $\mathbb{k}$ and $\mathbb{k}[t]/(f)$. In particular, the $\mathbb{k}$ -rational points are exactly those corresponding to ∞ and to linear polynomials.

III.D Graded maps and rational maps

Definition III.D.1 (Rational map). Let X and Y be schemes. A *rational map* from X to Y is an equivalence class of pairs (U, f) , where $U \subset X$ is a dense open subscheme and $f : U \rightarrow Y$ is a morphism. Two pairs (U, f) and (V, g) are equivalent if f and g agree on a dense open subscheme of $U \cap V$. We write

$$X \dashrightarrow Y.$$

Thus a formula defined only on an open subset represents a rational map only when that open subset is dense.

Definition III.D.2 (Birational map). Let X and Y be integral schemes. A rational map $\varphi : X \dashrightarrow Y$ is *birational* if it admits a rational inverse. Equivalently, there are non-empty open subschemes $U \subset X$ and $V \subset Y$ such that φ is represented by an isomorphism

$$U \xrightarrow{\sim} V.$$

The schemes X and Y are then said to be birational.

Example III.D.3. Let X and Y be schemes over a common base S , and let the S -morphism $f : U \rightarrow Y$ represent a rational map over S . Its *graph morphism* is

$$\Gamma_f = (\text{id}_U, f) : U \longrightarrow X \times_S Y.$$

It factors through $U \times_S Y$. When Y is projective over S , the resulting morphism $U \rightarrow U \times_S Y$ is a closed immersion (a special case of the closed-graph property for separated morphisms). The scheme-theoretic closure of its image in $X \times_S Y$,

$$\overline{\Gamma}_f \subset X \times_S Y$$

depends only on the rational map. The first projection is an isomorphism over U , while the second projection extends f . In this sense the graph closure eliminates indeterminacy.

The contravariance of Spec has a projective analogue, with an important difference: a homomorphism of graded rings need not define a morphism on the whole projective spectrum. However this analogue is defined as a rational map. Let

$$A = \bigoplus_{m \geq 0} A_m, \quad B = \bigoplus_{m \geq 0} B_m$$

be positively graded algebras, and let $\varphi : A \rightarrow B$ be a homogeneous homomorphism of positive degree d , meaning that

$$\varphi(A_m) \subset B_{dm} \quad \text{for every } m \geq 0.$$

Its *base ideal* is the homogeneous ideal

$$\mathfrak{b}_\varphi = \varphi(A_+)B \subset B,$$

and its *base locus* is $\mathbb{W}_+(\mathfrak{b}_\varphi) \subset \text{Proj}(B)$.

Proposition III.D.4 (The map defined by a graded homomorphism). *The homomorphism φ defines a morphism*

$$\text{Proj}(B) \setminus \mathbb{W}_+(\mathfrak{b}_\varphi) \longrightarrow \text{Proj}(A).$$

If its domain is dense in $\text{Proj}(B)$, this morphism determines a rational map

$$\text{Proj}(B) \dashrightarrow \text{Proj}(A).$$

Proof. The complement of the base locus is covered by the open subsets $D_+(\varphi(f))$, where $f \in A_+$ is homogeneous. Homogeneous localisation gives a homomorphism

$$(A[f^{-1}])_0 \longrightarrow (B[\varphi(f)^{-1}])_0,$$

and hence a morphism

$$D_+(\varphi(f)) \longrightarrow D_+(f).$$

These morphisms agree on pairwise overlaps and therefore glue. If their common domain is dense, Definition III.D.1 applies. $\square$

Thus the word “rational” requires a density hypothesis. Without it, the same construction still gives a perfectly good morphism on the displayed open subset, which could even be empty.

Example III.D.5 (Linear projection). Let

$$A = \mathbb{k}[u_1, \dots, u_n], \quad B = \mathbb{k}[x_0, x_1, \dots, x_n], \quad u_i \mapsto x_i.$$

The resulting rational map is

$$\mathbb{P}_{\mathbb{k}}^n \dashrightarrow \mathbb{P}_{\mathbb{k}}^{n-1}, \quad [x_0 : x_1 : \dots : x_n] \mapsto [x_1 : \dots : x_n].$$

Its base locus is the point $[1 : 0 : \dots : 0]$. For $n = 2$, restricting this projection to the conic $x_0x_2 = x_1^2$ gives the map studied in Example III.C.4; on the conic its apparent indeterminacy is removable.

Example III.D.6 (The standard quadratic transformation). The degree-two homomorphism

$$\mathbb{k}[u_0, u_1, u_2] \longrightarrow \mathbb{k}[x_0, x_1, x_2], \quad (u_0, u_1, u_2) \mapsto (x_1x_2, x_0x_2, x_0x_1)$$

defines the standard Cremona transformation

$$\text{Cr} : \mathbb{P}_{\mathbb{k}}^2 \dashrightarrow \mathbb{P}_{\mathbb{k}}^2, \quad [x_0 : x_1 : x_2] \mapsto [x_1x_2 : x_0x_2 : x_0x_1].$$

Its base locus consists of the three coordinate points. On the dense torus $x_0x_1x_2 \neq 0$ it is the inversion map $[x_0 : x_1 : x_2] \mapsto [x_0^{-1} : x_1^{-1} : x_2^{-1}]$, and applying it twice gives the identity.

Suppose now that A and B are graded algebras over a common ring R , that $\text{Proj}(B)$ is integral, and that the domain U of the rational map is dense. The preceding construction gives the graph model

$$\bar{\Gamma}_{\varphi} \subset \text{Proj}(B) \times_{\text{Spec}(R)} \text{Proj}(A).$$

For the standard Cremona transformation, one can show that this graph model is the blow-up of the three base points and is nonsingular; in that example it really is a resolution of indeterminacies.

Exercise III.D.7 (Projection and its graph). For the projection $\mathbb{P}_{\mathbb{k}}^2 \dashrightarrow \mathbb{P}_{\mathbb{k}}^1$ from $p = [1 : 0 : 0]$:

- (i) compute the base ideal and base locus;
- (ii) show that the graph closure in $\mathbb{P}_{\mathbb{k}}^2 \times \mathbb{P}_{\mathbb{k}}^1$ is defined by

$$x_1v - x_2u = 0,$$

where $[u : v]$ are the coordinates on $\mathbb{P}^1$;

- (iii) show that the first projection has one-point fibres away from p and fibre $\mathbb{P}_{\mathbb{k}}^1$ over p .

Solution. The graded homomorphism sends u to x_1 and v to x_2 , so its base ideal is (x_1, x_2) and its base locus is p . On the domain of the projection, the equality $[u : v] = [x_1 : x_2]$ is equivalent to $x_1v - x_2u = 0$. The same bihomogeneous equation defines its closure. If $[x_0 : x_1 : x_2] \neq p$, it determines the unique point $[u : v] = [x_1 : x_2]$. Over p the equation becomes $0 = 0$, so every $[u : v] \in \mathbb{P}_{\mathbb{k}}^1$ occurs.

Exercise III.D.8 (The standard Cremona transformation). For the map in Example III.D.6:

- (i) determine the base scheme and its underlying points;
- (ii) verify directly that Cr^2 is the identity wherever both sides are defined;

(iii) determine the image of each coordinate line away from the base points.

Solution. The base ideal is

$$(x_1x_2, x_0x_2, x_0x_1),$$

whose zero set consists of the three coordinate points. Applying the formula twice gives

$$[x_0x_1x_2x_0 : x_0x_1x_2x_1 : x_0x_1x_2x_2] = [x_0 : x_1 : x_2]$$

on the dense torus, hence wherever the two rational maps are defined. The line $x_i = 0$, with its two base points removed, is sent to the i -th coordinate point: for example, $x_0 = 0$ gives $[x_1x_2 : 0 : 0] = [1 : 0 : 0]$.

IV Invertible sheaves and projective space

We now develop the sheaf-theoretic language needed for the intrinsic description of morphisms to projective space.

IV.A Invertible sheaves

Definition IV.A.1. An $\mathcal{O}_X$ -module $\mathcal{L}$ is *invertible* if every point of X has an open neighbourhood U such that $\mathcal{L}|_U \simeq \mathcal{O}_U$. Equivalently, $\mathcal{L}$ is locally free of rank one. An invertible sheaf is also called a *line bundle*.

Remark IV.A.2. Choose a covering $X = \bigcup_i U_i$ and generators e_i of $\mathcal{L}|_{U_i}$. On an overlap there is a unique unit $g_{ij} \in \mathcal{O}_X(U_i \cap U_j)^\times$ such that

$$e_j = g_{ij}e_i.$$

For $i, j, k \in I$, the transition functions satisfy

$$g_{ii} = 1, \quad g_{ji} = g_{ij}^{-1}, \quad g_{ik} = g_{jk}g_{ij}.$$

Conversely, units satisfying these identities glue the modules $\mathcal{O}_{U_i}$ to an invertible sheaf.

Remark IV.A.3. The tensor product and dual are again invertible, with

$$\mathcal{L}^{-1} = \mathcal{H}om_{\mathcal{O}_X}(\mathcal{L}, \mathcal{O}_X), \quad \mathcal{L} \otimes \mathcal{L}^{-1} \simeq \mathcal{O}_X.$$

IV.B Twisting sheaves on projective space

Let $U_i = D_+(x_i)$ be the standard affine covering of $\mathbb{P}_B^n$. For $d \in \mathbb{Z}$, take a free generator $e_i^{(d)}$ on U_i and glue by

$$e_j^{(d)} = \left(\frac{x_j}{x_i}\right)^d e_i^{(d)} \quad \text{on } U_i \cap U_j.$$

The resulting invertible sheaf is denoted by $\mathcal{O}_{\mathbb{P}_B^n}(d)$. The transition functions immediately give

$$\mathcal{O}(d) \otimes \mathcal{O}(e) \simeq \mathcal{O}(d+e), \quad \mathcal{O}(d)^{-1} \simeq \mathcal{O}(-d).$$

If $F \in B[x_0, \dots, x_n]_d$ is homogeneous of degree $d \geq 0$, the local expressions

$$\frac{F}{x_i^d} e_i^{(d)}$$

agree on overlaps and define a global section of $\mathcal{O}(d)$. In fact, for $n \geq 1$,

$$\Gamma(\mathcal{O}_{\mathbb{P}_B^n}(d)) \simeq \begin{cases} B[x_0, \dots, x_n]_d, & d \geq 0, \\ 0, & d < 0. \end{cases}$$

For $\mathbb{P}_{\mathbb{C}}^1$ this recovers the computation of $\Gamma(\mathcal{O}_{\mathbb{P}_{\mathbb{C}}^1}(d))$ made in Chapter 1. In particular, $x_0, \dots, x_n$ are generating sections of $\mathcal{O}(1)$.

Exercise IV.B.1. Using the standard affine covering of $\mathbb{P}_B^1$, prove the displayed formula for $\Gamma(\mathcal{O}_{\mathbb{P}_B^1}(d))$. Show explicitly that the global sections of $\mathcal{O}(d)$ for $d \geq 0$ have basis

$$x_0^d, x_0^{d-1}x_1, \dots, x_1^d.$$

Solution. Write $t = x_1/x_0$ on U_0 and $s = x_0/x_1 = t^{-1}$ on U_1 . A global section is a pair $f(t)e_0^{(d)}, g(s)e_1^{(d)}$ satisfying

$$f(t) = t^d g(t^{-1}).$$

If $d \geq 0$, the common Laurent polynomial has exponents between 0 and d , giving the indicated $d+1$ homogeneous monomials. If $d < 0$, the two allowed ranges of exponents are disjoint and the section is zero.

IV.C Morphisms to projective space

The affine-target adjunction describes a morphism to affine space by global functions. For projective space the homogeneous coordinates are local and may be multiplied simultaneously by a unit. Invertible sheaves record precisely these changes of scale.

Proposition IV.C.1 (Morphisms to projective space). *Let X be a B -scheme. Morphisms of B -schemes $X \rightarrow \mathbb{P}_B^n$ are in natural bijection with equivalence classes of data*

$$(\mathcal{L}; s_0, \dots, s_n),$$

where $\mathcal{L}$ is an invertible sheaf and $s_0, \dots, s_n$ generate it (see §). Two such data are equivalent if an isomorphism of invertible sheaves carries each s_i to the corresponding section.

Proof. Given $(\mathcal{L}; s_0, \dots, s_n)$, let U_i be the open subset where s_i generates $\mathcal{L}$. On U_i the ratios s_j/s_i are regular functions and define a morphism

$$U_i \longrightarrow D_+(x_i) \subset \mathbb{P}_B^n.$$

On $U_i \cap U_j$, the two homogeneous tuples differ by multiplication by the unit s_j/s_i , so the local morphisms agree and glue.

Conversely, if $f : X \rightarrow \mathbb{P}_B^n$, take

$$\mathcal{L} = f^* \mathcal{O}_{\mathbb{P}_B^n}(1), \quad s_i = f^* x_i.$$

Since the x_i generate $\mathcal{O}(1)$, the sections s_i generate $\mathcal{L}$. The two constructions are inverse up to the stated equivalence. $\square$

Thus a tuple of global functions with no common zero describes only the special case $\mathcal{L} \simeq \mathcal{O}_X$. The line-bundle formulation is what makes the statement valid for an arbitrary, possibly non-affine source.

Example IV.C.2 (The Veronese morphism). The monomials

$$x_0^d, x_0^{d-1}x_1, \dots, x_1^d$$

generate $\mathcal{O}_{\mathbb{P}_{\mathbb{C}}^1}(d)$ and therefore define the degree- d Veronese morphism $\mathbb{P}_{\mathbb{C}}^1 \rightarrow \mathbb{P}_{\mathbb{C}}^d$. For $d = 2$ its image is the conic $z_0 z_2 = z_1^2$.

Chapter 4

First properties of schemes

I Connected and reduced schemes

Further reading. For connectedness, idempotents, reduction and the first topological properties of schemes, see [Liu02, Chapter 2], [Har77, Chapter II, §3], and the examples in [EH00, Chapter I].

Connectedness of a scheme is the connectedness of its underlying topological space. Reducedness, on the other hand, is an algebraic property: it requires the absence of nilpotent elements in all local rings.

I.A Connected schemes

Definition I.A.1. A scheme X is *connected* if its underlying topological space is connected.

Global idempotents detect connectedness even when the scheme is not affine.

Proposition I.A.2. A scheme X is connected if and only if $\Gamma(\mathcal{O}_X)$ has no non-trivial idempotents.

Proof. Let $X = U \amalg V$ be a decomposition into two non-empty open and closed subsets of X . Restriction gives a morphism $\Gamma(\mathcal{O}_X) \rightarrow \Gamma(\mathcal{O}_U) \times \Gamma(\mathcal{O}_V)$. Also, from sections $(u, v) \in \Gamma(\mathcal{O}_U) \times \Gamma(\mathcal{O}_V)$ we construct uniquely a global section $f \in \Gamma(\mathcal{O}_X)$ by gluing u and v , since the compatibility condition is void. Therefore, we have

$$\Gamma(\mathcal{O}_X) \simeq \Gamma(\mathcal{O}_U) \times \Gamma(\mathcal{O}_V).$$

The section corresponding to $(1, 0)$ is a non-trivial idempotent.

Conversely, let $e \in \Gamma(\mathcal{O}_X)$ satisfy $e^2 = e$. For every point $x \in X$, the stalk $\mathcal{O}_{X,x}$ is a local ring, so e_x is either 0 or 1. Therefore we have disjoint subsets of X :

$$U_0 = \{x \in X \mid e_x = 0\}, \quad U_1 = \{x \in X \mid e_x = 1\}$$

They are open, because equality of germs holds on a neighbourhood. They are thus also closed, because they are complement of each other in X . If e is non-trivial, both are non-empty, so X is disconnected. $\square$

Remark I.A.3 (The affine special case). For $X = \operatorname{Spec}(A)$ the proposition recovers the familiar equivalences

$$X \text{ disconnected} \iff A \text{ has a non-trivial idempotent} \iff A \simeq B \times C$$

with $B, C \neq 0$. The corresponding formula

$$\operatorname{Spec}(B \times C) \simeq \operatorname{Spec}(B) \amalg \operatorname{Spec}(C)$$

was established in Chapter 2.

I.B Reduced schemes

Reducedness is studied through nilpotent elements. The topological space underlying a scheme can always be equipped with a canonical reduced scheme structure obtained by killing nilpotents.

Definition I.B.1. A scheme X is *reduced* if, for every $x \in X$, the local ring $\mathcal{O}_{X,x}$ is reduced.

I.B.1 Reduced structure on a scheme

Recall that the *nilradical* of a ring A , denoted by $\text{Rad}(A)$, is the ideal of nilpotent elements of A . It is the intersection of all prime ideals of A .

Lemma I.B.2. Let X be a scheme.

- i) The scheme X is reduced if and only if $\mathcal{O}_X(U)$ is reduced for every open subset $U \subset X$.
- ii) There is a unique reduced closed subscheme $X_{\text{red}} \subset X$ sharing the same topological space as X .

Proof. We first introduce a quasi-coherent sheaf of ideals accounting for the radical of each local ring. For an open subset $U \subset X$, let

$$\mathcal{R}(U) = \{f \in \mathcal{O}_X(U) \mid f_x \in \text{Rad}(\mathcal{O}_{X,x}) \text{ for every } x \in U\}.$$

Since nilpotent elements form an ideal and nilpotence is preserved by restriction, $\mathcal{R}$ is a sheaf of ideals of $\mathcal{O}_X$. Let $U = \text{Spec}(A)$ be affine. We claim that

$$\mathcal{R}(U) = \text{Rad}(A).$$

If $f \in \text{Rad}(A)$, then $f^n = 0$ for some n , so the image of f in every localisation $A_{\mathfrak{p}}$ is nilpotent. Hence $f \in \mathcal{R}(U)$. Conversely, suppose that $f \in A$ has nilpotent image in $A_{\mathfrak{p}}$ for every $\mathfrak{p} \in \text{Spec}(A)$. For each $\mathfrak{p}$ there is an integer $n_{\mathfrak{p}} \geq 1$ such that

$$\left(\frac{f}{1}\right)^{n_{\mathfrak{p}}} = 0 \quad \text{in } A_{\mathfrak{p}}.$$

By the definition of localisation, there exists $s_{\mathfrak{p}} \in A \setminus \mathfrak{p}$ such that

$$s_{\mathfrak{p}} f^{n_{\mathfrak{p}}} = 0 \quad \text{in } A.$$

Since $s_{\mathfrak{p}} \notin \mathfrak{p}$, we have $\mathfrak{p} \in D(s_{\mathfrak{p}})$. Thus the open subsets $D(s_{\mathfrak{p}})$ cover $\text{Spec}(A)$. Since $\text{Spec}(A)$ is quasi-compact, finitely many of them suffice, so $\text{Spec}(A) = D(s_1) \cup \dots \cup D(s_r)$. Equivalently, $(s_1, \dots, s_r) = A$, so there are $a_1, \dots, a_r \in A$ such that $1 = a_1 s_1 + \dots + a_r s_r$. Let $N = \max\{n_1, \dots, n_r\}$. Since $s_i f^{n_i} = 0$, we also have $s_i f^N = 0$ for every i . Therefore

$$f^N = (a_1 s_1 + \dots + a_r s_r) f^N = 0.$$

Hence $f \in \text{Rad}(A)$, proving $\mathcal{R}(U) = \text{Rad}(A)$.

Next, we note that, for every $g \in A$, $\text{Rad}(A)_g = \text{Rad}(A_g)$. Indeed, if a/g^m belongs to $\text{Rad}(A_g)$, then

$$\left(\frac{a}{g^m}\right)^n = 0$$

for some integer n . Hence there is an integer $r \geq 0$ such that $g^r a^n = 0$ in A . Thus ga is nilpotent, since $(ga)^{n+r} = g^{n+r} a^{n+r} = 0$, and

$$\frac{a}{g^m} = \frac{ga}{g^{m+1}} \in \text{Rad}(A)_g.$$

The opposite inclusion is immediate, since localisation preserves nilpotent elements.

It follows that, on every affine open $U = \text{Spec}(A)$, the restriction of $\mathcal{R}$ is the quasi-coherent ideal sheaf associated with $\text{Rad}(A)$. Hence $\mathcal{R}$ is a quasi-coherent sheaf of ideals.

Now we prove i). Assume first that X is reduced, and let $U \subset X$ be open. Let $f \in \mathcal{O}_X(U)$ be nilpotent. For every $x \in U$, choose an affine open neighbourhood $V = \text{Spec}(A) \subset U$ of x . Since X is reduced, every local ring $\mathcal{O}_{X,y}$, $y \in V$, is reduced, hence $\mathcal{R}(V) = 0$. By the computation above, $\mathcal{R}(V) = \text{Rad}(A)$, so $A = \mathcal{O}_X(V)$ is reduced. Therefore the nilpotent section $f|_V$ is zero. Thus f vanishes on a neighbourhood of every point of U , and hence $f = 0$. Therefore $\mathcal{O}_X(U)$ is reduced.

Conversely, assume that $\mathcal{O}_X(U)$ is reduced for every open subset $U \subset X$. Given $x \in X$, choose an affine open neighbourhood $V = \text{Spec}(A)$ of x . Then $A = \mathcal{O}_X(V)$ is reduced. If x corresponds to $\mathfrak{p} \in \text{Spec}(A)$, we have $\mathcal{O}_{X,x} \simeq A_{\mathfrak{p}}$, which is reduced since localisation preserves reducedness. Hence every local ring of X is reduced, and therefore X is reduced.

Let us prove ii), the quotient sheaf $\mathcal{O}_X/\mathcal{R}$ defines a closed subscheme $X_{\text{red}} \subset X$. On an affine open subset $U = \text{Spec}(A)$ we have

$$X_{\text{red}} \cap U = \text{Spec}(A/\text{Rad}(A)).$$

Every prime ideal of A contains $\text{Rad}(A)$, so the canonical map

$$\text{Spec}(A/\text{Rad}(A)) \longrightarrow \text{Spec}(A)$$

is a homeomorphism. Thus X_{red} has the same underlying topological space as X . Moreover, $A/\text{Rad}(A)$ is reduced, so X_{red} is reduced.

Finally, let $Y \subset X$ be a reduced closed subscheme with the same underlying topological space. On an affine open subset $U = \text{Spec}(A)$, write $Y \cap U = \text{Spec}(A/\mathfrak{a})$. Since $Y \cap U$ has the same underlying closed subset as U , we have $\mathbb{W}(\mathfrak{a}) = \text{Spec}(A) = \mathbb{W}(0)$, and therefore $\sqrt{\mathfrak{a}} = \text{Rad}(A)$. On the other hand, $A/\mathfrak{a}$ is reduced, so $\mathfrak{a}$ is radical, namely $\mathfrak{a} = \sqrt{\mathfrak{a}}$. Hence $\mathfrak{a} = \text{Rad}(A)$. Thus Y and X_{red} agree on every affine open subset of X , and therefore $Y = X_{\text{red}}$. $\square$

Theorem I.B.3 (Universal property of the reduction). *Let Y be a reduced scheme and let $f : Y \rightarrow X$ be a morphism. Then there is a unique morphism $f_{\text{red}} : Y \rightarrow X_{\text{red}}$ such that*

$$f = i \circ f_{\text{red}},$$

where $i : X_{\text{red}} \hookrightarrow X$ is the canonical closed immersion.

Proof. Let $\mathcal{R} \subset \mathcal{O}_X$ be the nilradical sheaf introduced in Lemma I.B.2. Thus $\mathcal{O}_{X_{\text{red}}} = \mathcal{O}_X/\mathcal{R}$, and the canonical closed immersion $i : X_{\text{red}} \hookrightarrow X$ is the identity on the underlying topological spaces. Let

$$f : Y \longrightarrow X$$

be a morphism, where Y is reduced. We shall show that the morphism of structure sheaves

$$f^{\#} : \mathcal{O}_X \longrightarrow f_*\mathcal{O}_Y$$

vanishes on $\mathcal{R}$. Indeed, let $U \subset X$ be open and let $s \in \mathcal{R}(U)$. By definition, for every $x \in U$, the germ $s_x \in \mathcal{O}_{X,x}$ is nilpotent. Let $y \in f^{-1}(U)$, and put $x = f(y)$. The germ at y of the section $f^{\#}(s)$ is

$$(f^{\#}(s))_y = f_y^{\#}(s_x) \in \mathcal{O}_{Y,y}.$$

Since a ring homomorphism sends nilpotent elements to nilpotent elements, this germ is nilpotent in $\mathcal{O}_{Y,y}$. The scheme Y is reduced, so every local ring $\mathcal{O}_{Y,y}$ is reduced. Hence $f_y^\#(s_x) = 0$. Thus every germ of $f^\#(s)$ is zero, and therefore $f^\#(s) = 0$. We have proved that

$$\mathcal{R} \subseteq \ker(f^\#).$$

Consequently, $f^\#$ factors uniquely through the quotient sheaf:

$$\begin{array}{ccc} \mathcal{O}_X & \xrightarrow{f^\#} & f_*\mathcal{O}_Y \\ \downarrow i^\# & \nearrow f_{\text{red}}^\# & \\ \mathcal{O}_X/\mathcal{R} & & \end{array}$$

Here $f_{\text{red}}^\#$ is just defined as the map induced by the diagram. Since X_{red} and X have the same underlying topological space, we may regard the continuous map underlying f as a continuous map $f_{\text{red}} : Y \rightarrow X_{\text{red}}$. Together with the morphism of sheaves

$$f_{\text{red}}^\# : \mathcal{O}_{X_{\text{red}}} = \mathcal{O}_X/\mathcal{R} \rightarrow f_*\mathcal{O}_Y,$$

this defines a morphism of ringed spaces $f_{\text{red}} : Y \rightarrow X_{\text{red}}$.

We still have to check that this is a morphism of locally ringed spaces. Let $y \in Y$, and put $x = f(y)$. The induced homomorphism on stalks is

$$(f_{\text{red}}^\#)_y : \mathcal{O}_{X_{\text{red}},x} = \mathcal{O}_{X,x}/\mathcal{R}_x \rightarrow \mathcal{O}_{Y,y}.$$

We have $\mathcal{R}_x = \text{Rad}(\mathcal{O}_{X,x})$, so we have an inclusion-preserving bijection between prime ideals of $\mathcal{O}_{X,x}$ and prime ideals of $\mathcal{O}_{X,x}/\mathcal{R}_x$ by sending a prime ideal of $\mathcal{O}_{X,x}/\mathcal{R}_x$ to its pull-back in $\mathcal{O}_{X,x}$, and a prime ideal of $\mathcal{O}_{X,x}$ to its image in $\mathcal{O}_{X,x}/\mathcal{R}_x$. Under this correspondence, the maximal ideal $\mathfrak{m}_{X,x}$ is sent to $\mathfrak{m}_{X,x}/\mathcal{R}_x$ which is thus the maximal ideal of $\mathcal{O}_{X_{\text{red}},x}$. Since $f_y^\# : \mathcal{O}_{X,x} \rightarrow \mathcal{O}_{Y,y}$ is local, we have

$$(f_y^\#)^{-1}(\mathfrak{m}_{Y,y}) = \mathfrak{m}_{X,x}.$$

Passing to the quotient gives

$$(f_{\text{red}}^\#)_y^{-1}(\mathfrak{m}_{Y,y}) = \mathfrak{m}_{X,x}/\mathcal{R}_x.$$

Hence $(f_{\text{red}}^\#)_y$ is local. Thus f_{red} is a morphism of locally ringed spaces, i.e. a morphism of schemes. By construction, the composition $i \circ f_{\text{red}}$ has the same underlying continuous map as f and $f^\# = f_{\text{red}}^\# \circ i^\#$. In conclusion we have the factorisation $f = i \circ f_{\text{red}}$ as morphisms of schemes.

Finally, suppose that $g : Y \rightarrow X_{\text{red}}$ is another morphism such that $f = i \circ g$. Since i is the identity on the underlying topological spaces, the continuous map underlying g must coincide with that underlying f_{red} . Moreover, the morphism $g^\#$ must satisfy $f^\# = g^\# \circ i^\#$. The quotient morphism $i^\# : \mathcal{O}_X \rightarrow \mathcal{O}_X/\mathcal{R}$ is surjective, so such a factorisation is unique. Indeed, for all $y \in Y$, let $x = f(y)$ and consider an element t_x of $\mathcal{O}_{X,x}/\mathcal{R}_x$. Then t_x is represented by $s_x \in \mathcal{O}_{X,x}$, i.e. $i_x^\#(s_x) = t_x$ and $g_y^\#(t_x) = f_y^\#(s_x) = (f_{\text{red}}^\#)_y(t_x)$. Therefore $g^\# = f_{\text{red}}^\#$. In conclusion, we have $g = f_{\text{red}}$ as morphisms of schemes. This proves uniqueness. $\square$

I.B.2 The Zariski tangent space

The maps from the space of dual numbers to a given $\mathbb{k}$ -scheme are thought of as tangent directions at a given point of X . More precisely, consider the scheme of dual numbers

$$D_{\mathbb{k}} = \operatorname{Spec}(\mathbb{k}[\epsilon]/(\epsilon^2)).$$

Then (ϵ) is the unique prime ideal of $D_{\mathbb{k}}$. Note that $D_{\mathbb{k}}$ is not reduced. Its reduction is $(D_{\mathbb{k}})_{\text{red}} = \operatorname{Spec}(\mathbb{k})$.

Definition I.B.4 (Zariski tangent space). Let X be a $\mathbb{k}$ -scheme and let $x \in X(\mathbb{k})$. Write $i : \operatorname{Spec}(\mathbb{k}) \hookrightarrow D_{\mathbb{k}}$ for the inclusion given by reduction. The *Zariski tangent space* of X at x is the set

$$T_x X = \{v : D_{\mathbb{k}} \longrightarrow X \mid v \text{ is a } \mathbb{k}\text{-morphism and } v \circ i = x\}.$$

Thus a tangent vector is an infinitesimal $\mathbb{k}$ -valued path through x .

Let us introduce the space of derivations and identify it with the Zariski tangent space.

Definition I.B.5. Let $X = \operatorname{Spec}(A)$ be affine and let $x^\# : A \longrightarrow \mathbb{k}$ be the homomorphism corresponding to the $\mathbb{k}$ -rational point x . Put $\mathfrak{m} = \ker(x^\#)$. Via $x^\#$, the field $\mathbb{k}$ is an A -module. A $\mathbb{k}$ -derivation of A with values in $\mathbb{k}$ is a $\mathbb{k}$ -linear map $\theta : A \rightarrow \mathbb{k}$ such that

$$\theta(ab) = x^\#(a)\theta(b) + x^\#(b)\theta(a) \quad \text{for all } a, b \in A.$$

We denote the vector space of such derivations by

$$\operatorname{Der}_{\mathbb{k}}(A, \mathbb{k}).$$

Proposition I.B.6. Let $X = \operatorname{Spec}(A)$ be an affine $\mathbb{k}$ -scheme and let $x \in X(\mathbb{k})$, with corresponding homomorphism $x^\# : A \rightarrow \mathbb{k}$ and maximal ideal $\mathfrak{m} = \ker(x^\#)$. There are natural bijections

$$T_x X \simeq \operatorname{Der}_{\mathbb{k}}(A, \mathbb{k}) \simeq \operatorname{Hom}_{\mathbb{k}}(\mathfrak{m}/\mathfrak{m}^2, \mathbb{k}).$$

In particular, these identifications give $T_x X$ a natural $\mathbb{k}$ -vector-space structure.

Proof. A tangent vector is a $\mathbb{k}$ -algebra homomorphism $\varphi : A \longrightarrow \mathbb{k}[\epsilon]/(\epsilon^2)$ whose composite with the quotient map $\mathbb{k}[\epsilon]/(\epsilon^2) \longrightarrow \mathbb{k}$, sending ϵ to 0 is $x^\#$. It is therefore uniquely of the form

$$\varphi(a) = x^\#(a) + \theta(a)\epsilon$$

for a $\mathbb{k}$ -linear map $\theta : A \rightarrow \mathbb{k}$. The condition that φ preserve products amounts to

$$\begin{aligned} x^\#(ab) + \theta(ab)\epsilon &= (x^\#(a) + \theta(a)\epsilon)(x^\#(b) + \theta(b)\epsilon) \\ &= x^\#(a)x^\#(b) + (x^\#(a)\theta(b) + x^\#(b)\theta(a))\epsilon, \end{aligned}$$

for all $a, b \in A$, because $\epsilon^2 = 0$. Since $x^\#$ is a ring homomorphism, this is equivalent to

$$\theta(ab) = x^\#(a)\theta(b) + x^\#(b)\theta(a).$$

Thus tangent vectors are naturally identified with $\operatorname{Der}_{\mathbb{k}}(A, \mathbb{k})$.

Let now $\theta \in \operatorname{Der}_{\mathbb{k}}(A, \mathbb{k})$. If $a, b \in \mathfrak{m}$, then $x^\#(a) = x^\#(b) = 0$, and hence

$$\theta(ab) = 0.$$

Thus θ vanishes on $\mathfrak{m}^2$ and induces a $\mathbb{k}$ -linear map $\mathfrak{m}/\mathfrak{m}^2 \rightarrow \mathbb{k}$. Conversely, let

$$\ell : \mathfrak{m}/\mathfrak{m}^2 \longrightarrow \mathbb{k}$$

be $\mathbb{k}$ -linear. Every $a \in A$ has a unique decomposition

$$a = x^\#(a) + (a - x^\#(a)),$$

with $a - x^\#(a) \in \mathfrak{m}$. Define

$$\theta(a) = \ell(a - x^\#(a) \bmod \mathfrak{m}^2).$$

This is a $\mathbb{k}$ -derivation. Indeed,

$$\begin{aligned} ab - x^\#(ab) &= x^\#(a)(b - x^\#(b)) + x^\#(b)(a - x^\#(a)) \\ &\quad + (a - x^\#(a))(b - x^\#(b)), \end{aligned}$$

and the last term belongs to $\mathfrak{m}^2$. Passing to $\mathfrak{m}/\mathfrak{m}^2$ gives the Leibniz rule. The two constructions are inverse to one another. $\square$

The preceding proposition suggests the intrinsic definition at an arbitrary point.

Definition I.B.7 (Zariski tangent space at an arbitrary point). Let X be a scheme and let $x \in X$. We have $\mathfrak{m}_{X,x} = \mathfrak{m}_{X,x}$, the maximal ideal at x , and $\kappa(x) = \mathcal{O}_{X,x}/\mathfrak{m}_x$, the residue field at x . The *Zariski tangent space* of X at x is the $\kappa(x)$ -vector space

$$T_x X = \text{Hom}_{\kappa(x)}(\mathfrak{m}_x/\mathfrak{m}_x^2, \kappa(x)).$$

The vector space

$$\mathfrak{m}_x/\mathfrak{m}_x^2$$

is called the *cotangent space* of X at x .

If X is a $\mathbb{k}$ -scheme and $x \in X(\mathbb{k})$, then $\kappa(x) = \mathbb{k}$. Choosing an affine neighbourhood of x and applying Proposition I.B.6 shows that this intrinsic definition agrees with the definition by dual numbers.

Example I.B.8 (Equations and tangent vectors). Let

$$X = \mathbb{V}(f_1, \dots, f_r) \subset \mathbb{A}_{\mathbb{k}}^n$$

and let $a = (a_1, \dots, a_n) \in X(\mathbb{k})$. A tangent vector is represented by

$$x_i \longmapsto a_i + v_i \epsilon,$$

with $v = (v_1, \dots, v_n) \in \mathbb{k}^n$. For a monomial $x^\alpha = x_1^{\alpha_1} \cdots x_n^{\alpha_n}$, the relation $\epsilon^2 = 0$ gives

$$(a + v\epsilon)^\alpha = a^\alpha + \sum_{\alpha_i > 0} \alpha_i a_1^{\alpha_1} \cdots a_i^{\alpha_i - 1} \cdots a_n^{\alpha_n} v_i \epsilon.$$

By linearity, for every polynomial f one obtains the first-order Taylor formula

$$f(a + v\epsilon) = f(a) + \left(\sum_i \frac{\partial f}{\partial x_i}(a) v_i \right) \epsilon.$$

The tangent vector factors through X precisely when every f_j maps to zero. Since $a \in X$, its constant term already vanishes, and the remaining conditions are

$$\sum_i \frac{\partial f_j}{\partial x_i}(a) v_i = 0, \quad 1 \leq j \leq r.$$

Thus $T_a X$ is the kernel of the Jacobian matrix at a . For the parabola $y = x^2$ it is the line $v_y = 2av_x$ at (a, a^2) . For the node $xy = 0$, the tangent space at the origin is all of $\mathbb{k}^2$, reflecting the two crossing branches.

Exercise I.B.9 (Tangent spaces at a nonsingular and a singular point). For $\text{char}(\mathbb{k}) \neq 2, 3$ let

$$C = \mathbb{W}(y^2 - x^3) \subset \mathbb{A}_{\mathbb{k}}^2.$$

- (i) Compute $T_{(0,0)}C$.
- (ii) Compute $T_{(1,1)}C$ and give an equation for the resulting line.
- (iii) For a plane curve $\mathbb{W}(f)$, call a point *nonsingular* when (f_x, f_y) does not vanish there and *singular* otherwise. Explain the difference between the dimensions of the two tangent spaces above.

Solution. For $f = y^2 - x^3$, the Jacobian row is

$$(-3x^2, 2y).$$

At the origin it vanishes, so $T_{(0,0)}C = \mathbb{k}^2$ and the origin is singular. At $(1, 1)$ the tangent space is

$$T_{(1,1)}C = \{(u, v) \in \mathbb{k}^2 \mid -3u + 2v = 0\},$$

a one-dimensional line; the point is nonsingular. Thus at a nonsingular point the Zariski tangent space has the expected dimension of the curve, whereas at the cusp it jumps to dimension two.

Exercise I.B.10. Let $A = \mathbb{k}[x_1, x_2]$ and consider the ideals

$$\mathfrak{a} = (x_1, x_2), \quad \mathfrak{b} = (x_1, x_2)^2, \quad \mathfrak{c} = (x_1^2, x_1x_2).$$

- i) Compute $\sqrt{\mathfrak{a}}$, $\sqrt{\mathfrak{b}}$ and $\sqrt{\mathfrak{c}}$.
- ii) Determine the underlying closed subsets of $\text{Spec}(A/\mathfrak{a})$, $\text{Spec}(A/\mathfrak{b})$ and $\text{Spec}(A/\mathfrak{c})$.
- iii) Compute the reductions of these three affine schemes.
- iv) Exhibit a non-zero nilpotent element in $A/\mathfrak{b}$ and in $A/\mathfrak{c}$.

Solution. We have

$$\sqrt{\mathfrak{a}} = \mathfrak{a}, \quad \sqrt{\mathfrak{b}} = \mathfrak{a}, \quad \sqrt{\mathfrak{c}} = (x_1).$$

Indeed, $\mathfrak{b}$ contains x_1^2 and x_2^2 , while every element of $\mathfrak{c}$ is divisible by x_1 and $x_1^2 \in \mathfrak{c}$.

Consequently, the first two schemes are supported at the origin, whereas the third is supported on the line $\mathbb{W}(x_1)$. Their reductions are respectively

$$\text{Spec}(A/\mathfrak{a}), \quad \text{Spec}(A/\mathfrak{a}), \quad \text{Spec}(A/(x_1)).$$

The classes of x_1 and x_2 are non-zero nilpotents in $A/\mathfrak{b}$. The class of x_1 is a non-zero nilpotent in $A/\mathfrak{c}$.

II Irreducible schemes

Irreducibility is a property of the underlying topological space. It is stronger than connectedness.

II.A Irreducible components

Definition II.A.1. A non-empty topological space X is *irreducible* if, whenever $X = Z_1 \cup Z_2$ with Z_1, Z_2 closed subsets of X , then $X = Z_1$ or $X = Z_2$.

Remark II.A.2. For a non-empty topological space X , the following conditions are equivalent.

- i) The space X is irreducible.
- ii) Any two non-empty open subsets of X meet.
- iii) Every non-empty open subset of X is dense in X .

Proof. Taking complements transforms a decomposition of X into two proper closed subsets into a pair of disjoint non-empty open subsets. This proves the equivalence of i) and ii). A subset is dense if and only if it meets every non-empty open subset, which gives the equivalence of ii) and iii). $\square$

Definition II.A.3. An *irreducible component* of a topological space X is a maximal irreducible closed subset of X .

Every point of X is contained in an irreducible component. Indeed, the closure of a point is irreducible. More generally, the union of a chain of irreducible subsets is irreducible, and so is its closure. Zorn's lemma therefore gives a maximal irreducible closed subset containing any prescribed irreducible subset.

Lemma II.A.4. Let U be a non-empty open subset of a topological space X .

- i) If X is irreducible, then U is irreducible and dense in X .
- ii) If $(X_i)_{i \in I}$ are the irreducible components of X , then the non-empty sets $U \cap X_i$ are precisely the irreducible components of U .
- iii) If $X = X_1 \cup \dots \cup X_r$, where the X_i are irreducible closed subsets and no X_i is contained in another, then $X_1, \dots, X_r$ are the irreducible components of X .

Proof. The first assertion follows from Remark II.A.2.

If $U \cap X_i \neq \emptyset$, then it is irreducible and dense in X_i . If it were properly contained in an irreducible closed subset F of U , the closure of F in X would be an irreducible closed subset properly containing X_i , a contradiction. Conversely, if F is an irreducible component of U , then its closure in X is contained in some irreducible component X_i , and maximality gives $F = U \cap X_i$.

Finally, an irreducible subset contained in a finite union of closed subsets is contained in one of them. Hence every irreducible component of X is one of the X_i . The assumption that no X_i is contained in another shows that every X_i is maximal. $\square$

Example II.A.5. Let

$$X = \operatorname{Spec}(\mathbb{k}[x_1, x_2]/(x_1x_2)).$$

Then

$$X = \mathbb{V}(\bar{x}_1) \cup \mathbb{V}(\bar{x}_2).$$

Both closed subsets are isomorphic to $\mathbb{A}_{\mathbb{k}}^1$, hence irreducible, and neither is contained in the other. They are therefore the two irreducible components of X . Their intersection is the closed point defined by $(\bar{x}_1, \bar{x}_2)$.

Exercise II.A.6. Let

$$X = \mathbb{W}(x_1x_2(x_1 - x_2)) \subset \mathbb{A}_{\mathbb{k}}^2.$$

- i) Show that X is the union of three affine lines.
- ii) Prove that these lines are the irreducible components of X .
- iii) Determine the irreducible components of the open subset $U = X \setminus \{(0, 0)\}$.
- iv) Is U connected?

Solution. We have

$$X = \mathbb{W}(x_1) \cup \mathbb{W}(x_2) \cup \mathbb{W}(x_1 - x_2).$$

Each subset is an affine line and is therefore irreducible. No one of them is contained in another, so they are the irreducible components.

Their pairwise intersections are all equal to the origin. Hence removing the origin produces the disjoint union

$$U = (\mathbb{W}(x_1) \setminus \{0\}) \amalg (\mathbb{W}(x_2) \setminus \{0\}) \amalg (\mathbb{W}(x_1 - x_2) \setminus \{0\}).$$

These three subsets are the irreducible components of U , and U is disconnected.

II.B Irreducible affine schemes and minimal primes

Let A be a ring. A prime ideal $\mathfrak{p}$ is *minimal* if it contains no strictly smaller prime ideal. More generally, a prime ideal minimal among those containing an ideal $\mathfrak{a}$ is called a minimal prime over $\mathfrak{a}$.

Lemma II.B.1. *Let A be a ring and let $\mathfrak{a} \subset A$ be an ideal.*

- i) *The closed subset $\mathbb{W}(\mathfrak{a})$ is irreducible if and only if $\sqrt{\mathfrak{a}}$ is prime.*
- ii) *If $(\mathfrak{p}_i)_{i \in I}$ are the minimal primes over $\mathfrak{a}$, then the subsets $\mathbb{W}(\mathfrak{p}_i)$ are the irreducible components of $\mathbb{W}(\mathfrak{a})$.*
- iii) *The space $\text{Spec}(A)$ is irreducible if and only if A has a unique minimal prime, equivalently if and only if $\text{Rad}(A)$ is prime.*

Proof. Suppose first that $\sqrt{\mathfrak{a}}$ is prime and

$$\mathbb{W}(\mathfrak{a}) = \mathbb{W}(\mathfrak{b}_1) \cup \mathbb{W}(\mathfrak{b}_2) = \mathbb{W}(\mathfrak{b}_1\mathfrak{b}_2).$$

Then $\mathfrak{b}_1\mathfrak{b}_2 \subset \sqrt{\mathfrak{a}}$, so primality gives $\mathfrak{b}_1 \subset \sqrt{\mathfrak{a}}$ or $\mathfrak{b}_2 \subset \sqrt{\mathfrak{a}}$. In the first case,

$$\mathbb{W}(\mathfrak{a}) \subset \mathbb{W}(\mathfrak{b}_1),$$

and hence $\mathbb{W}(\mathfrak{a}) = \mathbb{W}(\mathfrak{b}_1)$; similarly in the second case.

Conversely, if $\sqrt{\mathfrak{a}}$ is not prime, choose $f_1, f_2 \notin \sqrt{\mathfrak{a}}$ with $f_1f_2 \in \sqrt{\mathfrak{a}}$. Then

$$\mathbb{W}(\mathfrak{a}) = \mathbb{W}(\mathfrak{a} + (f_1)) \cup \mathbb{W}(\mathfrak{a} + (f_2)),$$

and both closed subsets are proper. Thus $\mathbb{W}(\mathfrak{a})$ is reducible.

The second assertion follows because inclusion of prime ideals is reversed by $\mathbb{W}$: minimal primes over $\mathfrak{a}$ correspond exactly to maximal irreducible closed subsets of $\mathbb{W}(\mathfrak{a})$. The last assertion is the special case $\mathfrak{a} = (0)$. $\square$

Example II.B.2. The ring

$$A = \mathbb{k}[x_1, x_2]/(x_1^2)$$

is not reduced, but $\text{Spec}(A)$ is irreducible. Indeed,

$$\text{Rad}(A) = (\bar{x}_1),$$

and this ideal is prime because $A/(\bar{x}_1) \simeq \mathbb{k}[x_2]$ is an integral domain. Thus irreducibility does not imply reducedness.

Exercise II.B.3. For each of the following ideals of $\mathbb{k}[x_1, x_2]$, compute its radical, its minimal primes, and the irreducible components of its zero set:

$$\mathfrak{a}_1 = (x_1x_2), \quad \mathfrak{a}_2 = (x_1^2x_2), \quad \mathfrak{a}_3 = (x_1x_2(x_1 - x_2)).$$

Which of the quotient rings $\mathbb{k}[x_1, x_2]/\mathfrak{a}_i$ are reduced? Which spectra are irreducible?

Solution. We have

$$\sqrt{\mathfrak{a}_1} = \sqrt{\mathfrak{a}_2} = (x_1x_2) = (x_1) \cap (x_2),$$

so the minimal primes are (x_1) and (x_2) . Both zero sets have the two coordinate axes as irreducible components. The first quotient is reduced, while the second is not: the class of x_1x_2 is non-zero and has square zero modulo $(x_1^2x_2)$.

Moreover,

$$\sqrt{\mathfrak{a}_3} = \mathfrak{a}_3 = (x_1) \cap (x_2) \cap (x_1 - x_2),$$

so its zero set has three irreducible components. This quotient is reduced. None of the three spectra is irreducible.

II.C Generic points and specialisations

Definition II.C.1. Let X be a topological space and let $x, y \in X$. We say that y is a *specialisation* of x , or that x is a *generalisation* of y , if

$$y \in \overline{\{x\}}.$$

If $Z \subset X$ is an irreducible closed subset, a point $\eta \in Z$ is a *generic point* of Z if

$$\overline{\{\eta\}} = Z.$$

The generic points of the irreducible components of X are also called the generic points of X .

Lemma II.C.2. Let $X = \text{Spec}(A)$ and let $\mathfrak{p}, \mathfrak{q} \in X$. Then

$$\overline{\{\mathfrak{p}\}} = \mathbb{V}(\mathfrak{p}).$$

Consequently, $\mathfrak{q}$ is a specialisation of $\mathfrak{p}$ if and only if $\mathfrak{p} \subset \mathfrak{q}$. The generic points of X correspond to the minimal prime ideals of A .

Proof. The equality

$$\overline{\{\mathfrak{p}\}} = \mathbb{V}(\mathfrak{p})$$

was proved in Proposition II.A.6. Hence, by definition, $\mathfrak{q}$ is a specialisation of $\mathfrak{p}$ if and only if

$$\mathfrak{q} \in \mathbb{V}(\mathfrak{p}),$$

that is, if and only if $\mathfrak{p} \subset \mathfrak{q}$.

Finally, by Lemma II.B.1, ii), the irreducible components of $\text{Spec}(A)$ are precisely the subsets $\mathbb{V}(\mathfrak{p})$, where $\mathfrak{p}$ runs through the minimal prime ideals of A . Since $\overline{\{\mathfrak{p}\}} = \mathbb{V}(\mathfrak{p})$, their generic points are precisely the minimal prime ideals. $\square$

Proposition II.C.3. *Let X be a scheme.*

- i) *Every irreducible closed subset of X has a unique generic point. In particular, generic points are in bijection with irreducible components.*
- ii) *Let $x \in X$. The irreducible components of X containing x are in bijection with the irreducible components of $\text{Spec}(\mathcal{O}_{X,x})$.*

Proof. Let $Z \subset X$ be irreducible, equip it with its reduced induced scheme structure, and choose a non-empty affine open subset $U = \text{Spec}(A) \subset Z$. Since U is irreducible, A has a unique minimal prime $\mathfrak{p}$. The corresponding point η is dense in U , hence dense in Z . Uniqueness follows because two generic points of Z lie in every non-empty open subset of Z ; placing both in an affine open subset and using Lemma II.C.2 shows that they coincide.

For ii), choose an affine neighbourhood $U = \text{Spec}(A)$ of x , and let $\mathfrak{p}$ correspond to x . Prime ideals of $A_{\mathfrak{p}}$ correspond to prime ideals of A contained in $\mathfrak{p}$. Under this correspondence, minimal primes of $A_{\mathfrak{p}}$ correspond to minimal primes of A contained in $\mathfrak{p}$, namely to the generic points of the irreducible components of X passing through x . $\square$

Let X be a scheme and let $x \in X$. Consider $Y = X \setminus \{x\}$. We equip Y with the induced topology and the restricted sheaf $\mathcal{O}_Y = \mathcal{O}_X|_Y$.

Proposition II.C.4 (Generic points cannot be removed). *The point x is closed if and only if Y is a scheme.*

Proof. If x is closed, then Y is an open subscheme of X , hence again a scheme. Conversely, suppose that Y is a scheme. We claim that x must be closed. Indeed, the question is local on X , so choose an affine neighbourhood $V = \text{Spec}(A)$ of x . Then $Y \cap V = \text{Spec}(A) \setminus \{x\}$ is again a scheme.

Suppose that x is not closed. Let $\mathfrak{p} \subset A$ be the prime ideal corresponding to x . Since $\mathfrak{p}$ is not maximal, there exists a prime ideal $\mathfrak{q}$ of A with $\mathfrak{q} \supsetneq \mathfrak{p}$. The corresponding point belongs to Y , so it admits an affine neighbourhood

$$U = \text{Spec}(B) \subseteq Y \cap V.$$

Since U is an open neighbourhood of $\mathfrak{q}$ inside $\text{Spec}(A)$, the point $\mathfrak{q}$ corresponds to a minimal prime of B . Therefore the local ring $\mathcal{O}_{U,\mathfrak{q}}$ has a unique prime ideal.

On the other hand all the following rings are identified $\mathcal{O}_{U,\mathfrak{q}} = \mathcal{O}_{Y,\mathfrak{q}} = \mathcal{O}_{X,\mathfrak{q}} = A_{\mathfrak{q}}$. Moreover, $A_{\mathfrak{q}}$ has at least the two distinct prime ideals, namely $\mathfrak{p}A_{\mathfrak{q}}$ and $\mathfrak{q}A_{\mathfrak{q}}$, a contradiction. Hence Y is a scheme if and only if x is closed. $\square$

Example II.C.5. In $\mathbb{A}_{\mathbb{k}}^1 = \text{Spec}(\mathbb{k}[x_1])$, the point (0) is generic: its closure is all of $\mathbb{A}_{\mathbb{k}}^1$. Every closed point $(x_1 - a)$ is a specialisation of (0) because $(0) \subset (x_1 - a)$. A closed point has no proper specialisations.

Exercise II.C.6. Let

$$X = \text{Spec}(\mathbb{k}[x_1, x_2]/(x_1x_2))$$

and let o be the point defined by $(\bar{x}_1, \bar{x}_2)$.

- i) Determine the generic points of X and their closures.
- ii) Which points specialise to o ?
- iii) Compute the minimal primes of the local ring $\mathcal{O}_{X,o}$.
- iv) Verify directly the bijection in Proposition II.C.3, ii).

Solution. The minimal primes of the coordinate ring are $(\bar{x}_1)$ and $(\bar{x}_2)$. They are the generic points η_1 and η_2 of the two coordinate axes, with

$$\overline{\{\eta_1\}} = \mathbb{W}(\bar{x}_1), \quad \overline{\{\eta_2\}} = \mathbb{W}(\bar{x}_2).$$

Both ideals are contained in $(\bar{x}_1, \bar{x}_2)$, so both generic points specialise to o . Of course o also specialises to itself.

After localising at $(\bar{x}_1, \bar{x}_2)$, the minimal primes are $(\bar{x}_1)\mathcal{O}_{X,o}$ and $(\bar{x}_2)\mathcal{O}_{X,o}$. They correspond to the two irreducible components of X passing through o .

III Integral schemes

Integral schemes are the scheme-theoretic counterparts of integral domains.

Definition III..1. A scheme X is *integral* if it is reduced and irreducible. It is *integral at a point* $x \in X$ if the local ring $\mathcal{O}_{X,x}$ is an integral domain.

III.A Integral schemes and integral domains

Proposition III.A.1. For a scheme X , the following conditions are equivalent.

- i) The scheme X is integral.
- ii) Every non-empty affine open subset $U = \text{Spec}(A)$ of X has A an integral domain.
- iii) For every non-empty open subset $U \subset X$, the ring $\mathcal{O}_X(U)$ is an integral domain.

Proof. Assume first that X is integral, and let η be its generic point. We claim that, for every non-empty open subset $U \subset X$, the natural map

$$\mathcal{O}_X(U) \longrightarrow \mathcal{O}_{X,\eta}$$

is injective. Indeed, let $s \in \mathcal{O}_X(U)$ have zero germ at η . Then s vanishes on some non-empty open subset $V \subset U$. Let $W = \text{Spec}(A) \subset U$ be a non-empty affine open subset. Since U is irreducible, $V \cap W$ is non-empty; it contains a non-empty principal open subset $D(f) \subset W$. The ring A is a domain, and the restriction of $s|_W$ to $D(f)$ is zero. Hence some power of f annihilates $s|_W$, and therefore $s|_W = 0$. Since the non-empty affine open subsets cover U , we obtain $s = 0$.

On a non-empty affine open subset $\text{Spec}(A)$, the generic point corresponds to (0) , and hence

$$\mathcal{O}_{X,\eta} \simeq A_{(0)} = \text{Frac}(A)$$

is a field. Thus $\mathcal{O}_X(U)$ is a subring of a field and is an integral domain. This proves i) $\Rightarrow$ iii), and iii) $\Rightarrow$ ii) is immediate.

Finally, assume ii). The coordinate rings of an affine open covering are domains, so X is reduced. If X were reducible, it would contain two disjoint non-empty open subsets U and V . Choose non-empty affine open subsets $U_0 \subset U$ and $V_0 \subset V$. Their disjoint union is an affine open subset of X , with ring of sections

$$\Gamma(\mathcal{O}_{U_0 \amalg V_0}) \simeq \Gamma(\mathcal{O}_{U_0}) \times \Gamma(\mathcal{O}_{V_0}),$$

which is not a domain. This contradicts ii). Hence X is irreducible and therefore integral. $\square$

Example III.A.2. The cusp

$$X = \operatorname{Spec}(\mathbb{k}[x_1, x_2]/(x_2^2 - x_1^3))$$

is integral provided $x_2^2 - x_1^3$ is irreducible in $\mathbb{k}[x_1, x_2]$; for instance, this holds over every field. Indeed, regarding the polynomial as a polynomial in x_2 , a factorisation would force x_1^3 to be a square in $\mathbb{k}[x_1]$.

By contrast,

$$\operatorname{Spec}(\mathbb{k}[x_1, x_2]/(x_1x_2))$$

is reduced but reducible, while $\operatorname{Spec}(\mathbb{k}[x_1]/(x_1^2))$ is irreducible but non-reduced. Neither is integral.

Exercise III.A.3. Determine which of the following schemes are integral, reduced, or irreducible:

$$\begin{aligned} X_1 &= \operatorname{Spec}(\mathbb{k}[x_1, x_2]/(x_1x_2)), \\ X_2 &= \operatorname{Spec}(\mathbb{k}[x_1, x_2]/(x_1^2)), \\ X_3 &= \operatorname{Spec}(\mathbb{k}[x_1, x_2]/(x_1x_2 - 1)), \\ X_4 &= \operatorname{Spec}(\mathbb{k}[x_1, x_2]/((x_2 - x_1^2)^2)). \end{aligned}$$

Compute the reduction of each non-reduced scheme.

Solution. The scheme X_1 is reduced because (x_1x_2) is radical, but it is reducible, with components $\mathbb{V}(\bar{x}_1)$ and $\mathbb{V}(\bar{x}_2)$. Thus it is not integral.

The scheme X_2 is irreducible because its nilradical $(\bar{x}_1)$ is prime, but it is not reduced. Its reduction is $\operatorname{Spec}(\mathbb{k}[x_2])$.

For X_3 , eliminating x_2 after inverting x_1 gives

$$\mathbb{k}[x_1, x_2]/(x_1x_2 - 1) \simeq \mathbb{k}[x_1, x_1^{-1}],$$

which is an integral domain. Hence X_3 is integral.

Finally, X_4 is irreducible but non-reduced. Its reduction is the parabola

$$\operatorname{Spec}(\mathbb{k}[x_1, x_2]/(x_2 - x_1^2)) \simeq \mathbb{A}_{\mathbb{k}}^1.$$

III.B The function field

Lemma III.B.1. Let X be an integral scheme and let η be its generic point. Then $\mathcal{O}_{X,\eta}$ is a field. If $U = \operatorname{Spec}(A)$ is any non-empty affine open subset of X , then

$$\mathcal{O}_{X,\eta} = \operatorname{Frac}(A).$$

Proof. The generic point belongs to every non-empty open subset of X . On $U = \operatorname{Spec}(A)$, it corresponds to the prime ideal (0) , because A is an integral domain. Hence

$$\mathcal{O}_{X,\eta} = A_{(0)} = \operatorname{Frac}(A).$$

□

Definition III.B.2. Let X be an integral scheme and let η be the generic point of X . The field $K(X) = \mathcal{O}_{X,\eta} = \kappa(\eta)$ is called the *function field* of X .

Example III.B.3. Let A be an integral domain and put $K = \operatorname{Frac}(A)$. Then

$$K(\mathbb{A}_A^n) = K(x_1, \dots, x_n).$$

Likewise, using the affine chart $D_+(x_0)$, we get

$$K(\mathbb{P}_A^n) = K(y_1, \dots, y_n), \quad y_i = x_i/x_0,$$

Recall Definition III.D.2: a rational map between integral schemes is *birational* if it admits a rational inverse, equivalently if it restricts to an isomorphism between non-empty open subschemes. Such an isomorphism induces an isomorphism

$$K(Y) \xrightarrow{\sim} K(X).$$

More generally, a dominant rational map $X \dashrightarrow Y$ induces an inclusion $K(Y) \hookrightarrow K(X)$. Thus the map $t \mapsto t^2$ on the affine line is not birational: on function fields it gives the proper inclusion $\mathbb{k}(t^2) \subsetneq \mathbb{k}(t)$.

Example III.B.4. The map $\mathbb{A}_{\mathbb{k}}^1 \rightarrow \mathbb{W}(x_2 - x_1^2) \subset \mathbb{A}_{\mathbb{k}}^2$, defined by $t \mapsto (t, t^2)$, is an isomorphism. Hence the function field of the parabola is $\mathbb{k}(t)$.

The standard Cremona transformation

$$\mathbb{P}_{\mathbb{k}}^2 \dashrightarrow \mathbb{P}_{\mathbb{k}}^2, \quad [x_0 : x_1 : x_2] \mapsto [x_1x_2 : x_0x_2 : x_0x_1],$$

restricts to an automorphism of the open torus $x_0x_1x_2 \neq 0$. In the affine coordinates $y_i = x_i/x_0$, it induces the automorphism of $\mathbb{k}(y_1, y_2)$ defined by:

$$y_1 \mapsto y_1^{-1}, \quad y_2 \mapsto y_2^{-1}$$

Lemma III.B.5. Let X be an integral scheme and let $U \subset X$ be a non-empty open subset. Then

$$\mathcal{O}_X(U) = \bigcap_{x \in U} \mathcal{O}_{X,x} \subset K(X).$$

Proof. The injectivity of $\mathcal{O}_X(U) \rightarrow \mathcal{O}_{X,x}$ follows from integrality: a section whose germ vanishes at one point vanishes on a non-empty neighbourhood, hence on all of U . The second injection is checked on an affine neighbourhood.

It remains to prove the equality. The inclusion from left to right is clear. It is enough to treat $U = \text{Spec}(A)$. Let $f \in \text{Frac}(A)$ lie in $A_{\mathfrak{p}}$ for every prime $\mathfrak{p}$. Put

$$\mathfrak{b} = \{s \in A \mid sf \in A\}.$$

For every prime $\mathfrak{p}$, the condition $f \in A_{\mathfrak{p}}$ gives an element $s_{\mathfrak{p}} \in \mathfrak{b} \setminus \mathfrak{p}$. Thus $\mathfrak{b}$ is contained in no prime ideal, hence $\mathfrak{b} = A$. Therefore $1 \in \mathfrak{b}$ and $f \in A$. $\square$

Exercise III.B.6. Let $X = \mathbb{W}(x_1x_2 - 1) \subset \mathbb{A}_{\mathbb{k}}^2$ and $Y = \mathbb{W}(x_2 - x_1^2) \subset \mathbb{A}_{\mathbb{k}}^2$.

- i) Show that X and Y are integral.
- ii) Compute $K(X)$ and $K(Y)$.
- iii) Show that X is isomorphic to the principal open subset $D(x_1) \subset \mathbb{A}_{\mathbb{k}}^1$.
- iv) Are X and Y birational?

Solution. We have

$$\mathbb{k}[x_1, x_2]/(x_1x_2 - 1) \simeq \mathbb{k}[x_1, x_1^{-1}], \quad \mathbb{k}[x_1, x_2]/(x_2 - x_1^2) \simeq \mathbb{k}[x_1].$$

Both rings are integral domains. Their fraction fields are both isomorphic to $\mathbb{k}(x_1)$. The first isomorphism identifies X with $\text{Spec}(\mathbb{k}[x_1, x_1^{-1}]) = D(x_1) \subset \mathbb{A}_{\mathbb{k}}^1$. Since $D(x_1)$ is a non-empty open subset of $\mathbb{A}_{\mathbb{k}}^1 \simeq Y$, the two curves are birational.

IV Noetherianity and dimension

Most schemes studied in algebraic geometry satisfy finiteness conditions. We mentioned the condition of being of *finite type* over a base ring, notably of a base field $\mathbb{k}$, and the terminology *algebraic $\mathbb{k}$ -scheme*. Another basic condition is Noetherianity. Having these conditions in mind is important to develop a theory of dimension.

IV.A Noetherianity

Recall that a ring is Noetherian if every ideal is finitely generated, equivalently if every ascending chain of ideals is stationary.

Definition IV.A.1. A scheme X is *locally Noetherian* if it has an affine open covering $U_i = \text{Spec}(A_i)$ with all A_i Noetherian. It is *Noetherian* if it admits such a covering with finitely many members.

Example IV.A.2. Every finitely generated $\mathbb{k}$ -algebra is Noetherian by Hilbert's basis theorem. Hence every affine algebraic $\mathbb{k}$ -scheme is Noetherian, and every algebraic $\mathbb{k}$ -scheme is locally Noetherian. By contrast, the ring

$$\mathbb{k}[x_1, x_2, x_3, \dots]$$

is not Noetherian, because

$$(x_1) \subsetneq (x_1, x_2) \subsetneq (x_1, x_2, x_3) \subsetneq \dots$$

is a strictly increasing chain of ideals.

Exercise IV.A.3. Let X be a scheme.

- i) Show that X is locally Noetherian if and only if $\mathcal{O}_X(U)$ is Noetherian for every affine open subset $U \subset X$.
- ii) Show that every open subscheme of a locally Noetherian scheme is locally Noetherian.
- iii) Show that a quasi-compact open subscheme of a Noetherian scheme is Noetherian.

Solution. Only the forward implication in i) needs proof. Let $U = \text{Spec}(A)$ be an affine open subset and cover it by affine opens $U \cap U_i$. Refining, we may cover U by principal open subsets $D(f_j)$ such that each A_{f_j} is Noetherian. Since U is quasi-compact, finitely many suffice. The elements f_j generate the unit ideal. If $\mathfrak{a} \subset A$ is an ideal, each localisation $\mathfrak{a}_{f_j}$ is finitely generated. Choosing finitely many generators for all these localisations and clearing denominators gives a finitely generated ideal $\mathfrak{b} \subset \mathfrak{a}$ such that $\mathfrak{b}_{f_j} = \mathfrak{a}_{f_j}$ for every j . Since the $D(f_j)$ cover $\text{Spec}(A)$, we obtain $\mathfrak{a} = \mathfrak{b}$. Thus A is Noetherian.

Assertion ii) follows by restricting a Noetherian affine covering. For iii), a quasi-compact open subset has a finite affine covering by Noetherian rings, hence is Noetherian.

Proposition IV.A.4. Let $f : X \rightarrow Y$ be a morphism of finite type. If Y is Noetherian, then X is Noetherian.

Proof. Choose a finite affine covering $Y = \bigcup_{j=1}^r V_j$, with $V_j = \text{Spec}(A_j)$ and A_j Noetherian. Since f is of finite type, every $f^{-1}(V_j)$ has a finite affine covering by $U_{ij} = \text{Spec}(B_{ij})$, where B_{ij} is a finitely generated A_j -algebra. Hilbert's basis theorem implies that every B_{ij} is Noetherian. These finitely many affine opens cover X . $\square$

Proposition IV.A.5. A Noetherian scheme has only finitely many irreducible components.

Proof. It is enough to treat $X = \operatorname{Spec}(A)$ with A Noetherian. Replacing A by $A/\operatorname{Rad}(A)$ does not change the irreducible components, so assume that A is reduced.

Let $\mathcal{F}$ be the set of radical ideals which are not intersections of finitely many prime ideals. If $\mathcal{F}$ were non-empty, the ascending chain condition would give a maximal element $\mathfrak{a} \in \mathcal{F}$. It is not prime, so there are $u, v \notin \mathfrak{a}$ with $uv \in \mathfrak{a}$. The radical ideals

$$\sqrt{\mathfrak{a} + (u)} \quad \text{and} \quad \sqrt{\mathfrak{a} + (v)}$$

strictly contain $\mathfrak{a}$, hence each is an intersection of finitely many prime ideals. Moreover,

$$\sqrt{\mathfrak{a} + (u)} \cap \sqrt{\mathfrak{a} + (v)} = \sqrt{\mathfrak{a} + (uv)} = \mathfrak{a}.$$

Thus $\mathfrak{a}$ is also a finite intersection of prime ideals, a contradiction. Therefore (0) is a finite intersection of prime ideals. It follows that A has only finitely many minimal primes, hence X has only finitely many irreducible components. $\square$

IV.B Dimension

The dimension of a scheme is defined topologically, through chains of irreducible closed subsets, and agrees in the affine case with Krull dimension.

IV.B.1 Dimension of schemes and rings

Definition IV.B.1. Let X be a topological space. A chain of irreducible closed subsets of length n is a strictly increasing chain

$$Z_0 \subsetneq Z_1 \subsetneq \cdots \subsetneq Z_n.$$

The *dimension* of X , denoted by $\dim(X)$, is the supremum of the lengths of such chains. We put $\dim(\emptyset) = -\infty$. If $Z \subset X$ is an irreducible closed subset, its *codimension* in X is

$$\operatorname{codim}_X(Z) = \sup\{n \mid Z = Z_0 \subsetneq \cdots \subsetneq Z_n \text{ is a chain of irreducible closed subsets}\}.$$

The dimension of a scheme is the dimension of its underlying topological space, and its dimension at x is $\dim(\mathcal{O}_{X,x})$.

Definition IV.B.2. Let A be a ring. For a prime ideal $\mathfrak{p}$, its *height* is

$$\operatorname{ht}(\mathfrak{p}) = \sup\{n \mid \mathfrak{p}_0 \subsetneq \mathfrak{p}_1 \subsetneq \cdots \subsetneq \mathfrak{p}_n = \mathfrak{p} \text{ are prime ideals}\}.$$

The *Krull dimension* of A is

$$\dim(A) = \sup_{\mathfrak{p} \in \operatorname{Spec}(A)} \operatorname{ht}(\mathfrak{p}).$$

We set $\dim(0) = -\infty$.

Example IV.B.3. If A is a principal ideal domain which is not a field, then $\dim(A) = 1$. Its prime ideals are (0) and the ideals generated by irreducible elements. Hence the maximal chains have the form

$$(0) \subsetneq (p),$$

or, geometrically,

$$\mathbb{W}(p) \subsetneq \operatorname{Spec}(A).$$

In particular, $\dim(\operatorname{Spec}(\mathbb{Z})) = 1$.

Example IV.B.4. In $\mathbb{A}_{\mathbb{k}}^2$, the chain

$$\{(0,0)\} = \mathbb{V}(x_1, x_2) \subsetneq \mathbb{V}(x_1) \subsetneq \mathbb{A}_{\mathbb{k}}^2$$

has length 2. Our knowledge of prime ideals of $\mathbb{k}[x, y]$ implies that no longer chain exists. So $\dim(\mathbb{A}_{\mathbb{k}}^2) = 2$.

Lemma IV.B.5. *Let X be a topological space.*

- i) *If $Y \subset X$ is closed, then $\dim(Y) \leq \dim(X)$.*
- ii) *If X is irreducible of finite dimension and $Y \subsetneq X$ is closed, then $\dim(Y) < \dim(X)$.*
- iii) *If $(X_i)_{i \in I}$ are the irreducible components of X , then*

$$\dim(X) = \sup_{i \in I} \dim(X_i).$$

Proof. A chain of irreducible closed subsets of a closed subset Y is also such a chain in X , proving i). If X is irreducible and $Y \subsetneq X$, any chain in Y can be extended by appending X , proving ii). Finally, every irreducible closed subset of X is contained in an irreducible component, which proves iii). $\square$

Proposition IV.B.6. *Let A be a ring and $X = \text{Spec}(A)$.*

- i) *We have*

$$\dim(X) = \dim(A) = \dim(A/\text{Rad}(A)).$$

- ii) *For every $\mathfrak{p} \in \text{Spec}(A)$,*

$$\dim(A_{\mathfrak{p}}) = \text{ht}(\mathfrak{p}) = \text{codim}_X(\mathbb{V}(\mathfrak{p})).$$

Proof. The correspondence $\mathfrak{p} \mapsto \mathbb{V}(\mathfrak{p})$ reverses strict inclusions. Thus chains of prime ideals in A correspond bijectively, with the same length, to chains of irreducible closed subsets of X . This proves $\dim(A) = \dim(X)$. Passing to $A/\text{Rad}(A)$ does not change the prime spectrum.

Prime ideals of $A_{\mathfrak{p}}$ correspond to prime ideals of A contained in $\mathfrak{p}$. Hence chains in $\text{Spec}(A_{\mathfrak{p}})$ correspond to chains ending at $\mathfrak{p}$, proving $\dim(A_{\mathfrak{p}}) = \text{ht}(\mathfrak{p})$. After applying $\mathbb{V}$, these are exactly the chains of irreducible closed subsets beginning with $\mathbb{V}(\mathfrak{p})$. $\square$

Exercise IV.B.7. Compute the dimensions of the following rings and schemes:

$$\begin{aligned} A_1 &= \mathbb{k}[x_1]/(x_1^m), \quad m \geq 1, \\ A_2 &= \mathbb{k}[x_1, x_2]/(x_1x_2), \\ A_3 &= \mathbb{k}[x_1, x_2]/(x_1x_2 - 1). \end{aligned}$$

For $X = \text{Spec}(A_2)$, compute the dimension of the local ring at the origin and at the prime ideal $(\bar{x}_1)$.

Solution. The spectrum of A_1 has the same topology as $\text{Spec}(\mathbb{k})$, so $\dim(A_1) = 0$.

The irreducible components of $\text{Spec}(A_2)$ are two affine lines, hence $\dim(A_2) = 1$. At the origin, there are chains

$$(\bar{x}_1) \subsetneq (\bar{x}_1, \bar{x}_2) \quad \text{and} \quad (\bar{x}_2) \subsetneq (\bar{x}_1, \bar{x}_2),$$

so the local ring has dimension 1. At the generic point $(\bar{x}_1)$ of one component, the local ring has dimension 0. Finally,

$$A_3 \simeq \mathbb{k}[x_1, x_1^{-1}],$$

which is a localisation of the one-dimensional ring $\mathbb{k}[x_1]$. It has dimension 1.

IV.B.2 Dimension of algebraic schemes over a field

Let $\mathbb{k}$ be a field. Recall that an algebraic $\mathbb{k}$ -scheme is a $\mathbb{k}$ -scheme of finite type. In this subsection we assume $\mathbb{k}$ -schemes to be reduced when speaking of their irreducible components and function fields. We use Noether normalisation in the form proved in Theorem I.D.1, together with the standard fact that finite integral extensions preserve Krull dimension.

Theorem IV.B.8. *Let X be a reduced algebraic $\mathbb{k}$ -scheme. Then X has finitely many irreducible components $X_1, \dots, X_r$, and, for $1 \leq i \leq r$, we have*

$$\dim(X_i) = \text{trdeg}_{\mathbb{k}} K(X_i)$$

Lemma IV.B.9 (Integral extensions and chains of prime ideals). *Let $B \subset A$ be an integral extension of rings.*

- i) For every prime ideal $\mathfrak{p} \subset B$, there exists a prime ideal $\mathfrak{q} \subset A$ such that $\mathfrak{q} \cap B = \mathfrak{p}$. This is the lying-over property.*
- ii) Suppose that $\mathfrak{p} \subset \mathfrak{p}'$ are prime ideals of B , and let $\mathfrak{q} \subset A$ be a prime ideal such that $\mathfrak{q} \cap B = \mathfrak{p}$. Then there exists a prime ideal $\mathfrak{q}' \subset A$ such that $\mathfrak{q} \subset \mathfrak{q}'$, and $\mathfrak{q}' \cap B = \mathfrak{p}'$. This is the going-up property.*
- iii) If $\mathfrak{q} \subset \mathfrak{q}'$ are prime ideals of A and $\mathfrak{q} \cap B = \mathfrak{q}' \cap B$, then $\mathfrak{q} = \mathfrak{q}'$. This is the incomparability property.*

Proof. Let us prove i). Fix $\mathfrak{p} \in \text{Spec}(B)$ and put $S = B \setminus \mathfrak{p}$. Localisation gives an integral extension $B_{\mathfrak{p}} = S^{-1}B \subset S^{-1}A$. Choose a maximal ideal $\mathfrak{m} \subset S^{-1}A$ and let $\mathfrak{n} = \mathfrak{m} \cap B_{\mathfrak{p}}$. Then $B_{\mathfrak{p}}/\mathfrak{n} \subset (S^{-1}A)/\mathfrak{m}$ is an integral extension of integral domains. The ring on the right is a field, so Lemma I.C.10 shows that $B_{\mathfrak{p}}/\mathfrak{n}$ is a field. Hence $\mathfrak{n}$ is a maximal ideal of $B_{\mathfrak{p}}$. Since $B_{\mathfrak{p}}$ is local, we must have $\mathfrak{n} = \mathfrak{p}B_{\mathfrak{p}}$. If $\mathfrak{q} \subset A$ is the inverse image of $\mathfrak{m}$ under $A \rightarrow S^{-1}A$, then

$$\mathfrak{q} \cap B = \mathfrak{p}.$$

Now we prove ii). Suppose $\mathfrak{p} \subset \mathfrak{p}'$ and let $\mathfrak{q} \subset A$ lie over $\mathfrak{p}$. The induced inclusion $B/\mathfrak{p} \subset A/\mathfrak{q}$ is integral. By i), applied to the prime ideal $\mathfrak{p}'/\mathfrak{p} \subset B/\mathfrak{p}$, there exists a prime ideal

$$\overline{\mathfrak{q}'} \subset A/\mathfrak{q}$$

lying over $\mathfrak{p}'/\mathfrak{p}$. Let $\mathfrak{q}' \subset A$ be its inverse image. Then $\mathfrak{q} \subset \mathfrak{q}'$ and $\mathfrak{q}' \cap B = \mathfrak{p}'$. Finally, let us prove iii). Suppose $\mathfrak{q} \subset \mathfrak{q}'$ and put $\mathfrak{p} = \mathfrak{q} \cap B = \mathfrak{q}' \cap B$. Passing to quotients gives an integral extension of integral domains

$$B/\mathfrak{p} \subset A/\mathfrak{q}.$$

Set

$$S = (B/\mathfrak{p}) \setminus \{0\}.$$

After localisation we obtain an integral extension

$$\text{Frac}(B/\mathfrak{p}) \subset S^{-1}(A/\mathfrak{q}).$$

The ring on the right is an integral domain, so Lemma I.C.10 shows that it is a field.

Now $\mathfrak{q}'/\mathfrak{q} \subset A/\mathfrak{q}$ is disjoint from S , because its contraction to $B/\mathfrak{p}$ is zero. It therefore determines a prime ideal $S^{-1}(\mathfrak{q}'/\mathfrak{q})$ of the field $S^{-1}(A/\mathfrak{q})$, hence $S^{-1}(\mathfrak{q}'/\mathfrak{q}) = 0$. Since $A/\mathfrak{q}$ is a domain and the elements of S are non-zero, this implies $\mathfrak{q}'/\mathfrak{q} = 0$. Thus $\mathfrak{q} = \mathfrak{q}'$, as required. $\square$

Corollary IV.B.10. *Let $B \subset A$ be an integral extension. Then*

$$\dim(A) = \dim(B).$$

In particular, this holds whenever A is finitely generated as a B -module.

Proof. Let $\mathfrak{p}_0 \subsetneq \mathfrak{p}_1 \subsetneq \cdots \subsetneq \mathfrak{p}_r$ be a chain of prime ideals of B . By lying over, choose a prime $\mathfrak{q}_0 \subset A$ lying over $\mathfrak{p}_0$. Repeatedly applying going up gives a chain $\mathfrak{q}_0 \subsetneq \mathfrak{q}_1 \subsetneq \cdots \subsetneq \mathfrak{q}_r$ with $\mathfrak{q}_i \cap B = \mathfrak{p}_i$. Hence

$$\dim(B) \leq \dim(A).$$

Conversely, let $\mathfrak{q}_0 \subsetneq \mathfrak{q}_1 \subsetneq \cdots \subsetneq \mathfrak{q}_r$ be a chain of prime ideals of A . Contracting to B gives $\mathfrak{q}_0 \cap B \subseteq \mathfrak{q}_1 \cap B \subseteq \cdots \subseteq \mathfrak{q}_r \cap B$. By incomparability, all these inclusions are strict. Hence $\dim(A) \leq \dim(B)$. Therefore $\dim(A) = \dim(B)$. $\square$

Proof of Theorem IV.B.8. The scheme X is Noetherian, so it has finitely many irreducible components. Fix one of them, say X_i , endowed with its reduced induced scheme structure. Then X_i is an integral algebraic $\mathbb{k}$ -scheme.

Choose a finite affine open covering $X_i = U_1 \cup \cdots \cup U_r$, with $U_j = \text{Spec}(A_j)$, with every U_j non-empty. Since X_i is integral, every A_j is an integral domain, and Lemma III.B.1 gives

$$K(X_i) = \text{Frac}(A_j).$$

Fix j with $1 \leq j \leq r$. By Theorem I.D.1, there is a polynomial subring $B = \mathbb{k}[y_1, \dots, y_n] \subset A_j$ such that A_j is finite as a B -module. Hence $\dim(A_j) = \dim(B) = n$.

We claim that $\text{Frac}(A_j)$ is a finite extension of $\text{Frac}(B)$. Choose $a_1, \dots, a_s \in A_j$ which generate A_j as a B -module, and put

$$F = \text{Frac}(B), \quad L = \text{Frac}(A_j).$$

Inside L , consider the finite-dimensional F -vector space

$$V = Fa_1 + \cdots + Fa_s.$$

Since $a_\ell a_m \in A_j$, for every ℓ, m we may write

$$a_\ell a_m = \sum_q b_{\ell m q} a_q, \quad b_{\ell m q} \in B.$$

Thus V is closed under multiplication. Moreover $1 \in A_j \subset V$, so V is a subring of L . Since L is a field, V is an integral domain. A finite-dimensional algebra over a field which is an integral domain is a field: if $0 \neq v \in V$, multiplication by v gives an injective F -linear map

$$V \longrightarrow V, \quad w \longmapsto vw.$$

Since V is finite-dimensional, this map is surjective, and hence $vw = 1$ for some $w \in V$.

Therefore V is a field containing A_j . It contains $\text{Frac}(A_j) = L$, while $V \subset L$ by construction. Hence $V = L$, and

$$[\text{Frac}(A_j) : \text{Frac}(B)] < \infty.$$

In particular $\text{Frac}(A_j)$ is algebraic over

$$\text{Frac}(B) = \mathbb{k}(y_1, \dots, y_n),$$

and consequently

$$\text{trdeg}_{\mathbb{k}} K(X_i) = \text{trdeg}_{\mathbb{k}} \text{Frac}(A_j) = n.$$

We have proved that

$$\dim(U_j) = \dim(A_j) = \text{trdeg}_{\mathbb{k}} K(X_i)$$

for every j .

It remains to compare the dimension of X_i with those of the affine opens U_j . Clearly $\dim(U_j) \leq \dim(X_i)$ for every j . Conversely, let

$$Z_0 \subsetneq Z_1 \subsetneq \cdots \subsetneq Z_m$$

be a chain of irreducible closed subsets of X_i . Choose $x \in Z_0$ and an index j such that $x \in U_j$. Then every $Z_\ell \cap U_j$ is a non-empty open subset of the irreducible space Z_ℓ , hence is irreducible and dense in Z_ℓ . The inclusions

$$Z_0 \cap U_j \subsetneq Z_1 \cap U_j \subsetneq \cdots \subsetneq Z_m \cap U_j$$

are still strict: equality at some stage would imply, after taking closures in X_i , that $Z_\ell = Z_{\ell+1}$. Thus

$$m \leq \dim(U_j).$$

It follows that

$$\dim(X_i) = \max_j \dim(U_j) = \text{trdeg}_{\mathbb{k}} K(X_i),$$

as required. $\square$

Remark IV.B.11 (The base field is part of the data). There is no ambiguity in the formula above: an algebraic $\mathbb{k}$ -scheme includes its structure morphism to $\text{Spec}(\mathbb{k})$. When X is integral, this determines an embedding

$$\mathbb{k} \hookrightarrow K(X),$$

and it is with respect to this embedding that $\text{trdeg}_{\mathbb{k}} K(X)$ is computed.

If $K/\mathbb{k}$ is a field extension, the scheme X does not automatically become a K -scheme. There is a natural construction which produces a K -scheme from X , called *base change*; it will be introduced after fibred products have been constructed.

If an integral scheme X is already equipped with compatible structures over both $\mathbb{k}$ and K , then the tower formula gives

$$\text{trdeg}_{\mathbb{k}} K(X) = \text{trdeg}_{\mathbb{k}} K + \text{trdeg}_K K(X).$$

In particular, if $K/\mathbb{k}$ is algebraic, the two transcendence degrees coincide.

Example IV.B.12. The affine hypersurface

$$X = \mathbb{W}(x_1 x_2 - 1) \subset \mathbb{A}_{\mathbb{k}}^2$$

has function field $\mathbb{k}(x_1)$ and therefore dimension 1. More generally, a proper irreducible hypersurface in $\mathbb{A}_{\mathbb{k}}^n$ has dimension $n - 1$.

Corollary IV.B.13. For every $n \geq 0$,

$$\dim(\mathbb{A}_{\mathbb{k}}^n) = \dim(\mathbb{P}_{\mathbb{k}}^n) = n.$$

Every proper closed subscheme of $\mathbb{A}_{\mathbb{k}}^n$ or $\mathbb{P}_{\mathbb{k}}^n$ has dimension at most $n - 1$.

Exercise IV.B.14. Let

$$A = \mathbb{k}[x_1, x_2]/(x_2^2 - x_1^3).$$

i) Show that A is an integral domain.

- ii) Show that A is a finitely generated module over the polynomial subring $\mathbb{k}[\bar{x}_1]$, generated by 1 and $\bar{x}_2$.
- iii) Compute $\dim(A)$ and $\text{trdeg}_{\mathbb{k}} \text{Frac}(A)$.
- iv) Put $t = \bar{x}_2/\bar{x}_1 \in \text{Frac}(A)$. Show that $\bar{x}_1 = t^2$ and $\bar{x}_2 = t^3$, and deduce that $\text{Frac}(A) \simeq \mathbb{k}(t)$.

Solution. The polynomial $x_2^2 - x_1^3$ is irreducible: viewed as a polynomial in x_2 over $\mathbb{k}(x_1)$, it would be reducible only if x_1^3 were a square in $\mathbb{k}(x_1)$. Hence the quotient A is an integral domain.

The relation $\bar{x}_2^2 = \bar{x}_1^3$ reduces every power of $\bar{x}_2$ to a $\mathbb{k}[\bar{x}_1]$ -linear combination of 1 and $\bar{x}_2$. Thus A is finite over $\mathbb{k}[\bar{x}_1]$. Finite integral extensions preserve dimension, so

$$\dim(A) = 1.$$

Moreover $\text{Frac}(A)$ is algebraic over $\mathbb{k}(\bar{x}_1)$ and therefore has transcendence degree 1.

In the fraction field,

$$t^2 = \frac{\bar{x}_2^2}{\bar{x}_1^2} = \bar{x}_1, \quad t^3 = t\bar{x}_1 = \bar{x}_2.$$

Thus $\text{Frac}(A) = \mathbb{k}(t)$.

V Regular schemes

Further reading. For regular local rings, nonsingular varieties and the Jacobian criterion, see [Har77, Chapter I, §5] and [Har77, Chapter II, §8].

The purpose of this section is to give a precise meaning to the comparison between the dimension of a scheme and the dimension of its Zariski tangent space that already appeared in Example I.B.8.

V.A Regular local rings and the cotangent space

Let $(A, \mathfrak{m})$ be a Noetherian local ring and let $\kappa = A/\mathfrak{m}$ be its residue field. The vector space

$$\mathfrak{m}/\mathfrak{m}^2$$

is the *cotangent space* of A . By Nakayama's lemma, its dimension is the minimal number of generators of the maximal ideal.

Definition V.A.1. The *embedding dimension* of A is

$$\text{edim}(A) = \dim_{\kappa}(\mathfrak{m}/\mathfrak{m}^2).$$

The local ring A is *regular* if

$$\dim(A) = \text{edim}(A).$$

A locally Noetherian scheme X is *regular at* $x \in X$ if $\mathcal{O}_{X,x}$ is a regular local ring. It is a *regular scheme* if it is regular at every point.

The terminology is justified by a standard inequality from local algebra.

Proposition V.A.2. For every Noetherian local ring $(A, \mathfrak{m})$,

$$\dim(A) \leq \dim_{\kappa}(\mathfrak{m}/\mathfrak{m}^2).$$

Idea of proof. Choose elements $a_1, \dots, a_e \in \mathfrak{m}$ whose classes form a basis of $\mathfrak{m}/\mathfrak{m}^2$. Nakayama's lemma gives $\mathfrak{m} = (a_1, \dots, a_e)$. Krull's height theorem then gives $\dim(A) = \text{ht}(\mathfrak{m}) \leq e$. $\square$

Thus regularity says that the cotangent space has the smallest possible dimension.

For a scheme X and a point x , put

$$\mathfrak{m}_x = \mathfrak{m}_{X,x}, \quad \kappa(x) = \mathcal{O}_{X,x}/\mathfrak{m}_x.$$

By the definition of the intrinsic Zariski tangent space,

$$T_x X = \text{Hom}_{\kappa(x)}(\mathfrak{m}_x/\mathfrak{m}_x^2, \kappa(x)).$$

Consequently

$$\dim_x X = \dim(\mathcal{O}_{X,x}) \leq \dim_{\kappa(x)} T_x X,$$

and equality holds precisely when X is regular at x .

Example V.A.3 (Three familiar singularities). At the origin of the cusp $y^2 = x^3$ and of the node $xy = 0$, the local dimension is 1, whereas the Jacobian vanishes and the tangent space has dimension 2. Both points are therefore non-regular. For the dual numbers

$$A = \mathbb{k}[\epsilon]/(\epsilon^2),$$

the unique local ring has dimension 0, while $(\epsilon)/(\epsilon^2)$ has dimension 1. Thus a non-reduced point is not regular either.

V.B The Jacobian criterion for regularity

Let

$$X = \mathbb{V}(f_1, \dots, f_r) \subset \mathbb{A}_{\mathbb{k}}^n$$

and let x be a closed point. Assume that the finite extension $\kappa(x)/\mathbb{k}$ is separable; this holds, in particular, when $\mathbb{k}$ is perfect. Write

$$J_x = \left(\frac{\partial f_i}{\partial x_j}(x) \right)$$

for the Jacobian matrix with entries in $\kappa(x)$. As in Example I.B.8, its kernel is the relative Zariski tangent space at x . By the discussion following Definition I.B.4, separability identifies this space with the intrinsic tangent space

$$T_x X = \text{Hom}_{\kappa(x)}(\mathfrak{m}_x/\mathfrak{m}_x^2, \kappa(x)).$$

Theorem V.B.1 (Jacobian criterion for regularity). Put $d = \dim(\mathcal{O}_{X,x})$. Then the following conditions are equivalent:

- i) $\mathcal{O}_{X,x}$ is regular;
- ii) the Zariski tangent space $T_x X$ has dimension d ;
- iii) the Jacobian matrix J_x has rank $n - d$.

Proof. By definition, the local ring $\mathcal{O}_{X,x}$ is regular if and only if

$$d = \dim_{\kappa(x)}(\mathfrak{m}_x/\mathfrak{m}_x^2).$$

Since $T_x X$ is the dual of $\mathfrak{m}_x/\mathfrak{m}_x^2$, this is equivalent to

$$\dim_{\kappa(x)} T_x X = d.$$

On the other hand the computation with dual numbers gives

$$\dim_{\kappa(x)} T_x X = n - \text{rank}(J_x).$$

Thus the second condition is equivalent to $\text{rank}(J_x) = n - d$. □

Exercise V.B.2. Assume that $\mathbb{k}$ is algebraically closed and that $\text{char}(\mathbb{k}) \neq 2, 3$. Determine the regular locus of the cuspidal curve

$$C = \mathbb{W}(y^2 - x^3) \subset \mathbb{A}_{\mathbb{k}}^2.$$

Solution. For $f = y^2 - x^3$, the gradient is

$$(-3x^2, 2y).$$

It vanishes on the curve only at $(0, 0)$. Every closed point different from the origin is therefore regular by the Jacobian criterion. The generic point is regular as well, since its local ring is the function field of C , hence a field. Thus the regular locus is

$$C \setminus \{(0, 0)\}.$$

Chapter 5

Global properties of morphisms

I Fibred products

One important construction is that of products, or fibred products, in the category of schemes.

I.A The universal property of the fibred product

Definition I.A.1. Let $f : X \rightarrow S$ and $g : Y \rightarrow S$ be morphisms of schemes. A scheme W is the *fibred product* of f and g , denoted by $X \times_S Y$, if W is equipped with morphisms $F : W \rightarrow X$ and $G : W \rightarrow Y$ such that $f \circ F = g \circ G$ and, given any scheme T and morphisms $u : T \rightarrow X$ and $v : T \rightarrow Y$ with $f \circ u = g \circ v$, there is a unique morphism $w : T \rightarrow W$ such that $F \circ w = u$ and $G \circ w = v$. This is expressed in the next diagram.

$$\begin{array}{ccccc}
 & & & & \\
 & & & & \\
 T & & & & \\
 & \searrow v & & & \\
 & & W & \xrightarrow{F} & Y \\
 & \swarrow u & \downarrow G & & \downarrow g \\
 & & X & \xrightarrow{f} & S
 \end{array} \tag{I.1}$$

Remark I.A.2. The fibred product has the following immediate properties.

- i) It is unique up to a unique isomorphism compatible with the two projections. Indeed, if W_1 and W_2 both satisfy the universal property, it gives unique morphisms

$$W_1 \longrightarrow W_2, \quad W_2 \longrightarrow W_1$$

compatible with the projections. The two composites must be the identity by uniqueness.

- ii) Let

$$p_X : X \times_S Y \longrightarrow X$$

be the first projection. For every open subscheme $U \subset X$, the inverse image $p_X^{-1}(U)$ is canonically isomorphic to $U \times_S Y$.

- iii) Since every scheme has a unique morphism to $\text{Spec}(\mathbb{Z})$, the product of two schemes is their fibred product over $\text{Spec}(\mathbb{Z})$.

Exercise I.A.3. Prove item ii) of Remark I.A.2 directly from the universal property.

Solution. Let

$$W = X \times_S Y$$

and let $p_X : W \rightarrow X$ be the first projection. Put

$$W_U = p_X^{-1}(U).$$

The restrictions of the two projections give morphisms

$$W_U \longrightarrow U, \quad W_U \longrightarrow Y$$

over S .

Let T be a scheme equipped with morphisms $u : T \rightarrow U$ and $v : T \rightarrow Y$ over S . Composing u with $U \hookrightarrow X$, the universal property of W gives a unique morphism $w : T \rightarrow W$. Since $p_X \circ w$ has image in U , the morphism w factors uniquely through the open subscheme W_U . This is exactly the universal property of $U \times_S Y$.

I.B Constructing the fibred product

The next result is crucial in the understanding of schemes and families of schemes.

Theorem I.B.1. *For every scheme S , the category $(\text{Sch})_S$ admits finite products. Equivalently, fibred products of schemes exist.*

Proof. It is enough to construct the product of two S -schemes X and Y .

1. Assume first that

$$S = \text{Spec}(R), \quad X = \text{Spec}(A), \quad Y = \text{Spec}(B).$$

Set

$$W = \text{Spec}(A \otimes_R B).$$

The homomorphisms

$$A \longrightarrow A \otimes_R B, \quad a \longmapsto a \otimes 1, \quad B \longrightarrow A \otimes_R B, \quad b \longmapsto 1 \otimes b$$

give morphisms $W \rightarrow X$ and $W \rightarrow Y$ over S .

Let T be a scheme. By Theorem I.E.1, giving morphisms $T \rightarrow X$ and $T \rightarrow Y$ over S is equivalent to giving homomorphisms

$$A \longrightarrow \Gamma(\mathcal{O}_T), \quad B \longrightarrow \Gamma(\mathcal{O}_T)$$

which agree on R . By the universal property of the tensor product, this is equivalent to giving a unique homomorphism

$$A \otimes_R B \longrightarrow \Gamma(\mathcal{O}_T),$$

hence a unique morphism $T \rightarrow W$. Thus

$$X \times_S Y \simeq \text{Spec}(A \otimes_R B).$$

2. Assume that S and Y are affine, while X is arbitrary. Choose an affine open covering

$$X = \bigcup_{i \in I} U_i.$$

By the affine case, the fibred products

$$W_i = U_i \times_S Y$$

exist. If $U_{ij} = U_i \cap U_j$, then

$$W_i|_{U_{ij}} = U_{ij} \times_S Y = W_j|_{U_{ij}}$$

canonically. These identifications satisfy the cocycle condition, so the W_i glue to a scheme W with morphisms

$$W \longrightarrow X, \quad W \longrightarrow Y.$$

To verify the universal property, let $u : T \rightarrow X$ and $v : T \rightarrow Y$ be morphisms over S . The open subsets $T_i = u^{-1}(U_i)$ cover T , and the affine case gives compatible morphisms $T_i \rightarrow W_i$. They glue uniquely to a morphism $T \rightarrow W$. Hence $W \simeq X \times_S Y$.

3. Assume that S is affine and X, Y are arbitrary. Choose an affine open covering

$$Y = \bigcup_{j \in J} V_j.$$

By the preceding step, each $X \times_S V_j$ exists. Their restrictions to the overlaps $V_j \cap V_\ell$ are canonically isomorphic, so they glue to $X \times_S Y$.

4. Finally, let S be arbitrary. Choose an affine open covering

$$S = \bigcup_{k \in K} S_k$$

and put

$$U_k = X \times_S S_k = f^{-1}(S_k), \quad V_k = Y \times_S S_k = g^{-1}(S_k).$$

By the preceding step,

$$W_k = U_k \times_{S_k} V_k$$

exists. It also represents $U_k \times_S Y$: if $T \rightarrow U_k$ and $T \rightarrow Y$ have the same composite with S , then the second morphism factors uniquely through V_k .

On $S_k \cap S_\ell$, the corresponding restrictions of W_k and W_ℓ are canonically isomorphic. These isomorphisms satisfy the cocycle condition, so the W_k glue to a scheme W . The universal property is local on T , and therefore follows from the universal properties of the W_k . Thus $W \simeq X \times_S Y$.

□

Example I.B.2 (Scheme-theoretic intersections). Let A be a ring and let $\mathfrak{a}, \mathfrak{b} \subset A$ be ideals. The two closed subschemes

$$Z_1 = \operatorname{Spec}(A/\mathfrak{a}), \quad Z_2 = \operatorname{Spec}(A/\mathfrak{b})$$

of $\operatorname{Spec}(A)$ have fibred product

$$Z_1 \times_{\operatorname{Spec}(A)} Z_2 \simeq \operatorname{Spec}((A/\mathfrak{a}) \otimes_A (A/\mathfrak{b})) \simeq \operatorname{Spec}(A/(\mathfrak{a} + \mathfrak{b})).$$

Thus the fibred product recovers the scheme-theoretic intersection of two closed subschemes.

Exercise I.B.3. In $\mathbb{A}_{\mathbb{k}}^2 = \text{Spec}(\mathbb{k}[x, y])$, let

$$L = \mathbb{W}(y), \quad P = \mathbb{W}(y - x^2).$$

Compute $L \times_{\mathbb{A}_{\mathbb{k}}^2} P$. Compare it with the set-theoretic intersection of the line and the parabola.

Solution. Using the preceding example,

$$L \times_{\mathbb{A}_{\mathbb{k}}^2} P \simeq \text{Spec} \left(\frac{\mathbb{k}[x, y]}{(y) + (y - x^2)} \right) \simeq \text{Spec}(\mathbb{k}[x]/(x^2)).$$

The underlying topological space consists only of the origin, which is the set-theoretic intersection. Scheme-theoretically, however, the intersection is a double point: the class of x is nonzero and has square zero.

I.C Properties of the fibred product

First observe that the fibred product need not agree with the set-theoretic product of X and Y along S .

Remark I.C.1. The underlying set of a fibred product need not be the set-theoretic fibred product. Let

$$S = \text{Spec}(\mathbb{R}), \quad X = \text{Spec}(\mathbb{C}).$$

The set-theoretic fibred product of the one-point spaces $|X|$ with itself over $|S|$ has one point. On the other hand,

$$X \times_S X = \text{Spec}(\mathbb{C} \otimes_{\mathbb{R}} \mathbb{C}).$$

Using one copy of $\mathbb{C}$ as the coefficient field, we have

$$\mathbb{C} \otimes_{\mathbb{R}} \mathbb{C} \simeq \mathbb{C}[t]/(t^2 + 1) \simeq \mathbb{C} \times \mathbb{C},$$

where the last isomorphism follows from the factorisation

$$t^2 + 1 = (t - i)(t + i).$$

Thus $X \times_S X$ has two points.

Exercise I.C.2. Let X and Y be S -schemes and let Z be a Y -scheme. Prove the following properties.

1. $X \times_S S \simeq X$;
2. $X \times_S Y \simeq Y \times_S X$;
3. $(X \times_S Y) \times_Y Z \simeq X \times_S Z$.

Solution. All three isomorphisms follow from the universal property.

1. The morphisms

$$X \longrightarrow X, \quad X \longrightarrow S$$

give a morphism $X \rightarrow X \times_S S$. Conversely, the first projection gives $X \times_S S \rightarrow X$. The two composites are identities by uniqueness.

2. Exchanging the two projections gives a morphism

$$X \times_S Y \longrightarrow Y \times_S X.$$

The same construction in the other direction gives its inverse.

3. A morphism $T \rightarrow (X \times_S Y) \times_Y Z$ is equivalent to morphisms

$$T \rightarrow X, \quad T \rightarrow Y, \quad T \rightarrow Z$$

such that the map to Y is both the composite $T \rightarrow X \rightarrow S$ together with the structure data and the composite $T \rightarrow Z \rightarrow Y$. Equivalently, it is the datum of morphisms $T \rightarrow X$ and $T \rightarrow Z$ having the same composite with S . This is the universal property of $X \times_S Z$.

I.D Base change

Let S be a scheme and consider a scheme X over S and a morphism of schemes $g : S' \rightarrow S$. Then, using the fibred product, we obtain an S' -scheme $X' = X \times_S S' \rightarrow S'$. We say that X' is obtained by base change from X via g .

Remark I.D.1. Let B be a ring and let A be a positively graded B -algebra. Let C be another B -algebra. Then one can show that

$$\mathrm{Proj}(A \otimes_B C) \simeq \mathrm{Proj}(A) \times_{\mathrm{Spec}(B)} \mathrm{Spec}(C).$$

For instance, we defined the projective space $\mathbb{P}_{\mathbb{Z}}^n = \mathrm{Proj}(\mathbb{Z}[x_0, \dots, x_n])$. Then, given a scheme Y , by base change we obtain *the projective space over Y*

$$\mathbb{P}_Y^n = \mathbb{P}_{\mathbb{Z}}^n \times_{\mathrm{Spec}(\mathbb{Z})} Y.$$

Example I.D.2 (Extension of the ground field). Let $X = \mathrm{Spec}(A)$ be an affine $\mathbb{k}$ -scheme and let $K/\mathbb{k}$ be a field extension. The base change of X to K is

$$X_K = X \times_{\mathrm{Spec}(\mathbb{k})} \mathrm{Spec}(K) \simeq \mathrm{Spec}(A \otimes_{\mathbb{k}} K).$$

Thus base change replaces the coefficients in the equations defining X by coefficients in K .

Proposition I.D.3 (Dimension and extension of the ground field). *Let X be an integral algebraic $\mathbb{k}$ -scheme and let $K/\mathbb{k}$ be a field extension. Then*

$$\dim(X_K) = \dim(X).$$

Proof. Choose a finite covering of X by non-empty affine open subsets

$$U_i = \mathrm{Spec}(A_i).$$

By Proposition IV.B.8, all the U_i have the same dimension

$$d = \dim(X) = \mathrm{trdeg}_{\mathbb{k}} K(X).$$

For each i , Noether normalisation gives a finite inclusion

$$\mathbb{k}[y_1, \dots, y_d] \subset A_i.$$

Tensoring with K gives a finite inclusion

$$K[y_1, \dots, y_d] \subset A_i \otimes_{\mathbb{k}} K.$$

Finite integral extensions preserve Krull dimension, so

$$\dim(A_i \otimes_{\mathbb{k}} K) = d.$$

The schemes

$$(U_i)_K = \operatorname{Spec}(A_i \otimes_{\mathbb{k}} K)$$

form a finite open covering of X_K . As in the proof of Proposition IV.B.8, the dimension of a scheme is the supremum of the dimensions of the members of an open covering. Hence

$$\dim(X_K) = d = \dim(X).$$

□

Exercise I.D.4. Put $A = \mathbb{Z}[u]/(u^2 - 2)$ and let

$$X = \operatorname{Spec}(A)$$

with structure morphism $\operatorname{Spec}(\gamma_A) : X \rightarrow \operatorname{Spec}(\mathbb{Z})$. Compute the fibres over the generic point (0) and over the closed points (2), (3), and (7).

Solution. The generic fibre is

$$X_{(0)} \simeq \operatorname{Spec}(\mathbb{Q}[u]/(u^2 - 2)) = \operatorname{Spec}(\mathbb{Q}(\sqrt{2})).$$

Over 2, we obtain

$$X_{(2)} \simeq \operatorname{Spec}(\mathbb{F}_2[u]/(u^2)),$$

a non-reduced point.

Over 3, the polynomial $u^2 - 2 = u^2 + 1$ has no root in $\mathbb{F}_3$, so

$$X_{(3)} \simeq \operatorname{Spec}(\mathbb{F}_9)$$

is a single point with residue field $\mathbb{F}_9$.

Finally, 2 is a square modulo 7, since $3^2 = 2$. Hence

$$u^2 - 2 = (u - 3)(u + 3)$$

in $\mathbb{F}_7[u]$, and the Chinese remainder theorem gives

$$X_{(7)} \simeq \operatorname{Spec}(\mathbb{F}_7 \times \mathbb{F}_7),$$

the disjoint union of two $\mathbb{F}_7$ -rational points.

II Smoothness over a field

Further reading. For geometric regularity and smoothness over a field, see [Liu02].

Regularity is intrinsic: it only uses the local rings of a scheme. Smoothness over a field asks that regularity persist after extending the ground field. The base-change construction makes this precise.

Definition II.1. Let X be a scheme of finite type over $\mathbb{k}$. We say that X is *smooth over $\mathbb{k}$* if, for an algebraic closure $\bar{\mathbb{k}}$ of $\mathbb{k}$, the base change

$$X_{\bar{\mathbb{k}}} = X \times_{\operatorname{Spec}(\mathbb{k})} \operatorname{Spec}(\bar{\mathbb{k}})$$

is regular. We say that X is smooth at x if $X_{\bar{\mathbb{k}}}$ is regular at every point lying over x .

This is the notion usually called *geometric regularity*. For schemes of finite type over a field it is equivalent to the usual definition of a smooth structure morphism. The passage to an algebraic closure is essential: a regular scheme need not remain regular after extending the ground field.

Theorem II.2 (Regular versus smooth). *Let X be a scheme of finite type over a field $\mathbb{k}$.*

- i) If X is smooth over $\mathbb{k}$, then X is regular.*
- ii) If $\mathbb{k}$ is perfect, then X is smooth over $\mathbb{k}$ if and only if X is regular.*
- iii) The regular locus and the smooth locus are open subsets of X .*

In particular, regularity and smoothness agree over a field of characteristic zero or over an algebraically closed field.

We use this standard theorem without proof. Its second assertion says that, over a perfect field, regularity is preserved by extension of the ground field. The next example shows what can fail over a non-perfect field.

Example II.3 (Regular but not smooth). Assume that $\text{char}(\mathbb{k}) = p > 0$ and that $\mathbb{k}$ is not perfect. Choose $a \in \mathbb{k}$ which is not a p -th power and put

$$L = \mathbb{k}[t]/(t^p - a), \quad X = \text{Spec}(L).$$

The ring L is a field, so X is a regular zero-dimensional scheme. If $K = \mathbb{k}(a^{1/p})$, however, then

$$L \otimes_{\mathbb{k}} K \simeq K[u]/(u^p), \quad u = t - a^{1/p}.$$

Thus the base change X_K is non-reduced and is not regular. Hence X is not smooth over $\mathbb{k}$.

The Jacobian criterion for regularity proved in the previous chapter now immediately gives the corresponding criterion for smoothness over a perfect field.

Corollary II.4 (Jacobian criterion for smoothness). *Assume that $\mathbb{k}$ is perfect, let*

$$X = \mathbb{V}(f_1, \dots, f_r) \subset \mathbb{A}_{\mathbb{k}}^n,$$

let x be a closed point, and put $d = \dim(\mathcal{O}_{X,x})$. Then the following conditions are equivalent:

- i) X is smooth over $\mathbb{k}$ at x ;*
- ii) $\mathcal{O}_{X,x}$ is regular;*
- iii) the Zariski tangent space at x has dimension d ;*
- iv) the Jacobian matrix has rank $n - d$ at x .*

Proof. The equivalence of i) and ii) is Theorem II.2. The remaining equivalences are exactly Theorem V.B.1. $\square$

Exercise II.5. i) Use the Jacobian criterion to show that the projective conic

$$C = \mathbb{V}_+(xy - z^2) \subset \mathbb{P}_{\mathbb{k}}^2$$

is smooth over every field $\mathbb{k}$.

- ii) For the field $L/\mathbb{k}$ in Example II.3, verify directly that $\mathrm{Spec}(L)$ has zero-dimensional local ring and zero-dimensional cotangent space, but that its base change to $K = \mathbb{k}(a^{1/p})$ has a one-dimensional cotangent space.

Solution. The partial derivatives of $xy - z^2$ are

$$y, \quad x, \quad -2z.$$

After extending the ground field to an algebraic closure, if they all vanished at a point of the conic, then $x = y = 0$; the equation would force $z = 0$, which is not a projective point. This also works in characteristic 2, where the first two derivatives already suffice. Applying the affine Jacobian criterion on the standard charts shows that $C_{\mathbb{k}}$ is regular, hence C is smooth over $\mathbb{k}$.

The local ring of $\mathrm{Spec}(L)$ is the field L , whose maximal ideal and cotangent space are zero. After base change,

$$L \otimes_{\mathbb{k}} K \simeq K[u]/(u^p),$$

whose maximal ideal is (u) ; the quotient $(u)/(u^2)$ is one-dimensional over K . The base-changed point is therefore not regular.

III Fibres

III.A Fibres and residue fields

Let $f : X \rightarrow Y$ be a morphism of schemes and let $y \in Y$. The residue field at y is

$$\kappa(y) = \mathcal{O}_{Y,y}/\mathfrak{m}_{Y,y}.$$

There is a canonical morphism

$$i_y : \mathrm{Spec}(\kappa(y)) \longrightarrow Y$$

whose image is the point y . If $V = \mathrm{Spec}(A)$ is an affine open neighbourhood of y and y corresponds to the prime ideal $\mathfrak{p} \subset A$, then this morphism is induced by

$$A \longrightarrow A_{\mathfrak{p}} \longrightarrow A_{\mathfrak{p}}/\mathfrak{p}A_{\mathfrak{p}} = \kappa(\mathfrak{p}).$$

Notice that i_y is a closed immersion if and only if y is a closed point.

Definition III.A.1. The *fibre of f over y* is the $\kappa(y)$ -scheme

$$X_y = X \times_Y \mathrm{Spec}(\kappa(y)).$$

Proposition III.A.2. Let $y \in Y$. The projection $X_y \rightarrow X$ induces a homeomorphism

$$|X_y| \xrightarrow{\sim} f^{-1}(y),$$

where the right-hand side is equipped with the subspace topology.

Proof. The assertion is local on both X and Y . We may therefore assume that

$$Y = \mathrm{Spec}(A), \quad X = \mathrm{Spec}(B),$$

that f is induced by a ring homomorphism $\varphi : A \rightarrow B$, and that y corresponds to a prime ideal $\mathfrak{p} \subset A$. Then

$$X_y = \mathrm{Spec}(B \otimes_A \kappa(\mathfrak{p})).$$

Since

$$B \otimes_A \kappa(\mathfrak{p}) \simeq B_{\mathfrak{p}}/\mathfrak{p}B_{\mathfrak{p}},$$

its prime ideals correspond first to the prime ideals of $B_{\mathfrak{p}}$ containing $\mathfrak{p}B_{\mathfrak{p}}$, and hence to the prime ideals $\mathfrak{q} \subset B$ satisfying

$$\varphi^{-1}(\mathfrak{q}) \subset \mathfrak{p} \quad \text{and} \quad \varphi(\mathfrak{p}) \subset \mathfrak{q}.$$

These two conditions are equivalent to

$$\varphi^{-1}(\mathfrak{q}) = \mathfrak{p}.$$

Thus the points of X_y correspond exactly to the points of X mapping to y . The localisation and quotient correspondences identify closed subsets on both sides, so this bijection is a homeomorphism. $\square$

Example III.A.3 (A family of hyperbolas degenerating to two lines). Let

$$X = \operatorname{Spec}(\mathbb{k}[t, x, y]/(xy - t))$$

and let $f : X \rightarrow \mathbb{A}_{\mathbb{k}}^1 = \operatorname{Spec}(\mathbb{k}[t])$ be induced by the inclusion $\mathbb{k}[t] \hookrightarrow \mathbb{k}[t, x, y]/(xy - t)$. For $a \in \mathbb{k}$, the fibre is

$$X_a \simeq \operatorname{Spec}(\mathbb{k}[x, y]/(xy - a)).$$

If $a \neq 0$, then x is invertible and

$$X_a \simeq \operatorname{Spec}(\mathbb{k}[x, x^{-1}]) = \mathbb{G}_{m, \mathbb{k}}.$$

The special fibre is

$$X_0 = \mathbb{V}(xy),$$

the union of the two coordinate axes. Thus the geometry of the fibres may change in a family.

Exercise III.A.4. Let

$$X = \operatorname{Spec}(\mathbb{k}[t, x]/(x^2 - t))$$

and let $f : X \rightarrow \mathbb{A}_{\mathbb{k}}^1$ be the morphism induced by $t \mapsto x^2$. Compute the fibre over $a \in \mathbb{k}$. Show that the fibre over 0 is non-reduced.

Solution. The fibre over a is

$$X_a \simeq \operatorname{Spec}(\mathbb{k}[x]/(x^2 - a)).$$

For $a = 0$, this is

$$\operatorname{Spec}(\mathbb{k}[x]/(x^2)),$$

the double point, and is therefore non-reduced. For $a \neq 0$, the structure of the fibre depends on the factorisation of $x^2 - a$ over $\mathbb{k}$: it is the disjoint union of two rational points when a is a nonzero square and $\operatorname{char}(\mathbb{k}) \neq 2$, and otherwise it may be a single point with a quadratic residue field.

Example III.A.5 (The generic fibre of the affine-plane projection). Consider the projection

$$\pi : \mathbb{A}_{\mathbb{C}}^2 = \operatorname{Spec}(\mathbb{C}[t, x]) \longrightarrow \mathbb{A}_{\mathbb{C}}^1 = \operatorname{Spec}(\mathbb{C}[t]).$$

Let $\eta = (0)$ be the generic point of the base. Since $\kappa(\eta) = \mathbb{C}(t)$, the generic fibre is

$$(\mathbb{A}_{\mathbb{C}}^2)_{\eta} = \operatorname{Spec}(\mathbb{C}[t, x] \otimes_{\mathbb{C}[t]} \mathbb{C}(t)) = \operatorname{Spec}(\mathbb{C}(t)[x]).$$

Thus the ordinary affine line over the function field $\mathbb{C}(t)$ appears naturally as the generic member of the family of affine lines parametrised by $\mathbb{A}_{\mathbb{C}}^1$. Its points correspond to prime ideals of $\mathbb{C}[t, x]$ lying over the generic point of the base.

III.B Dimension of fibres

We now combine the construction of fibres with the dimension theory developed in the previous chapter. Throughout this subsection, an *integral algebraic $\mathbb{k}$ -variety* means an integral scheme of finite type over $\mathbb{k}$.

Let $f : X \rightarrow Y$ be a dominant morphism of integral schemes, and let η be the generic point of Y . Since

$$\kappa(\eta) = K(Y),$$

the fibre

$$X_\eta = X \times_Y \operatorname{Spec}(K(Y))$$

is called the *generic fibre* of f . Dominance implies that the generic point of X maps to η , and therefore induces an inclusion of function fields

$$K(Y) \hookrightarrow K(X).$$

Proposition III.B.1 (Dimension of the generic fibre). *Let $f : X \rightarrow Y$ be a dominant morphism of integral algebraic $\mathbb{k}$ -varieties, and let η be the generic point of Y . Then X_η is an integral $K(Y)$ -variety, its function field is naturally $K(X)$, and*

$$\dim(X_\eta) = \operatorname{trdeg}_{K(Y)} K(X) = \dim(X) - \dim(Y).$$

Proof. Choose non-empty affine open subsets

$$V = \operatorname{Spec}(A) \subset Y, \quad U = \operatorname{Spec}(B) \subset f^{-1}(V).$$

Since X and Y are integral and f is dominant, the induced morphism $A \rightarrow B$ is injective. Put $S = A \setminus \{0\}$. The part of the generic fibre lying over U is

$$U_\eta = \operatorname{Spec}(B \otimes_A K(Y)) = \operatorname{Spec}(S^{-1}B).$$

This is integral, and its fraction field is $\operatorname{Frac}(B) = K(X)$. Hence the generic fibre is integral and has function field $K(X)$, viewed as an extension of $K(Y)$. By the dimension–transcendence-degree formula,

$$\dim(X_\eta) = \operatorname{trdeg}_{K(Y)} K(X).$$

Finally, the tower formula for transcendence degrees gives

$$\operatorname{trdeg}_{\mathbb{k}} K(X) = \operatorname{trdeg}_{\mathbb{k}} K(Y) + \operatorname{trdeg}_{K(Y)} K(X),$$

and therefore

$$\dim(X_\eta) = \dim(X) - \dim(Y).$$

□

The dimension of a special fibre may be larger than the dimension of the generic fibre. Nevertheless, the generic value occurs on a non-empty open subset of the base.

Theorem III.B.2 (Generic constancy of fibre dimension). *Let $f : X \rightarrow Y$ be a dominant morphism of integral algebraic $\mathbb{k}$ -varieties. Then there exists a non-empty open subset $V \subset Y$ such that, for every $y \in V$, the fibre X_y is non-empty and*

$$\dim(X_y) = \dim(X) - \dim(Y).$$

Proof. Choose a non-empty affine open subset $V_0 = \text{Spec}(A) \subset Y$. Since f is of finite type, $f^{-1}(V_0)$ is quasi-compact; choose a finite covering by non-empty affine opens

$$U_i = \text{Spec}(B_i), \quad 1 \leq i \leq m.$$

Every U_i contains the generic point of the integral scheme X . Hence the maps $A \rightarrow B_i$ are injective and

$$\text{Frac}(B_i) = K(X).$$

Put

$$r = \text{trdeg}_{K(Y)} K(X) = \dim(X) - \dim(Y).$$

For completeness, the relative form of Noether normalisation used here is obtained by applying Chapter 1 over $K(Y) = \text{Frac}(A)$ to $B_i \otimes_A K(Y)$, then clearing the finitely many denominators in the chosen generators and in their monic integral equations. It gives a non-zero element $a_i \in A$ and elements $z_{i1}, \dots, z_{ir} \in (B_i)_{a_i}$ such that

$$C_i = A_{a_i}[z_{i1}, \dots, z_{ir}] \hookrightarrow (B_i)_{a_i}$$

is finite. Replacing all the a_i by their product, we may use one common principal open subset $D(a) \subset V_0$.

It remains to ensure that the displayed inclusions survive in every fibre. The quotient $(B_i)_a/C_i$ is a finite C_i -module. We use generic freeness in the following form: if A is a domain, C is a finite-type A -algebra and M is a finite C -module, then M_b is free, hence flat, over A_b for some non-zero $b \in A$. Applying this to the finitely many quotients and multiplying a by the resulting elements, every quotient is flat over A_a . Tensoring

$$0 \longrightarrow C_i \longrightarrow (B_i)_a \longrightarrow (B_i)_a/C_i \longrightarrow 0$$

with $\kappa(y)$ is therefore still exact for every $y \in D(a)$. We obtain an injective finite homomorphism

$$\kappa(y)[z_{i1}, \dots, z_{ir}] \hookrightarrow B_i \otimes_A \kappa(y).$$

Consequently $(U_i)_y$ is finite and dominant, hence finite and surjective, over $\mathbb{A}_{\kappa(y)}^r$. It is non-empty and has dimension r .

The schemes $(U_i)_y$ form a finite open covering of X_y . Thus X_y is non-empty and

$$\dim(X_y) = \max_i \dim((U_i)_y) = r$$

for every $y \in D(a)$, which proves the theorem. $\square$

Example III.B.3 (A projection). The projection

$$\pi : \mathbb{A}_{\mathbb{k}}^{m+n} \longrightarrow \mathbb{A}_{\mathbb{k}}^m$$

is dominant. Every fibre is isomorphic to $\mathbb{A}_{\kappa(y)}^n$, and hence

$$\dim(\pi^{-1}(y)) = n = (m+n) - m.$$

Here the generic dimension occurs over the whole base.

Example III.B.4 (A reducible special fibre). For the morphism

$$f : \text{Spec}(\mathbb{k}[t, x, y]/(xy - t)) \longrightarrow \mathbb{A}_{\mathbb{k}}^1,$$

the total space has dimension 2 and the base has dimension 1. Every fibre has dimension 1. For $a \neq 0$, the fibre $xy = a$ is isomorphic to $\mathbb{G}_{m, \mathbb{k}}$, whereas

$$X_0 = \mathbb{W}(xy)$$

is the union of the two coordinate axes. Fibre dimension is constant here, even though irreducibility is not.

Example III.B.5 (A jumping fibre). Consider

$$f : \mathbb{A}_{\mathbb{k}}^2 \longrightarrow \mathbb{A}_{\mathbb{k}}^2, \quad (x, y) \longmapsto (x, xy).$$

Equivalently, f is induced by

$$\mathbb{k}[u, v] \longrightarrow \mathbb{k}[x, y], \quad u \longmapsto x, \quad v \longmapsto xy.$$

This ring map is injective, so f is dominant. Over the principal open $D(u)$, the fibre over (a, b) consists of the single point

$$x = a, \quad y = b/a.$$

Thus the generic fibre has dimension $0 = 2 - 2$. However,

$$f^{-1}(0, 0) = \mathbb{V}(x) \simeq \mathbb{A}_{\mathbb{k}}^1,$$

so the fibre dimension jumps at the origin. If $a = 0$ and $b \neq 0$, the fibre is empty. This illustrates why the theorem only asserts constancy after restricting to a suitable non-empty open subset.

Remark III.B.6. Dimension does not detect nilpotent structure. For the power map

$$\mathbb{A}_{\mathbb{k}}^1 \longrightarrow \mathbb{A}_{\mathbb{k}}^1, \quad t \longmapsto t^n,$$

the fibre over 0 is the zero-dimensional but non-reduced scheme

$$\operatorname{Spec}(\mathbb{k}[t]/(t^n)).$$

Thus the scheme structure of a fibre contains information that its dimension alone forgets.

Exercise III.B.7 (The affine generic fibre). Let $A \subset B$ be finitely generated integral $\mathbb{k}$ -algebras and let

$$f : \operatorname{Spec}(B) \longrightarrow \operatorname{Spec}(A)$$

be the corresponding dominant morphism. Put $K = \operatorname{Frac}(A)$ and $S = A \setminus \{0\}$.

- i) Show that the generic fibre is $\operatorname{Spec}(S^{-1}B)$.
- ii) Show that it is integral and that its function field is $\operatorname{Frac}(B)$.
- iii) Deduce that

$$\dim(X_\eta) = \dim(B) - \dim(A).$$

Solution. The generic point of $\operatorname{Spec}(A)$ has residue field $K = S^{-1}A$. Hence

$$X_\eta = \operatorname{Spec}(B \otimes_A K) = \operatorname{Spec}(S^{-1}B).$$

Since B is a domain and S contains no zero element of B , the localisation $S^{-1}B$ is a domain. Its fraction field is the same as that of B . The last assertion follows from

$$\dim(S^{-1}B) = \operatorname{trdeg}_K \operatorname{Frac}(B) = \operatorname{trdeg}_{\mathbb{k}} \operatorname{Frac}(B) - \operatorname{trdeg}_{\mathbb{k}} K = \dim(B) - \dim(A).$$

Exercise III.B.8 (The jumping-fibre example). For the morphism

$$f : \mathbb{A}_{\mathbb{k}}^2 \longrightarrow \mathbb{A}_{\mathbb{k}}^2, \quad (x, y) \longmapsto (x, xy),$$

compute scheme-theoretically the fibre over $(a, b) \in \mathbb{A}_{\mathbb{k}}^2(\mathbb{k})$. Determine the locus on which the fibres are zero-dimensional and identify the fibre over $(0, 0)$.

Solution. The fibre over (a, b) is

$$\operatorname{Spec}(\mathbb{k}[x, y]/(x - a, xy - b)).$$

If $a \neq 0$, the equations give $x = a$ and $y = b/a$, so the fibre is one reduced point. If $a = 0$ and $b \neq 0$, the quotient is the zero ring and the fibre is empty. If $(a, b) = (0, 0)$, the quotient is

$$\mathbb{k}[x, y]/(x, xy) \simeq \mathbb{k}[y],$$

so the fibre is $\mathbb{A}_{\mathbb{k}}^1$. Thus the fibres are zero-dimensional on $D(u)$, while the dimension jumps at the origin.

Exercise III.B.9 (Dimension versus scheme structure). Assume that $n \geq 2$ and consider the morphism

$$f : \mathbb{A}_{\mathbb{k}}^1 \rightarrow \mathbb{A}_{\mathbb{k}}^1, \quad t \mapsto t^n.$$

Compute the fibre over $a \in \mathbb{k}$. Show that every non-empty fibre has dimension 0, but that the fibre over 0 is non-reduced.

Solution. The fibre over a is

$$\operatorname{Spec}(\mathbb{k}[t]/(t^n - a)).$$

Since this is finite-dimensional as a $\mathbb{k}$ -vector space, every non-empty fibre has dimension 0. At $a = 0$ one obtains

$$\operatorname{Spec}(\mathbb{k}[t]/(t^n)),$$

which is non-reduced because the class of t is non-zero and nilpotent.

Further reading. For the classical dimension-of-fibres theorem and its relation with transcendence degree, see [Har77, Chapter I, §7]. For a systematic scheme-theoretic treatment of fibres and dimension, see [Liu02].

IV Separated morphisms

The underlying topological space of a scheme is rarely Hausdorff. For instance, if an irreducible component of X contains two distinct points, then any neighbourhood of either point meets that component in a nonempty open subset and therefore contains its generic point. The two points cannot be separated by disjoint open neighbourhoods.

IV.A Separated morphisms and the diagonal

Exercise IV.A.1. Let X be a scheme.

- i) Show that if the underlying topological space of X is Hausdorff, then $\dim(X) = 0$.
- ii) Assume in addition that X is locally Noetherian. Show that the converse holds: if $\dim(X) = 0$, then the underlying topological space of X is discrete, hence Hausdorff.

Solution. i) In a Hausdorff space every point is closed. Let $U = \operatorname{Spec}(A)$ be an affine open subset of X . Every prime ideal of A is therefore a closed point of $\operatorname{Spec}(A)$, hence is maximal. Thus there is no strict inclusion of prime ideals, and $\dim(A) = 0$. Since this holds on every affine open subset, $\dim(X) = 0$.

- ii) Let $x \in X$ and choose a Noetherian affine neighbourhood $U = \text{Spec}(A)$. Since $\dim(A) = 0$, the Noetherian ring A is Artinian. Hence $\text{Spec}(A)$ is a finite discrete space. In particular, $\{x\}$ is open in U , and therefore open in X . Thus every point of X is open, so X is discrete and hence Hausdorff.

Even if schemes are rarely Hausdorff spaces, there is a notion adapted to schemes that plays the role of the Hausdorff condition. To understand it, let us start with the following observation. The idea is that, for a morphism $X \rightarrow Y$, the diagonal $\Delta_{X/Y} \subset X \times_Y X$ plays the role of the set of pairs of points of X that map to the same point in Y . Asking it to be closed is the scheme-theoretic analogue of the Hausdorff condition.

Remark IV.A.2. Let X be a topological space. We equip $X \times X$ with the product topology and consider $\Delta = \{(x_1, x_2) \in X \times X \mid x_1 = x_2\}$. Then X is Hausdorff if and only if Δ is closed in $X \times X$.

Indeed, let us assume that Δ is closed in $X \times X$. Let x_1, x_2 be distinct elements of X . Then (x_1, x_2) does not lie in Δ , so there is an open subset U of $X \times X$ containing (x_1, x_2) and disjoint from Δ . Since $X \times X$ is equipped with the product topology, there are open subsets U_1 and U_2 of X , with $x_1 \in U_1$ and $x_2 \in U_2$, such that $U_1 \times U_2 \subset U$. Then $U_1 \times U_2 \cap \Delta = \emptyset$, which implies $U_1 \cap U_2 = \emptyset$. So X is Hausdorff.

Conversely, assume that X is Hausdorff and let us check that $X \times X \setminus \Delta$ is open in $X \times X$. Let $(x_1, x_2) \in X \times X \setminus \Delta$, so that x_1 and x_2 are distinct elements of X . Then there are disjoint neighbourhoods U_1 and U_2 of x_1 and x_2 and $U_1 \times U_2$ is an open neighbourhood of (x_1, x_2) in $X \times X \setminus \Delta$.

Let $f : X \rightarrow Y$ be a morphism of schemes. Then we consider the fibred product $X \times_Y X$. Now, the identity of X gives, by the universal property of $X \times_Y X$, a unique morphism $X \rightarrow X \times_Y X$, which we call the diagonal morphism and denote by $\Delta_{X/Y}$. If $Y = \text{Spec}(\mathbb{Z})$, we write simply $\Delta_X : X \rightarrow X \times X$.

Definition IV.A.3. A morphism $f : X \rightarrow Y$ is separated if $\Delta_{X/Y}$ is a closed immersion. We also say that X is separated over Y . A scheme X is separated if it is separated over $\text{Spec}(\mathbb{Z})$.

Remark IV.A.4. Any morphism $f : X \rightarrow Y$ of affine schemes is separated. Indeed, let $X = \text{Spec}(B)$ and $Y = \text{Spec}(A)$, so that f corresponds to a morphism of rings $A \rightarrow B$. Then the diagonal morphism $\Delta_{X/Y} : X \rightarrow X \times_Y X$ corresponds to the morphism of rings $B \otimes_A B \rightarrow B$. Since this ring map is surjective, the associated morphism of schemes is a closed immersion.

Proposition IV.A.5 (Graphs). Let $f : X \rightarrow Y$ be a morphism of S -schemes. Its graph is the morphism

$$\Gamma_f = (\text{id}_X, f) : X \longrightarrow X \times_S Y.$$

It is a monomorphism. If Y is separated over S , then Γ_f is a closed immersion.

Proof. The graph is obtained by base change from the diagonal of Y :

$$\begin{array}{ccc} X & \xrightarrow{\Gamma_f} & X \times_S Y \\ f \downarrow & & \downarrow f \times \text{id}_Y \\ Y & \xrightarrow{\Delta_{Y/S}} & Y \times_S Y. \end{array}$$

The square is cartesian. Since a diagonal is always a monomorphism, so is its base change Γ_f . If $Y \rightarrow S$ is separated, then $\Delta_{Y/S}$ is a closed immersion, and closed immersions are preserved by base change. $\square$

Example IV.A.6. If $p(t) \in \mathbb{k}[t]$, the graph of the morphism

$$\mathbb{A}_{\mathbb{k}}^1 \longrightarrow \mathbb{A}_{\mathbb{k}}^1, \quad t \longmapsto p(t),$$

is the closed subscheme of $\mathbb{A}_{\mathbb{k}}^2 = \text{Spec}(\mathbb{k}[x, y])$ defined by

$$y - p(x) = 0.$$

IV.B Affine coverings of separated schemes

In a similar vein, we have the following lemma.

Lemma IV.B.1. *Let X be a scheme. Then X is separated if and only if, for any pair of open affine subsets U, V of X , $U \cap V$ is affine and the multiplication map $\mu_{U,V} : \mathcal{O}_X(U) \otimes \mathcal{O}_X(V) \rightarrow \mathcal{O}_X(U \cap V)$ is surjective. This happens if and only if there is an affine open covering $\{U_i \mid i \in I\}$ of X , such that all $i, j \in I$, $U_i \cap U_j$ is affine and the map μ_{U_i, U_j} is surjective.*

Proof. Let us take $U = \text{Spec}(A)$ and $V = \text{Spec}(B)$ affine open subsets of X . Then $U \times V$ is an open subset of $X \times X$ and $\Delta_X^{-1}(U \times V) = U \cap V$ as a topological space, as $\Delta_X(x) = (x, x)$. Since U and V are affine, we have $U \times V = \text{Spec}(A \otimes_{\mathbb{Z}} B)$ and the morphism of schemes Δ_X , restricted to $U \cap V$, corresponds to the ring homomorphism $\mu_{U,V} : A \otimes_{\mathbb{Z}} B \rightarrow \mathcal{O}_X(U \cap V)$.

If X is separated, Δ_X is a closed immersion so this map is surjective and $U \cap V$ is a closed subscheme of the affine $U \times V$, so $U \cap V$ is affine.

Let $\{U_i \mid i \in I\}$ be an open affine covering of X . If X is separated, we can apply the above argument to any $i, j \in I$ and see that μ_{U_i, U_j} is surjective.

Conversely, assume that for all $i, j \in I$ the map μ_{U_i, U_j} is surjective and recall that $U_i \cap U_j = \Delta_X^{-1}(U_i \times U_j)$. Since the morphism $U_i \cap U_j \rightarrow U_i \times U_j$ corresponds to the map $\mu_{U_i, U_j} : \mathcal{O}_X(U_i) \otimes \mathcal{O}_X(U_j) \rightarrow \mathcal{O}_X(U_i \cap U_j)$ the map Δ_X , restricted to $U_i \cap U_j$, is a closed immersion. Since the subsets $\{U_i \times U_j \mid (i, j) \in I^2\}$ cover $X \times X$, the map Δ_X is a closed immersion. $\square$

Exercise IV.B.2. Let $\mathbb{k}$ be a field.

1. Show that the affine line over $\mathbb{k}$ with two origins is not a separated scheme.
2. Construct a scheme X by gluing two affine planes $Z_1 = \mathbb{A}_{\mathbb{k}}^2$ and $Z_2 = \mathbb{A}_{\mathbb{k}}^2$ along $Z_{1,2} = Z_1 \setminus \{(0, 0)\}$ and $Z_{2,1} = Z_2 \setminus \{(0, 0)\}$ using the identity map.
3. Check that the scheme X constructed in the previous item has a covering by two affine subschemes U_1 and U_2 isomorphic to affine planes, but that $U_1 \cap U_2$ is not affine.

Solution. 1. Let $U_1, U_2 \simeq \text{Spec}(\mathbb{k}[x])$ be the two standard affine open subsets. Their intersection is $D(x) = \text{Spec}(\mathbb{k}[x, x^{-1}])$. The multiplication map

$$\mathbb{k}[x] \otimes_{\mathbb{k}} \mathbb{k}[x] \longrightarrow \mathbb{k}[x, x^{-1}]$$

has image contained in $\mathbb{k}[x]$, and in particular does not contain x^{-1} . It is not surjective, so the affine criterion for separatedness shows that the line with two origins is not separated.

2. Glue Z_1 and Z_2 by identifying their punctured planes through the identity map. Since there are only two charts, the cocycle condition is automatic. The gluing lemma produces a scheme X containing the images U_1, U_2 of the two affine planes as open subschemes.

3. We have

$$U_1 \cap U_2 \simeq \mathbb{A}_{\mathbb{k}}^2 \setminus \{(0, 0)\} = D(x) \cup D(y).$$

Its ring of global functions is

$$\mathbb{k}[x, y]_x \cap \mathbb{k}[x, y]_y = \mathbb{k}[x, y]$$

inside $\mathbb{k}(x, y)$. If the punctured plane were affine, the canonical morphism to the spectrum of its global functions would identify it with $\text{Spec}(\mathbb{k}[x, y]) = \mathbb{A}_{\mathbb{k}}^2$, which is impossible because the origin is missing. Hence $U_1 \cap U_2$ is not affine, and X is not separated.

Example IV.B.3. Let Y be a scheme. Then the structure morphism

$$\mathbb{P}_Y^n \longrightarrow Y$$

is separated. Indeed, we first treat $\mathbb{P}_{\mathbb{Z}}^n$. Put

$$A = \mathbb{Z}[x_0, \dots, x_n]$$

and let $U_i = D_+(x_i)$. Then

$$U_i = \operatorname{Spec}(A[1/x_i]_0)$$

is affine, and

$$U_i \cap U_j = \operatorname{Spec}(A[1/x_i, 1/x_j]_0).$$

The overlap ring is obtained from $A[1/x_i]_0$ by inverting x_j/x_i :

$$A[1/x_i, 1/x_j]_0 = A[1/x_i]_0 \left[\left(\frac{x_j}{x_i} \right)^{-1} \right].$$

The element $(x_j/x_i)^{-1} = x_i/x_j$ belongs to $A[1/x_j]_0$. Hence the multiplication map

$$A[1/x_i]_0 \otimes_{\mathbb{Z}} A[1/x_j]_0 \longrightarrow A[1/x_i, 1/x_j]_0$$

is surjective. The affine criterion therefore shows that $\mathbb{P}_{\mathbb{Z}}^n$ is separated. Since separatedness is preserved by base change and

$$\mathbb{P}_Y^n \simeq \mathbb{P}_{\mathbb{Z}}^n \times_{\operatorname{Spec}(\mathbb{Z})} Y,$$

the morphism $\mathbb{P}_Y^n \rightarrow Y$ is separated.

Exercise IV.B.4. Prove the following facts.

1. Closed immersions are separated.
2. Let $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ be separated morphisms. Show that $g \circ f$ is separated.
3. Let $X \rightarrow Y$ be a separated morphism of S -schemes and consider a morphism $S' \rightarrow S$. Set $X' = X \times_S S'$ and $Y' = Y \times_S S'$ and let $f' : X' \rightarrow Y'$ be the morphism induced by base change from S to S' . Then f' is separated.
4. Let X and Y be separated S -schemes. Then $X \times_S Y$ is a separated S -scheme.

Solution. 1. A closed immersion is a monomorphism, so its relative diagonal is an isomorphism. In particular, it is a closed immersion.

2. The morphism

$$X \times_Y X \longrightarrow X \times_Z X$$

is the inverse image of $\Delta_{Y/Z}$ under $f \times f$, hence is a closed immersion. The diagonal $\Delta_{X/Z}$ is the composite

$$X \xrightarrow{\Delta_{X/Y}} X \times_Y X \longrightarrow X \times_Z X.$$

It is therefore a closed immersion.

3. The diagonal of the base-changed morphism is the base change of $\Delta_{X/Y}$. Since closed immersions are preserved by base change, f' is separated.
4. The diagonal of $X \times_S Y$ identifies with the product of the two diagonals:

$$\Delta_{(X \times_S Y)/S} = \Delta_{X/S} \times_S \Delta_{Y/S}.$$

Equivalently, it is a composite of base changes of these diagonals. Hence it is a closed immersion.

IV.C Projective morphisms

Definition IV.C.1. A morphism of schemes $X \rightarrow Y$ is *projective* if it factors as $X \rightarrow \mathbb{P}_Y^n \rightarrow Y$, where $X \rightarrow \mathbb{P}_Y^n$ is a closed immersion and $\mathbb{P}_Y^n \rightarrow Y$ is the structure morphism.

From Example IV.B.3 and Exercise IV.B.4, we get the following result.

Proposition IV.C.2. *Let $f : X \rightarrow Y$ be a projective morphism, namely, the composition of a closed immersion $X \rightarrow \mathbb{P}_Y^n$ with the structure map $\mathbb{P}_Y^n \rightarrow Y$. Then f is separated.*

Proposition IV.C.3. *Let $f, g : X \rightarrow Y$ be morphisms of S -schemes. Assume that Y is separated over S , that X is reduced, and that there exists a dense open subset $U \subset X$ such that*

$$f|_U = g|_U.$$

Then $f = g$.

Proof. Let

$$H = (f, g) : X \longrightarrow Y \times_S Y.$$

Since Y is separated over S , the diagonal

$$\Delta_{Y/S} : Y \longrightarrow Y \times_S Y$$

is a closed immersion. The inverse image

$$E = H^{-1}(\Delta_{Y/S}(Y))$$

is therefore closed in X . It contains U , so its underlying topological space is all of X . Hence f and g agree on points.

It remains to compare their maps on structure sheaves. This is local on X and Y . Let $V = \operatorname{Spec}(A) \subset Y$ be affine and let $W = \operatorname{Spec}(B) \subset f^{-1}(V) = g^{-1}(V)$ be affine. The restrictions of f and g correspond to ring homomorphisms

$$\varphi, \psi : A \longrightarrow B.$$

The open subset $U \cap W$ is dense in W . For $a \in A$, put

$$b = \varphi(a) - \psi(a).$$

For every $\mathfrak{p} \in U \cap W$, equality of the two morphisms on U gives $b/1 = 0$ in $B_{\mathfrak{p}}$. In particular $b \in \mathfrak{p}$, so the closed subset $\mathbb{V}(b)$ contains the dense subset $U \cap W$. Therefore

$$\mathbb{V}(b) = \operatorname{Spec}(B),$$

and b belongs to the nilradical of B . Since X , hence W , is reduced, we obtain $b = 0$. Thus $\varphi = \psi$, and the local maps glue to give $f = g$. $\square$

Example IV.C.4 (Why reducedness is needed). Let

$$X = \operatorname{Spec}(\mathbb{k}[x, \varepsilon]/(\varepsilon^2, x\varepsilon))$$

and let $U = D(x)$. Since x becomes invertible on U , the relation $x\varepsilon = 0$ implies $\varepsilon|_U = 0$. The two morphisms

$$f, g : X \longrightarrow \mathbb{A}_{\mathbb{k}}^1$$

corresponding respectively to

$$t \longmapsto 0, \quad t \longmapsto \varepsilon$$

agree on the dense open subset U , but they are not equal on X .

V Proper morphisms

Proper schemes and morphisms are the algebraic analogue of the notion of compactness.

V.A Closed and universally closed morphisms

Definition V.A.1. Let $f : X \rightarrow Y$ be a morphism of schemes.

- i) The morphism f is *closed* if $f(F)$ is closed in Y for every closed subset $F \subset X$.
- ii) The morphism f is *universally closed* if, for every morphism $Y' \rightarrow Y$, the base-changed morphism

$$f' : X \times_Y Y' \longrightarrow Y'$$

is closed.

Note that not all closed maps are universally closed. For instance, if $\mathbb{k}$ is a field and X is a $\mathbb{k}$ -scheme, then the structure map $X \rightarrow \text{Spec}(\mathbb{k})$ is closed. But very often this map is not universally closed.

Example V.A.2. For $n \geq 1$, the affine space $\mathbb{A}_{\mathbb{k}}^n = \text{Spec}(\mathbb{k}[x_1, \dots, x_n]) \rightarrow \text{Spec}(\mathbb{k})$ is not universally closed, for we can base change along $\mathbb{A}_{\mathbb{k}}^1 = \text{Spec}(\mathbb{k}[t]) \rightarrow \text{Spec}(\mathbb{k})$ to get a morphism $\mathbb{A}_{\mathbb{k}}^{n+1} \rightarrow \mathbb{A}_{\mathbb{k}}^1$, which is not closed. Indeed, this morphism is given by the obvious morphism of rings $\mathbb{k}[t] \rightarrow \mathbb{k}[t, x_1, \dots, x_n]$, and the closed subset $\mathbb{W}(x_1 t - 1)$ is mapped to:

$$f'(\mathbb{W}(x_1 t - 1)) = \{\mathfrak{q} \cap \mathbb{k}[t] \mid \mathfrak{q} \in \text{Spec}(\mathbb{k}[t, x_1, \dots, x_n]), (x_1 t - 1) \in \mathfrak{q}\}.$$

We show that:

$$f'(\mathbb{W}(x_1 t - 1)) = D(t) \subset \mathbb{A}_{\mathbb{k}}^1.$$

Let $\mathfrak{p} = \mathfrak{q} \cap \mathbb{k}[t]$ be a prime ideal $f'(\mathbb{W}(x_1 t - 1))$. If $t \in \mathfrak{p}$, then $t \in \mathfrak{q}$, hence $1 = tx_1 - (x_1 t - 1) \in \mathfrak{q}$, which is not the case. So any $\mathfrak{p} \in f'(\mathbb{W}(x_1 t - 1))$ satisfies $t \notin \mathfrak{p}$, hence $f'(\mathbb{W}(x_1 t - 1)) \subset D(t)$.

Conversely, let $\mathfrak{p} \in D(t)$. Then, $\mathfrak{p}$ is obtained intersecting a prime ideal of $\mathbb{k}[t]_t$ with $\mathbb{k}[t]$. Under the isomorphism

$$\mathbb{k}[x_1, t]/(x_1 t - 1) \simeq \mathbb{k}[t, t^{-1}] \simeq \mathbb{k}[t]_t,$$

a prime ideal of $\mathbb{k}[t, t^{-1}]$ is sent to a prime ideal of $\mathbb{k}[t, x_1]/(x_1 t - 1)$, which pulls back in $\mathbb{k}[t, x_1]$ to a prime ideal containing $(x_1 t - 1)$. This further pulls-back to a prime ideal $\mathfrak{q}$ of $\mathbb{k}[t, x_1, \dots, x_n]$ which contains $(x_1 t - 1)$. So, $\mathfrak{p} = \mathfrak{q} \cap \mathbb{k}[t]$ and $\mathfrak{p} \in f'(\mathbb{W}(x_1 t - 1))$.

Example V.A.3. A closed immersion $i : X \hookrightarrow Y$ is universally closed. Indeed, after any base change $Y' \rightarrow Y$, the induced morphism

$$X \times_Y Y' \longrightarrow Y'$$

is again a closed immersion, hence a closed map.

Remark V.A.4. The terminology is motivated by a familiar topological fact. If K is a compact topological space, then the projection

$$\text{pr}_1 : X \times K \longrightarrow X$$

is closed for every topological space X . Indeed, if $F \subset X \times K$ is closed and $x \notin \text{pr}_1(F)$, then $\{x\} \times K$ is contained in the open subset $(X \times K) \setminus F$. By the tube lemma, there is an open neighbourhood U of x such that

$$U \times K \subset (X \times K) \setminus F.$$

Thus U is disjoint from $\text{pr}_1(F)$, proving that the latter is closed.

The condition “universally closed” asks for the analogous property after every base change.

Definition V.A.5. Let $f : X \rightarrow Y$ be a morphism of schemes. The morphism f is *proper* if it is

- i) of finite type;
- ii) separated;
- iii) universally closed.

Recall that f is of finite type if, for every affine open subset $V = \text{Spec}(A) \subset Y$, the inverse image $f^{-1}(V)$ admits a finite covering by affine open subsets

$$U_i = \text{Spec}(B_i)$$

such that each B_i is a finitely generated A -algebra.

An A -scheme X is *proper over A* if its structure morphism

$$X \longrightarrow \text{Spec}(A)$$

is proper. A scheme is *proper* if it is proper over $\text{Spec}(\mathbb{Z})$.

Proposition V.A.6. *Closed immersions are proper.*

Proof. Let $i : X \hookrightarrow Y$ be a closed immersion. It is universally closed because every base change of i is again a closed immersion. It is separated because its diagonal is a closed immersion.

Finally, if $V = \text{Spec}(A) \subset Y$ is affine, then

$$i^{-1}(V) = \text{Spec}(A/\mathfrak{a})$$

for some ideal $\mathfrak{a} \subset A$. The quotient $A/\mathfrak{a}$ is a finitely generated A -algebra, so i is of finite type. $\square$

Exercise V.A.7. Prove the following properties.

1. The composition of proper morphisms is proper.
2. The base change of a proper morphism is proper.
3. Suppose that $f : X \rightarrow S$ is proper and factors as

$$X \xrightarrow{f'} S' \longrightarrow S,$$

where $S' \rightarrow S$ is separated. Show that f' is proper.

4. Let X and Y be S -schemes. Assume that $X \rightarrow S$ is proper, that $Y \rightarrow S$ is separated and of finite type, and that there is a surjective S -morphism $X \rightarrow Y$. Show that $Y \rightarrow S$ is proper.

Solution. 1. Morphisms of finite type are stable under composition, as are separated morphisms. If $X \rightarrow Y$ and $Y \rightarrow Z$ are universally closed, then after any base change to $Z' \rightarrow Z$, the resulting composite is a composite of closed maps, hence is closed.

2. Finite type and separatedness are preserved by base change. Universal closedness is preserved by base change directly from its definition: a further base change of $X \times_Y Y' \rightarrow Y'$ is also a base change of $X \rightarrow Y$.

3. The graph

$$\Gamma_{f'} : X \longrightarrow X \times_S S'$$

is a closed immersion because $S' \rightarrow S$ is separated. The projection

$$X \times_S S' \longrightarrow S'$$

is proper, being a base change of $X \rightarrow S$. Since

$$f' = \text{pr}_{S'} \circ \Gamma_{f'},$$

the first item shows that f' is proper.

4. It remains only to prove universal closedness. Let $S' \rightarrow S$ be a base change and write

$$h' : X_{S'} \longrightarrow Y_{S'}.$$

The morphism h' is still surjective. If $F \subset Y_{S'}$ is closed, then $(h')^{-1}(F)$ is closed in $X_{S'}$. Since $X_{S'} \rightarrow S'$ is closed,

$$\text{im}((h')^{-1}(F) \rightarrow S')$$

is closed. By surjectivity of h' , this image is exactly the image of F in S' . Thus $Y_{S'} \rightarrow S'$ is closed. The assumptions already give finite type and separatedness.

Lemma V.A.8. *Properness is local on the target: if $Y = \bigcup_i V_i$ is an open covering, then $f : X \rightarrow Y$ is proper if and only if every restriction*

$$f^{-1}(V_i) \longrightarrow V_i$$

is proper.

Proof. Finite type and separatedness are local on the target. For universal closedness, perform an arbitrary base change $Y' \rightarrow Y$. The inverse images of the V_i form an open covering of Y' , and a subset of Y' is closed if and only if its intersection with each member of this covering is closed. Thus the base-changed morphism is closed if and only if all its restrictions over these open subsets are closed. $\square$

V.B The valuative criterion and projective morphisms

Definition V.B.1. A *valuation ring* is an integral domain R with fraction field K such that, for every $a \in K^\times$, either $a \in R$ or $a^{-1} \in R$. Equivalently, the ideals of R are totally ordered by inclusion.

Proposition V.B.2. *Every valuation ring R is local. Its maximal ideal is*

$$\mathfrak{m}_R = \{a \in R \mid a = 0 \text{ or } a^{-1} \notin R\},$$

the set of non-units of R .

Proof. The principal ideals of R are totally ordered. If a, b are non-units, assume after exchanging them that $aR \subset bR$, and write $a = bc$. Then $a + b = b(c + 1)$ is again a non-unit. A multiple of a non-unit is a non-unit as well. Hence the non-units form an ideal, which is necessarily the unique maximal ideal. $\square$

Example V.B.3 (Basic valuation rings). Every field is a valuation ring of itself; this is the trivial example. Every discrete valuation ring (Definition II.D.13) is a valuation ring. Indeed, if R has uniformiser π , every element of $K^\times$ has the form

$$u\pi^n, \quad u \in R^\times, \quad n \in \mathbb{Z},$$

so either it or its inverse belongs to R . In particular,

$$\mathbb{Z}_{(p)} \subset \mathbb{Q}, \quad \mathbb{k}[t]_{(t)} \subset \mathbb{k}(t), \quad \mathbb{k}[[t]] \subset \mathbb{k}((t))$$

are valuation rings. The first two are localisations of principal ideal domains, while the third records the order of vanishing of a Laurent series.

Remark V.B.4 (The geometric picture). The morphism $\text{Spec}(K) \rightarrow \text{Spec}(R)$ identifies $\text{Spec}(K)$ with the generic point of $\text{Spec}(R)$. The ring R is local, so it also has a unique closed point. For a discrete valuation ring these are the only two points; a general valuation ring may have a longer totally ordered chain of prime ideals. The valuative criterion asks whether a map defined at the generic point extends over this whole chain of specialisations.

Theorem V.B.5 (Valuative criterion for properness). *Let $f : X \rightarrow Y$ be a morphism of finite type. Then f is proper if and only if, for every valuation ring R with fraction field K , every commutative diagram*

$$\begin{array}{ccc} \text{Spec}(K) & \longrightarrow & X \\ \downarrow & \nearrow & \downarrow f \\ \text{Spec}(R) & \longrightarrow & Y \end{array}$$

admits a unique dotted arrow.

We use this standard criterion without proof. The existence of the lift expresses universal closedness, while its uniqueness expresses separatedness.

Exercise V.B.6. Let $R = \mathbb{k}[[\pi]]$ and $K = \mathbb{k}((\pi))$.

- i) Show that R is a valuation ring.
- ii) Describe $\text{Spec}(R)$ and identify the morphism $\text{Spec}(K) \rightarrow \text{Spec}(R)$.
- iii) Use the K -point $\pi^{-1} \in \mathbb{A}_{\mathbb{k}}^1(K)$ to see directly from the valuative criterion that $\mathbb{A}_{\mathbb{k}}^1 \rightarrow \text{Spec}(\mathbb{k})$ is not proper.

Solution. i) Every nonzero Laurent series can be written uniquely as

$$\pi^m u, \quad m \in \mathbb{Z},$$

with $u \in R^\times$. If $m \geq 0$, the element belongs to R ; if $m < 0$, its inverse belongs to R .

- ii) The ring R is a discrete valuation ring with maximal ideal (π) . Hence

$$\text{Spec}(R) = \{(0), (\pi)\}.$$

The morphism $\text{Spec}(K) \rightarrow \text{Spec}(R)$ identifies its unique point with the generic point (0) ; the point (π) is closed.

- iii) The element $\pi^{-1} \in K$ defines a morphism

$$\text{Spec}(K) \longrightarrow \mathbb{A}_{\mathbb{k}}^1.$$

An extension to $\text{Spec}(R)$ would correspond to an element of R whose image in K is π^{-1} , which is impossible. Hence the required lift does not exist.

Theorem V.B.7. *For every scheme Y , the projection*

$$\mathbb{P}_Y^n \longrightarrow Y$$

is proper.

Proof. The morphism is of finite type and was proved above to be separated. We check the valuative criterion. Let R be a valuation ring with fraction field K , and consider a diagram

$$\begin{array}{ccc} \mathrm{Spec}(K) & \longrightarrow & \mathbb{P}_Y^n \\ \downarrow & & \downarrow \\ \mathrm{Spec}(R) & \longrightarrow & Y. \end{array}$$

After base change by $\mathrm{Spec}(R) \rightarrow Y$, the upper morphism is a K -point of $\mathbb{P}_R^n$, hence is represented by homogeneous coordinates

$$[a_0 : \cdots : a_n], \quad a_i \in K,$$

not all zero.

The principal fractional ideals $a_i R$ are totally ordered. Choose a_j such that $a_j R$ contains all the $a_i R$. Then

$$a_i/a_j \in R \quad \text{for every } i,$$

and the j -th ratio is 1. Therefore

$$[a_0/a_j : \cdots : a_n/a_j]$$

defines an R -point of $\mathbb{P}_R^n$ extending the original K -point. Uniqueness follows from separatedness. The valuative criterion now implies that $\mathbb{P}_Y^n \rightarrow Y$ is proper. $\square$

Corollary V.B.8 (Projective morphisms are proper). *Let $f : X \rightarrow Y$ factor as*

$$X \hookrightarrow \mathbb{P}_Y^n \longrightarrow Y$$

with the first morphism a closed immersion. Then f is proper.

Proof. Closed immersions are proper, the projection $\mathbb{P}_Y^n \rightarrow Y$ is proper, and proper morphisms are stable under composition. $\square$

V.C Finite morphisms

Definition V.C.1. A morphism $f : X \rightarrow Y$ is *finite* if, for every affine open subset $V = \mathrm{Spec}(A) \subset Y$, the inverse image $f^{-1}(V)$ is affine, say

$$f^{-1}(V) = \mathrm{Spec}(B),$$

and B is a finitely generated A -module.

Proposition V.C.2 (Finite morphisms and projective embeddings). *Let $A \rightarrow B$ be a finite ring homomorphism. Then*

$$\mathrm{Spec}(B) \longrightarrow \mathrm{Spec}(A)$$

is projective.

More generally, every finite morphism is projective locally on its target.

Proof. Choose A -module generators

$$b_0 = 1, b_1, \dots, b_n$$

of B . Give $B[T]$ the grading for which B has degree 0 and T has degree 1. There is a surjective homomorphism of graded A -algebras

$$A[x_0, \dots, x_n] \longrightarrow B[T], \quad x_i \longmapsto b_i T.$$

Indeed, x_0 maps to T , and every element bT^d is an A -linear combination of the elements $b_i T^d = (b_i T) T^{d-1}$.

Taking Proj gives a closed immersion

$$\text{Proj}(B[T]) \hookrightarrow \mathbb{P}_A^n.$$

Since the irrelevant ideal of $B[T]$ is generated by T , the single chart $D_+(T)$ is all of $\text{Proj}(B[T])$, and

$$\text{Proj}(B[T]) \simeq \text{Spec}((B[T][T^{-1}])_0) = \text{Spec}(B).$$

This proves the assertion over an affine base. Applying the construction on affine open subsets of an arbitrary target proves the local statement. $\square$

Remark V.C.3. With the more general definition of a projective morphism using relative projective spectra, every finite morphism is projective globally:

$$X \simeq \text{Proj}_Y(f_* \mathcal{O}_X[T]).$$

With the present convention requiring one global closed immersion into a fixed $\mathbb{P}_Y^n$, the global conclusion follows whenever $f_* \mathcal{O}_X$ is generated by finitely many global sections; it is automatic when Y is affine.

Corollary V.C.4. *Every finite morphism is proper.*

Proof. Properness is local on the target. On every affine open subset of the target, the preceding proposition realizes the finite morphism as a projective morphism, hence as a proper morphism. $\square$

Proposition V.C.5 (Affine proper morphisms). *An affine morphism is proper if and only if it is finite.*

Proof. The implication “finite implies proper” was just proved. The converse is the standard algebraic theorem that an affine morphism of finite type which is universally closed is finite. We use this result without proof. $\square$

Example V.C.6 (The power map). Let $n \geq 1$. The homomorphism

$$\mathbb{k}[t] \longrightarrow \mathbb{k}[u], \quad t \longmapsto u^n$$

makes $\mathbb{k}[u]$ a free $\mathbb{k}[t]$ -module with basis

$$1, u, \dots, u^{n-1}.$$

Hence the induced morphism

$$\mathbb{A}_{\mathbb{k}}^1 \longrightarrow \mathbb{A}_{\mathbb{k}}^1$$

is finite, projective and proper.

Exercise V.C.7. Assume that $\mathbb{k}$ is algebraically closed and that $\text{char}(\mathbb{k}) \nmid n$. For the power map

$$f : \mathbb{A}_{\mathbb{k}}^1 \longrightarrow \mathbb{A}_{\mathbb{k}}^1, \quad u \longmapsto u^n,$$

compute the fibres over 0 and over $a \neq 0$.

Solution. The fibre over $a \in \mathbb{k}$ is

$$\mathrm{Spec}(\mathbb{k}[u]/(u^n - a)).$$

For $a = 0$, this is the non-reduced point

$$\mathrm{Spec}(\mathbb{k}[u]/(u^n)).$$

If $a \neq 0$, the polynomial $u^n - a$ has n distinct roots because $\mathbb{k}$ is algebraically closed and its derivative nu^{n-1} does not vanish at a nonzero root. Hence the fibre is the disjoint union of n reduced $\mathbb{k}$ -rational points.

Example V.C.8 (Normalisation of the cusp). The inclusion

$$\mathbb{k}[t^2, t^3] \subset \mathbb{k}[t]$$

is finite, since $\mathbb{k}[t]$ is generated by $1, t$ as a $\mathbb{k}[t^2, t^3]$ -module. It defines a finite, hence proper, morphism

$$\mathbb{A}_{\mathbb{k}}^1 \longrightarrow \mathrm{Spec}(\mathbb{k}[t^2, t^3]),$$

which is the normalisation morphism of the cuspidal cubic.

Theorem V.C.9 (Global functions on a proper scheme). *Let A be a ring and let X be a proper A -scheme. Then*

$$\Gamma(\mathcal{O}_X)$$

is integral over A . If X is a reduced proper k -scheme of finite type, where k is a field, then $\Gamma(\mathcal{O}_X)$ is a finite-dimensional k -vector space.

Proof. Put

$$B = \Gamma(\mathcal{O}_X)$$

and let $b \in B$. The section b determines an A -morphism

$$g : X \longrightarrow \mathbb{A}_A^1 = \mathrm{Spec}(A[t])$$

corresponding to the homomorphism

$$A[t] \longrightarrow B, \quad t \longmapsto b.$$

Let

$$C = \mathrm{im}(A[t] \rightarrow B)$$

and put $Z = \mathrm{Spec}(C)$. Then Z is a closed subscheme of $\mathbb{A}_A^1$, and g factors as

$$X \xrightarrow{\bar{g}} Z \longrightarrow \mathbb{A}_A^1.$$

The homomorphism $C \rightarrow B$ is injective, so $\bar{g}$ is dominant. Since $X \rightarrow \mathrm{Spec}(A)$ is proper and $Z \rightarrow \mathrm{Spec}(A)$ is separated, the morphism $\bar{g}$ is proper. It is therefore closed. Its image is both dense and closed in Z , hence $\bar{g}$ is surjective.

The scheme Z is separated and of finite type over A . Since it is the surjective image of the proper A -scheme X , the preceding exercise shows that Z is proper over A . But $Z \rightarrow \mathrm{Spec}(A)$ is affine, hence finite by the proposition above. Therefore C is a finitely generated A -module. In particular, the class of t in C , and hence b , is integral over A .

Now assume that X is reduced and of finite type over a field k . Let

$$X_1, \dots, X_r$$

be its irreducible components, endowed with their reduced induced scheme structures. They are proper integral k -schemes. Restriction gives an injective homomorphism

$$\Gamma(\mathcal{O}_X) \longrightarrow \prod_{i=1}^r \Gamma(\mathcal{O}_{X_i}).$$

Indeed, on an affine open subset this is the injectivity of a reduced ring into the product of its quotients by its minimal prime ideals.

For each i , the ring $\Gamma(\mathcal{O}_{X_i})$ embeds into the function field $k(X_i)$. By the first part of the theorem, all its elements are algebraic over k . The algebraic closure k_i of k inside the finitely generated field extension $k(X_i)/k$ is a finite extension of k . Hence

$$\Gamma(\mathcal{O}_{X_i}) \subset k_i$$

is finite-dimensional over k . The product of the k_i is finite-dimensional, and so is its k -subspace $\Gamma(\mathcal{O}_X)$. $\square$

Corollary V.C.10. *Let X be an integral proper k -scheme of finite type. Then $\Gamma(\mathcal{O}_X)$ is a finite field extension of k . In particular, if k is algebraically closed, then*

$$\Gamma(\mathcal{O}_X) = k.$$

Example V.C.11. For every field $\mathbb{k}$ and every $n \geq 1$,

$$\Gamma(\mathcal{O}_{\mathbb{P}_{\mathbb{k}}^n}) = \mathbb{k}.$$

This was computed directly in the preceding chapter; it also follows from properness and integrality.

Exercise V.C.12. Let $\mathbb{k}$ be algebraically closed, let X be an integral proper $\mathbb{k}$ -scheme of finite type, and let $Y = \text{Spec}(A)$ be an affine $\mathbb{k}$ -scheme. Show that every morphism

$$f : X \longrightarrow Y$$

is constant, in the sense that it factors through a $\mathbb{k}$ -rational point of Y .

Solution. By the theorem on morphisms to an affine scheme, the morphism f corresponds to a homomorphism of $\mathbb{k}$ -algebras

$$A \longrightarrow \Gamma(\mathcal{O}_X).$$

Since X is integral and proper and $\mathbb{k}$ is algebraically closed,

$$\Gamma(\mathcal{O}_X) = \mathbb{k}.$$

Thus the homomorphism factors as $A \rightarrow \mathbb{k}$, which is precisely a $\mathbb{k}$ -rational point $\text{Spec}(\mathbb{k}) \rightarrow Y$. Therefore f factors through that point.

Exercise V.C.13. Let $f : X \rightarrow Y$ be proper and let $y \in Y$. Show that the fibre

$$X_y \longrightarrow \text{Spec}(\kappa(y))$$

is proper.

Solution. The fibre morphism is obtained from f by base change along

$$\text{Spec}(\kappa(y)) \longrightarrow Y.$$

Properness is preserved by base change, so X_y is proper over $\kappa(y)$.

Chapter 6

A glimpse towards moduli spaces

The aim of this final chapter is not to develop the general theory of moduli spaces, but to show why the language introduced in this course is useful to prepare the ground for the study of moduli spaces. The first ingredient is just one more way to think of a morphism of schemes

$$X \longrightarrow T.$$

We view this morphism as a family of schemes parametrised by T . Namely, over a point $t \in T$, the element of the family is the fibre X_t .

A *moduli space* is a scheme whose T -points themselves parametrise families of geometric objects parametrised by T . The objects that one wants to parametrise, typically, are clear to the person embarking in the study of their moduli space: these could be compact Riemann surfaces of genus g , or vector bundles of a given rank on the projective plane, etc. Sometimes these objects are to be taken up to some equivalence relation (isomorphism of Riemann surfaces, of vector bundles, etc). So actually one starts by setting the *moduli problem*. This is about defining, for each scheme T (which could be over $\mathbb{Z}$, or over some other base scheme), the geometric objects parametrised by T , just as a set. This typically defines a functor, from schemes to sets. Then the question is: is that functor actually given by a space? Or, slightly more precisely, is that functor represented by a scheme?

I The Hilbert scheme functor

We will discuss briefly the idea of moduli space in one of the simplest situations, namely the Hilbert scheme.

I.A Families and the functor of points

Let us fix a ground field $\mathbb{k}$. For a $\mathbb{k}$ -scheme M , its functor of points is

$$\begin{aligned} h_M : (\mathrm{Sch})_{\mathbb{k}}^{\mathrm{op}} &\rightarrow (\mathrm{Set}) \\ T &\mapsto h_M(T) = \mathrm{Hom}_{(\mathrm{Sch})_{\mathbb{k}}}(T, M). \end{aligned}$$

Rational $\mathbb{k}$ -points correspond to $T = \mathrm{Spec}(\mathbb{k})$. Allowing arbitrary test schemes records more information. For instance, morphisms from

$$\mathrm{Spec}(\mathbb{k}[\varepsilon]/(\varepsilon^2))$$

encode infinitesimal tangent directions, as we have seen. Thus non-reduced test schemes are not an artificial complication: they are precisely what allows a moduli problem to remember first-order deformations.

Imagine now that we start from a functor $\mathcal{F}$. The idea is that this functor is supposed to assign to each scheme T the families of our favourite geometric objects parametrised by T . Thus we start with a functor

$$\mathcal{F} : (\text{Sch})_{\mathbb{k}}^{\text{op}} \rightarrow (\text{Set}).$$

We say that a scheme M represents $\mathcal{F}$ if $\mathcal{F} \simeq h_M$. This means that there are natural bijections

$$\text{Hom}_{(\text{Sch})_{\mathbb{k}}} (T, M) \simeq \mathcal{F}(T)$$

for every $\mathbb{k}$ -scheme T . This way, if $\mathcal{F}(T)$ is the set of families of objects parametrised by T , then a morphism $T \rightarrow M$ is one family of objects parametrised by T . Beware that, under this formalism, a morphism $f : T \rightarrow T'$ pulls a T' -family back to a T -family through $\mathcal{F}(f) : \mathcal{F}(T') \rightarrow \mathcal{F}(T)$. If the original family is represented by $\alpha' : T' \rightarrow M$, then the pull-back family is represented by $\alpha' \circ f : T \rightarrow M$.

It is also important to consider $T = M$. Under the representing bijection, the identity morphism of M gives a distinguished tautological element $\mathcal{U} \in \mathcal{F}(M)$. This element has the property that any family comes by pull-back of it. More precisely, for any $\mathbb{k}$ -scheme T , and for any $\mathcal{V} \in \mathcal{F}(T)$ we have a corresponding *moduli map* $\alpha_{\mathcal{V}} \in \text{Hom}_{(\text{Sch})_{\mathbb{k}}} (T, M)$, given by $\mathcal{V}$ under the bijection $\text{Hom}_{(\text{Sch})_{\mathbb{k}}} (T, M) \simeq \mathcal{F}(T)$. Then $\mathcal{F}(\alpha_{\mathcal{V}}) : \mathcal{F}(M) \rightarrow \mathcal{F}(T)$ sends $\mathcal{U}$ precisely to $\mathcal{V}$. So the slogan is: a T -family determines a moduli map $T \rightarrow M$ to the moduli space, in such a way that the family is the pull-back of the tautological family, which is therefore at the source of all families. Of course, a $\mathbb{k}$ -rational point of M is just a single geometric object, and all geometric objects (over $\mathbb{k}$) individually are in bijection with rational $\mathbb{k}$ -points. If one such object is defined over a field extension κ of $\mathbb{k}$, then it will be a κ -rational point of M .

I.B The Hilbert scheme functor of n points of the plane

For a systematic treatment of Hilbert schemes of points on surfaces and their matrix description, see [Nak99, Lectures 1–2]. See also [EH00, Chapter VI] for the general moduli philosophy.

Let T be a $\mathbb{k}$ -scheme and write

$$\mathbb{A}_T^2 = \mathbb{A}_{\mathbb{k}}^2 \times_{\text{Spec}(\mathbb{k})} T.$$

A family of length- n subschemes of the affine plane parametrised by T is a closed subscheme

$$Z \subset \mathbb{A}_T^2$$

such that the projection $p : Z \rightarrow T$ is finite and, locally on T , its coordinate algebra admits a basis of n elements over the coordinate ring of the base. More precisely, on an affine open subset $U = \text{Spec}(R) \subset T$, the family is given by an ideal

$$I \subset R[x, y]$$

such that $R[x, y]/I$ is a finite locally free R -module of rank n . Equivalently, after replacing U by a smaller affine open subset, this module is free of rank n . This condition is often expressed by saying that $p_* \mathcal{O}_Z$ is locally free of rank n . It implies that every fibre

$$Z_t = Z \times_T \text{Spec}(\kappa(t))$$

has coordinate ring of dimension n over $\kappa(t)$, hence length n . So we let $\mathcal{F}(T)$ be the set of these families. Again, if $T = \text{Spec}(R)$, then

$$\mathcal{F}(\text{Spec}(R)) = \{I \subset R[x, y] \mid R[x, y]/I \text{ is finite projective of constant rank } n.\}$$

Remark I.B.1. It is tempting to require only that every fibre have length n . Over reduced bases, and under suitable finiteness hypotheses, this often captures the same geometric idea. Over an arbitrary non-reduced base it is not sufficient: infinitesimal torsion may be invisible on all residue-field fibres. The local-basis condition is the down-to-earth form of the flatness requirement appearing in the general Hilbert functor. For projective families, its numerical shadow is the constancy of the Hilbert polynomial; for a zero-dimensional fibre of length n , that polynomial is the constant polynomial n .

Definition I.B.2. The *Hilbert functor of n points on the affine plane* assigns to a $\mathbb{k}$ -scheme T the set of closed subschemes $Z \subset \mathbb{A}_T^2$ satisfying the preceding condition. Pull-back along $T' \rightarrow T$ is given by

$$Z_{T'} = Z \times_T T' \subset \mathbb{A}_{T'}^2.$$

Theorem I.B.3 (Representability, stated without proof). *There exists an irreducible quasi-projective $\mathbb{k}$ -scheme*

$$\mathrm{Hilb}^n(\mathbb{A}_{\mathbb{k}}^2)$$

of dimension $2n$ representing the Hilbert functor of n points on $\mathbb{A}_{\mathbb{k}}^2$. It is smooth; the Barth–Nakajima tangent calculation below explains this last assertion. Thus morphisms

$$T \longrightarrow \mathrm{Hilb}^n(\mathbb{A}_{\mathbb{k}}^2)$$

are naturally the same as families of length- n subschemes of $\mathbb{A}_{\mathbb{k}}^2$ parametrised by T .

For $T = \mathrm{Spec}(\mathbb{k})$, a point of the Hilbert scheme is simply an ideal $I \subset \mathbb{k}[x, y]$ such that

$$\dim_{\mathbb{k}} \mathbb{k}[x, y]/I = n.$$

The open subset corresponding to n distinct reduced points is the unordered configuration space of n distinct points. Its boundary records collisions, and the resulting subschemes are usually non-reduced.

Example I.B.4. For $n = 1$, every colength-one ideal is a maximal ideal $(x - a, y - b)$, so

$$\mathrm{Hilb}^1(\mathbb{A}_{\mathbb{k}}^2) \simeq \mathbb{A}_{\mathbb{k}}^2.$$

For $n = 2$, the ideal

$$(y, x^2)$$

defines a non-reduced length-two subscheme supported at the origin. It may be viewed as two points which have collided while retaining an infinitesimal direction.

I.C Commuting matrices and a cyclic vector

Let $I \subset \mathbb{k}[x, y]$ have colength n and put

$$V = \mathbb{k}[x, y]/I.$$

Multiplication by x and y gives commuting endomorphisms

$$X, Y \in \mathrm{End}_{\mathbb{k}}(V), \quad [X, Y] = 0.$$

The class $v = 1 \bmod I$ is cyclic:

$$V = \mathbb{k}[X, Y]v.$$

Conversely, let V be an n -dimensional vector space, let $X, Y \in \mathrm{End}(V)$ commute, and let $v \in V$ be cyclic. Then

$$\mathbb{k}[x, y] \longrightarrow V, \quad f \longmapsto f(X, Y)v$$

is surjective, and its kernel is an ideal of colength n .

Proposition I.C.1 (Barth–Nakajima description). *The $\mathbb{k}$ -points of $\text{Hilb}^n(\mathbb{A}_{\mathbb{k}}^2)$ are in bijection with isomorphism classes of triples (X, Y, v) where*

$$X, Y \in \text{End}(\mathbb{k}^n), \quad [X, Y] = 0,$$

and $v \in \mathbb{k}^n$ is cyclic. Two triples are identified when they differ by a simultaneous change of basis by an element of $\text{GL}_n(\mathbb{k})$.

Proof. The two constructions just described are inverse. Starting from I , the map $f \mapsto f(X, Y)v$ is the quotient map $\mathbb{k}[x, y] \rightarrow \mathbb{k}[x, y]/I$, so its kernel is I . Starting from a cyclic triple, the induced identification $\mathbb{k}[x, y]/I \simeq V$ carries multiplication by x and y back to X and Y , and carries 1 to v .

It remains to account for the choice of a basis of V . Two identifications $V \simeq \mathbb{k}^n$ differ by an element of $\text{GL}_n(\mathbb{k})$, which simultaneously conjugates X and Y and sends v to the new coordinate vector. Conversely, such a change of basis does not change the kernel of $f \mapsto f(X, Y)v$. This proves the asserted bijection. $\square$

In families, the vector space V is replaced by a rank- n vector bundle $\mathcal{E}$ on T , the matrices by commuting endomorphisms $X, Y \in \text{End}(\mathcal{E})$, and the cyclic vector by a section $v \in \Gamma(T, \mathcal{E})$ which generates every fibre under X and Y . This realises the Hilbert scheme as the quotient of the cyclic locus

$$\{(X, Y, v) \mid [X, Y] = 0, \mathbb{k}[X, Y]v = \mathbb{k}^n\}$$

by GL_n . This is the simplest doorway to framed quiver representations and Nakajima quiver varieties.

Here is what is hidden in the word “quotient”. By Cayley–Hamilton, cyclicity can be tested using the finitely many vectors

$$X^a Y^b v, \quad 0 \leq a, b < n.$$

The condition that they span is the non-vanishing of suitable minors, hence defines an open locus. The GL_n -action on this locus is free: an endomorphism commuting with X and Y and fixing v fixes every $f(X, Y)v$, hence is the identity. A family with a rank- n bundle $\mathcal{E}$ becomes a family of matrices after passing to its frame bundle; changing the frame is precisely the GL_n -action. The geometric quotient therefore descends framed matrix data to families of ideals. The existence of this quasi-projective quotient is the non-elementary invariant-theoretic step in the Barth–Nakajima construction.

Example I.C.2. For $I = (y, x^2)$, use the basis $(1, x)$ of $\mathbb{k}[x, y]/I$. Multiplication gives

$$X = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \quad Y = 0, \quad v = \begin{pmatrix} 1 \\ 0 \end{pmatrix}.$$

The nilpotent Jordan block records the infinitesimal direction of the collision.

I.D Dual numbers and the Barth–Nakajima tangent calculation

Put $A = \mathbb{k}[x, y]$ and $D = \mathbb{k}[\varepsilon]/(\varepsilon^2)$. A tangent vector to $\text{Hilb}^n(\mathbb{A}_{\mathbb{k}}^2)$ at the point $[I]$ is a D -valued point specialising to $[I]$. In terms of ideals, it is an ideal

$$\tilde{I} \subset A \otimes_{\mathbb{k}} D = A[\varepsilon]/(\varepsilon^2)$$

such that $(A \otimes D)/\tilde{I}$ is free of rank n over D and $\tilde{I} \bmod \varepsilon = I$.

Proposition I.D.1 (First-order deformations of an ideal). *There is a natural isomorphism*

$$T_{[I]} \text{Hilb}^n(\mathbb{A}_{\mathbb{k}}^2) \simeq \text{Hom}_A(I, A/I).$$

More explicitly, if $f + \varepsilon g \in \tilde{I}$ lifts $f \in I$, the corresponding homomorphism is

$$\varphi_{\tilde{I}}(f) = g \bmod I.$$

Conversely, $\varphi \in \text{Hom}_A(I, A/I)$ determines

$$\tilde{I}_\varphi = \{f + \varepsilon g \mid f \in I, g \bmod I = \varphi(f)\}.$$

Proof. The local-freeness condition permits every $f \in I$ to be lifted to an element $f + \varepsilon g$ of $\tilde{I}$. If two such lifts are chosen, then $\varepsilon(g - g') \in \tilde{I}$. In the free D -module $Q = (A \otimes D)/\tilde{I}$, the kernel of multiplication by ε is εQ . Reducing modulo ε therefore shows that $g - g' \in I$. Thus $g \bmod I$ is well defined.

Addition of lifts proves additivity. Multiplying a lift by $a \in A$ shows that

$$\varphi_{\tilde{I}}(af) = a\varphi_{\tilde{I}}(f),$$

so $\varphi_{\tilde{I}}$ is A -linear. Conversely, A -linearity makes $\tilde{I}_\varphi$ an ideal. Choosing a $\mathbb{k}$ -linear section of $A \rightarrow A/I$ gives unique normal forms modulo $\tilde{I}_\varphi$ and identifies the quotient, as a D -module, with

$$D \otimes_{\mathbb{k}} (A/I).$$

It is therefore free of rank n . The two constructions are inverse. $\square$

We can now see the number $2n$ directly from the commuting-matrix description. Let (X, Y, v) be the cyclic triple corresponding to I and put $V = A/I$. A first-order deformation has the form

$$(X + \varepsilon P, Y + \varepsilon Q, v + \varepsilon w).$$

The commutation relation modulo ε^2 is

$$[P, Y] + [X, Q] = 0.$$

An infinitesimal change of basis $1 + \varepsilon a$ changes (P, Q, w) by

$$([a, X], [a, Y], av).$$

Consequently the tangent space is the middle cohomology of

$$\text{End}(V) \xrightarrow{d_0} \text{End}(V) \oplus \text{End}(V) \oplus V \xrightarrow{d_1} \text{End}(V),$$

where

$$d_0(a) = ([a, X], [a, Y], av), \quad d_1(P, Q, w) = [P, Y] + [X, Q].$$

Proposition I.D.2 (Barth–Nakajima tangent calculation). *For every colength- n ideal $I \subset \mathbb{k}[x, y]$,*

$$\dim_{\mathbb{k}} \text{Hom}_A(I, A/I) = 2n.$$

Thus every geometric tangent space of $\text{Hilb}^n(\mathbb{A}_{\mathbb{k}}^2)$ has dimension $2n$.

Proof. Cyclicity makes d_0 injective: if a commutes with X and Y and $av = 0$, then a vanishes on every vector $f(X, Y)v$, hence on all of V .

Use the non-degenerate trace pairing on $\text{End}(V)$. The orthogonal complement of $\text{im}(d_1)$ consists of the endomorphisms commuting with both X and Y . Evaluation at the cyclic vector gives an isomorphism

$$\{c \in \text{End}(V) \mid [c, X] = [c, Y] = 0\} \longrightarrow V, \quad c \longmapsto cv.$$

It is injective by cyclicity; it is surjective because every $u \in V$ has the form $u = f(X, Y)v$, and the endomorphism $f(X, Y)$ commutes with X and Y . Hence the common centraliser has dimension n , and

$$\text{rank}(d_1) = n^2 - n.$$

It follows that

$$\begin{aligned} \dim(\ker(d_1)/\text{im}(d_0)) &= (2n^2 + n) - (n^2 - n) - n^2 \\ &= 2n. \end{aligned}$$

The preceding proposition identifies this quotient with $\text{Hom}_A(I, A/I)$. Since the representing scheme is irreducible of dimension $2n$, its geometric tangent spaces all have the expected dimension. By the regularity and tangent-space criterion of Theorem II.4, applied after extending the ground field to an algebraic closure, the Hilbert scheme is regular there and hence smooth over $\mathbb{k}$. $\square$

Example I.D.3 (The double point). For $I = (y, x^2)$, the quotient A/I has basis $(1, x)$. The ideal I is generated by y and x^2 , with syzygy $(x^2, -y)$. This relation becomes zero in A/I , so the images of both generators under a homomorphism $I \rightarrow A/I$ may be chosen freely. Hence

$$T_{[I]} \text{Hilb}^2(\mathbb{A}_{\mathbb{k}}^2) \simeq (A/I)^{\oplus 2}$$

has dimension 4, just as it does at a pair of distinct points.

I.E Exercises

Exercise I.E.1. Assume that $\mathbb{k}$ is algebraically closed.

- i) Show that every ideal $I \subset \mathbb{k}[x]$ such that $\dim_{\mathbb{k}} \mathbb{k}[x]/I = n$ is generated by a unique monic polynomial

$$f(x) = x^n + a_1x^{n-1} + \cdots + a_n.$$

- ii) Deduce a bijection on $\mathbb{k}$ -points

$$\text{Hilb}^n(\mathbb{A}_{\mathbb{k}}^1)(\mathbb{k}) \simeq \mathbb{A}_{\mathbb{k}}^n(\mathbb{k}).$$

- iii) Explain how the factorisation

$$f(x) = \prod_i (x - \lambda_i)^{m_i}$$

records the support and multiplicities of the corresponding subscheme.

- iv) State the corresponding functorial result for a base $T = \text{Spec}(R)$ using monic polynomials in $R[x]$.

Solution. The ring $\mathbb{k}[x]$ is a principal ideal domain, so $I = (f)$ for a unique monic polynomial f . The quotient has dimension $\deg(f)$ with basis $1, x, \dots, x^{\deg(f)-1}$, hence $\deg(f) = n$. The coefficients $(a_1, \dots, a_n)$ give the required point of $\mathbb{A}_{\mathbb{k}}^n$. Over an algebraically closed field, the Chinese remainder theorem identifies the factors $(x - \lambda_i)^{m_i}$ with length- m_i subschemes supported at λ_i . Functorially, a monic polynomial

$$x^n + a_1x^{n-1} + \cdots + a_n \in R[x]$$

defines a quotient free over R with basis $1, x, \dots, x^{n-1}$; its coefficients define a morphism $T \rightarrow \mathbb{A}_{\mathbb{k}}^n$. This yields $\text{Hilb}^n(\mathbb{A}_{\mathbb{k}}^1) \simeq \mathbb{A}_{\mathbb{k}}^n$.

Exercise I.E.2. Let (X, Y, v) be a commuting cyclic triple on an n -dimensional vector space V . Show directly that

$$I = \{f \in \mathbb{k}[x, y] \mid f(X, Y)v = 0\}$$

is an ideal and that $\mathbb{k}[x, y]/I \simeq V$.

Solution. If $f \in I$ and $g \in \mathbb{k}[x, y]$, then commutativity gives

$$(gf)(X, Y)v = g(X, Y)f(X, Y)v = 0,$$

so $gf \in I$. The map $\mathbb{k}[x, y] \rightarrow V$, $f \mapsto f(X, Y)v$, is surjective because v is cyclic, and its kernel is I . The first isomorphism theorem gives $\mathbb{k}[x, y]/I \simeq V$.

Exercise I.E.3. Assume that $\mathbb{k}$ is algebraically closed. Describe the triples associated with

- i) two distinct points (a_1, b_1) and (a_2, b_2) of $\mathbb{A}_{\mathbb{k}}^2$;
- ii) the double point defined by (y, x^2) .

Solution. For two distinct points, take

$$X = \begin{pmatrix} a_1 & 0 \\ 0 & a_2 \end{pmatrix}, \quad Y = \begin{pmatrix} b_1 & 0 \\ 0 & b_2 \end{pmatrix}, \quad v = \begin{pmatrix} 1 \\ 1 \end{pmatrix}.$$

The vector v is cyclic because the two joint eigenvalues are distinct. For (y, x^2) , use the basis $(1, x)$ and obtain

$$X = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \quad Y = 0, \quad v = \begin{pmatrix} 1 \\ 0 \end{pmatrix}.$$

II What about more general moduli problems?

II.A Hilbert schemes of points on curves, surfaces and threefolds

The geometry of Hilbert schemes of points changes dramatically with the dimension of the ambient variety.

- On the affine line, every colength- n ideal is principal, and

$$\mathrm{Hilb}^n(\mathbb{A}_{\mathbb{k}}^1) \simeq \mathbb{A}_{\mathbb{k}}^n.$$

Equivalently, a length- n subscheme is determined by one monic polynomial of degree n , with repeated roots recording non-reduced points.

- On a smooth surface, the Hilbert scheme of n points is smooth and irreducible of dimension $2n$; this is the world studied in [Nak99].
- In dimension three, the geometry is much wilder. The scheme $\mathrm{Hilb}^n(\mathbb{A}_{\mathbb{k}}^3)$ is reducible for n sufficiently large. Classical constructions of Iarrobino already produce additional components for length 78; see [Iar72]. Thus the pleasant surface theorem is genuinely special to dimension two.

Remark II.A.1 (Partitions enter geometry). Over $\mathbb{C}$, the torus $(\mathbb{C}^\times)^2$ acts on $\mathbb{A}^2$ by scaling the two coordinates and hence acts on every $\mathrm{Hilb}^n(\mathbb{A}^2)$. Its fixed points are the monomial ideals of colength n . The monomials outside such an ideal form a Young diagram, so the fixed points are indexed by partitions $\lambda \vdash n$. A compatible cell decomposition then gives

$$\chi(\mathrm{Hilb}^n(\mathbb{C}^2)) = p(n),$$

where $p(n)$ is the number of partitions of n . Euler's identity packages all these Euler characteristics into

$$\sum_{n \geq 0} \chi(\mathrm{Hilb}^n(\mathbb{C}^2)) q^n = \sum_{n \geq 0} p(n) q^n = \prod_{m \geq 1} \frac{1}{1 - q^m}.$$

This elementary-looking product is closely related to the Dedekind eta function. Refined versions of the same generating series lead from Hilbert schemes to representation theory, modular forms and mathematical physics.

II.B Beyond Hilbert schemes of points

Can one construct schemes parametrising algebraic curves, vector bundles, coherent sheaves, or morphisms between schemes? Often the answer is yes, although the resulting moduli spaces may be singular, reducible, or even need to be replaced by more general objects such as algebraic stacks. The language developed in this course – valued points, families, fibres, nilpotents and dimension – is the natural starting point for these constructions.

Bibliography

- [EH00] DAVID EISENBUD AND JOE HARRIS, *The geometry of schemes*, Graduate Texts in Mathematics, vol. 197, Springer-Verlag, New York, 2000.
- [Har77] ROBIN HARTSHORNE, *Algebraic geometry*, Graduate Texts in Mathematics, vol. 52, Springer-Verlag, New York-Heidelberg, 1977.
- [Hol12] AUDUN HOLME, *A royal road to algebraic geometry*, Springer, Heidelberg, 2012.
- [Hul03] KLAUS HULEK, *Elementary algebraic geometry*, Student Mathematical Library, vol. 20, American Mathematical Society, Providence, RI, 2003, Translated from the 2000 German original by Helena Verrill.
- [Iar72] ANTHONY IARROBINO, *Reducibility of the family of 0-dimensional schemes on a variety*, *Inventiones Mathematicae* **15** (1972), 72–77.
- [Liu02] QING LIU, *Algebraic geometry and arithmetic curves*, Oxford Graduate Texts in Mathematics, vol. 6, Oxford University Press, Oxford, 2002, Translated from the French by Reinie Ern , Oxford Science Publications.
- [Nak99] HIRAKU NAKAJIMA, *Lectures on Hilbert schemes of points on surfaces*, University Lecture Series, vol. 18, American Mathematical Society, Providence, RI, 1999.