

RH: Interactive cues & phenology

## How interactive effects of temperature and photoperiod shape plant phenology responses to warming

E. M. WOLKOVICH<sup>1,2</sup>, C. J. CHAMBERLAIN<sup>1,2</sup>, D. M. BUONAIUTO<sup>1,2</sup>, A. K. ETTINGER<sup>3</sup>  
& I. MORALES-CASTILLA<sup>5</sup>

<sup>1</sup> *Forest & Conservation Sciences, Faculty of Forestry, University of British Columbia, 2424 Main Mall, Vancouver, BC V6T 1Z4*

<sup>2</sup> *Arnold Arboretum of Harvard University, 1300 Centre Street, Boston, Massachusetts, 02131, USA*

<sup>3</sup> *Organismic & Evolutionary Biology, Harvard University, 26 Oxford Street, Cambridge, Massachusetts, 02138, USA*

<sup>4</sup> *The Nature Conservancy, 74 Wall Street, Seattle, Washington USA*

<sup>5</sup> *Global Change Ecology and Evolution Group, Department of Life Sciences, University of Alcalá, Alcalá de Henares 28805, Spain*

**Corresponding author:** Lizzie, see <sup>3</sup> above ; E-mail: e.wolkovich@ubc.ca

### Abstract

Climate change has advanced plant phenology globally 4-6 days per °C on average, with some species shifting much more. Such shifts have been some of the most reported and predictable biological impacts of rising temperatures. Yet as climate change has marched on, phenological shifts have appeared muted over recent decades—failing to match simple predictions of an advancing spring with continued warming. The main hypothesis for these changing trends is that other cues of spring phenology—long-documented in lab environments—are playing a greater role in natural environments due to climate change. Here we argue that accurately linking shifts observed in long-term data to underlying phenological cues requires greater integration of controlled environment experiments with long-term data and insights from physiology. We highlight how seven decades of research in controlled environments can improve predictions for when, where, and how the interactive effects of other cues will impact simple linear predictions. At the same time a new generation of lab experiments could transform our fundamental understanding of phenology and improve forecasts of shifting phenology across different regions for decades to come.

**Keywords:** phenology, climate change, spring warming, chilling, forcing, daylength, photoperiod, non-linear responses, leafout, budburst

# 1 Main text

Shifts in spring plant phenology are one of the most reported and predictable changes with climate change. Decades of research have documented advancing budburst, leafout and flowering with climate change (Delpierre *et al.*, 2009; Yu *et al.*, 2010; Ellwood *et al.*, 2013; Jochner *et al.*, 2013; Hereford *et al.*, 2017), especially in temperate systems where long-term records highlight how humans have altered the timing of spring (Schwartz, 1997; Menzel *et al.*, 2003, 2006). Recently, however, these advances have appeared to slow (Fu *et al.*, 2015) or even reverse (Yu *et al.*, 2010)—failing to match simple predictions of an advancing spring with continued warming (Ellwood *et al.*, 2013). The main hypothesis for this failure is that spring warming—which most observational studies focus on—is no longer the only environmental cue that matters to predicting responses to warming (Chuine *et al.*, 2016; Gauzere *et al.*, 2019).

Despite the strong focus of spring phenology research on spring temperatures, increasing evidence suggests a more complicated underlying physiology for most plant species (e.g., Zohner *et al.*, 2016; Gauzere *et al.*, 2019; Ettinger *et al.*, 2020). For many species three major cues underlie spring phenology: forcing (warm temperatures, generally occurring in the late winter and early spring), chilling (cool temperatures, generally occurring in the fall and winter), and photoperiod (daylength).

Together, forcing, chilling and photoperiod may produce non-linear responses that many current methods do not predict, but observational studies have reported recently (Fu *et al.*, 2015). Predicting these non-linearities is a common goal in plant phenology research today (Güsewell *et al.*, 2017; Martinez-Luscher *et al.*, 2017; Gauzere *et al.*, 2019; Chen *et al.*, 2019; Keenan *et al.*, 2020), but has been slowed by data gaps and the underlying complexity of spring phenology.

The first step towards improved phenological predictions is robust measurements of chilling, forcing and photoperiod cues. Recently, efforts have focused on estimating these cues from long-term observational data (Luedeling *et al.*, 2009, 2013). Yet observational studies may struggle to robustly estimate any of the three major cues due to two types of statistical issues. First, in most observational data these cues correlated: forcing increases alongside longer photoperiods (Elmendorf & Ettinger, 2020), and changes to chilling and photoperiod cues with warming predict the same response—later spring phenology. Approaches that attempt to disentangle some of these correlations, such as leveraging trends across elevation or latitude, may run afoul of other correlations (Tansey *et al.*, 2017). Second, most observational studies focus on linear models of each cue, often without interactions between cues (Visser & Holleman, 2001; Polgar *et al.*, 2014; Asse *et al.*, 2018).

In contrast to the limitations of observational studies, one method is designed to measure the complexity of cues: controlled environment (e.g., growth chamber) studies. Such studies have been conducted for over 70 years and are designed to understand nonlinear responses to warming. In contrast to observational studies, controlled environment studies can manipulate all three cues and can tease out interactions between cues by experimentally decoupling them.

Despite the prevalence of controlled environment studies on spring phenological cues, they have rarely been reviewed. Perhaps more surprisingly, they are often poorly integrated into the current phenological literature on climate change, including in debates where they are critical (e.g., Fu *et al.*, 2015; Richardson *et al.*, 2018).

Here, we show how harnessing decades of controlled environment studies could transform our fundamental understanding of phenology and advance forecasting. We begin by outlining how the three major phenological cues—forcing, chilling and photoperiod—can produce non-linearities and how they will shift in coming decades with anthropogenic climate change. We then synthesize controlled environment studies to quantify how much of the cue-space (i.e., the possible range of each cue and interactions across cues) has been studied and how treatments compare to shifts in cues caused by climate change. Based on this, we outline a cross-disciplinary path forward where advances in physiology, greater integration of controlled environment studies with forecasting, and multi-species modeling can yield more robust predictions.

Given our aim to improve understanding of current trends and forecasts we focus on early vegetative phases (budburst and leafout), which are critical to plant growth and carbon storage, and which have shifted most with climate change (Cleland *et al.*, 2007; IPCC, 2014). Our review, therefore, concentrates on woody species phenology. Most of our conclusions and suggested approaches, however, could be adapted to non-woody species and other phenophases with similar underlying interactive cues.

## 1.1 How do phenological cues produce non-linear responses?

Forcing, chilling and photoperiod cues together determine the transition from dormancy to growth (usually first observed as budburst) each year for most temperate woody species (Chuine & Regniere, 2017; Ettinger *et al.*, 2020). Forcing and chilling cues are generally understood to be accumulation processes, where plants integrate chilling and forcing experienced over time to meet a threshold value at which they can budburst, leafout or flower (Chuine, 2000). In contrast, photoperiod is generally not considered an integral cue, but one evaluated daily (Singh *et al.*, 2017). In practice, these two types of processes are often abstracted into the effects of average experienced values (mean temperature or daylength), either over some temporal window in long-term data (e.g., Wolkovich *et al.*, 2012a; Fu *et al.*, 2015), or through controlled environments that hold temperatures and light regimes more constant (e.g., Worrall & Mergen, 1967; Heide, 1993a,b; Skuterud & Dietrichson, 1994).

Controlled environment studies show two major ways these cues can produce non-linear responses—each cue alone, or through interactions between cues (Fig. 1). Alone, each cue may be linear in the mid-range of values, while extremely high or low values of some cues may produce threshold responses (Fig. 1B). For example, at very low photoperiod (short days) plants often will budburst erratically (Heide, 1993a; Partanen *et al.*, 1998; Singh *et al.*, 2017; Rinne *et al.*, 2018), while at sufficiently long photoperiods maximum growth may occur, meaning photoperiods longer than some threshold will have no additional effect on budburst timing (e.g., Major, 1980). Similarly, extremely cool or very hot temperatures may limit forcing as plant developmental processes slow

(Parent & Tardieu, 2012). Such extreme values of cues, however, are likely uncommon in most natural environmental regimes.

More commonly, the interaction of cues produces non-linearities. For example, multiple studies now show that the threshold of forcing needed for budburst depends on the sum of chilling over the fall and winter and also by the photoperiod experienced in the spring (e.g., Zohner & Renner, 2014; Flynn & Wolkovich, 2018). Higher forcing is generally needed given lower chilling (Fig. 2) and shorter photoperiods (Basler & Körner, 2014; Fu *et al.*, 2019). This yields a sub-additive interaction of forcing x chilling and forcing x photoperiod, where high values of both cues together produce a more muted response than would be expected from examining either cue alone. This interaction produces a non-linearity when values of cues are correlated over time or space—for example, chilling declines correlated with greater forcing (Fig. 1A)—and may be critical to accurate forecasts with climate change (Fig. 1C-F).

## 1.2 How will chilling, forcing and photoperiod shift with climate change?

Translating when or if these non-linearities may be triggered by warming first requires understanding how climate change alters each of the three cues. Shifts in cues will vary substantially across space and time (Burrows *et al.*, 2011). Most notably to date, warming increases the forcing plants experience each day, with more rapid shifts—and thus greater shifts—at higher elevations and in the arctic (IPCC, 2014). Daily temperature minima (generally night-time temperatures) have warmed, and will continue to warm, more than maxima, making efforts to understand whether plants accumulate temperatures differently in the night or day critical (Prasad *et al.*, 2008; Shen *et al.*, 2018).

Warming across seasons also varies (Alexander *et al.*, 2006), meaning warming’s impact on forcing (generally accumulating in the late winter and spring) may not be equivalent to its impact on chilling (generally accumulating in the late fall through winter). Warming should translate into important shifts in chilling, which long-term observational studies have repeatedly suggested may already be occurring (Fu *et al.*, 2015; Piao *et al.*, 2017).

Research to date suggests chilling only accumulates in a certain range of temperatures with low (e.g.,  $<0^{\circ}\text{C}$ ) temperatures generally not contributing to accumulated chilling and higher temperatures (e.g.,  $>12^{\circ}\text{C}$ ) potentially decreasing previously accumulated chilling (see Fig. 3 and Richardson, 1974; Fishman *et al.*, 1987). Long-term studies generally focus on the warmer part of this chilling accumulation curve, suggesting that chilling should decrease with warming (Fu *et al.*, 2015; Piao *et al.*, 2017; Gauzere *et al.*, 2019). However, considering the cooler part of this curve, chilling could also increase with warming, which would yield earlier budburst, potentially far earlier than last frost dates (Guy, 2014). Thus, both increases and decreases of chilling are potential outcomes of warming (Fig. 3).

Shifts in chilling and forcing with warming have been studied far more than shifts in photoperiod (but see Saikkonen *et al.*, 2012; Way & Montgomery, 2015). While an environment’s photope-

riod does not shift with climate change, the relevant photoperiod a plant experiences at critical physiological points may change dramatically with warming. In particular, increases in chilling and/or forcing, which could alone produce much earlier budburst, may be offset by short photoperiods that delay budburst (Gauzere *et al.*, 2019). Similarly, long photoperiods can lead to budburst sooner than predicted solely by low chilling or forcing conditions (Nienstaedt, 1966; Myking & Heide, 1995; Partanen *et al.*, 1998).

These shifts—in forcing, chilling and photoperiod experienced near the time of an event—can produce non-linearities when they push a single cue across a critical threshold or inflection point (see threshold response in Fig. 1B). For example, if some species have a critical photoperiod for budburst and warming leads to forcing cues being met before the critical threshold, then we would expect incomplete or highly delayed budburst (Singh *et al.*, 2017; Rinne *et al.*, 2018). Alternatively, the threshold could be crossed in the opposite direction. For example, if pre-climate change conditions generally caused budburst to occur at the extreme values of some cues, but climate change has pushed budburst into values where responses are more linear. This is the mechanism often suggested for declining responses to warming in some temperate trees (Fu *et al.*, 2015; Piao *et al.*, 2017; Gauzere *et al.*, 2019), specifically that plants previously accumulated sufficient chilling such that forcing was the dominant cue—whereas warming has now reduced chilling such that more forcing is needed for budburst (producing an overall muted effect when estimated as change in days per °C, see Fu *et al.*, 2015, for one example). As this example highlights, however, changes in a single cue are unlikely to occur without additional effects on other cues.

We expect most non-linearities from climate change will come from the effects of interactive cues, either where changes in one cue trigger shifts in another cue or due to covarying shifts in cues (Fig. 1). While simple linear interactions between cues may not alone produce non-linearities, they quickly become non-linear when changes occur together (Fig. 1C-F). Predicting these non-linearities, however, requires a refined understanding of the interaction between cues and whether there are critical inflection points that may be crossed with continued warming. These complexities highlight how difficult predictions may be without careful efforts to tease out how each cue works alone and interactively.

### 1.3 Forecasting non-linear responses: Do we have the necessary data?

Controlled environment studies can help predict non-linear responses by allowing researchers to examine the effects of one cue with the others held constant, and examine interactive effects—given the appropriate study design. Such experiments may be especially useful for forecasting if they contain enough variation in treatments to capture precisely where non-linearities occur, and are designed across a range of levels relevant to current versus future conditions (Shen *et al.*, 2015). Indeed, one of the major advantages of experiments is that they allow treatments outside of the historical range of a species’ (or region’s) climate—an option observational data cannot provide.

To understand how valuable currently available data may be to forecasting, we reviewed controlled environment studies over the last seven decades, quantifying the range of treatments and how they compare to current and future conditions. We note that these studies were rarely conducted for climate change research, and most often done for reasons of fundamental or applied science (e.g., horticulture or forestry). Yet they are some of the best available data for how plants respond to the environment and thus a critical resource for climate change research today.

#### *How studies and their experimental treatments vary globally*

Controlled environment studies have been conducted with 226 woody species across the globe, with the majority of papers reporting research occurring in Europe (54 of 84 papers; and 93 of 136 studies across papers; a study is a unique experiment within a paper), followed by North America (22 papers and 32 studies, Fig. S1, Table S1). Most studies manipulate one cue though studies of two or three cues have occurred in almost every decade (Fig. S2). Across study designs, chilling was the most commonly studied cue (69% of 117 studies that manipulated forcing, chilling and/or photoperiod), followed by forcing and photoperiod (43% and 40%, respectively).

The levels of cues (e.g., 8 or 10 hours daylength) varied across latitude with a general trend toward examining more extreme values at higher latitudes (see Fig. S3). This trend appears related to latitudinal clines in the environment—higher latitudes experience colder temperatures and longer photoperiods, and see similar shifts in their controlled environment study designs. But this trend also introduces a bias in results, as any comparisons of studies from lower and higher latitudes are also comparing a different range of cues.

#### *How studies manipulate cues*

Studies can be broadly categorized as manipulating one, two or (rarely) three cues at once. Single cue studies were the most common (65 studies), with most manipulating chilling (41 studies), followed by 14 manipulating photoperiod and 10 manipulating forcing (Fig. 4). While valuable for defining potential non-linearities in one cue, single cue studies are difficult to use alone for forecasting as they prevent understanding interactions among cues or comparisons of which cues dominate phenological responses.

Multiple cue studies may be especially insightful for forecasting, especially if they test for interactions of cues. Of the studies manipulating at least one cue, roughly a third additionally manipulated another cue (37% or 43 studies). Study designs most often examined the interaction of cues (that is, whether the effect of one cue depends on the level of the other cue), with the studies almost evenly split across all possible two-way interactions (14 studies of photoperiod x chilling, followed by 13 studies of forcing x chilling, followed by 12 studies of photoperiod x forcing; 4 studies did not include an interaction).

Studies examining three cues directly were rare: we identified only three studies examining all three cues at once. Two of these were on *Picea abies* (Worrall & Mergen, 1967; Sogaard *et al.*, 2008), and the other on *Betula pendula* (Skuterud & Dietrichson, 1994). A slightly larger set of studies (5 studies from 4 papers) examined three cues indirectly—manipulating photoperiod

and forcing in controlled environments but equating chilling with sequential removal of tissue from the field—for 11 species (Schnabel & Wample, 1987; Heide, 1993a; Partanen *et al.*, 1998; Basler & Körner, 2014).

The limited number of studies examining multiple cues may impact forecasting because accurate models require understanding how—when combined—cues will determine future budburst with continued warming. Because the cues are all known to be interactive, estimates of any one cue are influenced by the level of each other cue. But knowing the level of each cue is difficult using current studies because they are often not reported (see ‘NA’ in Fig. 4), and many studies use field conditions to apply treatments (see Fig. 4, and ‘Paths forward’ below).

#### *How relevant are treatments to current and future conditions?*

The utility of controlled environment studies to forecasting also depends on how relevant treatments are to current and future conditions. Estimating such relevance is difficult as it depends on a species’ geographical range and projections considered. However, a simple analysis of two well studied species, *Fagus sylvatica* and *Betula pendula*, suggests experiments have generally bracketed the range of projected temperatures (Fig. 5), though there is a limited number of chilling studies that directly manipulate chilling temperature (Fig. 4). Indeed, we found no studies with multiple chill temperatures for *Fagus sylvatica*, even though it is one of the most well-studied species (Fig. 5C).

Experimental treatments were generally larger than expected shifts due to climate change. This makes sense from an experimental-statistical perspective: if the goal of an experiment is to identify if a cue is present then larger treatment differences should yield larger effect sizes and higher statistical power. But such large shifts may be risky to extrapolate from when predicting effects of warming.

## **1.4 Paths forward**

We argue that controlled environment experiments will be critical for accurate predictions of phenology given future warming. How accurate such predictions are will depend on the design of future experiments, breakthroughs in our physiological understanding of the major cues, and how well these two areas can be integrated with long-term data to improve models.

#### *Improving controlled environment studies*

We expect the most useful future experiments for forecasting will be designed to improve phenological models. In particular, experiments designed to identify threshold effects, optimal temperatures/photoperiods, and non-linearities from interactive cues may be most useful (Iler *et al.*, 2013). Identifying threshold effects and optimal temperatures or photoperiods generally requires many different levels of a single cue, which can make such experiments difficult to cross with other cues. Yet, understanding if findings are consistent across varying levels of other cues is critical for robust forecasting. Studies manipulating more than one cue also test for non-linearities due to interactive cues, as long as they are fully-crossed (i.e., every combination of levels is present in treatments). Such experiments can quickly require a large number of con-

255 trolled environments. However, as experimental results support that all cues are dependent on  
256 the level of other cues (Stearns & Olson, 1958; Flynn & Wolkovich, 2018) and long-term data  
257 hint at multiple cues (Fu *et al.*, 2015), we believe this should be a major research aim.

258  
259 Controlled environment studies may also be more readily applied to forecasting by exploring  
260 more realistic conditions. While identifying thresholds, optima and non-linearities may involve  
261 considering informative extremes in levels of cues, most changes in cues due to climate change  
262 are and will be on a (relatively) smaller scale (Fig. 5). Thus, experimentalists should examine  
263 cues within the current and projected future species' range. While species distribution models  
264 generally assume species will remain in the same climatic conditions (Elith & Leathwick, 2009),  
265 most evidence suggests species will lag in their spatial responses, meaning shifts in cues in the  
266 current range may be important to the fate of trailing edge populations (Bertrand *et al.*, 2011;  
267 Lenoir & Svenning, 2015; Savage & Vellend, 2015).

268  
269 Beyond the absolute level of cues, controlled environment studies need more work on what at-  
270 tributes of the design are more or less critical for replicating responses from the field. For exam-  
271 ple, controlled environment studies have shown differing day/night temperatures are important  
272 for some species (Heuvelink, 1989; Abrol & Ingram, 1996; Thingnaes *et al.*, 2003; Pressman  
273 *et al.*, 2006), but comparison studies have not been conducted for most species. Equally, a few  
274 studies have attempted to replicate certain aspects of the environment, such as fluctuating tem-  
275 peratures, ramped temperatures and the coincidence of temperature and sunrise (Erwin, 1998),  
276 but these are by no means widespread enough to understand how important these conditions  
277 are for extrapolation to models.

#### 278 *Understanding the physiology of phenology*

279 Even with all the suggested above improvements, controlled environment studies will still be fun-  
280 damentally limited in their utility for prediction without a deeper understanding of how major  
281 phenological cues act physiologically (Bahuguna & Jagadish, 2015). This problem is most acute  
282 in our understanding of dormancy, which divides chilling and forcing (Singh *et al.*, 2019; Chang  
283 *et al.*, 2021). Researchers may use the terms 'chilling' and 'forcing' for their treatments, but  
284 they rarely have physiological evidence that these are the actual conditions plants experience  
285 (Chuine *et al.*, 2016).

286  
287  
288 Chilling is generally defined as accumulated cool temperatures in the first phase of dormancy  
289 (endodormancy), after which plants enter ecodormancy, when accumulated forcing leads to bud-  
290 burst (Chuine *et al.*, 2016). Measuring endodormancy and its transition into ecodormancy is  
291 notoriously difficult (e.g., Junttila & Hänninen, 2012). Thus, in practice, many phenology stud-  
292 ies use the terms 'chilling' and 'forcing' to mean 'cool temperatures' (either in the fall and winter  
293 or applied in experimental conditions) and 'warm temperatures' (either in the spring or applied  
294 after sufficient chilling) and generally hope they correspond to endo- and eco-dormancy.

295  
296 Physiologists have long recognized this issue and recent breakthroughs provide new insights into  
297 what causes dormancy at the cellular level (van der Schoot *et al.*, 2014). Research suggests



endodormancy may break when enzymes sufficiently remove the sugar (callose) that blocks plasmodesmata in bud cells (reviewed in Chang *et al.*, 2021). Work thus far has relied generally on cellular staining methods tested on a very limited subset of species (Rinne *et al.*, 2011; Singh *et al.*, 2019), making extrapolation to other species difficult. Such results, however, hold promise for a much improved physiological understanding in the near future.

An improved understanding of endodormancy release could revolutionize models of chilling, and in turn, estimates of forcing. Forcing in controlled environment experiments is often defined simply as warm temperatures (or warm temperatures after cool temperatures). Future estimates could more accurately define forcing using only temperatures during ecodormancy, assuming tractable tests of endo- and eco-dormancy and the uptake of such tests in controlled environment studies. With these experiments in hand, researchers could quickly build improved models of chilling, and forcing and—for the first time—provide accurate estimates of how chilling has and will shift with climate change.

#### *Improving integration of controlled environment and physiological studies with long-term data*

Integrating long-term observational, physiological and controlled environment studies would help advance our understanding and climate change forecasting. With important exceptions (e.g., Gauzere *et al.*, 2017), studies of long-term observational phenology data have moved forward independently from advances in our physiological understanding and from controlled environment studies. Similarly, controlled environment studies generally do not use long-term data to help interpret results or define treatments.

Studies that have integrated results from both controlled environment experiments and with long-term observations provide a path forward (Caffarra *et al.*, 2011; Nagano *et al.*, 2012; Satake *et al.*, 2013; Ford *et al.*, 2016; Chuine & Regnier, 2017)—one that can happen while we await physiological breakthroughs in defining endo- and eco-dormancy. Experiments that test for thresholds and the presence of important interactions have helped re-design models (Caffarra *et al.*, 2011; Chuine & Regnier, 2017), while other studies have used experiments to test extremes combined with data from long-term provenance studies to understand how growth and phenology will combine to determine future ranges (Ford *et al.*, 2016). Such research underlies progress towards model development that relies continuously on a back-and-forth process between developing models based on both long-term data and experiments, then testing predictions with new experiments and newly-available observational data (i.e., more years and also data from new locations, Nagano *et al.*, 2012; Satake *et al.*, 2013). Such continual development, which takes extensive data, has been carried out on very few species (e.g., *Arabidopsis thaliana*, *Oryza sativa* (rice), *Arabidopsis halleri*, Wilczek *et al.*, 2009; Nagano *et al.*, 2012; Satake *et al.*, 2013).

#### *Building species-rich predictions*

Given the efforts and data involved in models for a single species, building up to multi-species predictions may appear daunting, but multi-species models are crucial for accurate forecasts that can apply to diverse regions and large-scale vegetation models. Addressing this issue requires, of

course, more data. Long-term data is generally more species-rich than controlled environment studies. For example long-term observational data in the PEP725 and NECTAR databases together have multi-site data on more than 2500 species (Wolkovich *et al.*, 2012b; Templ *et al.*, 2018), while our review of controlled environment studies found most studies focused on only one species, with data on a total of 226 species. Thus, more diverse controlled environment studies may be the current major data limitation. Beyond data, however, new modeling approaches can help integrate current and future data more powerfully.

Bayesian hierarchical models are specifically designed to synthesize across multiple sources of data. With the right information, they can attribute variation across studies to the species studied, the cues (i.e., chilling, forcing and photoperiod levels in studies) and remaining unmeasured variation in studies (i.e., unreported lab and chamber differences may be captured by including a parameter to estimate a ‘study’ effect). Such models are extremely powerful for building species-rich predictions, yet, like all models, they are more robust with more data. In particular, attributing variation due to study versus species requires the same species to be studied across several studies, which is currently not the case for most species, according to our literature review (81% of the 226 species appear in only one study; 14%, appear in two or three studies, and 4%, appear in more than three studies).

## 1.5 Right now: It’s your tomorrow

Research on phenology had been conducted for centuries before anthropogenic climate change caused earlier budburst and leafout across much of the globe (Lamb, 1948; Sparks & Carey, 1995). Decades of controlled environment studies contributed to our fundamental understanding of the drivers of spring plant phenology. Today, climate change requires leveraging these decades and centuries of research for more accurate predictions that can help humans adapt to warming.

We have outlined how researchers could better harness the power of controlled environment experiments to transform our fundamental understanding of phenology and advance forecasting. Controlled environment studies can critically rule out, or support, hypotheses to explain observed discrepancies in long-term data and open up new pathways to use long-term data to understand current trends. While understanding, modeling and predicting interactions among cues and their effects on phenology is challenging, it will yield more accurate predictions—with valuable implications to more realistically assess the effects of climate change on plant biodiversity, including agricultural and forest species.

## 2 References

- Abrol, Y.P. & Ingram, K.T. (1996) Effects of higher day and night temperatures on growth and yields of some crop plants. *Global climate change and agricultural production: Direct and indirect effects of changing hydrological, pedological and plant physiological processes* pp. 123–140.
- Alexander, L.V., Zhang, X., Peterson, T.C., Caesar, J., Gleason, B., Tank, A., Haylock, M., Collins, D., Trewin, B., Rahimzadeh, F., Tagipour, A., Kumar, K.R., Revadekar, J., Griffiths, G., Vincent, L., Stephenson, D.B., Burn, J., Aguilar, E., Brunet, M., Taylor, M., New, M., Zhai, P., Rusticucci, M. & Vazquez-Aguirre, J.L. (2006) Global observed changes in daily climate extremes of temperature and precipitation. *Journal of Geophysical Research-Atmospheres* **111**.
- Asse, D., Chuine, I., Vitasse, Y., Yoccoz, N.G., Delpierre, N., Badeau, V., Delestrade, A. & Randin, C.F. (2018) Warmer winters reduce the advance of tree spring phenology induced by warmer springs in the alps. *Agricultural and Forest Meteorology* **252**, 220–230.
- Bahuguna, R.N. & Jagadish, K.S. (2015) Temperature regulation of plant phenological development. *Environmental and Experimental Botany* **111**, 83–90.
- Basler, D. & Körner, C. (2014) Photoperiod and temperature responses of bud swelling and bud burst in four temperate forest tree species. *Tree Physiology* **34**, 377–388.
- Bertrand, R., Lenoir, J., Piedallu, C., Riofrío-Dillon, G., de Ruffray, P., Vidal, C., Pierrat, J.C. & Gégout, J.C. (2011) Changes in plant community composition lag behind climate warming in lowland forests. *Nature* **479**, 517.
- Burrows, M.T., Schoeman, D.S., Buckley, L.B., Moore, P., Poloczanska, E.S., Brander, K.M., Brown, C., Bruno, J.F., Duarte, C.M., Halpern, B.S., Holding, J., Kappel, C.V., Kiessling, W., O'Connor, M.I., Pandolfi, J.M., Parmesan, C., Schwing, F.B., Sydeman, W.J. & Richardson, A.J. (2011) The pace of shifting climate in marine and terrestrial ecosystems. *Science* **334**, 652–655, burrows, Michael T. Schoeman, David S. Buckley, Lauren B. Moore, Pippa Poloczanska, Elvira S. Brander, Keith M. Brown, Chris Bruno, John F. Duarte, Carlos M. Halpern, Benjamin S. Holding, Johnna Kappel, Carrie V. Kiessling, Wolfgang O'Connor, Mary I. Pandolfi, John M. Parmesan, Camille Schwing, Franklin B. Sydeman, William J. Richardson, Anthony J.
- Caffarra, A., Donnelly, A. & Chuine, I. (2011) Modelling the timing of *Betula pubescens* bud-burst. II. Integrating complex effects of photoperiod into process-based models. *Climate Research* **46**, 159–170.
- Cesaraccio, C., Spano, D., Snyder, R.L. & Duce, P. (2004) Chilling and forcing model to predict bud-burst of crop and forest species. *Agricultural and Forest Meteorology* **126**, 1–13.

- 411 Chang, C.Y.Y., Brautigam, K., Huner, N.P.A. & Ensminger, I. (2021) Champions of winter  
412 survival: cold acclimation and molecular regulation of cold hardiness in evergreen conifers.  
413 *New Phytologist* **229**, 675–691.
- 414 Chen, L., Huang, J.G., Ma, Q.Q., Hanninen, H., Tremblay, F. & Bergeron, Y. (2019) Long-term  
415 changes in the impacts of global warming on leaf phenology of four temperate tree species.  
416 *Global Change Biology* **25**, 997–1004.
- 417 Chuine, I. (2000) A unified model for budburst of trees. *Journal of Theoretical Biology* **207**, 337  
418 – 347.
- 419 Chuine, I., Bonhomme, M., Legave, J.M., García de Cortázar-Atauri, I., Charrier, G., Lacointe,  
420 A. & Améglio, T. (2016) Can phenological models predict tree phenology accurately in the  
421 future? The unrevealed hurdle of endodormancy break. *Global Change Biology* **22**, 3444–3460.
- 422 Chuine, I. & Regniere, J. (2017) Process-based models of phenology for plants and animals.  
423 *Annual Review of Ecology, Evolution, and Systematics* **48**, 159–182.
- 424 Cleland, E.E., Chuine, I., Menzel, A., Mooney, H.A. & Schwartz, M.D. (2007) Shifting plant  
425 phenology in response to global change. *Trends in Ecology & Evolution* **22**, 357–365.
- 426 Delpierre, N., Dufrene, E., Soudani, K., Ulrich, E., Cecchini, S., Boe, J. & Francois, C. (2009)  
427 Modelling interannual and spatial variability of leaf senescence for three deciduous tree species  
428 in France. *Agricultural and Forest Meteorology* **149**, 938–948.
- 429 Elith, J. & Leathwick, J.R. (2009) Species distribution models: ecological explanation and pre-  
430 diction across space and time. *Annual review of ecology, evolution, and systematics* **40**, 677–  
431 697.
- 432 Ellwood, E.R., Temple, S.A., Primack, R.B., Bradley, N.L. & Davis, C.C. (2013) Record-  
433 breaking early flowering in the eastern united states. *Plos One* **8**, ellwood, Elizabeth R.  
434 Temple, Stanley A. Primack, Richard B. Bradley, Nina L. Davis, Charles C.
- 435 Elmendorf, S.C. & Ettinger, A.K. (2020) Is photoperiod a dominant driver of secondary growth  
436 resumption? *Proceedings of the National Academy of Sciences of the United States of America*  
437 **117**, 32861–32864.
- 438 Erwin, J.E. (1998) Temperature and light effects on stem elongation. *Journal of the Japanese*  
439 *Society for Horticultural Science* **67**, 1113–1120.
- 440 Ettinger, A., Chamberlain, C., Morales-Castilla, I., Buonaiuto, D., Flynn, D., Savas, T., Samaha,  
441 J. & Wolkovich, E. (2020) Winter temperatures predominate in spring phenological responses  
442 to warming. *Nature Climate Change* pp. 1–6.
- 443 Fishman, S., Erez, A. & Couvillon, G. (1987) The temperature dependence of dormancy break-  
444 ing in plants: mathematical analysis of a two-step model involving a cooperative transition.  
445 *Journal of Theoretical Biology* **124**, 473–483.

- 446 Flynn, D.F.B. & Wolkovich, E.M. (2018) Temperature and photoperiod drive spring phenology  
447 across all species in a temperate forest community. *New Phytologist* **219**, 1353–1362.
- 448 Ford, K.R., Harrington, C.A., Bansal, S., Gould, Peter, J. & St. Clair, J.B. (2016) Will changes  
449 in phenology track climate change? A study of growth initiation timing in coast Douglas–fir.  
450 *Global Change Biology* **22**, 3712–3723.
- 451 Fu, Y.H., Piao, S., Zhou, X., Geng, X., Hao, F., Vitasse, Y. & Janssens, I.A. (2019) Short  
452 photoperiod reduces the temperature sensitivity of leaf-out in saplings of *Fagus sylvatica* but  
453 not in horse chestnut. *Global Change Biology* **25**, 1696–1703.
- 454 Fu, Y.S.H., Zhao, H.F., Piao, S.L., Peaucelle, M., Peng, S.S., Zhou, G.Y., Ciais, P., Huang, M.T.,  
455 Menzel, A., Uelas, J.P., Song, Y., Vitasse, Y., Zeng, Z.Z. & Janssens, I.A. (2015) Declining  
456 global warming effects on the phenology of spring leaf unfolding. *Nature* **526**, 104–107.
- 457 Gauzere, J., Delzon, S., Davi, H., Bonhomme, M., de Cortazar-Atauri, I.G. & Chuine, I. (2017)  
458 Integrating interactive effects of chilling and photoperiod in phenological process-based mod-  
459 els. A case study with two European tree species: *Fagus sylvatica* and *Quercus petraea*. *Agri-  
460 cultural and Forest Meteorology* **244**, 9–20.
- 461 Gauzere, J., Lucas, C., Ronce, O., Davi, H. & Chuine, I. (2019) Sensitivity analysis of tree phe-  
462 nology models reveals increasing sensitivity of their predictions to winter chilling temperature  
463 and photoperiod with warming climate. *Ecological Modelling* **441**, 108805.
- 464 Güsewell, S., Furrer, R., Gehrig, R. & Pietragalla, B. (2017) Changes in temperature sensitivity  
465 of spring phenology with recent climate warming in Switzerland are related to shifts of the  
466 preseason. *Global Change Biology* **23**, 5189–5202.
- 467 Guy, R.D. (2014) The early bud gets to warm. *New Phytologist* **202**, 7–9.
- 468 Heide, O. (1993a) Daylength and thermal time responses of budburst during dormancy release  
469 in some northern deciduous trees. *Physiologia Plantarum* **88**, 531–540.
- 470 Heide, O. (1993b) Dormancy release in beech buds (*Fagus sylvatica*) requires both chilling and  
471 long days. *Physiologia Plantarum* **89**, 187–191.
- 472 Hereford, J., Schmitt, J. & Ackerly, D.D. (2017) The seasonal climate niche predicts phenology  
473 and distribution of an ephemeral annual plant, *mollugo verticillata*. *Journal of Ecology* **105**,  
474 1323–1334.
- 475 Heuvelink, E. (1989) Influence of day and night temperature on the growth of young tomato  
476 plants. *Scientia Horticulturae* **38**, 11–22.
- 477 Iler, A.M., Høye, T.T., Inouye, D.W. & Schmidt, N.M. (2013) Nonlinear flowering responses to  
478 climate: are species approaching their limits of phenological change? *Philosophical Transac-  
479 tions of the Royal Society B: Biological Sciences* **368**, 20120489.
- 480 IPCC (2014) *Climate Change 2014: Impacts, Adaptation, and Vulnerability*. Cambridge Univer-  
481 sity Press, Cambridge, United Kingdom and New York, NY, USA.

- Jochner, S., Caffarra, A. & Menzel, A. (2013) Can spatial data substitute temporal data in phenological modelling? a survey using birch flowering. *Tree Physiology* **33**, 1256–1268.
- Junttila, O. & Hänninen, H. (2012) The minimum temperature for budburst in *Betula* depends on the state of dormancy. *Tree physiology* **32**, 337–345.
- Keenan, T.F., Richardson, A.D. & Hufkens, K. (2020) On quantifying the apparent temperature sensitivity of plant phenology. *New Phytologist* **225**, 1033–1040.
- Lamb, R.C. (1948) Effect of temperatures above and below freezing on the breaking of rest in the latham raspberry. *Proceedings of the American Society for Horticultural Science*, vol. 51, pp. 313–315, AMER SOC HORTICULTURAL SCIENCE 701 NORTH SAINT ASAPH STREET, ALEXANDRIA, VA 22314-1998.
- Lenoir, J. & Svenning, J.C. (2015) Climate-related range shifts—a global multidimensional synthesis and new research directions. *Ecography* **38**, 15–28.
- Luedeling, E., Guo, L., Dai, J., Leslie, C. & Blanke, M.M. (2013) Differential responses of trees to temperature variation during the chilling and forcing phases. *Agricultural and Forest Meteorology* **181**, 33–42.
- Luedeling, E., Zhang, M.H., McGranahan, G. & Leslie, C. (2009) Validation of winter chill models using historic records of walnut phenology. *Agricultural and Forest Meteorology* **149**, 1854–1864.
- Major, D.J. (1980) Photoperiod response characteristics controlling flowering of 9 crop species. *Canadian Journal of Plant Science* **60**, 777–784.
- Martinez-Luscher, J., Hadley, P., Ordidge, M., Xu, X.M. & Luedeling, E. (2017) Delayed chilling appears to counteract flowering advances of apricot in southern uk. *Agricultural and Forest Meteorology* **237**, 209–218.
- Menzel, A., Jakobi, G., Ahas, R., Scheifinger, H. & Estrella, N. (2003) Variations of the climatological growing season (1951–2000) in Germany compared with other countries. *International Journal of Climatology* **23**, 793–812.
- Menzel, A., Sparks, T.H., Estrella, N., Koch, E., Aasa, A., Ahas, R., Alm-Kuebler, K. *et al.* (2006) European phenological response to climate change matches the warming pattern. *Global Change Biology* **12**, 1969–1976.
- Myking, T. & Heide, O. (1995) Dormancy release and chilling requirement of buds of latitudinal ecotypes of *Betula pendula* and *B. pubescens*. *Tree Physiology* **15**, 697–704.
- Nagano, A.J., Sato, Y., Mihara, M., Antonio, B.A., Motoyama, R., Itoh, H., Nagamura, Y. & Izawa, T. (2012) Deciphering and prediction of transcriptome dynamics under fluctuating field conditions. *Cell* **151**, 1358–1369.
- Nienstaedt, H. (1966) Dormancy and dormancy release in white spruce. *Forest Science* **12**, 374–384.

- 518 Parent, B. & Tardieu, F. (2012) Temperature responses of developmental processes have not  
519 been affected by breeding in different ecological areas for 17 crop species. *New Phytologist*  
520 **194**, 760–774.
- 521 Partanen, J., Koski, V. & Hänninen, H. (1998) Effects of photoperiod and temperature on the  
522 timing of bud burst in Norway spruce (*Picea abies*). *Tree Physiology* **18**, 811–816.
- 523 Piao, S., Liu, Z., Wang, T., Peng, S., Ciais, P., Huang, M., Ahlstrom, A., Burkhardt, J.F.,  
524 Chevallier, F., Janssens, I.A. *et al.* (2017) Weakening temperature control on the interannual  
525 variations of spring carbon uptake across northern lands. *Nature climate change* **7**, 359.
- 526 Polgar, C.A., Primack, R.B., Dukes, J.S., Schaaf, C., Wang, Z. & Hoeppe, S.S. (2014) Tree leaf  
527 out response to temperature: comparing field observations, remote sensing, and a warming  
528 experiment. *International journal of biometeorology* **58**, 1251–1257.
- 529 Prasad, P.V.V., Pisipati, S.R., Ristic, Z., Bukovnik, U. & Fritz, A.K. (2008) Impact of nighttime  
530 temperature on physiology and growth of spring wheat. *Crop Science* **48**, 2372–2380.
- 531 Pressman, E., Shaked, R. & Firon, N. (2006) Exposing pepper plants to high day temperatures  
532 prevents the adverse low night temperature symptoms. *Physiologia Plantarum* **126**, 618–626.
- 533 Richardson, A.D., Hufkens, K., Milliman, T., Aubrecht, D.M., Furze, M.E., Seyednasrollah, B.,  
534 Krassovski, M.B., Latimer, J.M., Nettles, W.R., Heiderman, R.R., Warren, J.M. & Hanson,  
535 P.J. (2018) Ecosystem warming extends vegetation activity but heightens vulnerability to cold  
536 temperatures. *Nature* **560**, 368–+.
- 537 Richardson, E. (1974) A model for estimating the completion of rest for ‘Redhaven’ and ‘Elberta’  
538 peach trees. *HortScience* **9**, 331–332.
- 539 Rinne, P.L.H., Paul, L.K. & van der Schoot, C. (2018) Decoupling photo- and thermoperiod by  
540 projected climate change perturbs bud development, dormancy establishment and vernaliza-  
541 tion in the model tree populus. *Bmc Plant Biology* **18**.
- 542 Rinne, P.L.H., Welling, A., Vahala, J., Ripel, L., Ruonala, R., Kangasjarvi, J. & van der Schoot,  
543 C. (2011) Chilling of dormant buds hyperinduces FLOWERING LOCUS T and recruits GA-  
544 Inducible 1,3-beta-Glucanases to reopen signal conduits and release dormancy in *Populus*.  
545 *Plant Cell* **23**, 130–146.
- 546 Saikkonen, K., Taulavuori, K., Hyvönen, T., Gundel, P.E., Hamilton, C.E., Vänninen, I., Nissi-  
547 nen, A. & Helander, M. (2012) Climate change-driven species’ range shifts filtered by pho-  
548 toperiodism. *Nature Climate Change* **2**, 239.
- 549 Satake, A., Kawagoe, T., Saburi, Y., Chiba, Y., Sakurai, G. & Kudoh, H. (2013) Forecasting  
550 flowering phenology under climate warming by modelling the regulatory dynamics of flowering-  
551 time genes. *Nature Communications* **4**.
- 552 Savage, J. & Vellend, M. (2015) Elevational shifts, biotic homogenization and time lags in  
553 vegetation change during 40 years of climate warming. *Ecography* **38**, 546–555.

- 554 Schnabel, B.J. & Wample, R.L. (1987) Dormancy and cold hardiness in *Vitis vinifera* L. cv.  
555 White Riesling as influenced by photoperiod and temperature. *American Journal of Enology*  
556 *and Viticulture* **38**, 265–272.
- 557 Schwartz, M.D. (1997) Spring index models: An approach to connecting satellite and surface  
558 phenology. *Phenology in Seasonal Climates I* (eds. H. Lieth & M.D. Schwartz), pp. 22–38,  
559 Backhuys Publisher, Leiden, The Netherlands.
- 560 Shen, M., Piao, S., Dorji, T., Liu, Q., Cong, N., Chen, X., An, S., Wang, S., Wang, T. & Zhang,  
561 G. (2015) Plant phenological responses to climate change on the Tibetan Plateau: research  
562 status and challenges. *National Science Review* **2**, 454–467.
- 563 Shen, X.J., Liu, B.H., Henderson, M., Wang, L., Wu, Z.F., Wu, H.T., Jiang, M. & Lu, X.G.  
564 (2018) Asymmetric effects of daytime and nighttime warming on spring phenology in the  
565 temperate grasslands of china. *Agricultural and Forest Meteorology* **259**, 240–249.
- 566 Singh, R.K., Miskolczi, P., Maurya, J.P. & Bhalerao, R.P. (2019) A tree ortholog of short  
567 vegetative phase floral repressor mediates photoperiodic control of bud dormancy. *Current*  
568 *Biology* **29**, 128–+.
- 569 Singh, R.K., Svystun, T., AlDahmash, B., Jönsson, A.M. & Bhalerao, R.P. (2017) Photoperiod-  
570 and temperature-mediated control of phenology in trees—a molecular perspective. *New Phy-*  
571 *tologist* **213**, 511–524.
- 572 Skuterud, R. & Dietrichson, J. (1994) Budburst in detached birch shoots (*Betula pendula*) of  
573 different varieties winter-stored in darkness at three different temperatures. *Silva Fennica* **28**,  
574 223–224.
- 575 Søgaaard, G., Johnsen, Ø., Nilsen, J. & Junttila, O. (2008) Climatic control of bud burst in young  
576 seedlings of nine provenances of norway spruce. *Tree Physiology* **28**, 311–320.
- 577 Sønsteby, A. & Heide, O.M. (2014) Chilling requirements of contrasting black currant (*Ribes*  
578 *nigrum* L.) cultivars and the induction of secondary bud dormancy. *Scientia Horticulturae*  
579 **179**, 256–265.
- 580 Sparks, T.H. & Carey, P.D. (1995) The responses of species to climate over two centuries - an  
581 analysis of the Marsham phenological record, 1736–1947. *Journal of Ecology* **83**, 321–329.
- 582 Stearns, F. & Olson, J. (1958) Interactions of photoperiod and temperature affecting seed ger-  
583 mination in *Tsuga canadensis*. *American Journal of Botany* **45**, 53–58.
- 584 Tansey, C.J., Hadfield, J.D. & Phillimore, A.B. (2017) Estimating the ability of plants to plas-  
585 tically track temperature-mediated shifts in the spring phenological optimum. *Global Change*  
586 *Biology* **23**, 3321–3334.
- 587 Templ, B., Koch, E., Bolmgren, K., Ungersböck, M., Paul, A., Scheifinger, H., Rutishauser, T.,  
588 Busto, M., Chmielewski, F.M., Hájková, L., Hodzić, S., Kaspar, F., Pietragalla, B., Romero-  
589 Fresneda, R., Tolvanen, A., Vučetić, V., Zimmermann, K. & Züst, A. (2018) Pan European



590 Phenological database (PEP725): a single point of access for European data. *International*  
591 *Journal of Biometeorology* **62**, 1109–1113.

592 Thielges, B. & Beck, R. (1976) Control of bud break and its inheritance in *Populus deltoides*.  
593 *Tree Physiology and Yield Improvement* **14**, 253–259.

594 Thingnaes, E., Torre, S., Ernstsens, A. & Moe, R. (2003) Day and night temperature responses  
595 in arabidopsis: Effects on gibberellin and auxin content, cell size, morphology and flowering  
596 time. *Annals of Botany* **92**, 601–612.

597 van der Schoot, C., Paul, L.K. & Rinne, P.L.H. (2014) The embryonic shoot: a lifeline through  
598 winter. *Journal of Experimental Botany* **65**, 1699–1712.

599 Visser, M.E. & Holleman, L.J. (2001) Warmer springs disrupt the synchrony of oak and winter  
600 moth phenology. *Proceedings of the Royal Society of London. Series B: Biological Sciences*  
601 **268**, 289–294.

602 Way, D.A. & Montgomery, R.A. (2015) Photoperiod constraints on tree phenology, performance  
603 and migration in a warming world. *Plant, Cell & Environment* **38**, 1725–1736.

604 Weinberger, J.H. *et al.* (1950) Chilling requirements of peach varieties. *Proceedings. American*  
605 *Society for Horticultural Science*, vol. 56, pp. 122–28.

606 Wilczek, A.M., Roe, J.L., Knapp, M.C., Cooper, M.D., Lopez-Gallego, C., Martin, L.J., Muir,  
607 C.D., Sim, S., Walker, A., Anderson, J., Egan, J.F., Moyers, B.T., Petipas, R., Giakountis, A.,  
608 Charbit, E., Coupland, G., Welch, S.M. & Schmitt, J. (2009) Effects of genetic perturbation  
609 on seasonal life history plasticity. *Science* **323**, 930–934.

610 Wolkovich, E.M., Cook, B.I., Allen, J.M., Crimmins, T.M., Betancourt, J.L., Travers, S.E., Pau,  
611 S., Regetz, J., Davies, T.J., Kraft, N.J.B., Ault, T.R., Bolmgren, K., Mazer, S.J., McCabe,  
612 G.J., McGill, B.J., Parmesan, C., Salamin, N., Schwartz, M.D. & Cleland, E.E. (2012a)  
613 Warming experiments underpredict plant phenological responses to climate change. *Nature*  
614 **485**, 494–497.

615 Wolkovich, E.M., Cook, B.I. & Regetz, J. (2012b) NECTAR: Network of Ecological and Climato-  
616 logical Timings Across Regions, <http://knb.ecoinformatics.org/knb/metacat/nceas.988/knb>.

617 Worrall, J. & Mergen, F. (1967) Environmental and genetic control of dormancy in *Picea abies*.  
618 *Physiologia Plantarum* **20**, 733–745.

619 Yu, H.Y., Luedeling, E. & Xu, J.C. (2010) Winter and spring warming result in delayed spring  
620 phenology on the Tibetan Plateau. *Proceedings of the National Academy of Sciences of the*  
621 *United States of America* **107**, 22151–22156.

622 Zohner, C.M., Benito, B.M., Svenning, J.C. & Renner, S.S. (2016) Day length unlikely to con-  
623 strain climate-driven shifts in leaf-out times of northern woody plants. *Nature Climate Change*  
624 **6**, 1120–1123.

625 Zohner, C.M. & Renner, S.S. (2014) Common garden comparison of the leaf-out phenology  
626 of woody species from different native climates, combined with herbarium records, forecasts  
627 long-term change. *Ecology Letters* **17**, 1016–1025.






Figure 1: Interactions can produce nonlinearities, even in simple linear models if there are correlated shifts in cues. Much research focuses on how warming increases forcing (A), but it may also alter other cues, including photoperiod experienced near the time of the event, which is expected to shorten, and chilling, which may either increase or decrease (B). Shifts in forcing alongside shifts in a second cue (A) may produce non-linearities due to the interaction between cues (C-F showing the effect of: forcing-only in yellow, both cues without an interaction in lighter purple/blue, and both cues with an interaction in darker purple/blue); the overall change in budburst day predicted with warming depends on the sign (+/-), strength (weak/strong), and shape of the second cue (showing two simple examples: linear in C-D and threshold E-F).




Figure 2: A common example of how the level of one cue can modify the required level of another cue comes from experiments finding that the amount of chilling affects the amount of forcing needed for budburst. Here, we show this from experiments in black currant (*Ribes nigrum*) from Sønsteby & Heide (2014) and eastern cottonwood (*Populus deltoides*) from Thielges & Beck (1976). Forcing is estimated as growing degree days (GDD, sum of temperatures  $> 0^{\circ}\text{C}$ ) while chilling is estimated using the Utah Model (Richardson, 1974).




Figure 3: Current models of chilling suggest it may decrease or increase with winter warming. Here we show a common version of the Utah chilling model (top right inset and also turquoise line in main figure) with two conceptual scenarios of mean daily winter temperatures. When temperatures are generally below zero warming may increase accumulated chilling (left), while if pre-climate change temperatures are generally higher (near where chilling accumulates most per  $^{\circ}\text{C}$ ) then warming may decrease accumulated chilling (right).




Figure 4: Studies have manipulated a range of forcing temperatures (vertical axis) and photoperiods (horizontal axis) across two methods to manipulate chilling: a) experimentally or b) using multiple field sampling dates. This latter method, generally attributed to Weinberger *et al.* (1950), takes tissue (e.g., cuttings of adult dormant trees) from the field progressively across the fall and/or winter then exposes it to controlled environment conditions. These studies often equate tissue removed later as having received more chilling and thus often treat ‘time of cutting’ as interchangeable with ‘chilling,’ though forcing and photoperiod conditions also change. Gray squares indicate a combination of forcing x photoperiod not present for that method of chilling design (includes studies that did not clearly define chilling treatments), while a value of ‘NA’ indicates that we could not estimate a level of a particular cue because it was not clearly reported. Photoperiod treatments were generally only applied (or reported) during forcing conditions.




Figure 5: Controlled environment studies' treatments are comparable to predicted temperature changes in the future for minima (A, C) and maxima (B, D). (C-D). Points represent a site with spring phenology data for each of two species, *Fagus sylvatica* (A-B) and *Betula pendula*, (from the PEP725 database, Templ *et al.*, 2018) and show the predicted change (2080s average - 1970s average) in minimum (A,C) and maximum (B,D) temperatures 60 days before leafout. Inlay plots in the upper-left corner of each plot show a histogram of these same predicted changes in temperature overlaid with densities of the chilling (A, C) and forcing (B, D) differences in treatments across studies (green lines show the treatments for that exact species, while black lines show across all species; for *Fagus sylvatica* there are no chilling treatments of differing temperatures). For more details, see 'Comparing experimental treatments to forecasted trends' in the Supporting Information.