

# Chilling outweighs photoperiod and forcing cues for temperate trees in experiments, but not in natural systems

A.K. Ettinger, C. Chamberlain, I. Morales-Castilla, D. Buonaiuto, D. Flynn, T. Savas,  
J. Samaha & E. Wolkovich

May 6, 2019

## Abstract

Decades of research on woody species highlight how three major cues shape spring phenological events (e.g., budburst and leafout): forcing (warm temperatures, generally occurring in the late winter and early spring), daylength (photoperiod) and chilling (cool temperatures, generally occurring in the fall and late winter). How pervasive these cues are and whether some species are effectively governed by only one or two cues is a critical area of climate change biology research, as it would shape how complex responses to warming will be. Here we use a global meta-analysis of all published growth chamber studies to test for the relative effects of these three major cues across XX species. We find they almost all show these cues, making climate change responses complex.

## Text so far...

For decades, plant phenology has been one of the most reported and consistent biological imprints of climate change (IPCC, 2014), with many temperate plants leafing and flowering earlier with rising temperatures (Miller-Rushing and Primack, 2008; Menzel et al., 2006). Understanding such shifts is important as phenology shapes a suite of ecosystem services, including pollination and carbon sequestration, and scales up to impact projections of climate change itself.

As research interest in phenology has progressed, critical discrepancies and uncertainties in our understanding have emerged. Though responses to warming are consistent on average, they show high variation across species and sites (Wolkovich et al., 2012). Furthermore, long-term observational studies provide increasing evidence that sensitivities of phenology to temperature are weakening in recent decades (Rutishauser et al., 2008; Yu et al., 2010; Fu et al., 2015). In Europe, recent work from many of the most well-studied tree species shows declining responses to temperature, suggesting that the long-term trend towards ever-earlier springs may be stalling (Fu et al., 2015). The authors, and others, suggest that responses to other environmental cues underlie these declining temperature sensitivities.

Fundamental research in phenology outlines three major cues known to shape spring phenology (Chuine, 2000): chilling (cool temperatures, generally occurring in the fall and late winter), forcing (warm temperatures, generally occurring in the late winter and early spring), and daylength (photoperiod). These cues are thought to provide multiple routes to budburst each spring depending on the environment. For example, in some species a cool winter resulting in high chilling will require a lower amount of forcing to trigger budburst, compared to a warmer winter that results in significantly lower chilling (Harrington and Gould, 2015), while in other species daylength may help trigger budburst given low chilling and/or forcing (CITES). Research suggests that all three cues may underlie spring phenology for many temperate woody species (Flynn and

Wolkovich, 2018; Basler and Körner, 2014; Caffarra et al., 2011a), but there is strong debate, with some research suggesting some cues—such as photoperiod— may be effectively absent in some species, but dominate in others (Zohner et al., 2016; Koerner and Basler, 2010a).

Given the declining response to temperature observed in long-term observational studies (Fu et al., 2015), a number of papers have tried to tease out evidence that chilling or daylength cues are playing an increasingly important role in recent years (Basler and Körner, 2014; Zohner et al., 2016; Laube et al., 2014). This work must overcome the fundamental challenge that all three cues are strongly correlated in nature. During the transition from winter to spring at most temperate latitudes, air temperatures increase (i.e., forcing increases) at the same time that daylength is increasing; likewise, winters with low amounts of chilling are often correlated with warmer springs, and thus higher forcing. Identifying which of these cues most strongly affects spring phenology is critical for forecasting future phenological changes. For example, if forcing is the dominant cue (as many observational studies to date have assumed, (Bradley et al., 1999; Menzel et al., 2006; Harrington and Gould, 2015)), then we can expect additional spring advancement as temperatures continue to warm. However, if unfulfilled chilling limits budburst, then we may see delays in spring phenology with additional global warming, which will reduce chilling in many areas (Fraga et al., 2019).

In contrast to observational studies, controlled environment experiments can break correlations between chilling, forcing, and photoperiod to reveal which cues underlie budburst phenology. These experiments—most often conducted in growth chambers or similar systems to control temperature and light—have been conducted for decades as a major method to understand the fundamental drivers of spring phenology. To date, controlled environment experiments have identified contrasting effects of the three major budburst cues. Some studies have found that photoperiod is likely to constrain species responses to climatic warming (Basler and Körner, 2012; Caffarra et al., 2011b; Caffarra and Donnelly, 2011), whereas others report that photoperiod is not a strong cue for most species (Zohner et al., 2016; Laube et al., 2014) and that chilling is more important to current and future trends.

Here, we leverage nearly 40 years of controlled environment studies to understand how chilling, forcing, and photoperiod contribute to budburst timing in woody species. Using a meta-analytic approach we reviewed XX papers from controlled environment studies, then extracted data from any papers that met XX conditions, yielding data from 74 studies across 39 years and 223 species (reference map of studies). This database includes only studies for which we could identify forcing, photoperiod, and chilling treatments quantitatively. As chilling was only rarely reported, we estimated chilling (when possible) using local climate data (see Supplemental Materials). We used a Bayesian hierarchical model to estimate the effects of chilling, forcing, and photoperiod. This model estimates both species-level responses (generally yielding more accurate estimates for well-studied species, such as *Fagsyl*, *Betpen*) and the distribution from which they are drawn, yielding a higher-level estimate of the overall response across species (see Supplemental Materials- mention species complex).

Across studies, all cues—chilling, forcing, and photoperiod—each advance budburst phenology (Fig. 1). Using a standardized scale to allow comparisons of the three cues we found that chilling was the strongest cue (-8.18 days/standard unit or -0.25 days per chill portion, Fig. 2), followed by forcing (-4.85 days/standard unit or -0.85 days per degree of warming, Fig. 2), and photoperiod (-3.17 days/standard unit or -0.53 days per hour of daylength). While photoperiod had the smallest effect among the three cues, our results contrast with the extensive literature suggesting photoperiod is a weak or non-existent cue for many species (Zohner et al., 2016; Koerner and Basler, 2010a)— instead we found it was surprisingly large and consistent across species. Only *Fagus sylvatica*, a species well-known for having a large response to photoperiod deviating far from the overall estimate (Figure 1). Species also showed fairly consistent responses to chilling (Figure 1, though two species delayed budburst with chilling *Tilia codata*, *Salix* complex). Responses to forcing, on the other hand, were the most variable across species ( $\text{sigma} = 6$ ).

As temperature is radically altered by anthropogenic climate change, our finding that different ends of the temperature spectrum—chilling and forcing—have the strongest effects on budburst suggests that under-

standing these cues will be critical for forecasting. Many previous studies attribute advances in budburst to increased forcing (Basler and Körner, 2014; Bradley et al., 1999; Menzel et al., 2006; Harrington and Gould, 2015), and forcing sensitivity in our model (-0.85 days per degree of warming) is consistent with what previous experiments (CITES), and observational studies (CITES) have observed. Our results, however, suggest chilling has a greater effect on budburst than forcing (Fig. 1). This has not been widely suggested previously, perhaps because little work has directly manipulated chilling, and the few studies that have were designed to compare chilling and photoperiod effects (e.g., Basler and Körner, 2014; Caffarra et al., 2011a; Laube et al., 2014; Zohner et al., 2016).

The paucity of studies directly manipulating chilling—which our results suggest has the greatest effect on budburst—suggests a major gap in current research. While many studies (X out of Y here) directly manipulated forcing, far fewer directly manipulated chilling (Z out of Y). Instead many studies (J out of Y) estimate chilling effects through sequential removal of tissue from the field and exposure to ‘forcing’ conditions (cite Weinberger), with the assumption that tissue collected later experienced more chilling. The challenge with this method is that photoperiod and other factors have also changed during this time. Indeed, we found estimates varied in XX way when derived from direct manipulations of chilling versus the sequential ‘Weinberger’ method. Given the limited manipulation of chilling in studies, we estimated chilling for all studies by combining chilling from the field (i.e., chilling before plants were sampled) and experimental chilling into two widely used metrics of chilling: Utah (citation) and Chill portions (citation). We found the effects of chilling and other cues remain qualitatively the consistent across the two chilling units, though chilling and photoperiod estimates were slightly lower using chill portions compared to Utah (cite supplemental table comparing estimates with both units).

An additional important limitation in most current experiments is the rarity of studies designed to test interactions. While interactions between these cues are widely expected (cites) and, when examined, often found (cites), we were unable to estimate interactions in our meta-analysis because very few studies design experiments to test for interactions between chilling, forcing, and photoperiod (cite table with number of interactions from coding challenge!). The few that do incorporate interactions generally use the Weinberger method, which is not designed to robustly tease out of the effects of multiple cues (cites, Tables, figs). Our estimated effects average over interactions (cite Gelman), but identifying them in future research will be critical to understanding and predicting budburst. For example, the most commonly observed interaction between chilling and forcing—that lower amounts of chilling increase forcing requirements for budburst (cite)—is the hypothesized cause of declining sensitivities in European trees (cites). As more data becomes available, it would allow additional tests of important interactions, such as how responses vary across latitudes (see Supp).

A simple interpretation of our overall model supports hypotheses that chilling and photoperiod cues may underlie declining sensitivities to warming in long-term data observed recently across Europe (cite papers, ref our pep fig?). Under these hypotheses warming increases forcing and thus advances budburst, but such advances become muted if warming also causes declines in chilling and shorter photoperiods experienced near budburst (cite our photoperiod paper? or something else...). Our model supports this superficially as it predicts increased forcing advances budburst while less chilling and shorter photoperiods both delay budburst (Fig. 3), but these estimates integrate across experimental conditions (see FIG). A more useful understanding the implications of our findings requires interpreting them in the context of observed environmental conditions.

Reinterpreting our model using the climate and phenology data that has led to observations of declining temperature sensitivities across Europe suggests chilling and photoperiod cues are highly unlikely to underlie the observed declines. Our model predicts such declines only at extreme warming for most sites (Supplemental Materials for details). [Would be good to say here quickly in one sentence WHY we find this ... ] In contrast to the common hypothesis that chilling declines with warming we found that chilling often increased with small amounts of warming, though this varied with current local climate (Fig. 3 A-D, supp heat map fig).

We also found that patterns of budburst advancement with warming vary considerably depending on the current climate (e.g. how much advancement will continue with warming depends on how much chilling is currently experienced and whether that will increase or decrease with warming). The decline in sensitivity

observed with high warming was due to declines in chilling; photoperiod had little effect on budburst day of year, even for the photosensitive species *F. sylvatica* (Fig. 3 E,F). Portions of Europe have experienced more dramatic warming in winter versus summer (citatopm), but even if warming *only* occurs in the winter, the OSPREE model suggests that delays due to decreased chilling will occur at warming levels of 4°C or more (CHECK THIS!). Given the modest amounts of warming experienced at these sites (XX on average), the OSPREE model predicts a general advance of budburst for Germany.

We compared our model's predictions with observations from the PEP database, and found a different response than predicted by the OSPREE model. The PEP data show declining sensitivities to warming in recent decades (Fig. 4). (Compare advances in our models to PEP725 data?). Our results suggest few sites with delays before 3-4 degrees warming (CHECK) and Germany has warmed X amount. Why do the OSPREE model and PEP observations not match up? Possibly because conditions are different between experiments and natural systems: although forcing conditions are roughly similar between experimental and observed conditions, experimental chilling is generally lower than field chilling. Lizzie- could you write the below points up in a few sentences, probably to stick in here:

1. Speeding up a biological process given sampling time resolution could lead to declining estimates of sensitivities, even if unchanged
2. Say something about what to do about this and how to figure out if this is the issue or it's cues.

Our results suggest most or all studied species are responsive to chilling, forcing, and photoperiod, and we expect climate change to continue to have dramatic effects on spring phenology, especially because the two temperature-derived cues (chilling and forcing) strongly affect budburst (Laube et al., 2014). However, the relative importance of chilling versus forcing (i.e., the extent to which a chilling threshold will be reached and cause delays in budburst with additional warming) will vary spatially. Our results comparing experiments to observations are only for one region, but highlight the critical nature of accurate forecasts of shifts in forcing and chilling at local scales. To do this, we desperately need to better understand chilling (i.e., the process of dormancy release), so that we can accurately predict it in the future. Current common models for chilling (i.e., Utah which was developed for XX species, chill portions which was developed for XX species) are *hypotheses* for how chilling may accumulate to affect the process of dormancy release, but are likely to be inaccurate for many species. Our model highlights how the choice of chill units can affect model estimates and associated forecasts (reference supplemental figure with Utah units vs chill portions in main manuscript). Alongside this, we need improved fundamental understanding of the way these three cues interact. This requires larger studies across diverse species. An improved understanding of interactive cues, however, is unlikely to alter our fundamental predictions of an increasing advance for many temperate trees in the future, even those with strong chilling or forcing cues (Gauzere et al., 2017), unless cues are changing very asynchronously.

## Figures




Figure 1: Budburst model estimates  
for model fit to centered data with Chill Portions.




Figure 2: Effects of chilling, forcing, and photoperiod, across the experimental conditions in the OS-PREE database. Make this part of a 2-panel figure with 1?




Figure 3: Implications of global warming on budburst of *Betula pendula* (A,B) and *Fagus sylvatica* (C,D) as predicted by the OSPREE model. We show the maximum and minimum latitudes at which each species occurred in the PEP database for Germany as examples; these locations differ in current climate. For all sites effects of potential shifts in photoperiod with advancing budburst were minimal (E,F), even the photosensitive species *Fagus sylvatica*




Figure 4: Implications of global warming on budburst of *Betula pendula* (A,B) and *Fagus sylvatica* (C,D) as predicted by the OSPREE model. We show the maximum and minimum latitudes at which each species occurred in the PEP database for Germany as examples; these locations differ in current climate. For all sites effects of potential shifts in photoperiod with advancing budburst were minimal (E,F), even the photosensitive species *Fagus sylvatica*

## Supplemental Materials

### Supplemental methods

1. Forecasting with the OSPREE model: We selected sites in Germany where temperature and budburst have been monitored since the 1950s. We extracted mean temperature data from 1950 through 1980 (pre warming time period) and used these values as baseline data in our model. We then investigated model predictions of budburst given different levels of warming (from 1-7 °C), including altered chilling and forcing as well as potential declines in photoperiod due to advancing phenology. We did this for two common European species: *Betula pendula* (silver birch) and *Fagus sylvatica* at all lat/longs included in the PEP database for Germany.
2. To understand how experimental temperature, photoperiod, and budburst sensitivity compares to past and current conditions in nature, we used data from the PEP database (cite). We summarized forcing, chilling, and budburst day for two common species: *Betula pendula* (silver birch) and *Fagus sylvatica* (European beech) during a pre-warming time-period (1950-1980) and post-warming period (1981-2014?).

### Supplemental figures/tables:



1. Map of study locations, shading or symbol coding for number of cues (Lizzie)
2. Map of species forecasting to justify sites
3. Tables, yes.
4. Heat maps for the main data, including by actual study design and by calculated chilling (our calculations)
5. Photoperiod x latitude effects figure
6. Equation of our model

## Reference list

A few categories:

Papers about contrasting results over what cues matter from growth chamber studies: Basler and Körner (2012, 2014); Caffarra et al. (2011*a*); Caffarra and Donnelly (2011); Caffarra et al. (2011*b*); Heide and Prestud (2005); Koerner and Basler (2010*b*); Laube et al. (2014); Vitasse and Basler (2013); Zohner et al. (2016). Get Nanninga *et al.* 2017: 'Increased exposure to chilling advances the time to budburst in North American tree species' and maybe Malyshev *et al.* 2018 'Temporal photoperiod sensitivity and forcing requirements for budburst in temperate tree seedlings.'

Papers about declining sensitivities (Ailene will update this list): Rutishauser et al. (2008); Fu et al. (2015). Also look for a Wang *et al.* article 'Impacts of global warming on phenology of spring leaf unfolding remain stable in the long run.' Vitasse paper on declining variation across elevation gradient. See Yu et al. (2010), but this is not temperate trees.

Papers about chilling units paper (Lizzie gets a list): Fu 2012 from OSPREE. Harrington and Gould (2015)Luedeling et al. (2011); Luedeling and Brown (2011); Luedeling et al. (2013)

## References

- Basler, D., and C. Körner. 2012. Photoperiod sensitivity of bud burst in 14 temperate forest tree species. *Agricultural and Forest Meteorology* 165:73–81.
- . 2014. Photoperiod and temperature responses of bud swelling and bud burst in four temperate forest tree species. *Tree physiology* 34:377–388.
- Bradley, N. L., A. C. Leopold, J. Ross, and W. Huffaker. 1999. Phenological changes reflect climate change in wisconsin. *Proceedings of the National Academy of Sciences* 96:9701–9704.
- Caffarra, A., and A. Donnelly. 2011. The ecological significance of phenology in four different tree species: effects of light and temperature on bud burst. *International journal of Biometeorology* 55:711–721.
- Caffarra, A., A. Donnelly, and I. Chuine. 2011*a*. Modelling the timing of *Betula pubescens* budburst. II. Integrating complex effects of photoperiod into process-based models. *Climate Research* 46:159–170. Times Cited: 2 Caffarra, Amelia Donnelly, Alison Chuine, Isabelle.

- Caffarra, A., A. Donnelly, I. Chuine, and M. B. Jones. 2011*b*. Modelling the timing of *betula pubescens* bud-burst. i. temperature and photoperiod: A conceptual model. *Climate Research* 46:147.
- Chuine, I. 2000. A unified model for budburst of trees. *Journal of Theoretical Biology* 207:337–347.
- Flynn, D. F. B., and E. M. Wolkovich. 2018. Temperature and photoperiod drive spring phenology across all species in a temperate forest community. *New Phytologist* 0.
- Fraga, H., J. G. Pinto, and J. A. Santos. 2019. Climate change projections for chilling and heat forcing conditions in european vineyards and olive orchards: a multi-model assessment. *Climatic Change* pages 1–15.
- Fu, Y. S. H., H. F. Zhao, S. L. Piao, M. Peaucelle, S. S. Peng, G. Y. Zhou, P. Ciais, M. T. Huang, A. Menzel, J. P. Uelas, Y. Song, Y. Vitasse, Z. Z. Zeng, and I. A. Janssens. 2015. Declining global warming effects on the phenology of spring leaf unfolding. *Nature* 526:104–107.
- Gauzere, J., S. Delzon, H. Davi, M. Bonhomme, I. G. de Cortazar-Atauri, and I. Chuine. 2017. Integrating interactive effects of chilling and photoperiod in phenological process-based models. a case study with two european tree species: *Fagus sylvatica* and *quercus petraea*. *Agricultural and Forest Meteorology* 244:9–20.
- Harrington, C. A., and P. J. Gould. 2015. Tradeoffs between chilling and forcing in satisfying dormancy requirements for pacific northwest tree species. *Frontiers in Plant Science* 6:120.
- Heide, O., and A. Prestrud. 2005. Low temperature, but not photoperiod, controls growth cessation and dormancy induction and release in apple and pear. *Tree Physiology* 25:109–114.
- IPCC. 2014. *Climate Change 2014: Impacts, Adaptation, and Vulnerability*. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.
- Koerner, C., and D. Basler. 2010*a*. Phenology under global warming. *Science* 327:1461–1462.
- . 2010*b*. Warming, photoperiods, and tree phenology response. *Science* 329:278–278.
- Laube, J., T. H. Sparks, N. Estrella, J. Höfler, D. P. Ankerst, and A. Menzel. 2014. Chilling outweighs photoperiod in preventing precocious spring development. *Global Change Biology* 20:170–182.
- Luedeling, E., and P. H. Brown. 2011. A global analysis of the comparability of winter chill models for fruit and nut trees. *International Journal of Biometeorology* 55:411–421.
- Luedeling, E., E. H. Girvetz, M. A. Semenov, and P. H. Brown. 2011. Climate change affects winter chill for temperate fruit and nut trees. *Plos One* 6.
- Luedeling, E., L. Guo, J. H. Dai, C. Leslie, and M. M. Blanke. 2013. Differential responses of trees to temperature variation during the chilling and forcing phases. *Agricultural and Forest Meteorology* 181:33–42.
- Menzel, A., T. H. Sparks, N. Estrella, E. Koch, A. Aasa, R. Ahas, K. Alm-Kuebler, et al. 2006. European phenological response to climate change matches the warming pattern. *Global Change Biology* 12:1969–1976.
- Miller-Rushing, A. J., and R. B. Primack. 2008. Global warming and flowering times in Thoreau’s Concord: A community perspective. *Ecology* 89:332–341.
- Rutishauser, T., J. Luterbacher, C. Defila, D. Frank, and H. Wanner. 2008. Swiss spring plant phenology 2007: Extremes, a multi-century perspective, and changes in temperature sensitivity. *Geophysical Research Letters* 35:L05703. Rutishauser, This Luterbacher, Juerg Defila, Claudio Frank, David Wanner, Heinz.
- Vitasse, Y., and D. Basler. 2013. What role for photoperiod in the bud burst phenology of european beech. *European Journal of Forest Research* 132:1–8.

- Wolkovich, E. M., B. I. Cook, J. M. Allen, T. M. Crimmins, J. L. Betancourt, S. E. Travers, S. Pau, J. Regetz, T. J. Davies, N. J. B. Kraft, T. R. Ault, K. Bolmgren, S. J. Mazer, G. J. McCabe, B. J. McGill, C. Parmesan, N. Salamin, M. D. Schwartz, and E. E. Cleland. 2012. Warming experiments underpredict plant phenological responses to climate change. *Nature* 485:494–497. Wolkovich, E. M. Cook, B. I. Allen, J. M. Crimmins, T. M. Betancourt, J. L. Travers, S. E. Pau, S. Regetz, J. Davies, T. J. Kraft, N. J. B. Ault, T. R. Bolmgren, K. Mazer, S. J. McCabe, G. J. McGill, B. J. Parmesan, C. Salamin, N. Schwartz, M. D. Cleland, E. E.
- Yu, H. Y., E. Luedeling, and J. C. Xu. 2010. Winter and spring warming result in delayed spring phenology on the Tibetan Plateau. *Proceedings of the National Academy of Sciences of the United States of America* 107:22151–22156. Yu, Haiying Luedeling, Eike Xu, Jianchu.
- Zohner, C. M., B. M. Benito, J. C. Svenning, and S. S. Renner. 2016. Day length unlikely to constrain climate-driven shifts in leaf-out times of northern woody plants. *Nature Climate Change* .