

Limiting cues: How spring warming, winter chilling and daylength will shape climate change responses

THE LAB AS IT WAS IN 2017^{1,2}

¹ *Arnold Arboretum of Harvard University, 1300 Centre Street, Boston, Massachusetts, 02131, USA*

² *Organismic & Evolutionary Biology, Harvard University, 26 Oxford Street, Cambridge, Massachusetts, 02138, USA*

³ *Forest & Conservation Sciences, Faculty of Forestry, University of British Columbia, 2424 Main Mall, Vancouver, BC V6T 1Z4*

1 Overview of OSPREE database

We built the Observed Spring Phenology Responses in Experimental Environments (OSPREE) database, by searching both ISI Web of Science and Google Scholar the following terms:

1. TOPIC = (budburst OR leaf-out) AND (photoperiod or daylength) AND temperature*, which yielded 85 publications
2. TOPIC = (budburst OR leaf-out) AND dorman*, which yielded 193 publications

We extracted data (using ImageJ for figures, transcribing values from tables and extracting date, location and other methods from the text) from papers on woody plants that test for photoperiod and/or temperature effects on budburst, leafout, or flowering. Ettinger *et al.* (2020) used a subset of these data (studies on budburst for which we could estimate forcing, chilling and photoperiod treatments), while here we present the full database, capturing data from 84 papers (see Table S1 for a list), of which 21 are focused on crops (*Actinidia deliciosa*, *Malus domestica*, *Vitis vinifera*, *Ribes nigrum*, *Vaccinium ashei*, *Vaccinium corymbosum*, *Prunus persica*). Papers often reported more than one experiment, which we refer to as a ‘study.’ OSPREE includes 136 studies (with the earliest study was conducted in 1947 (Lamb, 1948)), which spanned a variety of plant materials, though studies on ‘seedlings’ (51 studies) and ‘cuttings’ (55 studies) were most common.

2 Trends in experimental treatments over space

Cue levels studied varied across latitude, with a general trend toward examining more extreme values at higher latitudes. Forcing and chilling treatments decline 0.1°C per 1° of latitude (for forcing, min is -0.1, for max it is -0.06, see Fig S3; for chilling it is -0.06 for min and -0.09 for max); and the maximum studied photoperiod increases with latitude (0.09 hr per ° latitude, see Fig S3).

3 Comparing experimental treatments to forecasted trends

To compare the magnitude of experimental treatments to forecasted changes in temperature we calculated treatment differences as the differences within varying forcing and chilling treatments within a single study (e.g., a study with a 1 and 4°C chilling treatment would yield a value of 3°C). We did this across all studies (136 total) and for the 19 studies of *Fagus sylvatica* and 17 studies of *Betula pendula*. We calculated forecasted changes in minimum and maximum average daily temperatures over a 60 day window using RCP8.5 from the NCAR Large Ensemble (LENS, a multi-member ensemble of a single general circulation model, GCM, the Community Earth System Model Kay *et al.*, 2015).

4 References

- Ashby, W. (1962) Germination capacity in American Basswood *Tilia americana*. *Transactions of the Illinois State Academy of Science* **55**, 120–3.
- Basler, D. & Körner, C. (2012) Photoperiod sensitivity of bud burst in 14 temperate forest tree species. *Agricultural and Forest Meteorology* **165**, 73–81.
- Basler, D. & Körner, C. (2014) Photoperiod and temperature responses of bud swelling and bud burst in four temperate forest tree species. *Tree Physiology* **34**, 377–388.
- Biasi, L., Zanette, F. & Carvalho, R. (2012) Dormancy dynamics of grape and kiwifruit buds in a region of low chill occurrence. *XXVIII International Horticultural Congress on Science and Horticulture for People (IHC2010): International Symposium on Plant* 932, pp. 507–512.
- Boyer, J. & South, D. (1986) Dormancy, chilling requirements, and storability of container-grown loblolly pine seedlings .
- Caffarra, A. & Donnelly, A. (2011) The ecological significance of phenology in four different tree species: effects of light and temperature on bud burst. *International Journal of Biometeorology* **55**, 711–721.
- Caffarra, A., Donnelly, A., Chuine, I. & Jones, M.B. (2011) Modelling the timing of *Betula pubescens* bud-burst. I. Temperature and photoperiod: A conceptual model. *Climate Research* **46**, 147.
- Calmé, S., Bigras, F.J., Margolis, H.A. & Hébert, C. (1994) Frost tolerance and bud dormancy of container-grown yellow birch, red oak and sugar maple seedlings. *Tree Physiology* **14**, 1313–1325.
- Campbell, R.K. & Sugano, A.I. (1975) Phenology of bud burst in Douglas-fir related to provenance, photoperiod, chilling, and flushing temperature. *Botanical Gazette* pp. 290–298.
- Cannell, M. & Smith, R. (1983) Thermal time, chill days and prediction of budburst in *Picea sitchensis*. *Journal of applied Ecology* pp. 951–963.
- Charrier, G., Bonhomme, M., Lacoite, A. & Améglio, T. (2011) Are budburst dates, dormancy and cold acclimation in walnut trees (*Juglans regia* L.) under mainly genotypic or environmental control? *International journal of biometeorology* **55**, 763–774.
- Chavarria, G., Herter, F.G., Raseira, M.d.C.B., Rodrigues, A.C., Reisser, C. & Silva, J.B.d. (2009) Mild temperatures on bud breaking dormancy in peaches. *Ciência Rural* **39**, 2016–2021.
- Cook, C. & Jacobs, G. (2000) Progression of apple (*malus × domestica* Borkh.) bud dormancy in two mild winter climates. *The Journal of Horticultural Science and Biotechnology* **75**, 233–236.

- Cook, N.C., Bellen, A., Cronjé, P.J., De Wit, I., Keulemans, W., Van den Putte, A. & Steyn, W. (2005) Freezing temperature treatment induces bud dormancy in ‘granny smith’ apple shoots. *Scientia horticultrae* **106**, 170–176.
- Cronjé, P., Jacobs, G., Sadie, A. & Cook, N. (2003) Quantification of the dormancy progression in terminal apple buds. changes in growth rate and water status. *Advances in horticultural science* pp. 105–110.
- Dantec, C.F., Vitasse, Y., Bonhomme, M., Louvet, J.M., Kremer, A. & Delzon, S. (2014) Chilling and heat requirements for leaf unfolding in European beech and sessile oak populations at the southern limit of their distribution range. *International journal of biometeorology* **58**, 1853–1864.
- De Vries, D., Smeets, L. & Dubois, L.A. (1982) Interaction of temperature and light on growth and development of hybrid tea-rose seedlings, with reference to breeding for low-energy requirements. *Scientia Horticulturae* **17**, 377–382.
- Ettinger, A., Chamberlain, C., Morales-Castilla, I., Buonaiuto, D., Flynn, D., Savas, T., Samaha, J. & Wolkovich, E. (2020) Winter temperatures predominate in spring phenological responses to warming. *Nature Climate Change* pp. 1–6.
- Falusi, M. & Calamassi, R. (1990) Bud dormancy in beech (*Fagus sylvatica* L.). Effect of chilling and photoperiod on dormancy release of beech seedlings. *Tree Physiology* **6**, 429–438.
- Falusi, M. & Calamassi, R. (1996) Geographic variation and bud dormancy in beech seedlings (*Fagus sylvatica* L.). *Annales des Sciences forestières*, vol. 53, pp. 967–979, EDP Sciences.
- Falusi, M. & Calamassi, R. (1997) Bud dormancy in *Fagus sylvatica* L. II. The evolution of dormancy in seedlings and one-node cuttings. *Plant Biosystems-An International Journal Dealing with all Aspects of Plant Biology* **131**, 143–148.
- Falusi, M. & Calamassi, R. (2003) Dormancy of *Fagus sylvatica* L. buds III. Temperature and hormones in the evolution of dormancy in one-node cuttings. *Plant Biosystems-An International Journal Dealing with all Aspects of Plant Biology* **137**, 185–191.
- Fu, Y.H., Campioli, M., Deckmyn, G. & Janssens, I.A. (2013) Sensitivity of leaf unfolding to experimental warming in three temperate tree species. *Agricultural and Forest Meteorology* **181**, 125–132.
- Gansert, D. (2002) *Betula ermanii*, a dominant subalpine and subarctic treeline tree species in japan: ecological traits of deciduous tree life in winter. *Arctic, Antarctic, and Alpine Research* pp. 57–64.
- Ghelardini, L., Santini, A., Black-Samuelsson, S., Myking, T. & Falusi, M. (2010) Bud dormancy release in elm (*ulmus* spp.) clones—a case study of photoperiod and temperature responses. *Tree physiology* **30**, 264–274.
- Gianfagna, T. & Mehlenbacher, S. (1985) Importance of heat requirement for bud break and time of flowering in apple. *HortScience* **20**, 909–911.

- Gömöry, D., Foffová, E., Longauer, R. & Krajmerová, D. (2015) Memory effects associated with early-growth environment in Norway spruce and European larch. *European Journal of Forest Research* **134**, 89–97.
- Granhus, A., Fløistad, I.S. & SØgaard, G. (2009) Bud burst timing in *Picea abies* seedlings as affected by temperature during dormancy induction and mild spells during chilling. *Tree physiology* **29**, 497–503.
- Guak, S., Olszyk, D.M., Fuchigami, L.H. & Tingey, D.T. (1998) Effects of elevated CO₂ and temperature on cold hardiness and spring bud burst and growth in douglas-fir (*Pseudotsuga menziesii*). *Tree Physiology* **18**, 671–679.
- Guerriero, P., Scalabrelli, G. & Grazzini, G. (1990) Chilling effect on inhibition removal in kiwifruit dormant lateral buds. *I International Symposium on Kiwifruit 282*, pp. 79–86.
- Gunderson, C.A., Edwards, N.T., Walker, A.V., O'Hara, K.H., Champion, C.M. & Hanson, P.J. (2012) Forest phenology and a warmer climate—growing season extension in relation to climatic provenance. *Global Change Biology* **18**, 2008–2025.
- Hawerroth, F.J., Herter, F.G., Petri, J.L., Marafon, A.C. & Leonetti, J.F. (2013) Evaluation of winter temperatures on apple budbreak using grafted twigs. *Revista Brasileira de Fruticultura* **35**, 713–721.
- Hawkins, C.D. & Dhar, A. (2012) Spring bud phenology of 18 *Betula papyrifera* populations in British Columbia. *Scandinavian Journal of Forest Research* **27**, 507–519.
- Heide, O. (1993a) Daylength and thermal time responses of budburst during dormancy release in some northern deciduous trees. *Physiologia Plantarum* **88**, 531–540.
- Heide, O. (1993b) Dormancy release in beech buds (*Fagus sylvatica*) requires both chilling and long days. *Physiologia Plantarum* **89**, 187–191.
- Heide, O. (2003) High autumn temperature delays spring bud burst in boreal trees, counterbalancing the effect of climatic warming. *Tree Physiology* **23**, 931–936.
- Heide, O. & Prestrud, A. (2005) Low temperature, but not photoperiod, controls growth cessation and dormancy induction and release in apple and pear. *Tree Physiology* **25**, 109–114.
- Heide, O.M. (2008) Interaction of photoperiod and temperature in the control of growth and dormancy of *Prunus* species. *Scientia Horticulturae* **115**, 309–314.
- Heide, O.M. (2011) Temperature rather than photoperiod controls growth cessation and dormancy in *Sorbus* species. *Journal of Experimental Botany* **62**, 5397–5404.
- Heide, O.M. & Sønsteby, A. (2012) Floral initiation in black currant cultivars (*Ribes nigrum* L.): Effects of plant size, photoperiod, temperature, and duration of short day exposure. *Scientia Horticulturae* **138**, 64–72.
- Heide, O.M. & Sonsteby, A. (2015) Simultaneous dormancy induction interferes with short day floral induction in black currant (*Ribes nigrum* L.). *Scientia Horticulturae* **185**, 228–232.

- Howe, G.T., Hackett, W.P., Furnier, G.R. & Klevorn, R.E. (1995) Photoperiodic responses of a northern and southern ecotype of black cottonwood. *Physiologia Plantarum* **93**, 695–708.
- Jones, H., Hillis, R., Gordon, S. & Brennan, R. (2012) An approach to the determination of winter chill requirements for different *Ribes* cultivars. *Plant Biology* **15**, 18–27.
- Junttila, O. & Hänninen, H. (2012) The minimum temperature for budburst in *Betula* depends on the state of dormancy. *Tree physiology* **32**, 337–345.
- Karlsson, P., Bylund, H., Neuvonen, S., Heino, S. & Tjus, M. (2003) Climatic response of budburst in the mountain birch at two areas in northern fennoscandia and possible responses to global change. *Ecography* **26**, 617–625.
- Kay, J.E., Deser, C., Phillips, A., Mai, A., Hannay, C., Strand, G., Arblaster, J.M., Bates, S.C., Danabasoglu, G., Edwards, J., Holland, M., Kushner, P., Lamarque, J.F., Lawrence, D., Lindsay, K., Middleton, A., Munoz, E., Neale, R., Oleson, K., Polvani, L. & Vertenstein, M. (2015) The Community Earth System Model (CESM) Large Ensemble Project A community resource for studying climate change in the presence of internal climate variability. *Bulletin of the American Meteorological Society* **96**, 1333–1349.
- Lamb, R.C. (1948) Effect of temperatures above and below freezing on the breaking of rest in the latham raspberry. *Proceedings of the American Society for Horticultural Science*, vol. 51, pp. 313–315, AMER SOC HORTICULTURAL SCIENCE 701 NORTH SAINT ASAPH STREET, ALEXANDRIA, VA 22314-1998.
- Laube, J., Sparks, T.H., Estrella, N., Höfler, J., Ankerst, D.P. & Menzel, A. (2014a) Chilling outweighs photoperiod in preventing precocious spring development. *Global Change Biology* **20**, 170–182.
- Laube, J., Sparks, T.H., Estrella, N. & Menzel, A. (2014b) Does humidity trigger tree phenology? Proposal for an air humidity based framework for bud development in spring. *New Phytologist* **202**, 350–355.
- Li, C., Welling, A., Puhakainen, T., Viherä-Aarnio, A., Ernsten, A., Junttila, O., Heino, P. & Palva, E.T. (2005) Differential responses of silver birch (*Betula pendula*) ecotypes to short-day photoperiod and low temperature. *Tree physiology* **25**, 1563–1569.
- Linkosalo, T. & Lechowicz, M.J. (2006) Twilight far-red treatment advances leaf bud burst of silver birch (*Betula pendula*). *Tree Physiology* **26**, 1249–1256.
- Man, R. & Lu, P. (2010) Effects of thermal model and base temperature on estimates of thermal time to bud break in white spruce seedlings. *Canadian Journal of Forest Research* **40**, 1815–1820.
- Manson, P. & Snelgar, W. (1991) Effect of time of budburst and apical shoot growth on flower production in kiwifruit. *New Zealand Journal of Crop and Horticultural Science* **19**, 441–445.
- Morin, X., Roy, J., Sonié, L. & Chuine, I. (2010) Changes in leaf phenology of three European oak species in response to experimental climate change. *New Phytologist* **186**, 900–910.

- Myking, T. (1997) Effects of constant and fluctuating temperature on time to budburst in *Betula pubescens* and its relation to bud respiration. *Trees* **12**, 107–112.
- Myking, T. (1998) Interrelations between respiration and dormancy in buds of three hardwood species with different chilling requirements for dormancy release. *Trees* **12**, 224–229.
- Myking, T. & Heide, O. (1995) Dormancy release and chilling requirement of buds of latitudinal ecotypes of *Betula pendula* and *B. pubescens*. *Tree Physiology* **15**, 697–704.
- Nienstaedt, H. (1966) Dormancy and dormancy release in white spruce. *Forest Science* **12**, 374–384.
- Nishimoto, N. & Fujisaki, M. (1994) Chilling requirement of buds of some deciduous fruits grown in southern japan and the means to break dormancy. *Dormancy and the related Problems of Deciduous Fruit Trees* 395 pp. 153–160.
- Okie, W.R. & Blackburn, B. (2011) Interactive effects of light and chilling on peach flower and leaf budbreak. *HortScience* **46**, 1056–1062.
- Pagter, M., Andersen, U.B. & Andersen, L. (2015) Winter warming delays dormancy release, advances budburst, alters carbohydrate metabolism and reduces yield in a temperate shrub. *AoB plants* **7**, plv024.
- Partanen, J., Hänninen, H. & Häkkinen, R. (2005) Bud burst in Norway spruce (*Picea abies*): preliminary evidence for age-specific rest patterns. *Trees* **19**, 66–72.
- Partanen, J., Koski, V. & Hänninen, H. (1998) Effects of photoperiod and temperature on the timing of bud burst in Norway spruce (*Picea abies*). *Tree Physiology* **18**, 811–816.
- Partanen, J., Leinonen, I. & Repo, T. (2001) Effect of accumulated duration of the light period on bud burst in Norway spruce (*Picea abies*) of varying ages. *Silva Fennica* **35**, 111–117.
- Pettersen, H. (1972) Effect of temperature and daylength on shoot growth and bud formation in azaleas. *Journal of the American Society for Horticultural Science* **97**, 17.
- Pop, E.W., Oberbauer, S.F. & Starr, G. (2000) Predicting vegetative bud break in two arctic deciduous shrub species, *salix pulchra* and *betula nana*. *Oecologia* **124**, 176–184.
- Ramos, A. & Rallo, L. (1999) Effect of the bearing condition of the tree, chilling and defoliation on the forced budburst of olive cuttings at different temperatures. *III International Symposium on Olive Growing 474* (eds. I.T. Metzidakis & D.G. Voyiatzis), pp. 251–254.
- Rinne, P., Hänninen, H., Kaikuranta, P., Jalonen, J. & Repo, T. (1997) Freezing exposure releases bud dormancy in *Betula pubescens* and *B. pendula*. *Plant, Cell & Environment* **20**, 1199–1204.
- Rinne, P., Saarelainen, A. & Junttila, O. (1994) Growth cessation and bud dormancy in relation to ABA level in seedlings and coppice shoots of *Betula pubescens* as affected by a short photoperiod, water stress and chilling. *Physiologia Plantarum* **90**, 451–458.

- Ruesink, J. (1998) Long day treatment prevents flower bud formation in pieris. *Gartenbauwissenschaft (Germany)* .
- Sanz-Pérez, V. & Castro-Díez, P. (2010) Summer water stress and shade alter bud size and budburst date in three Mediterranean *Quercus* species. *Trees* **24**, 89–97.
- Sanz-Perez, V., Castro-Diez, P. & Valladares, F. (2009) Differential and interactive effects of temperature and photoperiod on budburst and carbon reserves in two co-occurring Mediterranean oaks. *Plant Biology* **11**, 142–51.
- Schnabel, B.J. & Wample, R.L. (1987) Dormancy and cold hardiness in *Vitis vinifera* L. cv. White Riesling as influenced by photoperiod and temperature. *American Journal of Enology and Viticulture* **38**, 265–272.
- Skre, O., Taulavuori, K., Taulavuori, E., Nilsen, J., Igeland, B. & Laine, K. (2008) The importance of hardening and winter temperature for growth in mountain birch populations. *Environmental and experimental botany* **62**, 254–266.
- Skuterud, R. & Dietrichson, J. (1994) Budburst in detached birch shoots (*Betula pendula*) of different varieties winter-stored in darkness at three different temperatures. *Silva Fennica* **28**, 223–224.
- Søgaard, G., Johnsen, Ø., Nilsen, J. & Junttila, O. (2008) Climatic control of bud burst in young seedlings of nine provenances of norway spruce. *Tree Physiology* **28**, 311–320.
- Sønsteby, A. & Heide, O. (2013) Variation in seasonal timing of flower bud initiation in black currant (*ribes nigrum* l.) cultivars of contrasting geographic origin. *The Journal of Horticultural Science and Biotechnology* **88**, 403–408.
- Sønsteby, A. & Heide, O.M. (2014) Chilling requirements of contrasting black currant (*Ribes nigrum* L.) cultivars and the induction of secondary bud dormancy. *Scientia Horticulturae* **179**, 256–265.
- Spiers, J. & Draper, A. (1974) Effect of chilling on bud break in rabbiteye blueberry cultivars. *Journal American Society for Horticultural Science* .
- Swartz, H. & Powell Jr, L. (1981) The effect of long chilling requirement on time of bud break in apple. *Symposium on Growth Regulators in Fruit Production 120*, pp. 173–178.
- Thielges, B. & Beck, R. (1976) Control of bud break and its inheritance in *Populus deltoides*. *Tree Physiology and Yield Improvement* **14**, 253–259.
- Viherä-Aarnio, A., Häkkinen, R. & Junttila, O. (2006) Critical night length for bud set and its variation in two photoperiodic ecotypes of *Betula pendula*. *Tree Physiology* **26**, 1013–1018.
- Webb, D.P. (1977) Root regeneration and bud dormancy of sugar maple, silver maple, and white ash seedlings: effects of chilling. *Forest Science* **23**, 474–483.
- Worrall, J. & Mergen, F. (1967) Environmental and genetic control of dormancy in *Picea abies*. *Physiologia Plantarum* **20**, 733–745.

- Yazdaniha, A. (1967) *Effects of Chilling, Chemicals and Pruning on the Rest Period of Peach Trees*. Master's thesis, Utah State University, <http://digitalcommons.usu.edu/etd/3596>.
- Zohner, C.M., Benito, B.M., Svenning, J.C. & Renner, S.S. (2016) Day length unlikely to constrain climate-driven shifts in leaf-out times of northern woody plants. *Nature Climate Change* **6**, 1120–1123.

Supplemental Tables

Table S1: **Dataset names and references for papers in the OS-PREE database.**

Dataset	Reference
ashby62	(Ashby, 1962)
basler12	(Basler & Körner, 2012)
basler14	(Basler & Körner, 2014)
biasi12	(Biasi <i>et al.</i> , 2012)
boyer	(Boyer & South, 1986)
caffarra11a	(Caffarra & Donnelly, 2011)
caffarra11b	(Caffarra <i>et al.</i> , 2011)
calme94	(Calmé <i>et al.</i> , 1994)
campbell75	(Campbell & Sugano, 1975)
cannell83	(Cannell & Smith, 1983)
charrier11	(Charrier <i>et al.</i> , 2011)
chavarria09	(Chavarria <i>et al.</i> , 2009)
cook00b	(Cook & Jacobs, 2000)
cook05	(Cook <i>et al.</i> , 2005)
cronje03	(Cronjé <i>et al.</i> , 2003)
dantec14	(Dantec <i>et al.</i> , 2014)
devries82	(De Vries <i>et al.</i> , 1982)
falusi03	(Falusi & Calamassi, 2003)
falusi90	(Falusi & Calamassi, 1990)
falusi96	(Falusi & Calamassi, 1996)
falusi97	(Falusi & Calamassi, 1997)
fu13	(Fu <i>et al.</i> , 2013)
gansert02	(Gansert, 2002)
ghelardini10	(Ghelardini <i>et al.</i> , 2010)
gianfagna85	(Gianfagna & Mehlenbacher, 1985)
gomory15	(Gömöry <i>et al.</i> , 2015)
granhus09	(Granhus <i>et al.</i> , 2009)
guak98	(Guak <i>et al.</i> , 1998)
guerriero90	(Guerriero <i>et al.</i> , 1990)
gunderson12	(Gunderson <i>et al.</i> , 2012)
hawerorth13	(Hawerorth <i>et al.</i> , 2013)
hawkins12	(Hawkins & Dhar, 2012)
heide03	(Heide, 2003)
heide05	(Heide & Prestrud, 2005)
heide08	(Heide, 2008)
heide11	(Heide, 2011)
heide12	(Heide & Sønsteby, 2012)
heide15	(Heide & Sonsteby, 2015)
heide93	(Heide, 1993a)
heide93a	(Heide, 1993b)
howe95	(Howe <i>et al.</i> , 1995)
jones12	(Jones <i>et al.</i> , 2012)
junttila12	(Junttila & Hänninen, 2012)
karlsson03	(Karlsson <i>et al.</i> , 2003)
lamb37	(Lamb, 1948)
laube14a	(Laube <i>et al.</i> , 2014a)
laube14b	(Laube <i>et al.</i> , 2014b)

Table S1: **Dataset names and references for papers in the OS-PREE database.**

Dataset	Reference
li05	(Li <i>et al.</i> , 2005)
linkosalo06	(Linkosalo & Lechowicz, 2006)
man10	(Man & Lu, 2010)
manson91	(Manson & Snelgar, 1991)
morin10	(Morin <i>et al.</i> , 2010)
myking95	(Myking & Heide, 1995)
myking97	(Myking, 1997)
myking98	(Myking, 1998)
nienstaedt66	(Nienstaedt, 1966)
nishimoto95	(Nishimoto & Fujisaki, 1994)
okie11	(Okie & Blackburn, 2011)
pagter15	(Pagter <i>et al.</i> , 2015)
partanen01	(Partanen <i>et al.</i> , 2001)
partanen05	(Partanen <i>et al.</i> , 2005)
partanen98	(Partanen <i>et al.</i> , 1998)
pettersen71	(Pettersen, 1972)
pop2000	(Pop <i>et al.</i> , 2000)
ramos99	(Ramos & Rallo, 1999)
rinne94	(Rinne <i>et al.</i> , 1994)
rinne97	(Rinne <i>et al.</i> , 1997)
ruesink98	(Ruesink, 1998)
sanx-perez09	(Sanz-Perez <i>et al.</i> , 2009)
sanx-perez10	(Sanz-Pérez & Castro-Díez, 2010)
schnabel87	(Schnabel & Wample, 1987)
skre08	(Skre <i>et al.</i> , 2008)
skuterud94	(Skuterud & Dietrichson, 1994)
sogaard08	(Søgaard <i>et al.</i> , 2008)
sonsteby13	(Sønsteby & Heide, 2013)
sonsteby14	(Sønsteby & Heide, 2014)
spiers74	(Spiers & Draper, 1974)
swartz81	(Swartz & Powell Jr, 1981)
thielges75	(Thielges & Beck, 1976)
viheraaarnio06	(Viherä-Aarnio <i>et al.</i> , 2006)
webb78	(Webb, 1977)
worrall67	(Worrall & Mergen, 1967)
yazdaniha64	(Yazdaniha, 1967)
zohner16	(Zohner <i>et al.</i> , 2016)




Figure S1: We reviewed seven decades of controlled environment studies, from Lamb (1948) to Zohner *et al.* (2016), conducted across the globe generally on 1-3 species in each experiment (size of circles and exact number of species given after each study). See Table S1 for references for each ‘Dataset.’




Figure S2: Prevalence of number of cues (chilling, forcing, photoperiod) manipulated in studies over time over time.




Figure S3: Experimental treatments correlate with latitude. Here we show the average latitude of a study (averaged over all latitudes from which tissue was taken) versus the minimum (upper left) and maximum (upper right) forcing treatments and the maximum (lower left) photoperiod treatment and range of photoperiod treatments (lower right, range calculated as maximum versus minimum treatments in a study). Colors represent unique datasets (see Table S1).