# Describing phenology with beta, beta-binomial, and beta-hypergeometric distributions

**Geoffrey Legault** 

University of British Columbia

# Characterizing phenology

# CONCEPTS & SYNTHESIS

EMPHASIZING NEW IDEAS TO STIMULATE RESEARCH IN ECOLOGY


Ecological Monographs, 89(2), 2019, e01352 © 2019 by the Ecological Society of America

Phenology as a process rather than an event: from individual reaction norms to community metrics

BRIAN D. INOUYE D, 1,2,3,5 JOHAN EHRLÉN, 2,4 AND NORA UNDERWOOD 1,2,3

- Phenology is usually characterized with discrete measures such as first flowering
- ► This approach can obscure many important features
- More useful to describing phenology as a distribution

# Phenology as a distribution



Source: Inouye et al. 2019

# Advantages of phenology as a distribution

- More comprehensive (accurate?) description
- Use more of the available data/information
- Fit the distribution to data
- Parameters of the distribution will (hopefully) be interpretable
- Given a distribution, we can estimate measures like mean, variance, correlation
- May allow comparisons over time/environments

## Objective

Find an appropriate, general, and above all interpretable distribution for describing phenology

#### Where to start

▶ With the available data, of course

e.g., percent budburst over time


#### Where to start


With the available data, of course
 e.g., percent budburst over time

Probability theory

#### Where to start


- With the available data, of course
  e.g., percent budburst over time
- Probability theory
- Beta distribution






- Only two parameters (that's nice)
- Effects of  $\alpha$  and  $\beta$  are fairly interpretable

- Only two parameters (that's nice)
- Effects of  $\alpha$  and  $\beta$  are fairly interpretable
- Problem 1: x ranges between 0 and 1
- Problem 2: Area under the curve is 1







- 2 shape parameters, 2 "interval" parameters
- Parameters are still fairly interpretable


- 2 shape parameters, 2 "interval" parameters
- Parameters are still fairly interpretable
- Problem 1: x ranges between 0 and 1


Solved!

- 2 shape parameters, 2 "interval" parameters
- Parameters are still fairly interpretable
- Problem 1: x ranges between 0 and 1 Solved!
- Problem 2: Area under the curve is 1
- ► Problem 3: Assumes data are continuous









- Compound distribution
- ▶ Binomial random variable, B(n, p), p is beta-distributed

▶ PMF of binomial random variable:

$$P(X = k; p, n) = \binom{n}{k} p^k (1 - p)^{n-k}$$

PMF of binomial random variable:

$$P(X = k; p, n) = \binom{n}{k} p^k (1 - p)^{n-k}$$

▶ PDF of a beta random variable:

$$f_P(p)=rac{p^{lpha-1}(1-p)^{eta-1}}{B(lpha,eta)}$$

PMF of binomial random variable:

$$P(X = k; p, n) = \binom{n}{k} p^k (1 - p)^{n-k}$$

▶ PDF of a beta random variable:

$$f_P(p) = \frac{p^{\alpha-1}(1-p)^{\beta-1}}{B(\alpha,\beta)}$$

PMF of compound distribution:

$$Pr(k; \boldsymbol{n}, \alpha, \beta) = \binom{n}{k} \frac{1}{B(\alpha, \beta)} \int_0^1 p^{k+\alpha-1} (1-p)^{n-k+\beta-1} dp$$

PMF of binomial random variable:

$$P(X = k; p, n) = \binom{n}{k} p^k (1 - p)^{n-k}$$

▶ PDF of a beta random variable:

$$f_P(p) = \frac{p^{\alpha-1}(1-p)^{\beta-1}}{B(\alpha,\beta)}$$

PMF of compound distribution:

$$Pr(k; n, \alpha, \beta) = \binom{n}{k} \frac{1}{B(\alpha, \beta)} \int_0^1 p^{k+\alpha-1} (1-p)^{n-k+\beta-1} dp$$

- Problem: This uses the <u>standardized</u> beta distribution (range 0 to 1)
- We probably want the <u>beta-hypergeometric</u> (which doesn't exist yet, but soon will!)