Supplemental materials for Chilling outweighs photoperiod and forcing cues for temperate trees in experiments, but not in natural systems

A.K. Ettinger, C. Chamberlain, I. Morales-Castilla, D. Buonaiuto, D. Flynn, T. Savas, J. Samaha & E. Wolkovich

May 8, 2019

Supplemental Methods

- 1. Equation of our model
- 2. Forecasting with the OSPREE model: We selected sites in Germany where temperature and budburst have been monitored since the 1950s. We extracted mean temperature data from 1950 through 1980 (pre warming time period) and used these values as baseline data in our model. We then investigated model predictions of budburst given different levels of warming (from 1-7 °C), including altered chilling and forcing as well as potential declines in photoperiod due to advancing phenology. We did this for two common European species: Betula pendula (silver birch) and Fagus sylvatica at all lat/longs included in the PEP database for Germany.
- 3. To understand how experimental temperature, photoperiod, and budburst sensitivity compares to past and current conditions in nature, we used data from the PEP database (cite). We summarized forcing, chilling, and budburst doy for two common species: Betula pendula (silver birch) and Fagus sylvatica (European beech) during a pre-warming time-period (1950-1980) and post-warming period (1981-2014?).

Supplemental figures/tables:

- 1. Map of study locations, shading or symbol coding for number of cues (Lizzie)
- 2. Map of species forecasting to justify sites
- 3. Heat maps for the main data, including by actual study design and by calculated chilling (our calculations)
- 4. Photoperiod x latitude effects figure

Reference list

A few categories:

Papers about contrasting results over what cues matter from growth chamber studies: Basler and Körner (2012, 2014); Caffarra et al. (2011a); Caffarra and Donnelly (2011); Caffarra et al. (2011b); Heide and Prestrud (2005); Koerner and Basler (2010); Laube et al. (2014); Vitasse and Basler (2013); Zohner et al. (2016). Get Nanninga et al. 2017: 'Increased exposure to chilling advances the time to budburst in North American tree species' and maybe Malyshev et al. 2018 'Temporal photoperiod sensitivity and forcing requirements for budburst in temperate tree seedlings.'

Papers about declining sensitivities (Ailene will update this list): Rutishauser et al. (2008); Fu et al. (2015). Also look for a Wang et al. article 'Impacts of global warming on phenology of spring leaf unfolding remain stable in the long run.' Vitasse paper on declining variation across elevation gradient. See Yu et al. (2010), but this is not temperate trees.

Papers about chilling units paper (Lizzie gets a list): Fu 2012 from OSPREE. Harrington and Gould (2015)Luedeling et al. (2011); Luedeling and Brown (2011); Luedeling et al. (2013)

References

- Basler, D., and C. Körner. 2012. Photoperiod sensitivity of bud burst in 14 temperate forest tree species. Agricultural and Forest Meteorology 165:73–81.
- ———. 2014. Photoperiod and temperature responses of bud swelling and bud burst in four temperate forest tree species. Tree physiology 34:377–388.
- Caffarra, A., and A. Donnelly. 2011. The ecological significance of phenology in four different tree species: effects of light and temperature on bud burst. International journal of Biometeorology 55:711–721.
- Caffarra, A., A. Donnelly, and I. Chuine. 2011a. Modelling the timing of *Betula pubescens* budburst. II. Integrating complex effects of photoperiod into process-based models. Climate Research 46:159–170. Times Cited: 2 Caffarra, Amelia Donnelly, Alison Chuine, Isabelle.
- Caffarra, A., A. Donnelly, I. Chuine, and M. B. Jones. 2011b. Modelling the timing of betula pubescens bud-burst. i. temperature and photoperiod: A conceptual model. Climate Research 46:147.
- Fu, Y. S. H., H. F. Zhao, S. L. Piao, M. Peaucelle, S. S. Peng, G. Y. Zhou, P. Ciais, M. T. Huang, A. Menzel, J. P. Uelas, Y. Song, Y. Vitasse, Z. Z. Zeng, and I. A. Janssens. 2015. Declining global warming effects on the phenology of spring leaf unfolding. Nature 526:104–107.
- Harrington, C. A., and P. J. Gould. 2015. Tradeoffs between chilling and forcing in satisfying dormancy requirements for pacific northwest tree species. Frontiers in Plant Science 6:120.
- Heide, O., and A. Prestrud. 2005. Low temperature, but not photoperiod, controls growth cessation and dormancy induction and release in apple and pear. Tree Physiology 25:109–114.
- Koerner, C., and D. Basler. 2010. Warming, photoperiods, and tree phenology response. Science 329:278–278.
- Laube, J., T. H. Sparks, N. Estrella, J. Höfler, D. P. Ankerst, and A. Menzel. 2014. Chilling outweighs photoperiod in preventing precocious spring development. Global Change Biology 20:170–182.
- Luedeling, E., and P. H. Brown. 2011. A global analysis of the comparability of winter chill models for fruit and nut trees. International Journal of Biometeorology 55:411–421.
- Luedeling, E., E. H. Girvetz, M. A. Semenov, and P. H. Brown. 2011. Climate change affects winter chill for temperate fruit and nut trees. Plos One 6.

- Luedeling, E., L. Guo, J. H. Dai, C. Leslie, and M. M. Blanke. 2013. Differential responses of trees to temperature variation during the chilling and forcing phases. Agricultural and Forest Meteorology 181:33–42.
- Rutishauser, T., J. Luterbacher, C. Defila, D. Frank, and H. Wanner. 2008. Swiss spring plant phenology 2007: Extremes, a multi-century perspective, and changes in temperature sensitivity. Geophysical Research Letters 35:L05703. Rutishauser, This Luterbacher, Juerg Defila, Claudio Frank, David Wanner, Heinz.
- Vitasse, Y., and D. Basler. 2013. What role for photoperiod in the bud burst phenology of european beech. European Journal of Forest Research 132:1–8.
- Yu, H. Y., E. Luedeling, and J. C. Xu. 2010. Winter and spring warming result in delayed spring phenology on the Tibetan Plateau. Proceedings of the National Academy of Sciences of the United States of America 107:22151–22156. Yu, Haiying Luedeling, Eike Xu, Jianchu.
- Zohner, C. M., B. M. Benito, J. C. Svenning, and S. S. Renner. 2016. Day length unlikely to constrain climate-driven shifts in leaf-out times of northern woody plants. Nature Climate Change .