



Supplementary Material

Phylogenetic estimates of species-level phenology improve  
ecological forecasting

May 5, 2023

Authors:

The Wolkovich Lab in 2019 & collaborators <sup>1,2,3,4</sup>

*Author affiliations:*

<sup>1</sup>Forest & Conservation Sciences, Faculty of Forestry, University of British Columbia, 2424 Main Mall, Vancouver, BC V6T 1Z4;

<sup>2</sup>Arnold Arboretum of Harvard University, 1300 Centre Street, Boston, Massachusetts, USA;

<sup>3</sup>Organismic & Evolutionary Biology, Harvard University, 26 Oxford Street, Cambridge, Massachusetts, USA;

<sup>4</sup>Edificio Ciencias, Campus Universitario 28805 Alcalá de Henares, Madrid, Spain

\*Corresponding author: ignacio.moralesc@uah.es

## Extended Methods

### Interpretation of $\lambda_j$ and $\sigma_j^2$ on slopes and intercepts

Most current phylogenetic regression approaches aimed at controlling for phylogenetic non-independence of analysis units (i.e. species, see ?) assume the  $\lambda$  scaling parameter is constant across the full set of predictors in the model. Thus,  $\lambda$  is estimated as a single parameter based on one single residual term VCV matrix. While useful for correcting for phylogenetic non-independence this approach does not allow the phylogeny to differentially affect different predictors (i.e. environmental cues in our example). In models with multiple cues, species responses to all cues are estimated as similarly phylogenetically structured, but this may not be the case. For example, in a PGLS model with three cues, it would be possible to have a high (i.e. close to 1) value of  $\lambda$ , due to either a strong phylogenetic signal in the response, but no phylogenetic structuring in the cues, or one or more predictors being strongly phylogenetically structured. In the latter case, phylogenetic structuring of responses to cues could be correlated (i.e., responses to cues evolving in a correlated fashion) or uncorrelated (i.e., independent evolution of responses to cues). Discerning these different situations is not trivial as they would inform whether responses to predictors configure in a structured fashion along the evolutionary process. However, most current approaches act as a black box regarding this information; they simply inform whether or not model residuals are phylogenetically structured (i.e. in PGLS) or the amount of model variance attributable to the phylogeny and independent from other sources of variation (i.e., in PMM, see ?).

Because we are specifically interested in estimating the phylogenetic structure of each cue, our approach explicitly partitions variance into specific components relative to the model intercept and predictor (cue) slopes (see equation ??). The multivariate normal distributions of the intercept and slope terms include each a variance term (see equation ??), modelled with a  $\lambda$  scaling parameter. The interpretation of  $\lambda$ s in our models are analogous to Pagel’s  $\lambda$  (?) parameter (?), constrained to range from 0 to 1, with values of 0 indicating absence of phylogenetic relatedness, and values of 1 indicating *Brownian Motion* evolution (BM). Estimated  $\lambda$ s are not fully equivalent to computing phylogenetic signal of the slopes of each cue separately (i.e., fitting a multilevel regression model with species as a grouping factor on intercepts, and subsequently estimating phylogenetic signal for model slopes). Instead, they are a relative metric of phylogenetic relatedness allowing us to compare among responses known to interact with each other and estimated simultaneously. This approach has the further benefit of adjusting our partial pooling (‘random effect’ of species) based on evolutionary distance, more strongly pooling closely related species, and only weakly pooling distantly related species (see Gaussian process models in ?).

A traditional interpretation of  $\sigma^2$ s under Brownian Motion evolution, is an ‘evolutionary rate’

or phenotypic accumulation over time (?). In PGLS,  $\sigma_\epsilon^2$  is estimated for the model error term, which is distributed as a multivariate normal with VCV matrix given by  $\sigma_\epsilon^2 \mathbf{\Sigma}_i$ . Here, similar to our approach to  $\lambda$ , we estimate four  $\sigma^2$  values, corresponding to each model parameter. In our particular case (i.e., modelling a phenological response to three environmental cues),  $\sigma_\alpha^2$  for the intercept could be interpreted as the phenological variation across species accumulated along evolution independently from the cues. The  $\sigma_{\beta_1}^2$ ,  $\sigma_{\beta_2}^2$ , and  $\sigma_{\beta_3}^2$ , corresponding to model slopes, would represent the phylogenetic variance linked to species responses to each of the modelled cues (i.e., forcing, chilling, and photoperiod, respectively). This is, the variability in how species shift their phenology responding to temperature and light, accumulated along the evolutionary process and considered in concert.

## 0.1 Forecasting for two species

We forecasted estimated shifts in species phenologies from our phylogenetic and non-phylogenetic models for two species with overlapping European ranges to show the impact of differences across models in a well (*Betula pendula*) versus poorly sampled (*Acer campestre*) species. For this, we fit our models using natural (not-scored units) then applied a 2°warming to historical data from XXXX-XXXX for each gridcell in the range (ranges extracted from XXX) using XXX climate data, recalculated chilling units and projected the change in budburst date.

## Supporting Figures and Tables

Table S1: Estimated sensitivities of 191 tree species to three environmental cues: chilling (b.chill), forcing (b.force) and photoperiod (b.photo), along with their corresponding 2.5% (low) and 97.5% (up) Credible Intervals. Values correspond to the full model accounting for phylogenetic relationships.

Species	b.chill	low	up	b.force	low	up	b.photo	low	up
<i>Populus deltoides</i>	-15.16	-25.05	-5.82	-6.43	-13.73	1.82	-0.97	-5.39	3.23
<i>Populus koreana</i>	-8.65	-18.98	1.84	-7.47	-15.61	1.11	-0.87	-5.03	3.22
<i>Populus tremula</i>	-1.94	-7.35	3.60	-7.41	-10.54	-4.23	0.05	-2.11	2.28
<i>Populus grandidentata</i>	-14.08	-21.56	-6.90	-9.98	-15.90	-4.64	-1.18	-5.12	2.65
<i>Euonymus europaeus</i>	-3.61	-15.78	8.88	-6.23	-17.68	5.35	-1.15	-5.80	3.39
<i>Euonymus latifolius</i>	1.12	-9.12	11.97	-6.32	-17.69	5.22	-1.01	-5.32	3.41
<i>Nothofagus antarctica</i>	-10.16	-22.40	1.95	-6.62	-17.37	4.52	-1.63	-6.10	2.79
<i>Carya cordiformis</i>	-15.52	-25.34	-6.01	-4.51	-14.91	6.16	-2.65	-7.00	1.57
<i>Carya laciniosa</i>	-17.48	-27.47	-7.95	-4.03	-14.28	6.79	-1.72	-6.04	2.60
<i>Alnus incana</i>	-1.73	-5.59	2.12	-10.18	-14.56	-5.96	-0.52	-2.82	1.82
<i>Alnus glutinosa</i>	-5.85	-12.14	0.49	-10.25	-15.37	-5.24	-0.60	-3.06	1.93
<i>Alnus maximowiczii</i>	-5.35	-14.90	4.33	-7.32	-16.89	2.55	-1.05	-5.26	3.15
<i>Betula nana</i>	-5.76	-14.88	3.60	-5.15	-13.97	3.61	-0.92	-4.90	3.08
<i>Betula pendula</i>	-4.33	-5.44	-3.23	-1.38	-3.58	0.76	-0.42	-1.41	0.59
<i>Betula pubescens</i>	-2.13	-3.47	-0.83	-6.64	-9.29	-4.11	-0.75	-1.85	0.33
<i>Betula populifolia</i>	-5.30	-14.54	4.00	-5.48	-14.46	3.60	-0.86	-4.88	3.16
<i>Betula papyrifera</i>	-27.28	-33.48	-21.06	-5.69	-10.02	-1.32	1.25	-2.24	5.13
<i>Betula alleghaniensis</i>	-11.23	-18.21	-4.10	-6.99	-10.69	-3.42	-1.47	-5.48	2.47
<i>Betula lenta</i>	-3.33	-10.33	3.85	-4.84	-12.37	2.89	-0.98	-5.16	2.98
<i>Corylus cornuta</i>	-6.20	-17.60	5.56	-7.24	-15.02	0.13	-1.35	-5.69	2.86
<i>Ostrya carpinifolia</i>	-5.52	-14.52	3.75	-4.92	-14.08	3.93	-0.92	-4.91	3.01
<i>Ostrya virginiana</i>	-11.38	-20.74	-2.10	-4.75	-14.01	4.43	-0.75	-4.80	3.43
<i>Carpinus laxiflora</i>	-8.99	-19.72	1.53	-4.74	-13.68	3.95	-0.91	-5.22	3.32
<i>Carpinus betulus</i>	-12.30	-18.99	-5.82	-2.59	-7.41	2.38	-0.59	-4.06	3.02
<i>Carpinus monbeigiana</i>	-5.66	-14.75	3.82	-4.73	-13.83	4.11	-0.89	-4.97	3.20
<i>Rosa majalis</i>	-4.26	-14.32	5.85	-7.15	-17.85	3.38	-0.94	-5.20	3.37
<i>Rosa hugonis</i>	-6.88	-18.69	5.08	-7.23	-17.86	3.75	-1.03	-5.49	3.50
<i>Aronia melanocarpa</i>	-2.38	-10.96	6.08	-5.07	-12.57	2.62	-0.74	-4.72	3.33
<i>Photinia villosa</i>	-2.92	-11.68	5.92	-5.86	-14.58	2.79	-0.76	-4.64	3.30
<i>Spiraea japonica</i>	-3.74	-12.85	5.42	-6.66	-16.49	3.07	-0.71	-4.74	3.41
<i>Spiraea chamaedryfolia</i>	-4.02	-13.48	5.21	-6.72	-16.37	3.13	-0.73	-4.89	3.58
<i>Spiraea canescens</i>	-5.16	-16.13	5.65	-6.78	-16.48	2.85	-0.79	-5.14	3.64
<i>Prunus tenella</i>	-3.89	-15.10	7.59	-7.19	-16.37	2.29	-0.93	-5.27	3.29
<i>Prunus serrulata</i>	-4.63	-15.76	6.60	-7.08	-15.83	2.23	-0.78	-4.88	3.43
<i>Prunus pensylvanica</i>	-3.85	-14.58	7.26	-7.01	-14.53	0.66	-0.91	-5.02	3.29
<i>Prunus serotina</i>	-7.40	-16.97	2.22	-6.26	-14.41	2.36	-0.83	-4.90	3.31
<i>Prunus padus</i>	-1.76	-7.09	3.46	-10.44	-14.27	-6.74	-0.96	-2.96	1.03
<i>Prinsepia uniflora</i>	-5.04	-16.35	6.67	-6.97	-16.85	3.27	-0.72	-4.89	3.49
<i>Prinsepia sinensis</i>	-5.07	-17.10	6.91	-7.03	-16.88	3.01	-0.80	-5.04	3.55

Continued on next page

Table S1: Estimated sensitivities of 191 tree species to three environmental cues: chilling (b.chill), forcing (b.force) and photoperiod (b.photo), along with their corresponding 2.5% (low) and 97.5% (up) Credible Intervals. Values correspond to the full model accounting for phylogenetic relationships.

Species	b.chill	low	up	b.force	low	up	b.photo	low	up
<i>Oemleria cerasiformis</i>	-4.93	-15.87	6.42	-6.84	-16.51	3.03	-0.75	-4.99	3.60
<i>Ulmus minor</i>	-16.20	-20.78	-11.59	-10.39	-14.29	-6.49	-2.57	-6.32	0.98
<i>Ulmus glabra</i>	-18.61	-25.10	-12.12	-10.25	-17.35	-2.78	-1.49	-5.16	2.38
<i>Ulmus macrocarpa</i>	-17.08	-23.91	-10.39	-10.51	-18.27	-2.82	-1.49	-5.27	2.54
<i>Ulmus pumila</i>	-12.48	-17.10	-7.94	-10.68	-14.62	-6.65	-1.81	-5.35	1.76
<i>Ulmus parvifolia</i>	-15.59	-20.21	-10.92	-9.42	-13.49	-5.28	-2.39	-6.08	1.18
<i>Ulmus laevis</i>	-16.26	-25.01	-7.62	-9.63	-17.46	-1.30	-1.64	-5.64	2.43
<i>Ulmus americana</i>	-13.53	-22.09	-4.68	-9.81	-17.57	-1.43	-1.59	-5.51	2.44
<i>Ulmus villosa</i>	-15.74	-20.40	-11.06	-11.92	-15.90	-8.04	-1.94	-5.59	1.65
<i>Caragana pygmaea</i>	-5.09	-16.95	6.79	-4.36	-14.86	6.22	-1.14	-5.42	3.32
<i>Robinia pseudoacacia</i>	-5.49	-15.64	4.60	-0.16	-5.50	5.18	-0.98	-5.30	3.51
<i>Amorpha fruticosa</i>	-1.98	-12.24	8.75	-3.88	-13.42	5.70	-1.04	-5.35	3.42
<i>Cercis chinensis</i>	-0.91	-10.52	9.36	-5.45	-16.53	5.76	-1.86	-6.37	2.54
<i>Cercis canadensis</i>	-4.02	-13.75	5.85	-5.15	-15.95	6.47	-1.45	-6.00	3.10
<i>Tilia japonica</i>	-2.93	-11.73	6.09	-8.03	-16.42	0.99	-1.26	-5.59	3.04
<i>Tilia cordata</i>	-3.70	-8.16	0.93	-9.91	-14.05	-5.82	-2.55	-6.15	0.92
<i>Tilia dasystyla</i>	-7.53	-17.00	1.52	-7.95	-16.41	1.18	-1.28	-5.47	2.95
<i>Tilia platyphyllos</i>	-4.30	-13.23	4.64	-7.95	-16.40	1.33	-1.27	-5.57	3.13
<i>Hibiscus syriacus</i>	0.01	-9.30	10.00	-7.50	-17.14	2.41	-1.26	-5.54	3.08
<i>Aesculus flava</i>	-14.90	-24.42	-5.70	-1.49	-10.89	7.74	-1.45	-5.58	2.77
<i>Aesculus parviflora</i>	-12.63	-21.95	-3.69	-2.00	-11.47	7.49	-2.00	-6.23	2.02
<i>Aesculus hippocastanum</i>	-8.33	-16.34	-0.34	1.57	-4.06	7.72	-1.33	-5.34	2.74
<i>Toona sinensis</i>	-3.77	-13.34	6.19	-4.78	-15.53	6.37	-1.31	-5.70	2.94
<i>Orixa japonica</i>	-5.76	-15.83	4.15	-4.75	-15.49	6.38	-1.21	-5.45	3.25
<i>Ptelea trifoliata</i>	-4.44	-14.20	5.45	-4.60	-15.22	6.67	-1.23	-5.57	3.14
<i>Ribes divaricatum</i>	-11.44	-23.49	0.20	-6.57	-18.06	4.68	-1.25	-5.82	3.30
<i>Ribes glaciale</i>	-7.39	-17.37	2.48	-6.69	-18.49	4.88	-0.96	-5.39	3.52
<i>Ribes alpinum</i>	-9.60	-21.69	2.07	-6.65	-18.27	4.94	-1.07	-5.56	3.50
<i>Hamamelis virginiana</i>	-8.19	-20.19	3.97	-9.99	-19.60	-0.73	-1.60	-6.28	3.08
<i>Hamamelis vernalis</i>	-12.75	-22.86	-2.86	-7.46	-18.49	3.77	-1.08	-5.47	3.33
<i>Hamamelis japonica</i>	-8.81	-18.66	1.17	-7.47	-18.61	3.42	-1.11	-5.53	3.31
<i>Sinowilsonia henryi</i>	-4.90	-15.08	5.12	-7.03	-18.68	4.55	-1.22	-5.58	3.34
<i>Corylopsis sinensis</i>	-7.56	-19.20	4.40	-7.10	-18.43	3.99	-1.33	-5.75	3.21
<i>Corylopsis spicata</i>	-7.91	-20.00	4.03	-7.06	-18.29	4.06	-1.42	-5.81	3.03
<i>Liquidambar styraciflua</i>	-7.91	-17.96	1.79	-6.68	-17.99	4.77	-1.51	-5.92	2.83
<i>Liquidambar orientalis</i>	-2.27	-12.08	7.89	-6.81	-18.31	4.47	-1.21	-5.61	3.23
<i>Cercidiphyllum japonicum</i>	-9.19	-20.49	2.23	-6.65	-17.89	4.96	-1.24	-5.71	3.26
<i>Cercidiphyllum magnificum</i>	-9.91	-19.98	-0.03	-6.58	-17.94	4.98	-1.30	-5.65	3.18
<i>Parrotia persica</i>	-8.06	-20.72	4.46	-6.62	-17.88	4.72	-1.01	-5.32	3.49
<i>Paeonia rockii</i>	-6.41	-18.74	6.05	-6.63	-18.19	5.11	-1.17	-5.66	3.44

Continued on next page

Table S1: Estimated sensitivities of 191 tree species to three environmental cues: chilling (b.chill), forcing (b.force) and photoperiod (b.photo), along with their corresponding 2.5% (low) and 97.5% (up) Credible Intervals. Values correspond to the full model accounting for phylogenetic relationships.

Species	b.chill	low	up	b.force	low	up	b.photo	low	up
<i>Syringa villosa</i>	-5.04	-15.49	5.54	-4.17	-12.82	4.14	-0.89	-5.19	3.46
<i>Syringa vulgaris</i>	-4.55	-13.61	4.73	-1.79	-6.08	2.68	-0.93	-5.22	3.18
<i>Syringa josikaea</i>	-4.34	-14.73	6.34	-4.07	-12.75	4.47	-0.96	-5.36	3.41
<i>Syringa reticulata</i>	-4.51	-15.06	5.98	-4.12	-12.75	4.24	-0.88	-5.17	3.50
<i>Ligustrum tschonoskii</i>	-4.45	-14.78	5.75	-4.13	-12.79	4.31	-0.88	-5.20	3.49
<i>Forsythia ovata</i>	-4.34	-14.99	6.24	-4.68	-14.36	4.83	-0.88	-5.23	3.38
<i>Forsythia suspensa</i>	-4.40	-13.57	5.01	-4.73	-14.22	4.57	-0.83	-5.14	3.55
<i>Cephalanthus occidentalis</i>	-2.38	-12.22	7.91	-5.37	-16.10	5.42	-1.11	-5.46	3.35
<i>Viburnum buddleifolium</i>	-10.26	-21.58	0.26	-4.75	-13.53	4.42	-0.23	-4.45	4.30
<i>Viburnum carlesii</i>	-7.53	-18.46	2.95	-5.33	-14.31	3.30	-0.86	-5.13	3.60
<i>Viburnum cassinoides</i>	-4.06	-11.31	3.10	-4.63	-10.55	1.19	-1.07	-4.93	2.89
<i>Viburnum lantanoides</i>	-8.43	-15.76	-1.23	-7.04	-12.94	-1.20	-1.27	-5.27	2.77
<i>Viburnum plicatum</i>	-9.13	-20.55	1.66	-5.00	-13.89	4.25	-0.49	-4.90	4.04
<i>Viburnum opulus</i>	-6.20	-15.54	3.42	-5.27	-14.19	3.75	-0.80	-5.10	3.44
<i>Viburnum betulifolium</i>	-7.78	-18.78	3.06	-5.35	-14.50	3.79	-0.99	-5.29	3.27
<i>Weigela coraeensis</i>	-4.55	-14.12	4.95	-4.97	-15.03	5.46	-0.70	-4.94	3.65
<i>Weigela florida</i>	-5.05	-16.05	6.17	-5.02	-15.07	5.15	-0.72	-5.01	3.64
<i>Weigela maximowiczii</i>	-5.99	-15.36	3.50	-4.92	-15.24	5.49	-0.67	-4.88	3.66
<i>Heptacodium miconioides</i>	-4.27	-15.74	7.38	-5.05	-14.97	4.95	-0.78	-5.20	3.75
<i>Symphoricarpos albus</i>	-6.84	-16.49	2.74	-3.29	-8.35	1.96	-0.74	-5.02	3.60
<i>Lonicera maximowiczii</i>	-4.60	-16.20	6.94	-5.01	-14.15	4.06	-0.71	-4.96	3.80
<i>Lonicera alpigena</i>	-4.83	-16.11	6.80	-4.91	-14.25	4.31	-0.61	-4.86	3.79
<i>Lonicera canadensis</i>	-4.56	-16.34	7.44	-5.48	-13.30	1.79	-0.87	-5.28	3.53
<i>Lonicera caerulea</i>	-4.96	-17.41	7.31	-5.19	-14.44	3.80	-0.77	-5.21	3.86
<i>Eleutherococcus sieboldianus</i>	-3.08	-12.87	6.67	-5.51	-16.16	5.23	-0.75	-5.07	3.58
<i>Eleutherococcus setchuenensis</i>	-3.74	-15.08	7.72	-5.47	-16.29	5.56	-0.86	-5.32	3.72
<i>Eleutherococcus senticosus</i>	-3.20	-12.73	6.71	-5.45	-16.37	5.49	-0.72	-4.90	3.73
<i>Ilex mucronata</i>	-2.88	-10.53	4.87	-6.00	-12.28	0.16	-1.91	-6.17	2.25
<i>Rhododendron prinophyllum</i>	-3.02	-14.99	9.25	-11.17	-20.26	-2.67	-0.97	-5.49	3.70
<i>Rhododendron canadense</i>	0.00	-9.69	9.99	-10.20	-19.93	-0.41	-0.78	-4.94	3.73
<i>Rhododendron dauricum</i>	-1.90	-13.20	9.95	-10.15	-20.05	0.24	-0.66	-5.05	3.89
<i>Rhododendron mucronulatum</i>	-2.54	-14.20	9.05	-10.38	-20.08	-0.54	-0.78	-5.24	3.88
<i>Kalmia angustifolia</i>	-2.93	-14.96	9.56	-12.80	-21.36	-4.83	-1.53	-6.00	2.93
<i>Vaccinium myrtilloides</i>	-1.69	-13.89	10.77	-9.51	-17.61	-1.70	-0.80	-5.17	3.86
<i>Lyonia ligustrina</i>	-3.56	-15.92	9.13	-11.82	-20.72	-3.04	-0.99	-5.57	3.76
<i>Nyssa sylvatica</i>	-6.22	-19.01	6.72	-8.05	-17.75	1.31	-1.44	-6.02	3.16
<i>Cornus alba</i>	-4.43	-14.38	5.63	2.72	-4.19	9.86	-1.48	-5.75	2.74

Continued on next page

Table S1: Estimated sensitivities of 191 tree species to three environmental cues: chilling (b.chill), forcing (b.force) and photoperiod (b.photo), along with their corresponding 2.5% (low) and 97.5% (up) Credible Intervals. Values correspond to the full model accounting for phylogenetic relationships.

Species	b.chill	low	up	b.force	low	up	b.photo	low	up
<i>Cornus kousa</i>	-2.92	-12.91	6.78	-2.84	-12.93	7.21	-1.30	-5.77	3.02
<i>Cornus mas</i>	-4.83	-14.88	5.25	-2.17	-8.45	4.40	-1.15	-5.51	3.30
<i>Hydrangea arborescens</i>	-8.72	-18.61	0.95	-5.11	-16.13	6.25	-0.91	-5.30	3.50
<i>Hydrangea involucrata</i>	-8.38	-17.92	1.10	-5.18	-16.24	6.18	-0.96	-5.14	3.46
<i>Hydrangea serrata</i>	-7.90	-19.38	3.71	-5.24	-16.55	6.05	-0.95	-5.30	3.43
<i>Deutzia gracilis</i>	-5.33	-16.70	6.09	-5.43	-16.20	5.60	-0.87	-5.24	3.65
<i>Deutzia scabra</i>	-3.84	-13.80	6.13	-5.30	-16.00	5.63	-0.87	-5.16	3.50
<i>Decaisnea fargesii</i>	-8.07	-21.12	4.95	-6.02	-18.26	5.96	-1.05	-5.68	3.54
<i>Berberis dielsiana</i>	-5.03	-18.26	8.40	-6.23	-18.11	6.06	-0.98	-5.72	3.80
<i>Liriodendron tulipifera</i>	-12.23	-26.09	1.05	-6.28	-18.58	5.93	-1.57	-6.32	3.20
<i>Acer pseudoplatanus</i>	-10.59	-16.87	-4.06	-9.06	-12.19	-5.96	-1.30	-4.58	2.05
<i>Acer saccharinum</i>	-7.95	-12.05	-3.92	-3.33	-10.91	4.47	-1.20	-5.57	3.07
<i>Acer rubrum</i>	-15.89	-22.63	-9.17	-0.34	-5.01	4.46	-0.42	-4.11	3.53
<i>Acer barbinerve</i>	-9.88	-20.06	0.71	-4.03	-12.65	4.64	-1.19	-5.29	2.92
<i>Acer negundo</i>	-11.88	-21.14	-2.52	-2.89	-10.66	5.24	-1.30	-5.54	2.85
<i>Acer pensylvanicum</i>	-9.42	-16.45	-2.35	-5.80	-11.67	-0.04	-1.97	-5.94	1.83
<i>Acer platanoides</i>	-10.35	-19.35	-1.41	-3.77	-12.18	4.69	-1.21	-5.25	2.80
<i>Acer campestre</i>	-11.25	-20.38	-2.26	-3.71	-12.17	5.09	-1.10	-5.10	3.04
<i>Acer tataricum</i>	-9.05	-18.10	0.14	-2.60	-10.40	5.71	-1.18	-5.31	3.01
<i>Acer ginnala</i>	-9.18	-19.83	1.23	-3.93	-12.38	4.92	-1.22	-5.43	3.05
<i>Amelanchier laevis</i>	-5.13	-15.80	6.01	-5.88	-14.63	2.85	-0.91	-5.08	3.31
<i>Amelanchier florida</i>	-6.02	-15.05	2.89	-5.86	-14.55	3.07	-0.72	-4.73	3.36
<i>Buddleja davidii</i>	-3.03	-15.36	9.46	-5.45	-15.85	5.04	-0.75	-5.13	3.84
<i>Buddleja alternifolia</i>	-2.79	-14.91	9.62	-5.45	-15.69	5.26	-0.83	-5.19	3.75
<i>Buddleja albiflora</i>	-3.35	-15.38	8.66	-5.48	-16.42	4.90	-0.79	-5.21	3.92
<i>Celtis laevigata</i>	-12.47	-22.42	-2.60	-7.60	-18.51	3.47	-1.60	-5.81	2.64
<i>Celtis occidentalis</i>	-7.70	-19.65	3.89	-7.48	-18.18	3.43	-1.40	-5.77	3.03
<i>Celtis caucasica</i>	-10.86	-20.81	-1.13	-7.54	-18.07	2.98	-1.29	-5.60	3.02
<i>Cladrastis lutea</i>	-9.69	-19.88	0.48	-4.68	-15.83	6.43	-1.93	-6.40	2.32
<i>Elaeagnus ebbingei</i>	-7.93	-20.04	4.44	-8.29	-19.52	2.52	-1.09	-5.42	3.40
<i>Fagus crenata</i>	-13.85	-22.96	-4.61	-7.94	-16.59	0.62	-3.49	-8.02	1.21
<i>Fagus engleriana</i>	-13.94	-22.98	-5.04	-7.93	-16.88	0.24	-3.39	-7.90	1.36
<i>Fagus grandifolia</i>	-13.77	-21.02	-6.50	-8.83	-14.77	-2.89	-4.52	-8.88	-0.31
<i>Fagus orientalis</i>	-16.38	-25.68	-7.38	-8.06	-16.89	0.26	-4.00	-8.52	0.42
<i>Fagus sylvatica</i>	-14.31	-15.89	-12.74	-2.91	-4.92	-0.92	-9.37	-11.91	-6.71
<i>Fraxinus excelsior</i>	-6.70	-15.49	1.96	-5.66	-13.56	1.70	-1.34	-5.59	2.76
<i>Fraxinus ornus</i>	-11.44	-20.97	-2.69	-4.07	-12.57	4.71	-0.88	-4.99	3.28
<i>Fraxinus nigra</i>	-6.30	-16.53	4.27	-7.89	-15.98	-0.64	-1.49	-5.81	2.64
<i>Fraxinus pennsylvanica</i>	-4.76	-13.81	4.38	-4.04	-11.90	4.08	-1.06	-5.25	3.18
<i>Fraxinus americana</i>	-7.23	-11.63	-2.76	-3.87	-11.48	4.02	-0.98	-5.27	3.39
<i>Fraxinus latifolia</i>	-7.66	-16.43	1.07	-4.63	-13.19	3.81	-1.45	-5.61	2.74

Continued on next page



Table S1: Estimated sensitivities of 191 tree species to three environmental cues: chilling (b.chill), forcing (b.force) and photoperiod (b.photo), along with their corresponding 2.5% (low) and 97.5% (up) Credible Intervals. Values correspond to the full model accounting for phylogenetic relationships.

Species	b.chill	low	up	b.force	low	up	b.photo	low	up
<i>Fraxinus chinensis</i>	-5.94	-14.92	2.99	-4.40	-12.35	3.60	-0.95	-5.04	3.16
<i>Juglans cinerea</i>	-11.11	-20.91	-1.25	-2.91	-12.33	7.19	-1.45	-5.75	2.70
<i>Juglans ailantifolia</i>	-11.61	-21.18	-2.29	-2.85	-12.14	7.07	-1.51	-5.90	2.77
<i>Parrotiopsis jaquemontiana</i>	-5.83	-15.63	3.89	-7.40	-18.44	3.63	-1.11	-5.55	3.32
<i>Pyrus ussuriensis</i>	-6.05	-16.71	4.34	-5.98	-14.58	2.52	-0.73	-4.74	3.31
<i>Pyrus elaeagnifolia</i>	-6.92	-15.89	1.86	-5.85	-14.48	3.13	-0.80	-4.83	3.31
<i>Quercus faginea</i>	-20.99	-32.47	-10.85	-12.86	-18.58	-7.15	-3.69	-8.03	0.37
<i>Quercus bicolor</i>	-15.08	-23.84	-6.01	-9.69	-16.58	-1.46	-2.31	-6.48	1.89
<i>Quercus alba</i>	-14.33	-24.51	-3.88	-12.84	-19.68	-5.59	-2.69	-6.74	1.39
<i>Quercus coccifera</i>	-21.14	-32.27	-10.79	-12.91	-20.09	-5.50	-2.50	-6.73	1.85
<i>Quercus rubra</i>	-18.25	-24.21	-12.29	-11.54	-15.11	-7.89	-2.56	-6.10	1.07
<i>Quercus ellipsoidalis</i>	-17.47	-27.04	-7.96	-13.15	-20.62	-5.69	-2.63	-6.89	1.63
<i>Quercus velutina</i>	-14.69	-25.24	-3.81	-13.13	-20.18	-6.08	-2.47	-6.65	1.80
<i>Quercus shumardii</i>	-17.04	-25.91	-8.41	-12.12	-19.62	-4.18	-2.40	-6.36	1.79
<i>Quercus ilex</i>	-23.29	-34.81	-13.14	-17.73	-24.27	-11.90	-2.57	-6.69	1.78
<i>Quercus petraea</i>	-16.47	-22.58	-10.32	-12.45	-15.88	-9.04	-2.64	-6.62	1.38
<i>Quercus pubescens</i>	-14.74	-25.34	-3.62	-16.42	-24.65	-9.56	-2.33	-6.61	2.28
<i>Quercus robur</i>	-13.18	-19.40	-6.83	-11.66	-15.22	-8.02	0.28	-3.49	4.35
<i>Rhamnus cathartica</i>	-4.58	-12.27	3.22	-11.90	-22.38	-2.51	-1.93	-6.12	2.18
<i>Rhamnus alpina</i>	-11.53	-21.67	-2.00	-8.90	-19.55	1.61	-1.14	-5.42	3.13
<i>Rhamnus frangula</i>	-6.98	-18.76	4.88	-10.67	-19.83	-1.62	-1.29	-5.82	3.19
<i>Salix gracilistyla</i>	-7.38	-18.54	3.62	-6.15	-15.42	3.21	-1.03	-5.34	3.26
<i>Salix smithiana</i>	-5.00	-11.20	1.20	-4.71	-10.10	0.96	-1.98	-6.28	2.08
<i>Salix repens</i>	-9.73	-19.17	-0.53	-6.02	-14.86	3.35	-1.16	-5.36	3.15
<i>Sambucus tigranii</i>	-8.12	-20.14	3.49	-5.33	-15.53	5.07	-0.83	-5.23	3.56
<i>Sambucus pubens</i>	-8.30	-20.17	3.35	-5.33	-15.75	5.11	-0.79	-5.26	3.70
<i>Sorbus aucuparia</i>	-5.67	-15.28	4.26	-5.60	-10.21	-0.98	-0.80	-4.24	2.58
<i>Sorbus torminalis</i>	-8.15	-17.44	0.37	-6.21	-14.71	2.47	-0.83	-4.46	2.95
<i>Sorbus aria</i>	-8.86	-18.00	0.08	-5.67	-14.08	2.86	-0.86	-4.74	3.09
<i>Sorbus decora</i>	-6.40	-15.18	2.22	-5.78	-14.19	2.78	-0.85	-4.88	3.17
<i>Sorbus commixta</i>	-5.84	-14.68	3.21	-5.88	-14.37	2.63	-0.77	-4.82	3.39
<i>Spirea alba</i>	-2.58	-14.70	10.38	-5.51	-13.84	2.80	-0.99	-5.28	3.46
<i>Stachyurus praecox</i>	-6.22	-18.40	6.27	-6.02	-16.95	5.21	-1.08	-5.46	3.37
<i>Stachyurus sinensis</i>	-6.14	-16.90	4.62	-5.65	-17.01	5.99	-0.97	-5.47	3.62

Table S2: Estimated sensitivities of 191 tree species to three environmental cues: chilling (b.chill), forcing (b.force) and photoperiod (b.photo), along with their corresponding 2.5% (low) and 97.5% (up) Credible Intervals. Values correspond to the non phylogenetic model.

Species	b.chill	low	up	b.force	low	up	b.photo	low	up
<i>Populus deltoides</i>	-18.11	-28.51	-7.77	-6.23	-15.95	4.03	-1.26	-5.84	3.32
<i>Populus koreana</i>	-7.92	-20.17	4.43	-7.42	-18.36	3.81	-1.10	-5.60	3.35
<i>Populus tremula</i>	-0.58	-6.21	5.09	-7.44	-10.70	-4.20	0.08	-2.10	2.32
<i>Populus grandidentata</i>	-15.62	-23.79	-7.77	-11.33	-17.49	-5.03	-1.32	-5.36	2.76
<i>Euonymus europaeus</i>	-8.27	-20.53	4.41	-7.49	-17.98	2.90	-1.18	-5.65	3.40
<i>Euonymus latifolius</i>	0.39	-10.11	10.84	-7.43	-18.03	3.43	-0.92	-5.14	3.40
<i>Nothofagus antarctica</i>	-10.50	-23.34	2.22	-7.46	-17.74	3.05	-1.38	-5.87	3.14
<i>Carya cordiformis</i>	-14.04	-24.40	-3.93	-7.85	-18.59	2.95	-2.62	-7.25	1.74
<i>Carya laciniosa</i>	-17.27	-27.99	-6.99	-6.61	-17.22	4.39	-1.16	-5.49	3.25
<i>Alnus incana</i>	-1.59	-5.55	2.37	-10.38	-14.59	-6.15	-0.55	-2.95	1.88
<i>Alnus glutinosa</i>	-6.51	-13.31	0.43	-10.42	-15.83	-5.14	-0.61	-3.13	1.88
<i>Alnus maximowiczii</i>	-6.18	-16.56	4.39	-7.15	-17.62	3.58	-1.21	-5.73	3.36
<i>Betula nana</i>	-6.39	-16.31	3.94	-7.15	-17.95	3.49	-1.13	-5.50	3.18
<i>Betula pendula</i>	-4.32	-5.43	-3.24	-1.34	-3.51	0.83	-0.43	-1.41	0.55
<i>Betula pubescens</i>	-2.11	-3.42	-0.79	-6.88	-9.54	-4.20	-0.78	-1.88	0.31
<i>Betula populifolia</i>	-5.25	-15.53	5.25	-7.38	-18.12	3.16	-1.08	-5.38	3.30
<i>Betula papyrifera</i>	-29.62	-35.81	-23.48	-5.80	-10.59	-0.96	1.37	-2.37	5.36
<i>Betula alleghaniensis</i>	-9.92	-17.11	-2.60	-6.93	-10.75	-3.03	-2.03	-6.13	1.95
<i>Betula lenta</i>	-2.29	-9.47	5.25	-5.99	-14.69	2.79	-1.23	-5.54	3.13
<i>Corylus cornuta</i>	-6.37	-19.26	7.06	-9.25	-17.67	-1.12	-1.62	-5.98	2.70
<i>Ostrya carpinifolia</i>	-4.10	-14.33	6.22	-7.40	-18.07	3.12	-1.08	-5.33	3.33
<i>Ostrya virginiana</i>	-13.26	-23.48	-3.08	-6.97	-17.59	3.71	-0.85	-5.04	3.54
<i>Carpinus laxiflora</i>	-9.40	-22.06	3.28	-7.33	-18.04	3.19	-1.13	-5.57	3.27
<i>Carpinus betulus</i>	-13.38	-20.38	-6.35	-2.12	-7.22	3.01	-0.62	-4.38	3.08
<i>Carpinus monbeigiana</i>	-4.43	-14.44	6.14	-7.23	-17.98	3.60	-1.05	-5.41	3.16
<i>Rosa majalis</i>	-4.90	-14.82	5.20	-7.49	-18.17	2.80	-0.91	-5.30	3.54
<i>Rosa hugonis</i>	-8.24	-20.58	4.29	-7.48	-17.98	2.93	-1.01	-5.44	3.50
<i>Aronia melanocarpa</i>	-2.26	-12.02	8.08	-5.30	-13.53	3.41	-1.01	-5.18	3.30
<i>Photinia villosa</i>	-3.69	-13.75	6.68	-7.29	-17.69	3.12	-0.99	-5.24	3.28
<i>Spiraea japonica</i>	-5.62	-15.79	4.68	-7.23	-17.59	3.41	-0.90	-5.24	3.44
<i>Spiraea chamaedryfolia</i>	-6.08	-16.17	4.00	-7.35	-17.96	3.50	-0.90	-5.25	3.53
<i>Spiraea canescens</i>	-8.77	-21.15	3.53	-7.52	-17.64	3.02	-1.05	-5.46	3.37
<i>Prunus tenella</i>	-6.68	-18.91	6.02	-7.50	-18.67	3.50	-1.11	-5.55	3.30
<i>Prunus serrulata</i>	-8.11	-20.34	4.47	-7.26	-17.51	2.93	-0.94	-5.47	3.52
<i>Prunus pensylvanica</i>	-6.31	-18.71	6.34	-6.83	-14.73	1.34	-1.14	-5.66	3.25
<i>Prunus serotina</i>	-10.58	-21.02	-0.14	-5.49	-14.91	4.23	-1.09	-5.41	3.21
<i>Prunus padus</i>	-2.17	-7.46	3.09	-10.81	-14.58	-6.99	-1.02	-3.02	0.93
<i>Prinsepia uniflora</i>	-7.80	-20.33	4.82	-7.62	-18.52	3.05	-0.96	-5.31	3.54
<i>Prinsepia sinensis</i>	-8.08	-21.71	5.47	-7.73	-18.30	2.92	-1.03	-5.45	3.47
<i>Oemleria cerasiformis</i>	-7.71	-19.80	4.48	-7.38	-17.76	2.99	-0.89	-5.29	3.44
<i>Ulmus minor</i>	-15.63	-20.50	-10.78	-9.99	-14.32	-5.72	-2.53	-6.44	1.32
<i>Ulmus glabra</i>	-17.95	-25.56	-10.49	-8.23	-17.95	1.42	-0.95	-5.10	3.26

Continued on next page

Table S2: Estimated sensitivities of 191 tree species to three environmental cues: chilling (b.chill), forcing (b.force) and photoperiod (b.photo), along with their corresponding 2.5% (low) and 97.5% (up) Credible Intervals. Values correspond to the non phylogenetic model.

Species	b.chill	low	up	b.force	low	up	b.photo	low	up
<i>Ulmus macrocarpa</i>	-16.04	-23.67	-8.80	-8.41	-18.61	1.26	-0.98	-5.12	3.21
<i>Ulmus pumila</i>	-11.49	-16.11	-6.71	-10.23	-14.42	-6.11	-1.42	-5.28	2.43
<i>Ulmus parvifolia</i>	-15.06	-19.83	-10.22	-8.70	-12.86	-4.46	-2.26	-6.03	1.52
<i>Ulmus laevis</i>	-14.85	-25.33	-4.77	-7.03	-17.89	3.92	-1.09	-5.44	3.11
<i>Ulmus americana</i>	-10.02	-20.18	0.50	-7.25	-17.66	3.23	-1.08	-5.43	3.28
<i>Ulmus villosa</i>	-15.17	-19.92	-10.37	-11.93	-16.26	-7.68	-1.64	-5.41	2.16
<i>Caragana pygmaea</i>	-6.71	-18.99	6.14	-7.62	-17.95	2.61	-1.00	-5.50	3.49
<i>Robinia pseudoacacia</i>	-6.86	-17.21	3.40	-0.60	-5.78	4.61	-0.75	-5.06	3.72
<i>Amorpha fruticosa</i>	-2.88	-13.18	7.81	-6.49	-15.80	3.00	-0.89	-5.06	3.46
<i>Cercis chinensis</i>	-1.56	-11.77	9.06	-7.60	-17.89	2.95	-1.74	-6.21	2.63
<i>Cercis canadensis</i>	-6.58	-16.76	3.84	-7.05	-17.67	3.67	-1.16	-5.52	3.17
<i>Tilia japonica</i>	-4.13	-14.09	6.20	-7.27	-17.82	3.35	-0.92	-5.13	3.38
<i>Tilia cordata</i>	-4.07	-8.65	0.70	-10.14	-14.18	-6.13	-2.80	-6.50	0.72
<i>Tilia dasystyla</i>	-11.94	-21.87	-2.13	-7.08	-17.24	3.43	-0.97	-5.30	3.48
<i>Tilia platyphyllos</i>	-6.54	-16.83	3.82	-7.15	-17.61	3.34	-0.95	-5.30	3.35
<i>Hibiscus syriacus</i>	-0.59	-11.02	10.10	-7.26	-17.56	3.34	-1.01	-5.34	3.55
<i>Aesculus flava</i>	-16.49	-27.49	-6.45	-6.64	-16.97	4.11	-1.19	-5.56	3.16
<i>Aesculus parviflora</i>	-12.49	-22.79	-2.37	-7.72	-18.11	3.24	-2.03	-6.34	2.29
<i>Aesculus hippocastanum</i>	-6.13	-14.54	2.27	0.38	-5.66	6.53	-1.15	-5.18	2.96
<i>Toona sinensis</i>	-4.10	-14.20	6.30	-7.25	-17.75	3.20	-1.17	-5.46	3.17
<i>Orixa japonica</i>	-6.50	-16.39	3.67	-7.19	-18.04	3.52	-1.13	-5.59	3.42
<i>Ptelea trifoliata</i>	-5.10	-15.40	5.25	-7.18	-17.65	3.19	-1.13	-5.31	3.00
<i>Ribes divaricatum</i>	-13.67	-26.45	-1.51	-7.57	-17.79	2.87	-1.34	-5.84	3.10
<i>Ribes glaciale</i>	-6.68	-16.53	3.71	-7.52	-18.12	3.24	-0.93	-5.24	3.38
<i>Ribes alpinum</i>	-10.17	-22.66	2.37	-7.55	-18.28	2.77	-1.09	-5.59	3.36
<i>Hamamelis virginiana</i>	-8.61	-21.51	4.26	-10.91	-20.31	-2.12	-1.71	-6.07	2.63
<i>Hamamelis vernalis</i>	-14.59	-24.87	-4.51	-6.98	-17.43	3.98	-0.99	-5.36	3.38
<i>Hamamelis japonica</i>	-9.15	-19.59	1.08	-7.27	-17.57	2.99	-1.06	-5.29	3.44
<i>Sinowilsonia henryi</i>	-5.24	-15.18	4.75	-7.49	-18.03	2.98	-1.18	-5.49	3.06
<i>Corylopsis sinensis</i>	-8.51	-21.07	3.86	-7.55	-18.23	3.10	-1.28	-5.87	3.26
<i>Corylopsis spicata</i>	-9.18	-22.03	3.41	-7.43	-17.82	3.06	-1.39	-5.73	2.99
<i>Liquidambar styraciflua</i>	-11.04	-21.08	-0.87	-7.33	-17.56	2.99	-1.53	-5.94	2.77
<i>Liquidambar orientalis</i>	-1.35	-11.39	9.46	-7.50	-17.89	2.68	-1.10	-5.42	3.36
<i>Cercidiphyllum japonicum</i>	-9.68	-22.12	2.60	-7.37	-17.85	2.64	-1.21	-5.57	3.14
<i>Cercidiphyllum magnificum</i>	-10.65	-20.75	-0.52	-7.40	-18.14	3.44	-1.26	-5.51	3.06
<i>Parrotia persica</i>	-9.39	-21.92	2.89	-7.29	-17.70	3.04	-0.96	-5.35	3.61
<i>Paeonia rockii</i>	-7.59	-19.89	4.62	-7.50	-17.90	2.67	-1.11	-5.54	3.45
<i>Syringa villosa</i>	-7.83	-20.07	4.59	-7.56	-17.94	3.00	-0.96	-5.47	3.67
<i>Syringa vulgaris</i>	-6.30	-16.36	3.99	-1.65	-6.09	2.84	-1.10	-5.37	3.20
<i>Syringa josikaea</i>	-6.60	-19.15	6.33	-7.45	-18.02	2.72	-1.06	-5.44	3.39
<i>Syringa reticulata</i>	-6.89	-19.47	5.72	-7.39	-17.73	3.23	-0.99	-5.25	3.46

Continued on next page

Table S2: Estimated sensitivities of 191 tree species to three environmental cues: chilling (b.chill), forcing (b.force) and photoperiod (b.photo), along with their corresponding 2.5% (low) and 97.5% (up) Credible Intervals. Values correspond to the non phylogenetic model.

Species	b.chill	low	up	b.force	low	up	b.photo	low	up
Ligustrum tschonoskii	-6.63	-19.20	6.28	-7.60	-18.30	3.08	-0.97	-5.36	3.50
Forsythia ovata	-6.77	-19.55	5.87	-7.44	-17.72	2.73	-1.05	-5.50	3.29
Forsythia suspensa	-6.12	-16.36	4.09	-7.35	-17.43	3.19	-0.91	-5.26	3.49
Cephalanthus occidentalis	-4.96	-14.92	5.38	-7.24	-17.89	3.24	-1.20	-5.44	3.08
Viburnum buddleifolium	-14.54	-27.32	-2.15	-6.42	-16.75	4.49	-0.33	-4.81	4.32
Viburnum carlesii	-8.95	-21.76	3.97	-7.51	-18.21	3.24	-1.24	-5.64	3.15
Viburnum cassinoides	-3.71	-11.38	3.89	-5.17	-11.27	1.16	-1.47	-5.57	2.67
Viburnum lantanoides	-9.24	-16.57	-1.67	-8.47	-14.62	-2.48	-1.61	-5.72	2.45
Viburnum plicatum	-12.20	-25.12	0.40	-6.78	-16.97	3.77	-0.68	-5.11	3.94
Viburnum opulus	-6.92	-17.13	3.34	-7.13	-17.59	3.54	-1.07	-5.31	3.32
Viburnum betulifolium	-9.67	-21.96	2.29	-7.35	-18.04	2.93	-1.35	-5.71	3.06
Weigela coraeensis	-6.56	-16.50	3.63	-7.28	-17.73	3.12	-0.97	-5.39	3.55
Weigela florida	-7.68	-20.09	4.64	-7.44	-17.87	2.82	-1.02	-5.43	3.48
Weigela maximowiczii	-8.92	-19.08	1.20	-7.17	-17.88	3.39	-0.93	-5.25	3.48
Heptacodium miconioides	-6.82	-19.04	5.69	-7.43	-18.06	3.16	-1.06	-5.51	3.45
Symphoricarpos albus	-9.52	-19.78	0.77	-3.21	-8.44	2.01	-0.99	-5.31	3.30
Lonicera maximowiczii	-7.39	-19.31	5.44	-7.68	-18.06	2.94	-0.99	-5.42	3.58
Lonicera alpigena	-7.78	-20.33	4.51	-7.44	-18.17	3.04	-0.91	-5.30	3.73
Lonicera canadensis	-5.93	-18.81	7.24	-6.98	-14.79	0.81	-1.10	-5.56	3.39
Lonicera caerulea	-8.21	-21.52	5.31	-7.69	-18.25	2.78	-1.07	-5.56	3.48
Eleutherococcus sieboldianus	-5.41	-15.51	4.71	-7.38	-17.99	3.12	-0.93	-5.20	3.20
Eleutherococcus setchuenensis	-7.57	-20.49	5.35	-7.35	-17.75	2.96	-1.11	-5.57	3.36
Eleutherococcus senticosus	-5.49	-15.94	5.09	-7.28	-17.96	3.46	-0.85	-5.16	3.48
Ilex mucronata	-3.79	-11.32	3.87	-6.56	-12.55	-0.46	-1.95	-6.21	2.13
Rhododendron prinophyllum	-8.09	-20.84	4.67	-9.26	-18.33	-0.45	-1.15	-5.60	3.14
Rhododendron canadense	-2.62	-12.72	7.74	-7.23	-17.42	3.14	-0.97	-5.29	3.32
Rhododendron dauricum	-6.80	-19.00	5.68	-7.19	-17.73	3.19	-0.78	-5.16	3.84
Rhododendron mucronulatum	-7.40	-20.33	5.50	-7.54	-18.22	3.36	-0.93	-5.24	3.42
Kalmia angustifolia	-7.32	-19.73	5.25	-12.11	-20.86	-4.08	-1.88	-6.36	2.60
Vaccinium myrtilloides	-5.98	-18.81	6.75	-7.41	-15.38	0.43	-0.96	-5.29	3.52
Lyonia ligustrina	-8.15	-21.11	4.80	-10.43	-20.28	-1.38	-1.13	-5.67	3.42
Nyssa sylvatica	-8.31	-20.67	4.49	-9.52	-18.51	-0.80	-1.39	-5.88	3.14
Cornus alba	-6.41	-16.04	3.37	1.92	-5.09	9.28	-1.53	-5.67	2.62
Cornus kousa	-4.68	-14.61	5.57	-7.39	-17.85	2.97	-1.23	-5.54	3.10
Cornus mas	-6.61	-16.98	3.82	-3.77	-10.09	2.70	-1.00	-5.49	3.49
Hydrangea arborescens	-10.58	-20.66	-0.37	-7.14	-17.55	3.55	-1.00	-5.24	3.33
Hydrangea involucrata	-10.15	-20.14	-0.07	-7.34	-18.13	3.41	-1.09	-5.52	3.30
Hydrangea serrata	-9.89	-22.57	2.80	-7.40	-17.99	3.12	-1.05	-5.61	3.29

Continued on next page

Table S2: Estimated sensitivities of 191 tree species to three environmental cues: chilling (b.chill), forcing (b.force) and photoperiod (b.photo), along with their corresponding 2.5% (low) and 97.5% (up) Credible Intervals. Values correspond to the non phylogenetic model.

Species	b.chill	low	up	b.force	low	up	b.photo	low	up
Deutzia gracilis	-7.60	-20.50	4.96	-7.50	-18.00	2.96	-0.97	-5.22	3.39
Deutzia scabra	-5.07	-15.20	5.25	-7.38	-17.61	2.97	-0.96	-5.33	3.45
Decaisnea fargesii	-9.88	-22.56	2.67	-7.29	-18.05	3.42	-1.04	-5.49	3.53
Berberis dielsiana	-6.59	-19.04	6.31	-7.55	-18.15	3.04	-1.03	-5.46	3.38
Liriodendron tulipifera	-13.64	-26.59	-1.43	-7.48	-17.99	2.87	-1.54	-6.05	2.97
Acer pseudoplatanus	-9.70	-16.44	-2.92	-9.89	-12.97	-6.82	-1.04	-4.56	2.52
Acer saccharinum	-7.59	-11.67	-3.57	-5.92	-14.96	3.57	-1.10	-5.65	3.47
Acer rubrum	-16.72	-23.81	-9.65	-0.57	-5.33	4.18	-0.11	-3.91	3.89
Acer barbinerve	-8.41	-20.80	4.09	-7.50	-17.88	3.24	-1.09	-5.44	3.39
Acer negundo	-11.74	-22.34	-1.06	-5.75	-15.29	4.11	-1.25	-5.65	3.01
Acer pensylvanicum	-8.87	-16.22	-1.40	-7.87	-13.96	-1.91	-2.06	-6.16	1.94
Acer platanoides	-9.84	-20.42	0.22	-7.16	-17.34	2.95	-1.08	-5.44	3.22
Acer campestre	-10.97	-21.18	-0.77	-7.11	-17.69	3.50	-0.95	-5.30	3.32
Acer tataricum	-7.79	-18.03	2.46	-5.40	-14.64	4.07	-1.15	-5.40	3.14
Acer ginnala	-7.24	-19.58	5.24	-7.49	-18.33	3.43	-1.12	-5.68	3.44
Amelanchier laevis	-8.10	-21.11	5.19	-7.37	-17.58	3.14	-1.20	-5.72	3.31
Amelanchier florida	-8.42	-18.53	1.88	-7.19	-17.84	3.50	-0.98	-5.22	3.36
Buddleja davidii	-6.63	-18.75	5.87	-7.44	-17.88	3.29	-0.89	-5.27	3.52
Buddleja alternifolia	-6.14	-18.70	6.71	-7.50	-17.98	2.95	-0.99	-5.35	3.43
Buddleja albiflora	-6.94	-19.44	5.67	-7.57	-17.97	2.86	-0.92	-5.35	3.61
Celtis laevigata	-12.78	-22.93	-2.58	-7.23	-17.80	3.21	-1.47	-5.78	2.81
Celtis occidentalis	-6.21	-18.44	6.36	-7.09	-17.45	3.03	-1.26	-5.69	3.24
Celtis caucasica	-10.47	-20.58	-0.58	-7.13	-17.53	3.27	-1.10	-5.50	3.24
Cladrastis lutea	-12.14	-22.62	-1.82	-7.49	-17.74	3.05	-1.87	-6.32	2.42
Elaeagnus ebbingei	-7.84	-20.50	4.78	-7.59	-18.36	3.17	-0.94	-5.27	3.50
Fagus crenata	-9.96	-20.36	0.39	-7.46	-17.89	2.82	-1.39	-5.68	2.90
Fagus engleriana	-10.08	-20.42	-0.09	-7.30	-17.89	3.21	-1.29	-5.55	3.13
Fagus grandifolia	-12.35	-20.21	-4.58	-9.72	-16.07	-3.46	-3.14	-7.46	0.95
Fagus orientalis	-14.05	-24.38	-3.82	-7.67	-17.96	2.80	-2.12	-6.56	2.13
Fagus sylvatica	-14.21	-15.81	-12.65	-2.65	-4.64	-0.66	-9.51	-12.05	-7.03
Fraxinus excelsior	-8.32	-18.52	1.93	-8.54	-17.44	0.30	-1.54	-5.89	2.75
Fraxinus ornus	-16.69	-27.23	-6.40	-6.19	-16.38	4.61	-0.79	-5.22	3.63
Fraxinus nigra	-8.33	-21.58	4.78	-11.81	-20.23	-3.87	-1.66	-6.24	2.85
Fraxinus pennsylvanica	-5.37	-15.49	5.02	-6.09	-15.47	3.72	-1.14	-5.50	3.25
Fraxinus americana	-7.61	-12.04	-3.03	-5.48	-14.63	4.13	-1.11	-5.73	3.57
Fraxinus latifolia	-10.54	-20.86	-0.45	-7.26	-17.74	3.42	-1.68	-6.06	2.61
Fraxinus chinensis	-7.27	-17.55	3.05	-6.59	-15.87	3.19	-0.98	-5.29	3.43
Juglans cinerea	-8.76	-19.59	2.18	-6.49	-15.82	2.86	-1.05	-5.29	3.25
Juglans ailantifolia	-9.49	-20.35	1.28	-6.42	-15.56	2.82	-1.17	-5.49	3.22
Parrotiopsis jaquemontiana	-5.61	-15.90	4.64	-7.37	-17.60	2.86	-1.06	-5.36	3.28
Pyrus ussuriensis	-8.49	-21.12	4.01	-7.50	-17.79	3.05	-1.01	-5.40	3.50

Continued on next page

Table S2: Estimated sensitivities of 191 tree species to three environmental cues: chilling (b.chill), forcing (b.force) and photoperiod (b.photo), along with their corresponding 2.5% (low) and 97.5% (up) Credible Intervals. Values correspond to the non phylogenetic model.

Species	b.chill	low	up	b.force	low	up	b.photo	low	up
<i>Pyrus elaeagnifolia</i>	-9.48	-19.57	0.58	-7.28	-18.02	3.53	-1.07	-5.26	3.22
<i>Quercus faginea</i>	-18.47	-33.56	-5.41	-11.03	-17.86	-4.45	-3.11	-7.94	1.24
<i>Quercus bicolor</i>	-8.94	-19.24	1.46	-6.42	-15.74	3.46	-1.21	-5.63	3.23
<i>Quercus alba</i>	-9.45	-22.53	3.34	-11.13	-20.45	-2.23	-1.97	-6.64	2.55
<i>Quercus coccifera</i>	-18.01	-32.31	-4.98	-8.90	-18.92	0.86	-1.25	-5.73	3.27
<i>Quercus rubra</i>	-16.88	-23.01	-10.62	-10.78	-14.60	-6.94	-2.17	-6.10	1.72
<i>Quercus ellipsoidalis</i>	-13.69	-25.01	-2.52	-7.55	-17.79	2.46	-1.22	-5.78	3.27
<i>Quercus velutina</i>	-9.87	-23.30	3.02	-11.53	-21.17	-2.22	-1.60	-6.14	2.87
<i>Quercus shumardii</i>	-12.36	-22.59	-2.46	-7.18	-17.39	3.45	-1.31	-5.42	3.02
<i>Quercus ilex</i>	-22.78	-36.98	-10.07	-18.72	-25.67	-12.04	-1.40	-5.94	3.15
<i>Quercus petraea</i>	-14.19	-21.05	-7.52	-12.70	-16.43	-9.14	-1.64	-5.91	2.50
<i>Quercus pubescens</i>	-5.83	-19.07	7.80	-15.90	-26.67	-6.44	-0.98	-5.62	3.61
<i>Quercus robur</i>	-10.11	-16.66	-3.51	-11.38	-15.34	-7.53	1.99	-1.51	5.79
<i>Rhamnus cathartica</i>	-4.04	-11.63	3.79	-11.39	-21.57	-1.84	-1.89	-6.25	2.37
<i>Rhamnus alpina</i>	-13.28	-23.56	-3.29	-6.84	-17.38	3.86	-0.92	-5.18	3.57
<i>Rhamnus frangula</i>	-7.45	-20.31	5.55	-9.97	-19.23	-0.83	-1.15	-5.51	3.37
<i>Salix gracilistyla</i>	-7.77	-20.03	5.07	-7.52	-17.74	2.88	-0.98	-5.36	3.50
<i>Salix smithiana</i>	-4.74	-11.00	1.55	-4.61	-10.32	0.93	-2.27	-6.68	2.05
<i>Salix repens</i>	-11.68	-21.88	-1.57	-7.11	-17.62	3.43	-1.11	-5.42	3.22
<i>Sambucus tigranii</i>	-10.43	-22.84	1.55	-7.40	-17.95	3.27	-1.05	-5.56	3.38
<i>Sambucus pubens</i>	-10.68	-23.03	1.36	-7.42	-17.88	3.17	-1.03	-5.39	3.43
<i>Sorbus aucuparia</i>	-8.39	-20.03	3.44	-6.35	-11.41	-1.50	-1.11	-4.81	2.58
<i>Sorbus torminalis</i>	-10.89	-21.41	-0.75	-8.01	-18.33	2.39	-1.09	-5.06	2.89
<i>Sorbus aria</i>	-13.07	-23.69	-2.74	-6.98	-17.30	3.44	-1.08	-5.35	3.24
<i>Sorbus decora</i>	-8.57	-18.81	1.94	-7.34	-17.48	2.93	-1.09	-5.39	3.30
<i>Sorbus commixta</i>	-7.51	-17.66	2.70	-7.38	-17.75	3.10	-1.01	-5.41	3.50
<i>Spirea alba</i>	-5.95	-18.47	6.94	-6.51	-14.74	1.51	-1.05	-5.49	3.46
<i>Stachyurus praecox</i>	-7.90	-20.29	4.77	-7.55	-17.73	2.86	-1.05	-5.54	3.43
<i>Stachyurus sinensis</i>	-7.56	-18.17	3.53	-7.03	-17.41	3.35	-0.99	-5.43	3.44




Figure S1: Non-phylogenetic phenological sensitivity to three environmental cues, chilling (a), forcing (b) and photoperiod (c) measured in change in days to budburst per standardized unit (z-transformation) of the cues across 191 tree species. Sensitivity estimates are computed by commonly used hierarchical model where phylogenetic distances are not accounted for ( $\lambda = 0$ ). The same phylogenetic tree is shown in each panel, colored according to an estimation of ancestral character states, being the states at the tips the species' sensitivities to a cue. Species sensitivities are shown along with 50% Credible Intervals in the diagrams. Note that the color scale varies in each panel. Total tree depth is 81. My.




Figure S2: Correlations among estimated sensitivities to the environmental cues comparing forcing vs. chilling (a,d), forcing vs. photoperiod (b,e) and chilling vs. photoperiod (c,f). Upper panels show correlations among estimated sensitivities by the phylogenetic model and lower panels show results for the non-phylogenetic model.






Figure S3: Comparison of uncertainty around estimated sensitivities to chilling (a), forcing (b) and photoperiod (c) of individual species between the phylogenetic model with estimated  $\lambda$  ( $\lambda_{\text{est}}$ ), and the non-phylogenetic model with  $\lambda = 0$  ( $\lambda_0$ ). The non-phylogenetic model increases uncertainty.




Figure S4: Bias in estimation of sensitivity to chilling (a), forcing (b) and photoperiod (c). Histograms show the difference between the phylogenetic model with estimated  $\lambda$  against the non-phylogenetic model with  $\lambda = 0$ . Positive values indicate that sensitivities estimated by the non-phylogenetic model are smaller than those estimated by the phylogenetic model.




Figure S5: Marginal plots contrasting the posterior distributions (in red) of estimated sensitivities to chilling (a), forcing (b) and photoperiod (c) against prior distributions (blue). Lower panels show marginal plots for the posterior distributions of phylogenetic parameter  $\lambda$  fitted for chilling (d), forcing (e) and photoperiod (f).




Figure S6: Estimates of sigmas using our phylogenetic model and simulated data with responses to forcing and photoperiod generated from a delta model of evolution (fastBM in phytools package)), with delta set to 2 for forcing and to 0.5 for photoperiod (intercept and chilling generated from lambda model with a lambda of 1).

## 1 Stan code