

# Fixed-Shape Ellipse by Three Points

Danilo Tedeschi  
Dra. Marina Andretta


Universidade de São Paulo

25 de Outubro de 2019

This research has been funded by Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES). This is part of a Master's research project on the problem: *Planar Maximal Covering by Ellipses with Rotation*.

# Introduction


The shape of an ellipse is given by its major-axis and minor-axis,  $(a, b) \in \mathbb{R}^2$ , with  $a > b > 0$ .



**Figura:** An ellipse with shape parameters  $a$  and  $b$ .

# Introduction

Here, the shape will be fixed and the center and angle of rotation are free.



**Figura:** A fix-shape ellipse at different centers and with different angles of rotation.

# Introduction

## Problem definition

Given three points  $u, v, w \in \mathbb{R}^2$ , and the shape  $(a, b) \in \mathbb{R}^2$  of an ellipse:





Figura: An instance of the problem.

# Introduction

## Problem definition

A solution is given by the ellipse's center  $q \in \mathbb{R}^2$  and the angle of rotation  $\theta \in [0, \pi)$ , such that  $u, v, w$  lie on its border. **We want to find every solution!**



**Figura:** Every solution for the instance shown previously.

# Introduction

The equation of an ellipse is given by:


$$\frac{\left( \begin{bmatrix} x - q_x \\ y - q_y \end{bmatrix}^T \begin{bmatrix} \cos \theta \\ \sin \theta \end{bmatrix} \right)^2}{a^2} + \frac{\left( \begin{bmatrix} x - q_x \\ q_y - y \end{bmatrix}^T \begin{bmatrix} \sin \theta \\ \cos \theta \end{bmatrix} \right)^2}{b^2} = 1.$$

- ▶ Fixing the points  $u, v, w$ , we get 3 equations and 3 unknowns  $(q_x, q_y, \theta)$ .
- ▶ Finding every solution is difficult.

# Transforming the problem

Let's make the problem simpler by transforming it into a circle problem.

An ellipse with shape  $(a, b)$  can be transformed into a circle of radius  $b$  through scaling the  $x$ -axis by  $\frac{b}{a}$ :



**Figura:** Turning an ellipse with shape  $(a, b)$  into a circle of radius  $b$ .

# Transforming the problem

Let's rotate the points instead of rotating the ellipse:



Figura: Three points at their initial location.



# Transforming the problem

Firstly, we rotate leaving one point fixed at  $(0,0)$ :



Figura: After rotation.

# Transforming the problem

Then, we scale by  $\frac{b}{a}$  and check the radius of the circle:




Figura: After scaling by  $\frac{b}{a}$ .

## Transforming the problem

Then, we apply the inverse transformation.





Figura: After scaling by  $\frac{a}{b}$ .

# Transforming the problem

If the radius is  $b$ , the angle of rotation is a solution:



**Figura:** A solution is obtained after scaling by  $\frac{a}{b}$  and rotating by  $-\theta$ .

# Transforming the problem

Formally, we can transform the problem by:

- ▶ Translate the points so  $u = (0, 0)$ .
- ▶ Rotate by  $\theta$  and scale the  $x$ -axis by  $\frac{b}{a}$ .
- ▶ Find the  $\theta$ 's which produce a circle with radius  $b$ .

This transformation is expressed by:

$$\varphi(p, \theta) = \begin{bmatrix} \frac{b}{a} & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix} \begin{bmatrix} p_x \\ p_y \end{bmatrix},$$

for  $p = u, v, w$ .

**This is an one variable problem on a closed interval!**

# Fixed-shape ellipse by three points

There is a known formula for the radius of a circumscribed circle [JY60, p. 189]:

$$R = \frac{\|\varphi(v, \theta)\|_2 \|\varphi(w, \theta)\|_2 \|\varphi(v, \theta) - \varphi(w, \theta)\|_2}{4A(\theta)}$$

- ▶  $R$  is the radius.
- ▶  $A(\theta)$  is the area of the triangle defined by the points  $\varphi(u, \theta), \varphi(v, \theta), \varphi(w, \theta)$ .

# Fixed-shape ellipse by three points

We define the function  $\xi : [0, \pi) \mapsto \mathbb{R}$ :

$$\xi(\theta) = 16b^2 A(\theta)^2 - \|\varphi(v, \theta)\|_2^2 \|\varphi(w, \theta)\|_2^2 \|\varphi(v, \theta) - \varphi(w, \theta)\|_2^2$$

- It can be written as

$$\sum_{0 \leq j, k \leq \text{Deg}(\xi)} \alpha_{j,k} \cos^j \theta \sin^k \theta.$$

- Degree 6, at most 12 roots in  $[0, 2\pi)$  [Pow81, p. 150]
- The roots of  $\xi$  are solutions of our problem.

# Fixed-shape ellipse by three points

An example with no roots.




Figura: An example of  $\xi$ .



# Fixed-shape ellipse by three points

An example with two roots.




Figura: Another example of  $\xi$ .

# Fixed-shape ellipse by three points

An example with four roots.




Figura: An example of  $\xi$  with 4 roots.

# Polynomial Interpolation

It is a way to approximate a function by a simpler one (a polynomial).

- ▶ A degree  $n$  of the interpolation is determined.
- ▶  $n + 1$  points are chosen, such that the polynomial has to pass through.
- ▶ Can be calculated using Lagrange's formula.
- ▶ We can find every root of a polynomial by determining the eigenvalues of a matrix called The Companion Matrix [HJ86, p. 195].
- ▶ Depending on the points, the interpolation can be a bad approximation. It can get worse even if  $n$  is increased (Runge's Phenomenon) [Pow81, p. 37].

# Chebyshev Interpolation

## Chebyshev Polynomial

$T_n : [-1, 1] \mapsto [-1, 1]$  is the  $n$ -degree Chebyshev polynomial [MH03]:

$$T_n(\cos t) = \cos(nt)$$

Also, it can be defined recursively:

$$T_0(x) = 1$$

$$T_1(x) = x$$

$$T_n(x) = 2xT_{n-1}(x) - T_{n-2}(x)$$

# Chebyshev Interpolation




Figura: Chebyshev Polynomials of degree 1, 2, 3, 4.

# Chebyshev Interpolation

Interpolation on the roots of  $T_n$ , also known as Chebyshev Nodes:

$$x_k = \cos\left(\pi \frac{2k-1}{2n}\right)$$

The interpolation of a function  $f : [-1, 1] \mapsto \mathbb{R}$  can be written directly using Chebyshev polynomial as basis:

$$f(x) \approx \sum_{k=0}^n a_k T_k(x)$$

- ▶ This can be done in  $\mathcal{O}(n^3)$  [Boy13].
- ▶ A simple change of coordinates lets the interpolation to be done on any closed interval!

# Chebyshev Interpolation

Why is it good?

- ▶ Numerically stable! Way better than polynomials in the power format [Gau79].
- ▶ No Runge's Phenomenon, the interpolation converges to  $f$ .
  - ▶  $\mathcal{O}(n^{-m})$  if  $f$  is  $m$  times differentiable [GO77, p. 28].
  - ▶  $\mathcal{O}(C^n)$ , for  $C < 1$ , if  $f$  is analytical in a neighborhood of  $[-1, 1]$  [BT04].
- ▶ Very used in practice:
  - ▶ Present in external libraries like NumPy for Python.
  - ▶ Matlab tool Chebfun: allows functions to be treated as vectors.

## Example

$\xi$  and its approximation of degree 8.




Figura: An example of degree 8 approximation.



## Example

$\xi$  and its approximation of degree 10.




Figura: An example of degree 10 approximation.

## Example

$\xi$  and its approximation of degree 12.




Figura: An example of degree 12 approximation.

# Chebyshev Interpolation

Choosing the degree of the interpolation:

- ▶ There is no guaranteed way to choose it.
- ▶ A good rule is to examine the last coefficient (the last coefficient rule-of-thumb [Boy01, p .50]).
- ▶ For a predefined  $\epsilon$ , choose  $n$ , such that:

$$|a_n| \leq \epsilon$$

- ▶ There are other ways like checking the error on a Lobatto grid [BG07].

# Chebyshev Interpolation

For  $n = 32$ , a precision of  $10^{-10}$  is expected.




Figura:  $|a_n|$  for the interpolation of  $\xi$  for an instance.

# Chebyshev Interpolation

## Roots

The roots of a Chebyshev polynomial can be found though determining the eigenvalues of a Chebyshev companion matrix [Boy13]. For  $n = 5$ , we have:

$$\begin{bmatrix} 0 & 1 & 0 & 0 & 0 \\ \frac{1}{2} & 0 & \frac{1}{2} & 0 & 0 \\ 0 & \frac{1}{2} & 0 & \frac{1}{2} & 0 \\ 0 & 0 & \frac{1}{2} & 0 & \frac{1}{2} \\ -\frac{a_0}{2a_5} & -\frac{a_1}{2a_5} & -\frac{a_2}{2a_5} & -\frac{a_3}{2a_5} & -\frac{a_4}{2a_5} \end{bmatrix}$$

- ▶ This is a Hessenberg matrix.
- ▶ Its eigenvalues can be found by a QR decomposition in  $\mathcal{O}(n^3)$ .

# Chebyshev Interpolation

## Roots

The largest error on roots that were found for  $n = 32$  is around  $10^{-14}$  for an instance:




Figura:  $|\xi(\hat{\theta})|$ , where  $\hat{\theta}$  is a root of  $f_n$ .

# Chebyshev Interpolation

## Roots

The experiments were made using Python with the NumPy library.  
The running time is really low, even for  $n = 64$ .




Figura: The running time to find the roots of  $f_n$ .

## Example

An example with 4 solutions.













Figura: An example with 4 solutions.



# References I

-  John P. Boyd and Daniel H. Gally, *Numerical experiments on the accuracy of the chebyshev–frobenius companion matrix method for finding the zeros of a truncated series of chebyshev polynomials*, Journal of Computational and Applied Mathematics **205** (2007), no. 1, 281 – 295.
-  John P. Boyd, *Chebyshev and Fourier spectral methods*, second ed., Dover Books on Mathematics, Dover Publications, Mineola, NY, 2001.
-  John Boyd, *Finding the zeros of a univariate equation: Proxy rootfinders, chebyshev interpolation, and the companion matrix*, SIAM Review **55** (2013).
-  Zachary Battles and Lloyd N Trefethen, *An extension of MATLAB to continuous functions and operators*, SIAM Journal on Scientific Computing **25** (2004), no. 5, 1743–1770.

# References II

-  Walter Gautschi, *The condition of polynomials in power form*, Mathematics of Computation - Math. Comput. **33** (1979).
-  D. Gottlieb and S.A. Orszag, *Numerical analysis of spectral methods: Theory and applications*, CBMS-NSF Regional Conference Series in Applied Mathematics, Society for Industrial and Applied Mathematics, 1977.
-  Roger A. Horn and Charles R. Johnson (eds.), *Matrix analysis*, Cambridge University Press, New York, NY, USA, 1986.
-  R.A. Johnson and Y.W. Young, *Advance euclidean geometry (modern geometry): An elementary treatise on the geometry of the triangle and the circle*, Dover books on advanced mathematics, Dover, 1960.
-  J. C. Mason and D. C. Handscomb, *Chebyshev polynomials*, Chapman & Hall/CRC, Boca Raton, FL, 2003. MR MR1937591 (2004h:33001)

# References III



M. J. D. (Michael James David) Powell, *Approximation theory and methods*, Cambridge [England] ; New York : Cambridge University Press, 1981 (English), Includes index.