

Finite difference scheme for a Kirchhoff thin plate with free edges

Daniel Tang

November 25, 2018

The vibrations in a thin, isotropic, homogeneous plate can be described as

$$\frac{\partial^2 u}{\partial t^2} = -\kappa^2 \Delta u$$

where

$$\kappa = \frac{EH^2}{12\rho(1-\nu^2)}$$

where E is the Youngs modulus of the material, H is the thickness of the plate, ρ is its density and ν is its Poisson ratio.

For free edges, we have the boundary conditions (along a left-hand edge):

$$\frac{\partial^2 u}{\partial x^2} + \nu \frac{\partial^2 u}{\partial y^2} = 0$$

and

$$\frac{\partial^3 u}{\partial x^3} + (2-\nu) \frac{\partial^3 u}{\partial xy^2} = 0$$

while at a corner

$$\frac{\partial^2 u}{\partial xy} = 0$$

Boundary conditions for other edge orientations can be inferred by symmetry.

Using the notation ∂_x , ∂_{xx} etc to mean the second order, central finite difference approximation of $\frac{\partial}{\partial x}$, $\frac{\partial^2}{\partial x^2}$ etc. and ∂_{x+} and ∂_{x-} to mean the second order, forward and backward finite difference respectively, we can express the thin plate equation as a finite difference scheme:

$$\partial_{tt}u = -\kappa(\partial_{xx} + \partial_{yy})(\partial_{xx} + \partial_{yy})u$$

The boundary conditions along a left-hand edge can be expressed

$$\partial_{xx}u + \nu\partial_{yy}u = 0 \tag{1}$$

$$\partial_{x-}(\partial_{xx}u + (2-\nu)\partial_{yy}u) = 0 \tag{2}$$

and at a bottom-left corner as

$$\partial_{x-y-}u = 0 \tag{3}$$

0.1 Laplacian finite difference at an edge

At a left-hand edge, from equation 1 we have

$$\partial_{xx} = -\nu\partial_{yy}$$

so

$$\Delta = \partial_{xx} + \partial_{yy} = (1 - \nu)\partial_{yy} \quad (4)$$

Rearranging equation 2 gives

$$\partial_{x-}(\Delta + (1 - \nu)\partial_{yy}) = 0$$

so, at a left-hand edge

$$\partial_{x-}\Delta = (1 - \nu)\partial_{x-yy} \quad (5)$$

but

$$\partial_{xx}\Delta = \frac{\partial_{x+}\Delta - \partial_{x-}\Delta}{Dx} = \frac{\partial_{x+}\Delta - (1 - \nu)\partial_{x-yy}}{Dx} \quad (6)$$

where Dx is the grid spacing in the x direction.

Differentiating equation 1 with respect to y gives

$$\partial_{xxyy} = \frac{\partial_{x+yy} - \partial_{x-yy}}{Dx} = -\nu\partial_{yyyy}$$

so

$$\partial_{x-yy} = \partial_{x+yy} + Dx\nu\partial_{yyyy}$$

substituting back into equation 6

$$\partial_{xx}\Delta = \frac{\partial_{x+}\Delta - (1 - \nu)(\partial_{x+yy} + Dx\nu\partial_{yyyy})}{Dx}$$

but, differentiating equation 4 by y gives

$$\partial_{yy}\Delta = (1 - \nu)\partial_{yyyy}$$

so, at a left-hand edge

$$\Delta\Delta = \frac{\partial_{x+}\Delta - (1 - \nu)\partial_{x+yy}}{Dx} - (1 - \nu)^2\partial_{yyyy} \quad (7)$$

0.2 Laplacian finite difference at a corner

At the bottom-left corner we have, from equation 1,

$$\partial_{xx} = -\nu\partial_{yy}$$

and

$$\partial_{yy} = -\nu\partial_{xx}$$

Since ν is positive, this implies $\partial_{xx} = 0$ and $\partial_{yy} = 0$ so $\Delta = 0$

At the corner, we cannot differentiate past the corner, so we need to calculate ∂_{x-yy} in a different way:

$$\partial_{x-yy} = \frac{\partial_{x-}(\partial_{y+} - \partial_{y-})}{Dy}$$

so, from equation 3

$$\partial_{x-yy} = \frac{\partial_{x-y+}}{Dy}$$

However, along the left-hand edge

$$\frac{\partial_{x+} - \partial_{x-}}{Dx} = -\nu \partial_{yy}$$

so

$$\partial_{x-} = Dx\nu\partial_{yy} + \partial_{x+}$$

so

$$\partial_{x-yy} = \frac{\partial_{y+}(Dx\nu\partial_{yy} + \partial_{x+})}{Dy}$$

substituting this into equation 6

$$\partial_{xx}\Delta = \frac{Dy\partial_{x+}\Delta - (1-\nu)\partial_{y+}(Dx\nu\partial_{yy} + \partial_{x+})}{DxDy}$$

by symmetry

$$\partial_{yy}\Delta = \frac{Dx\partial_{y+}\Delta - (1-\nu)\partial_{x+}(Dy\nu\partial_{xx} + \partial_{y+})}{DxDy}$$

so, at the bottom-left corner

$$\Delta\Delta = \frac{Dx\partial_{y+}\Delta + Dy\partial_{x+}\Delta - (1-\nu)(\partial_{y+}(Dx\nu\partial_{yy} + \partial_{x+}) + \partial_{x+}(Dy\nu\partial_{xx} + \partial_{y+}))}{DxDy} \quad (8)$$