Species Distribution Modeling of *Xantusia vigilis*

Presenters: Daniel Furman and Sarah Halvorsen Advisors: Stephen Adolph, PhD and Tanja Srebotnjak, PhD November 23, 2019

Background






www.inaturalist.org

Research Questions

1. What is the set of climate conditions that defines *X. vigilis's* geographical distribution?

2. How will the distribution shift as climate changes?

X. vigilis and climate




Rapid climate changes could alter the suitability of an ecosystem for *X. vigilis*.


Xantusia Fecundity and Rainfall at Fort Tejon (Zweifel and Lowe, 1966)

Species Distribution Models


- Estimates geographical niche of an organism
- Correlates organism locations with climate variables




Occurrence Points





Background Points






Maxent Distribution






Maxent Distribution









Random Forest Distribution







General Logistic Regression Distribution




Model Overlap







Three Model Overlap

Variable Name	Average (1970-2000)
Precipitation driest month	2.59 mm
Min temp coldest month	-1.62 °C
Elevation	1003 m




Known Geographic Range







Conservative Range vs. Accepted Range



Projection



Future prediction is
50.3%
of current distribution
(using 2070 CNRM-CM5,
4.5 rcp climate model)



Conclusions



Conclusions

- All models show high agreement with previous range estimates.
- Models show climatic suitability, but tell us nothing about X.
 vigilis's abundance or long-term movements.
- Future Work:
 - More rigorous testing of model parameters
 - More projections to future climates
 - Test X. vigilis's response to new environmental conditions

Acknowledgments and References

We would like to express our gratitude to professors Stephen Adolph and Tanja **Srebotnjak at Harvey Mudd College** for their guidance and advice throughout the project, and our fellow undergraduate **Kyra Clark**. We would also like to thank students **Daniela Elizondo**, Deyana Marsh, and Steven Witkin for their help revising our final work. Furthermore, we would like to acknowledge Drs. Hijmans' and **Elith**'s tutorial for modelling species' distributions in R^H. Their tutorial gave us a fundamental understanding of SDMs and provided our project with a basic foundation to start our work. Finally, we would like to thank Professor **Steven Williams** of James Cook University for suggestions and guidance.

- A. Bezy, R. L. 1989. Night Lizards: The evolution of habitat specialists. *Terra* 28: 29-34.
- B. Bezy, R. L. 2014. Desert night lizard, *Xantusia vigilis* Baird, 1859. *Sonoran Herpetologist* 27: 7-10.
- C. Routson, C.C. *et al.* 2019. Mid-latitude net precipitation decreased with Arctic warming during the Holocene. *Nature* 568: 83-87. Available at https://doi.org/10.1038/s41586-019-1060-3
- D. Zweifel, R.G., Lowe, C.H. 1966. The ecology of a population of *Xantusia vigilis*, the desert night lizard. *Am. Mus. Novit.* 2247.: 1-57.
- E. Fick, S.E. and R.J. Hijmans, 2017. Worldclim 2: New 1-km spatial resolution climate surfaces for global land areas. *International Journal of Climatology. / http://www.worldclim.org/*
- F. R Core Team (2019). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. https://www.R-project.org/.
- G. Bezy, R. L. 1982. *Xantusia vigilis. Catalogue of American Amphibians and Reptiles* 302: 1-4.
- H. Cole et al. 2011. Past and Ongoing Shifts in Joshua Tree Distribution. *Ecological Applications*. 21: 137-149
- Hijmans, R. J. and Elith J. 2017. Species Distribution Modeling in R. (and references within) https://cran.r-project.org/web/packages/dismo/vignettes/sdm.pdf

Overlap with Y. brevifolia: Environmental Space

- Method: Principal component analysis
- Convex hull volumes

o X. vigilis: 531

Y. brevifolia: 517

