

Horn Clauses in Hybrid-Dynamic First-Order Logic

Daniel Găină

Institute of Mathematics for Industry, Kyushu University, Japan

Department of Mathematics and Statistics, La Trobe University, Australia

daniel@imi.kyushu-u.ac.jp

Ionuț Tuțu

Simion Stoilow Institute of Mathematics of the Romanian Academy, Romania

Department of Computer Science, Royal Holloway University of London, UK

ittutu@gmail.com

Abstract

We propose a hybrid-dynamic first-order logic as a formal foundation for specifying and reasoning about reconfigurable systems. As the name suggests, the formalism we develop extends (many-sorted) first-order logic with features that are common to hybrid logics and to dynamic logics. This provides certain key advantages for dealing with reconfigurable systems, such as: (a) a signature of nominals, including operation and relation symbols, that allows references to specific possible worlds / system configurations – as in the case of hybrid logics; (b) distinguished signatures of rigid and flexible symbols, where the rigid symbols are interpreted uniformly across possible worlds; this supports a rigid form of quantification, which ensures that variables have the same interpretation regardless of the possible world where they are evaluated; (c) hybrid terms, which increase the expressive power of the logic in the context of rigid symbols; and (d) modal operators over dynamic-logic actions, which are defined as regular expressions over binary nominal relations. We then study Horn clauses in this hybrid-dynamic logic, and develop a series of results that lead to an initial-semantics theorem for arbitrary sets of clauses. This shows that a significant fragment of hybrid-dynamic first-order logic has good computational properties, and can serve as a basis for defining executable languages for reconfigurable systems. Lastly, we set out the foundations of logic programming in this fragment by proving a hybrid-dynamic variant of Herbrand’s theorem, which reduces the semantic entailment of a logic-programming query by a program to the search of a suitable answer substitution.

2012 ACM Subject Classification Theory of computation → Modal and temporal logics; Theory of computation → Logic and verification

Keywords and phrases Hybrid logic, Dynamic logic, Horn clause, Herbrand’s theorem

1 Introduction

The dynamic-reconfiguration paradigm is a most promising approach in the development of highly complex and integrated systems of interacting ‘components’, which now often evolve dynamically, at run time, in response to internal or external stimuli. More than ever, we are witnessing a continuous increase in the number of applications with reconfigurable features, many of which have aspects that are safety- or security-critical. This calls for suitable formal-specification and verification technologies, and there is already a significant body of research on this topic; hybrid(ized) logics [2, 13], first-order dynamic logic [11], and modal μ -calculus [10] are three prominent examples, among many others.

In this paper, we focus on those reconfigurable systems whose states or configurations can be presented explicitly, based on some kind of context-independent data types, and for which we distinguish the computations performed at the local/configuration level from the dynamic evolution of the configurations. This suggests a two-layered approach to the design and analysis of reconfigurable systems, involving a *local view*, which amounts to describing the structural properties of configurations, and a *global view*, which corresponds to a specialized language for specifying and reasoning about the way system configurations evolve.

For that purpose, we propose a new modal logical system for the reconfiguration paradigm that is obtained by enriching first-order logic (regarded as a base logic, or parameter for the whole construction) with both hybrid and dynamic features. More specifically, we model reconfigurable systems as Kripke structures (or transition systems), where:

- from a local perspective, we consider a dedicated first-order signature for configurations, and hence capture configurations as first-order structures for that signature; and
- from a global perspective, we consider a second first-order signature for the possible worlds of the Kripke structure; the terms over that signature are *nominals* used to identify configurations, and the binary nominal relations are regarded as *modalities*, which capture the transitions, or accessibility relations, between configurations.

Concerning the syntax of the logic proposed, sentences are build from equations and relational atoms over the two first-order signatures mentioned above (one pertaining to data, and the other to possible worlds) by using Boolean connectives, quantifiers, standard hybrid-logic operators such as *retrieve* and *store*, and dynamic-logic operators such as *necessity* over structured actions, which are defined as regular expressions over modalities.

The construction is reminiscent of the hybridization of institutions from [13, 5] and of the hybrid-dynamic logics presented in [1, 12], but it departs fundamentally from any of those studies due to the fact that the possible worlds of the Kripke structures that we consider here have an algebraic structure. This special feature of the logic that we put forward is extremely important for dealing with reconfigurable systems whose states are obtained from initial configurations by applying constructor operations; see, e.g. [8]. In this context, we advance a general notion of Horn clause, which allows the use of implications, universal quantifiers, as well as the hybrid- and dynamic-logic operators listed above. We study congruences over Kripke structures, and show how these can be used in quotienting initial Kripke models with respect to Horn clauses. Initial semantics has received a lot of attention in the formal-specification literature, where it has traditionally been linked to formalisms that support execution by means of rewriting. Here, we explore the key role of initiality in logic programming – where the initial models are typically referred to as least Herbrand models – and we prove a hybrid-dynamic version of Herbrand’s theorem. This a fundamental result, because it establishes a connection between the semantic entailment of a logic-programming query and the existence of an answer substitution for that query – which is a syntactic construct, and can subsequently be obtained by proof-theoretic means.

The paper is structured as follows: Section 2 is devoted to the definition of the new logic, which we call hybrid-dynamic first-order logic with user-defined sharing (this last attribute is meant to indicate the fact that users have control over the symbols that should be interpreted the same across the possible worlds of a Kripke structure); then, in Section 3 we introduce Horn clauses and prove the main initiality results of the paper (for atomic sentences and for Horn clauses), while in Section 4 we focus on queries and Herbrand’s theorem; lastly, in Sections 5 and 6 we discuss related work and potential future research.

2 Hybrid-Dynamic First-Order Logic with user-defined Sharing

The hybrid-dynamic first-order logic (HDFOLS) that we propose in this paper is rooted in the same ideas that underlie hybrid first-order logic [2] and hybrid first-order logic with rigid symbols [5, 4]. Its signatures contain distinguished subsets of so-called *rigid symbols*, which may be sorts, operation or relation symbols, and are meant to have the same interpretation across the possible worlds of a given Kripke structure. This provides support for the most common form of quantification in modal logic, where the semantics of variables is rigid. We

develop these ideas further by considering first-order structures of possible worlds.

We present HDFOLS from an institutional perspective [9], meaning that we focus on signatures and signature morphisms (though, for the purpose of this paper, inclusions would suffice), Kripke structures and homomorphisms, sentences, and the (local) satisfaction relation and condition that relate the syntax and the semantics of HDFOLS. However, other than the notations used, the text requires no prior knowledge of institution theory, and should be accessible to readers with a general background in modal and first-order model theory.

Signatures The *signatures* of HDFOLS are tuples of the form $\Delta = (\Sigma^n, \Sigma^r \subseteq \Sigma)$, where:

1. $\Sigma^n = (S^n, F^n, P^n)$ is a first-order signature of *nominals* such that $S^n = \{\star\}$ is a singleton,
2. $\Sigma^r = (S^r, F^r, P^r)$ is a first-order signature of so-called *rigid symbols*, and
3. $\Sigma = (S, F, P)$ is a first-order signature of both *rigid* and *flexible* symbols.

We denote by F^f and P^f the sub-families of F and P , respectively, that consist of *flexible symbols* (obtained by removing the rigid symbols). Throughout the paper, we generally denote by Δ and Δ' signatures of the form $(\Sigma^n, \Sigma^r \subseteq \Sigma)$ and $(\Sigma'^n, \Sigma'^r \subseteq \Sigma')$, respectively.

Signature morphisms A *signature morphism* $\varphi: \Delta \rightarrow \Delta'$ consists of a pair of first-order signature morphisms $\varphi^n: \Sigma^n \rightarrow \Sigma'^n$ and $\varphi^r: \Sigma^r \rightarrow \Sigma'^r$ such that $\varphi(\Sigma^r) \subseteq \Sigma'^r$.

Kripke structures The *models* of a signature Δ are pairs (W, M) , where:

1. W is a Σ^n -model, for which we denote by $|W|$ the carrier set of the only sort \star in Σ^n , and
2. $M = \{M_w\}_{w \in |W|}$ is a family of Σ -models, indexed by *possible worlds* $w \in |W|$, such that the rigid symbols have the same interpretation across the worlds; that is, $M_{w_1, \varsigma} = M_{w_2, \varsigma}$ for all possible worlds $w_1, w_2 \in |W|$ and all symbols ς in Σ^r .

Kripke homomorphisms A *morphism* $h: (W, M) \rightarrow (W', M')$ is also a pair, given by a Σ^n -homomorphism $h: W \rightarrow W'$ and, for each $w \in |W|$, a Σ -homomorphism $h_w: M_w \rightarrow M'_{h(w)}$, such that $h_{w_1, s} = h_{w_2, s}$ for all possible worlds $w_1, w_2 \in |W|$ and all rigid sorts $s \in S^r$.

Actions As in dynamic logic, HDFOLS supports structured actions obtained from atoms using sequential composition, union, and iteration. The set A^n of *actions* over Σ^n is defined inductively, according to the following grammar: $\alpha ::= \lambda \in P_{**}^n \mid \alpha \circ \alpha \mid \alpha + \alpha \mid \alpha^*$.

Actions are interpreted in Kripke structures as *accessibility relations* between possible worlds. This is done by extending the interpretation of binary modalities (symbols in P_{**}^n) as follows: $W_{\alpha_1 \circ \alpha_2} = W_{\alpha_1} \circ W_{\alpha_2}$ (diagrammatic composition of relations), $W_{\alpha_1 + \alpha_2} = W_{\alpha_1} \cup W_{\alpha_2}$ (union of relations), and $W_{\alpha^*} = (W_\alpha)^*$ (reflexive and transitive closure of a relation).

Hybrid terms For every Σ^n -model W , the family $T^W = \{T_w^W\}_{w \in |W|}$ of hybrid terms over W is defined inductively as the least family of (sorted) sets satisfying the following rules:¹

$$\begin{array}{lll}
 (1) \frac{w \in |W| \quad \tau \in T_{w, ar}^W}{\sigma(\tau) \in T_{w, s}^W} & (2) \frac{w \in |W| \quad \tau \in T_{w, ar}^W}{\sigma(w, \tau) \in T_{w, s}^W} & (3) \frac{w_0, w \in |W| \quad \tau \in T_{w_0, s}^W}{\tau \in T_{w, s}^W} \\
 \text{where } \sigma \in F_{ar \rightarrow s}^r & \text{where } \sigma \in F_{ar \rightarrow s}^f & \text{where } s \in S^r
 \end{array}$$

Notice that flexible operation symbols receive a possible world $w \in |W|$ as an extra argument while the rigid operation symbols keep their initial arity. Also, hybrid terms of rigid sorts are shared across the worlds. These are important considerations for Proposition 2 below.

Given a world $w \in |W|$, the S -sorted set T_w^Δ can be regarded as a Σ -model by interpreting every rigid operation symbol $\sigma: ar \rightarrow s$ as the function that maps hybrid terms $\tau \in T_{w, ar}^W$ to $\sigma(\tau)$, every flexible operation symbol $\sigma: ar \rightarrow s$ as the function that maps hybrid terms $\tau \in T_{w, ar}^W$ to $\sigma(w, \tau)$, and every relation symbol (rigid or flexible) as the empty set. A proof of the following lemma, as well as of Proposition 2, can be found in the Appendix.

¹ For brevity, we often use τ or t to denote not only terms, but also tuples of terms.

135 ▶ **Lemma 1** (Hybrid-term model and its freeness). *For every Σ^n -model W , (W, T^W) is a Δ -*
 136 *model. Moreover, for any Δ -model (W', M') and first-order Σ^n -homomorphism $f: W \rightarrow W'$*
 137 *there exists a unique Δ -homomorphism $h: (W, T^W) \rightarrow (W', M')$ that agrees with f on W . ◀*

138 **Standard term model** When W is the first-order term model T_{Σ^n} , by Lemma 1 we obtain
 139 the standard hybrid-term model over Δ , which we denote by $(T_{\Sigma^n}, \{T_k^\Delta\}_{k \in T_{\Sigma^n}})$.

140 The initiality of the standard term model provides a straightforward interpretation of
 141 hybrid terms in Δ -models (W, M) : for every hybrid term $t \in T_k^\Delta$, we denote by $(W, M)_t$ or
 142 $M_{h(k),t}$ the image of t under h_k , where h is the unique homomorphism $(T_{\Sigma^n}, T^\Delta) \rightarrow (W, M)$.

143 ▶ **Proposition 2** (Reachability of the hybrid-term model). *If W is a reachable Σ^n -model,*
 144 *meaning that the first-order homomorphism $T_{\Sigma^n} \rightarrow W$ is surjective, then (W, T^W) is reachable*
 145 *too; i.e. the unique homomorphism $h: (T_{\Sigma^n}, \{T_k^\Delta\}_{k \in T_{\Sigma^n}}) \rightarrow (W, \{T_w^W\}_{w \in W})$ is surjective. ◀*

146 **Sentences** The atomic sentences ρ defined over a HDFOLS-signature Δ are given by:

$$147 \quad \rho ::= k_1 = k_2 \mid \lambda(k') \mid t_1 =_{k,s} t_2 \mid \varpi(t) \mid \pi(k, t)$$

148 where $k, k_i \in T_{\Sigma^n}$ are nominal terms, k' is a tuple of terms corresponding to the arity of
 149 $\lambda \in P^n$, $t_i \in T_{k,s}^\Delta$ and $t \in T_{k,ar}^\Delta$ are (tuples of) hybrid terms,² $\varpi \in P_{ar}^r$, and $\pi \in P_{ar}^f$. We
 150 refer to these sentences, in order, as *nominal equations*, *nominal relations*, *hybrid equations*,
 151 *rigid hybrid relations*, and *non-rigid/flexible hybrid relations*, respectively. When there is no
 152 danger of confusion, we may drop one or both subscripts k, s from the notation $t_1 =_{k,s} t_2$.

153 *Full sentences* over Δ are built from atomic sentences according to the following grammar:

$$154 \quad \gamma ::= \rho \mid \mathbf{a}(k_1, k_2) \mid @_k \gamma \mid \neg \gamma \mid \bigwedge \Gamma \mid \downarrow z \cdot \gamma' \mid \forall X \cdot \gamma'' \mid [\mathbf{a}] \gamma \mid (\sigma) \gamma$$

155 where $k, k_i \in T_{\Sigma^n}$ are nominal terms, $\mathbf{a} \in A^n$ is an action, Γ is a finite set of sentences, z is a
 156 nominal variable, γ' is a sentence over the signature $\Delta[z]$ obtained by adding z as a new
 157 constant to F^n , X is a block of nominal and/or rigid variables, γ'' is a sentence over the
 158 signature $\Delta[X]$ obtained by adding the elements of X as a new constants to F^n and F^r , and
 159 σ is a unary operation symbol in F^n . Other than the first two kinds of sentences (*atoms* and
 160 *action relations*), we refer to the sentence-building operators, in order, as *retrieve*, *negation*,
 161 *conjunction*, *store*, *universal quantification*, *necessity*, and *next*, respectively.

162 We denote by $\text{Sen}^{\text{HDFOLS}}(\Delta)$ the set of all HDFOLS-sentences over Δ .

163 **The local satisfaction relation** Given a Kripke structure (W, M) for Δ and a possible world
 164 $w \in |W|$, we define the *satisfaction of Δ -sentences at w* by structural induction as follows:

- 165 1. *For atomic sentences:*
 - 166 – $(W, M) \models^w k_1 = k_2$ iff $W_{k_1} = W_{k_2}$ for all nominal equations $k_1 = k_2$;
 - 167 – $(W, M) \models^w \lambda(k)$ iff $W_k \in W_\lambda$ for all nominal relations $\lambda(k)$;
 - 168 – $(W, M) \models^w t_1 =_k t_2$ iff $M_{w',t_1} = M_{w',t_2}$, where $w' = W_k$, for all equations $t_1 =_k t_2$;
 - 169 – $(W, M) \models^w \varpi(t)$ iff $(W, M)_t \in M_{w,\varpi}$ for all rigid relations $\varpi(t)$;
 - 170 – $(W, M) \models^w \pi(k, t)$ iff $(W, M)_t \in M_{w',\pi}$, where $w' = W_k$, for flexible relations $\pi(k, t)$.
- 171 2. *For full sentences:*
 - 172 – $(W, M) \models^w \mathbf{a}(k_1, k_2)$ iff $(W_{k_1}, W_{k_2}) \in W_{\mathbf{a}}$ for all action relations $\mathbf{a}(k_1, k_2)$;
 - 173 – $(W, M) \models^w @_k \gamma$ iff $(W, M) \models^{w'} \gamma$, where $w' = W_k$;
 - 174 – $(W, M) \models^w \neg \gamma$ iff $(W, M) \not\models^w \gamma$;
 - 175 – $(W, M) \models^w \bigwedge \Gamma$ iff $(W, M) \models^w \gamma$ for all $\gamma \in \Gamma$;

² Note that, if the arity ar is rigid, then the sets $\{T_{k,ar}^\Delta\}_{k \in T_{\Sigma^n}}$ coincide.

- 176 $(W, M) \models^w \downarrow z \cdot \gamma$ iff $(W, M)^{z \leftarrow w} \models^w \gamma$,
- 177 where $W^{z \leftarrow w}$ is the unique $\Delta[z]$ -expansion³ of W that interprets the variable z as w ;
- 178 $(W, M) \models^w \forall X \cdot \gamma$ iff $(W', M') \models^w \gamma$ for all $\Delta[X]$ -expansions (W', M') of (W, M) ;
- 179 $(W, M) \models^w [\mathbf{a}] \gamma$ iff $(W, M) \models^{w'} \gamma$ for all worlds $w' \in |W|$ such that $(w, w') \in W_{\mathbf{a}}$;
- 180 $(W, M) \models^w (\sigma) \gamma$ iff $(W, M) \models^{w'} \gamma$, where $w' = W_{\sigma}(w)$.

181 \triangleright **Fact 3.** The next properties follow easily from the definition of the satisfaction relation.

182 1. The satisfaction of atoms and of action relations ρ does not depend on the possible worlds.

183 More specifically, $(W, M) \models^w \rho$ iff $(W, M) \models^{w'} \rho$ for all possible worlds $w, w' \in |W|$.

184 2. The satisfaction of atoms and of action relations ρ is preserved by homomorphisms. That

185 is, if $(W, M) \models \rho$ and there is a homomorphism $(W, M) \rightarrow (W', M')$, then $(W', M') \models \rho$.

186 To state the *satisfaction condition* – and thus finalize the presentation of HDFOLS – let

187 us first notice that every signature morphism $\varphi: \Delta \rightarrow \Delta'$ induces appropriate *translations*

188 *of sentences* and *reductions of models*, as follows: every Δ -sentence γ is translated to a

189 Δ' -sentence $\varphi(\gamma)$ by replacing (usually in an inductive manner) the symbols in Δ with

190 symbols from Δ' according to φ ; and every Δ' -model (W', M') is reduced to a Δ -model

191 $(W', M') \downarrow_{\varphi}$ that interprets every symbol x in Δ as $(W', M')_{\varphi(x)}$. When φ is an inclusion,

192 we usually denote $(W', M') \downarrow_{\varphi}$ by $(W', M') \downarrow_{\Delta}$ – in this case, the model reduct simply forgets

193 the interpretation of those symbols in Δ' that do not belong to Δ .

194 The following satisfaction condition can be proved by induction on the structure of

195 Δ -sentences. Its argument is virtually identical with several other variants presented in the

196 literature (see, e.g. [4]), hence we choose to present the result without a proof.

197 \triangleright **Proposition 4** (Local satisfaction condition for signature morphisms). *For every signature*

198 *morphism $\varphi: \Delta \rightarrow \Delta'$, Δ' -model (W', M') , world $w' \in |W'|$, and Δ -sentence γ , we have:*

$$199 \quad (W', M') \models^{w'} \varphi(\gamma) \quad \text{if and only if} \quad (W', M') \downarrow_{\varphi} \models^{w'} \gamma.^4 \quad \blacktriangleleft$$

200 **Substitutions** Consider two signature extensions $\Delta[X]$ and $\Delta[Y]$ with blocks of variables,

201 and let $X = X^n \cup X^r$ and $Y = Y^n \cup Y^r$ be the partitions of X and Y into blocks of

202 nominal and rigid variables. A Δ -substitution $\theta: X \rightarrow Y$ consists of a pair of functions

203 $\theta^n: X^n \rightarrow T_{\Sigma^n[Y^n]}$ and $\theta^r: X^r \rightarrow T_k^{\Delta[Y]}$, where k is a nominal term.⁵

204 Similarly to signature morphisms, Δ -substitutions $\theta: X \rightarrow Y$ determine translations of

205 $\Delta[X]$ -sentences into $\Delta[Y]$ -sentences, and reductions of $\Delta[Y]$ -models to $\Delta[X]$ -models. The

206 proofs of the next two propositions are similar to the ones given in [7] for hybrid substitutions.

207 \triangleright **Proposition 5** (Local satisfaction condition for substitutions). *For every Δ -substitution*

208 *$\theta: X \rightarrow Y$, $\Delta[Y]$ -model (W, M) , world $w \in |W|$, and $\Delta[X]$ -sentence γ , we have:*

$$209 \quad (W, M) \models^w \theta(\gamma) \quad \text{if and only if} \quad (W, M) \downarrow_{\theta} \models^w \gamma. \quad \blacktriangleleft$$

210 Proposition 5, together with Proposition 6 below, has an important technical role in the

211 initial-model and Herbrand's-theorem proofs presented in the later sections of the paper.

212 \triangleright **Proposition 6** (Substitutions generated by expansions of reachable models). *If (W, M) is*

213 *a reachable Δ -model, then for every $\Delta[X]$ -expansion (W', M') of (W, M) there exists a*

214 *substitution $\theta: X \rightarrow \mathcal{O}$ such that $(W, M) \downarrow_{\theta} = (W', M')$.* \blacktriangleleft

³ In general, by a $\Delta[X]$ -expansion of (W, M) we understand a $\Delta[X]$ -model (W', M') that interprets all symbols in Δ in the same way as (W, M) . In particular, (W', M') has the same carrier sets as (W, M) .

⁴ Note that, by the definition of model reducts, (W', M') and $(W', M') \downarrow_{\varphi}$ have the same possible worlds.

⁵ Since the sorts of the variables are rigid, it does not matter which nominal term k we choose.

Expressive power Fact 3 highlights one of the main distinguishing features of HDFOLS: the satisfaction of atomic sentences, whether they involve flexible symbols or not, does not depend on the possible world where the sentences are evaluated. This contrasts the standard approach in hybrid logic, where each nominal is regarded as an atomic sentence satisfied precisely at the world that corresponds to the interpretation of that nominal. In HDFOLS, the dependence of the satisfaction of sentences on possible worlds is explicit rather than implicit, and is achieved through the *store* operator. These two approaches are often inter-definable: every nominal sentence k is semantically equivalent to $\downarrow z \cdot z = k$;⁶ moreover, when the logical system admits existential quantification over nominal variables, every *store* sentence $\downarrow z \cdot \gamma$ is semantically equivalent to $\exists z \cdot (z \wedge \gamma)$. Following the lines of [7, Section 4.3], the first part of this correspondence can be used to show that even without considering action relations, HDFOLS is strictly more expressive than other standard hybrid logics constructed from the same base logic such as the hybrid first-order logic with rigid symbols [5, 4].

As it is often the case, other sentence-building operators can be derived from those presented on page 4 through standard constructions: for instance, *implication* can be defined based on conjunction and negation, *existential quantification* based on universal quantification and negation, and the modal-logic *possibility* operator based on necessity and negation.

Necessity and *next* can be derived as well, since all sentences $[a]\gamma$ and $(\sigma)\gamma$ are semantically equivalent to $\downarrow z \cdot \forall z' \cdot a(z, z') \Rightarrow @_{z'} \gamma$ and $\downarrow z \cdot @_{\sigma(z)} \gamma$, respectively. However, we have chosen to present them independently, as there are logical systems that do not support the *store* operator or *universal quantifiers*. The results in this paper are modular; their proofs are not based on semantic abbreviations such as those listed above, nor on the existence of certain operators – with one exception: the main theorems rely on the existence of *retrieve*.

3 Initiality

Hybrid-dynamic *Horn clauses* are obtained from atomic sentences by repeated applications of (1) retrieve (2) implication such that the condition is a conjunction of atomic sentences or action relations, (3) store, (4) universal quantification, (5) necessity, or (6) next. We denote by HDCLS the Horn-clause fragment of HDFOLS – i.e. with the same signatures and models as HDFOLS, and with sentences defined as Horn clauses. This restriction is crucial, because it entails that any set of sentences in HDCLS has an initial model, as we prove next.

3.1 The basic level

We first deal with the initiality property for the basic level of atoms. The following result can also be found in [7], though in a simplified setting where Σ^n consists only of constants. A detailed proof, adapted to hybrid-dynamic first-order logic can be found in the Appendix.

► **Proposition 7** (Initiality of nominal equations and relations). *Every set Γ_n of nominal equations or relations admits a reachable initial model $(W^{\Gamma_n}, M^{\Gamma_n})$.* ◀

In order to extend Proposition 7 to arbitrary sets of HDCLS-sentences, we use a notion of congruence on a Kripke structure and the universal property of its corresponding quotient.

► **Definition 8** (Congruence). *Let (W, M) be a Kripke structure over a HDFOLS-signature Δ . A Δ -congruence $\equiv = \{\equiv_w\}_{w \in |W|}$ on (W, M) is a family of Σ -congruences \equiv_w on M_w , for each $w \in |W|$, such that $(\equiv_{w_1, s}) = (\equiv_{w_2, s})$ for all $w_1, w_2 \in |W|$ and $s \in S^r$.*

⁶ Formally, in hybrid logic, $(W, M) \models^w k$ if and only if $w = W_k$.

The following construction and universal property are straightforward generalizations of their first-order counterparts, and have been studied in several other papers in the literature (see, e.g. [6]). For that reason, we include them for further reference without a proof.

► **Proposition 9** (Quotient model). *Every HDCLS-congruence \equiv on a Δ -model (W, M) determines a quotient-model homomorphism $(_/\equiv): (W, M) \rightarrow (W, M/\equiv)$ that acts as an identity on possible worlds, and for which $(M/\equiv)_w$ is the quotient Σ -model M_w/\equiv_w .*

Moreover, $(_/\equiv)$ has the following universal property: for any Kripke homomorphism $h: (W, M) \rightarrow (W', M')$ such that $\equiv \subseteq \ker(h)$,⁷ there exists a unique homomorphism $h': (W, M/\equiv) \rightarrow (W', M')$ such that $(_/\equiv) \circ h' = h$.⁸

► **Proposition 10.** *Let Γ be a set of nominal or hybrid equations over Δ , and W a reachable Σ^n -model such that $\Gamma \models \Gamma_W$,⁹ where $\Gamma_W = \{k_1 = k_2 \in \text{Sen}(\Sigma^n) \mid W_{k_1} = W_{k_2}\}$. Then Γ generates a congruence \equiv on (W, T^W) defined by $\tau_1 \equiv_w \tau_2$ iff there exist a nominal term k and hybrid terms t_1 and t_2 such that $\Gamma \models t_1 =_k t_2$, $w = W_k$, and $\tau_i = T_{w, t_i}^W$.*

Proof. We first show that for every model (W', M') of Γ there exists a homomorphism $(W, T^W) \rightarrow (W', M')$. Since W is reachable, the unique homomorphism $q: T_{\Sigma^n} \rightarrow W$ is surjective. If $(W', M') \models \Gamma$, then $W' \models \Gamma_W$. Therefore, the unique homomorphism $f: T_{\Sigma^n} \rightarrow W'$ satisfies $\ker(q) \subseteq \ker(f)$. By the universal property of first-order model quotients, there exists a morphism $h: W \rightarrow W'$ such that $q \circ h = f$, which can then be uniquely extended to a Kripke homomorphism $h: (W, T^W) \rightarrow (W', M')$ as per Lemma 1.

The above observation allows us to prove that the definition of \equiv_w does not depend on the choice of k . That is, if $w = W_k = W_{k'}$ and $\Gamma \models t_1 =_k t_2$, then there exist $t'_1, t'_2 \in T_{k'}^\Delta$ such that $T_{w, t_i}^W = T_{w, t'_i}^W$ and $\Gamma \models t'_1 =_{k'} t'_2$. It suffices to consider the following list of inferences:

- 1 $T_{w, t_1}^W = T_{w, t'_1}^W$ for some $t'_1, t'_2 \in T_{k'}^\Delta$ by Proposition 2
- 2 $(W', M')_{t_i} = h(T_{w, t_i}^W) = h(T_{w, t'_i}^W) = (W', M')_{t'_i}$ since there exists a morphism
for all Δ -models (W', M') such that $(W', M') \models \Gamma$ $h: (W, T^W) \rightarrow (W', M')$
- 3 $(W', M')_{t_1} = (W', M')_{t_2}$ since $\Gamma \models t_1 =_k t_2$
for all Δ -models (W', M') such that $(W', M') \models \Gamma$
- 4 $(W', M')_{t'_1} = (W', M')_{t'_2}$ from 2 and 3
for all Δ -models (W', M') such that $(W', M') \models \Gamma$
- 5 $\Gamma \models t'_1 =_{k'} t'_2$ by the definition of \models

The fact that \equiv_w is both reflexive and symmetric is a straightforward consequence of the reachability of T^W . Therefore, we focus on the transitivity and the compatibility of \equiv_w with the operations in F . For transitivity, suppose $\tau_1 \equiv_w \tau_2$ and $\tau_2 \equiv_w \tau_3$. Then:

- 1 $\Gamma \models t_1 =_k t_2$, $w = W_k$, and $\tau_i = T_{w, t_i}^W$ for $i \in \{1, 2\}$, by the definition of \equiv_w
for some nominal term k and hybrid terms $t_1, t_2 \in T_k^\Delta$
- 2 $\Gamma \models t'_2 =_k t'_3$, $w = W_k$, and $\tau_i = T_{w, t'_i}^W$ for $i \in \{2, 3\}$, by the definition of \equiv_w
for some nominal term k and hybrid terms $t'_2, t'_3 \in T_k^\Delta$
- 3 $(W', M')_{t_2} = h(T_{w, t_2}^W) = h(T_{w, t'_2}^W) = (W', M')_{t'_2}$ since there exists a morphism
for all Δ -models (W', M') such that $(W', M') \models \Gamma$ $h: (W, T^W) \rightarrow (W', M')$
- 4 $\Gamma \models t_2 =_k t'_2$ by the definition of \models

⁷ This means that $h_{w,s}(a_1) = h_{w,s}(a_2)$ for all $a_1, a_2 \in M_{w,s}$ such that $a_1 \equiv_{w,s} a_2$.

⁸ We use the diagrammatic notation for function composition, where $((_/\equiv) \circ h')(a) = h'(a/\equiv)$.

⁹ By the definition of *semantic entailment*, this means that all models satisfying Γ satisfy Γ_W as well.

- 5 $\Gamma \models t_1 =_k t'_3$ from 1, 4, and 2
 6 $\tau_1 \equiv_w \tau_3$ by the definition of \equiv_w

281 For the compatibility of \equiv_w with the operations in F , assume $\sigma \in F_{ar \rightarrow s}$ and $\tau_1, \tau_2 \in T_{w, ar}^W$
 282 such that $\tau_1 \equiv_w \tau_2$. There is virtually no distinction between the case where σ is rigid and
 283 the case where σ is flexible, hence it suffices to make explicit the proof for $\sigma \in F_{ar \rightarrow s}^r$:

- 1 $\Gamma \models t_1 =_k t_2$, $w = W_k$, and $\tau_i = T_{w, t_i}^W$ for $i \in \{1, 2\}$, by the definition of \equiv_w
 for some nominal k and hybrid terms $t_1, t_2 \in T_k^\Delta$
 2 $\Gamma \models \sigma(t_1) =_k \sigma(t_2)$ by the general properties of \models
 3 $T_{w, \sigma(t_1)}^W \equiv_w T_{w, \sigma(t_2)}^W$ by the definition of \equiv_w
 4 $(T_w^W)_\sigma(\tau_1) \equiv_w (T_w^W)_\sigma(\tau_2)$ since $T_{w, \sigma(t_i)}^W = (T_w^W)_\sigma(\tau_i)$

284 Finally, we need to show that the relations $\{\equiv_w\}_{w \in |W|}$ coincide on rigid sorts, that is
 285 $(\equiv_{w, s}) = (\equiv_{w', s})$ for all $w, w' \in |W|$ and $s \in S^r$. Suppose $\tau_1 \equiv_{w, s} \tau_2$. It follows that:

- 1 $\Gamma \models t_1 =_{k, s} t_2$, $w = W_k$, and $\tau_i = T_{w, t_i}^W$ for $i \in \{1, 2\}$, by the definition of $\equiv_{w, s}$
 for some nominal k and hybrid terms $t_1, t_2 \in T_{k, s}^\Delta$
 2 $w' = W_{k'}$ for some nominal k' since W is reachable
 3 $\Gamma \models t_1 =_{k', s} t_2$ and $t_1, t_2 \in T_{k', s}^\Delta$ since s is rigid
 4 $\tau_1 \equiv_{w', s} \tau_2$ by the definition of $\equiv_{w', s}$ \blacktriangleleft

286 Unlike first-order congruences and the Δ -congruences presented in Definition 8, the
 287 general notion of Kripke congruence is significantly more complex when it involves the
 288 formation of a quotient model for the possible worlds. In this paper, we consider congruences
 289 over hybrid-term models, for which the quotients can be obtained in two stages: first on
 290 possible worlds, then on their local models. This constitutes a step forward in understanding
 291 Kripke congruences and their implications. Therefore, Proposition 10 is an interesting result
 292 in its own right, since it deals with the congruence generated by a set of equations. We use
 293 it next as a basis for proving the existence of initial models for atomic sentences.

294 **► Theorem 11 (Atomic initiality).** *Every set Γ of atomic sentences over a HDCLS-signature*
 295 *$\Delta = (\Sigma^n, \Sigma^r \subseteq \Sigma)$ admits a reachable initial model (W^Γ, M^Γ) .*

296 **Proof.** Let Γ_n be the subset of nominal equations and relations in Γ . By Proposition 7, there
 297 exists a reachable initial model $(W^{\Gamma_n}, M^{\Gamma_n})$ of Γ_n . For notational convenience, we denote the
 298 interpretation of a (nominal or hybrid) term t in $(W^{\Gamma_n}, M^{\Gamma_n})$ by $[t]$.

299 For each $w \in |W^{\Gamma_n}|$, we define a relation \equiv_w on $M_w^{\Gamma_n}$ as follows: $\tau_1 \equiv_w \tau_2$ if and only
 300 if there exist a nominal k and hybrid terms $t_1, t_2 \in T_k^\Delta$ such that $w = [k]$, $\tau_i = [t_i]$, and
 301 $\Gamma \models t_1 =_k t_2$. Since W^{Γ_n} is reachable, by Proposition 10, \equiv is a Δ -congruence on $(W^{\Gamma_n}, M^{\Gamma_n})$.

302 By Proposition 9, the congruence \equiv determines a quotient-model homomorphism $(_/\equiv)$
 303 between $(W^{\Gamma_n}, M^{\Gamma_n})$ and $(W^{\Gamma_n}, M^{\Gamma_n}/\equiv)$. Let (W^Γ, M^Γ) be the Δ -model obtained from
 304 $(W^{\Gamma_n}, M^{\Gamma_n}/\equiv)$ by interpreting, for every $w \in |W^{\Gamma_n}|$, each relation symbol

305 \blacksquare $\pi \in P^r$ as $M_{w, \pi}^\Gamma = \{[t]/\equiv_w \in M_w^{\Gamma_n}/\equiv_w \mid t \in T_k^\Delta, [k] = w, \text{ and } \Gamma \models \pi(t)\}$, and

306 \blacksquare $\pi \in P^f$ as $M_{w, \pi}^\Gamma = \{[t]/\equiv_w \in M_w^{\Gamma_n}/\equiv_w \mid t \in T_k^\Delta, [k] = w, \text{ and } \Gamma \models \pi(k, t)\}$.

307 Trivially, since $(W^{\Gamma_n}, M^{\Gamma_n})$ is reachable and $(_/\equiv)$ is surjective, then (W^Γ, M^Γ) is reachable
 308 as well. Therefore, all we need to prove is that (W^Γ, M^Γ) is an initial model of Γ .

309 The fact that (W^Γ, M^Γ) is a model of Γ follows in a straightforward manner from the very
 310 construction of (W^Γ, M^Γ) . We focus on the initiality property. Let (W, M) be a Δ -model
 311 satisfying Γ . Since $\Gamma_n \subseteq \Gamma$, we have $(W, M) \models \Gamma_n$, and thus, by Proposition 7, there exists a
 312 unique homomorphism $h_n: (W^{\Gamma_n}, M^{\Gamma_n}) \rightarrow (W, M)$. Moreover, for every $\tau_1 \equiv_w \tau_2$ we have:

- 1 $\Gamma \models t_1 =_k t_2$, $w = [k]$, and $\tau_i = [t_i]$ for $i \in \{1, 2\}$, by the definition of \equiv_w
for some nominal term k and hybrid terms $t_1, t_2 \in T_k^\Delta$
- 2 $(W, M) \models t_1 =_k t_2$ since $(W, M) \models \Gamma$
- 3 $h_n(\tau_1) = h_n(\tau_2)$ since $h_n(\tau_i) = h_n([t_i]) = (W, M)_{t_i}$

313 This shows that $\equiv \subseteq \ker(h_n)$. By Proposition 9, there exists a unique Kripke homomorph-
314 ism $h: (W^{\Gamma_n}, M^{\Gamma_n}/\equiv) \rightarrow (W, M)$ such that $(_/\equiv) \circ h = h_n$. To finalize the proof, we need to
315 show that h preserves the interpretation of the relation symbols in Δ .

316 We consider only the case where π is a rigid relation symbol. Non-rigid relation symbols
317 can be treated in a similar manner. Assume $a \in M_{w,\pi}^\Gamma$ for some $w \in |W^\Gamma|$. It follows that:

- 1 $\Gamma \models \pi(t)$, $w = [k]$, and $a = [t]$ by the definition of $M_{w,\pi}^\Gamma$
for some nominal term k and hybrid term $t \in T_k^\Delta$
- 2 $M_{W_k,t} \in M_{W_k,\pi}$ since $(W, M) \models \Gamma$
- 3 $M_{h(w),t} \in M_{h(w),\pi}$ since $h(w) = h([k]) = W_k$
- 4 $h_w(a) \in M_{h(w),\pi}$ since $h_w(a) = h_w([t]) = M_{h(w),t}$ ◀

318 ▶ **Corollary 12.** *Under the notations and hypotheses of Theorem 11, we have that $\Gamma \models \rho$ if*
319 *and only if $(W^\Gamma, M^\Gamma) \models \rho$, for all atomic sentences or action relations ρ over Δ .*

320 **Proof.** Since (W^Γ, M^Γ) is a model of Γ , the ‘only if’ part holds trivially by the definition of
321 the semantic entailment. For the ‘if’ part, let ρ be an atomic sentence or an action relation
322 over Δ such that $(W^\Gamma, M^\Gamma) \models \rho$, and (W, M) an arbitrary Δ -model that satisfies Γ . Since
323 (W^Γ, M^Γ) is an initial model of Γ , there exists a unique homomorphism $(W^\Gamma, M^\Gamma) \rightarrow (W, M)$.
324 And because the satisfaction of both atomic sentences and action relations is preserved by
325 homomorphisms (by Fact 3), it follows that $(W, M) \models \rho$. Therefore, $\Gamma \models \rho$. ◀

326 3.2 The upper level

327 In this subsection, we lift the initiality property from the basic level to the level of Horn
328 clauses. To that end, let us denote by HDFOLS_0 the atomic fragment of HDFOLS .

329 ▶ **Proposition 13.** *Let Γ be a set of Horn clauses over Δ and $(W^{\Gamma_0}, M^{\Gamma_0})$ an initial model*
330 *of the set $\Gamma_0 = \{\rho \in \text{Sen}^{\text{HDFOLS}_0}(\Delta) \mid \Gamma \models \rho\}$. For each nominal k and Horn clause γ over Δ ,*

$$331 \quad \Gamma \models @_k \gamma \quad \text{implies} \quad (W^{\Gamma_0}, M^{\Gamma_0}) \models @_k \gamma.$$

332 **Proof.** We prove the result by induction on the structure of the clauses γ built over Δ .
333 $[\gamma = \rho \in \text{Sen}^{\text{HDFOLS}_0}(\Delta)]$ If $\Gamma \models @_k \rho$, then we obtain:

- 1 $@_k \rho \models \rho$ since ρ is atomic
- 2 $\Gamma \models \rho$ since $\Gamma \models @_k \rho \models \rho$
- 3 $\rho \in \Gamma_0$ by the definition of Γ_0
- 4 $(W^{\Gamma_0}, M^{\Gamma_0}) \models \rho$ by Corollary 12
- 5 $(W^{\Gamma_0}, M^{\Gamma_0}) \models @_k \rho$ since $@_k \rho \models \rho$

334 $[@_{k'} \gamma]$ This case is trivial, because $@_k @_{k'} \gamma \models @_{k'} \gamma$.

335 $[\bigwedge H \Rightarrow \gamma]$ Suppose $\Gamma \models @_k (\bigwedge H \Rightarrow \gamma)$ and $(W^{\Gamma_0}, M^{\Gamma_0}) \models^w H$, where $w = W_k^{\Gamma_0}$. Then:

- 1 $(W^{\Gamma_0}, M^{\Gamma_0}) \models H$ by Fact 3
- 2 $\Gamma_0 \models H$ by Corollary 12
- 3 $\Gamma \models H$ since $\Gamma \models \Gamma_0$

- 4 $\Gamma \models @_k \gamma$ since $H \cup \{ @_k (\bigwedge H \Rightarrow \gamma) \} \models @_k \gamma$
5 $(W^{\Gamma_0}, M^{\Gamma_0}) \models @_k \gamma$ by the induction hypothesis
6 $(W^{\Gamma_0}, M^{\Gamma_0}) \models^w \gamma$ since $w = W_k^{\Gamma_0}$
7 $(W^{\Gamma_0}, M^{\Gamma_0}) \models^w \bigwedge H \Rightarrow \gamma$ from 1–6, by the definition of \models
8 $(W^{\Gamma_0}, M^{\Gamma_0}) \models @_k (\bigwedge H \Rightarrow \gamma)$ by the definition of \models
- 336 $[\downarrow z \cdot \gamma]$ This case is straightforward, because $@_k \downarrow z \cdot \gamma \models @_k \theta_{z \leftarrow k}(\gamma)$, where $\theta_{z \leftarrow k}: \{z\} \rightarrow \emptyset$
337 is the Δ -substitution that maps the nominal variable z to the (ground) term k .
338 $[\forall X \cdot \gamma]$ Suppose $\Gamma \models @_k \forall X \cdot \gamma$ and let (W, M) be a $\Delta[X]$ -expansion of $(W^{\Gamma_0}, M^{\Gamma_0})$. Since
339 $(W^{\Gamma_0}, M^{\Gamma_0})$ is reachable, by Proposition 6, there exists a Δ -substitution $\theta: X \rightarrow \emptyset$ such
340 that $(W^{\Gamma_0}, M^{\Gamma_0}) \upharpoonright_{\theta} = (W, M)$. It follows that:
- 1 $\Gamma \models @_k \theta(\gamma)$ since $\Gamma \models @_k \forall X \cdot \gamma$
2 $(W^{\Gamma_0}, M^{\Gamma_0}) \models @_k \theta(\gamma)$ by the induction hypothesis
3 $(W^{\Gamma_0}, M^{\Gamma_0}) \models^w \theta(\gamma)$, where $w = W_k^{\Gamma_0}$ by the definition of \models
4 $(W, M) \models^w \gamma$ by the local satisfaction condition for θ
5 $(W^{\Gamma_0}, M^{\Gamma_0}) \models^w \forall X \cdot \gamma$ by 4, since (W, M) is an arbitrary $\Delta[X]$ -expansion of $(W^{\Gamma_0}, M^{\Gamma_0})$
6 $(W^{\Gamma_0}, M^{\Gamma_0}) \models @_k \forall X \cdot \gamma$ by the definition of \models
- 341 $[\mathbf{a}\gamma]$ Suppose $\Gamma \models @_k [\mathbf{a}\gamma]$ and let $(w, w') \in W_a^{\Gamma_0}$ such that $w = W_k^{\Gamma_0}$. Since $(W^{\Gamma_0}, M^{\Gamma_0})$
342 is reachable, there exists a nominal term k' such that $w' = W_{k'}^{\Gamma_0}$. It follows that:
- 1 $(W^{\Gamma_0}, M^{\Gamma_0}) \models \mathbf{a}(k, k')$ since $(w, w') \in W_a^{\Gamma_0}$
2 $\Gamma_0 \models \mathbf{a}(k, k')$ by Corollary 12
3 $\Gamma \models \mathbf{a}(k, k')$ since $\Gamma \models \Gamma_0$
4 $\Gamma \models @_{k'} \gamma$ since $\{ @_k [\mathbf{a}\gamma], \mathbf{a}(k, k') \} \models @_{k'} \gamma$
5 $(W^{\Gamma_0}, M^{\Gamma_0}) \models @_{k'} \gamma$ by the induction hypothesis
6 $(W^{\Gamma_0}, M^{\Gamma_0}) \models^{w'} \gamma$ since $w' = W_{k'}^{\Gamma_0}$
7 $(W^{\Gamma_0}, M^{\Gamma_0}) \models @_k [\mathbf{a}\gamma]$ from 1–6, by the definition of \models
- 343 $[(\sigma)\gamma]$ This case is trivial, because $@_k (\sigma)\gamma \models @_{\sigma(k)} \gamma$. \blacktriangleleft

344 Proposition 13 shows that the semantic consequences of a set Γ of Horn clauses are
345 satisfied by the initial models of the set Γ_0 of atomic sentences. This is a very useful insight
346 into the initiality property, which can be explored for other logical systems as well.

347 **► Theorem 14** (Horn-clause initiality). *Let Γ be a set of Horn clauses over a signature Δ and*
348 *$(W^{\Gamma_0}, M^{\Gamma_0})$ a reachable initial model of Γ_0 . Then $(W^{\Gamma_0}, M^{\Gamma_0})$ is also an initial model of Γ .*

349 **Proof.** Since $\Gamma \models \Gamma_0$, it follows that for every model (W, M) of Γ there exists a unique homo-
350 morphism $(W^{\Gamma_0}, M^{\Gamma_0}) \rightarrow (W, M)$. Therefore, what remains to prove is that $(W^{\Gamma_0}, M^{\Gamma_0})$ is a
351 model of Γ . Let $w \in |W^{\Gamma_0}|$ and $\gamma \in \Gamma$. Since $(W^{\Gamma_0}, M^{\Gamma_0})$ is reachable, there exists a nominal
352 term k such that $w = W_k^{\Gamma_0}$. Moreover, since $\Gamma \models \gamma$, we have $\Gamma \models @_k \gamma$. By Lemma 13, we
353 deduce that $(W^{\Gamma_0}, M^{\Gamma_0}) \models @_k \gamma$, which is equivalent to $(W^{\Gamma_0}, M^{\Gamma_0}) \models^w \gamma$. \blacktriangleleft

354 4 Herbrand's theorem

355 A query for a signature Δ is a sentence of the form $\exists X \cdot \bigwedge E$ such that X is a set of variables
356 and E is a finite set of atomic sentences or action relations. The following result, which is

fundamental for the development of logic programming for HDFOLS, states that establishing the (semantic) entailment of a query by a program – understood here as a set of Horn clauses – is equivalent to finding a (syntactic) *correct-answer substitution* for that query.

► **Theorem 15** (Herbrand's theorem). *Consider a HDFOLS-signature Δ , a set Γ of Horn clauses, and a query $\exists X \cdot \bigwedge E$ over Δ . The following statements are equivalent:*

1. $\Gamma \models \exists X \cdot \bigwedge E$.
2. $(W^\Gamma, M^\Gamma) \models \exists X \cdot \bigwedge E$, where (W^Γ, M^Γ) is an initial model of Γ .
3. $\Gamma \models \theta(E)$ for some Δ -substitution $\theta: X \rightarrow \emptyset$.

Proof. Note that, by Theorem 14, Γ has indeed a *reachable* initial reachable model (W^Γ, M^Γ) .

[1 \Rightarrow 2] By the fact that $(W^\Gamma, M^\Gamma) \models \Gamma$.

[2 \Rightarrow 3] Let w be an arbitrary possible world of (W^Γ, M^Γ) . It follows that:

1. $(W, M) \models^w E$ for some $\Delta[X]$ -expansion (W, M) of (W^Γ, M^Γ) by the definition of \models
2. $(W, M) = (W^\Gamma, M^\Gamma) \upharpoonright_\theta$ for some Δ -substitution $\theta: X \rightarrow \emptyset$ since (W^Γ, M^Γ) is reachable
3. $(W^\Gamma, M^\Gamma) \models \theta(E)$ by the local sat. cond. for θ
4. $(W', M') \models \theta(E)$ for all models (W', M') such that $(W', M') \models \Gamma$ by Fact 3, based on the universal property of (W^Γ, M^Γ)
5. $\Gamma \models \theta(E)$ by the definition of \models

[3 \Rightarrow 1] Let (W, M) be a model of Γ and $w \in |W|$. It follows that:

1. $(W, M) \models^w \theta(E)$ since $\Gamma \models \theta(E)$
2. $(W, M) \upharpoonright_\theta \models^w E$ by the local sat. cond. for θ
3. $(W, M) \models^w \exists X \cdot \bigwedge E$ since $(W, M) \upharpoonright_\theta \upharpoonright_\Delta = (W, M)$
4. $\Gamma \models \exists X \cdot \bigwedge E$ by the definition of \models ◀

5 Related work

This work builds on previous results presented by the first author in [7], and is related to developments in the initial semantics for hybridized institutions reported in [4] and in the foundations of logic-independent logic programming from [3, 14, 15]. Clearly, those related studies are all logic-independent, whereas here we made a deliberate choice to focus on a concrete logical system – still, many of the results developed in this paper can be lifted to an institution-independent setting. The hybrid-dynamic logic that we propose has a number of distinguishing features that take it outside the scope of the frameworks presented in [7, 4, 3]. More specifically, since possible worlds are no longer plain, but have an algebraic structure, one cannot employ the strategy used in [7] (or [4], for that matter) for proving Theorem 11. In fact, the nominal framework discussed therein can be regarded as an instance of the one we consider here for HDFOLS, where all nominals are constants. That restriction leads to a much simpler construction of the quotient models (and proof of the existence of initial models), where the quotienting on possible worlds can be done by means of signature morphisms whose action on nominal constants is surjective.

Compared to [4], the initiality results presented in this paper are based on constructive arguments that avoid the heavier model-theoretic infrastructure of quasi-varieties and inclusion systems. Moreover, those arguments hold as well for some classes of constrained models (e.g. models with reachable possible worlds) that do not form a quasi-variety.

Besides these technical advancements, it is also worth noting that the Horn clauses and queries examined in this paper exceed the expressive power of the clauses and queries considered in [7]. For instance, the logical implications used in clauses are no longer restricted to atoms; instead, the hypothesis of an implication can involve both atomic sentences and action relations (which in general do not have an initial model), and the conclusion can be any Horn clause. Moreover, queries can involve action relations too, which in general are not basic sentences. For that reason, the variant of Herbrand’s theorem presented here does not fit into the framework advanced in [3]. A more general approach to Herbrand’s theorem can be found in [15], but the result presented there relies on a much more complex institution-theoretic infrastructure, which is outside the scope of this paper.

6 Conclusions and further work

In this work, we have presented a new hybrid-dynamic logic for the specification and analysis of reconfigurable systems. From a semantic perspective, this allows us to capture reconfigurable systems as Kripke structures whose possible worlds (1) have an algebraic structure, which provides support for operations on system configurations, and (2) are labelled with constrained first-order models that capture the local/inner structure and behaviour of configurations. The model constraints that we consider here are given by rigid declarations of first-order symbols (sorts, or operation or relation symbols), which ensure a uniform interpretation of those symbols across the possible worlds. This kind of modelling is also reflected in the syntax of the logic, where we use nominal and hybrid terms to refer to possible worlds and to the elements of the first-order structures associated to those worlds. Terms are used to form nominal and hybrid equations, as well as relational atoms, from which we build complex sentences using Boolean connectives, quantifiers, and operators that are specific to hybrid logics and to dynamic logics. To the best of our knowledge, there is currently no other logical system that supports such an approach to the reconfiguration paradigm.

We have shown that every set of atomic hybrid-dynamic sentences admits a reachable initial model (Kripke structure), and then generalized this result to Horn clauses, which in this case have a highly complex structure, involving implications and universal quantifiers (as in first-order logic), but also hybrid-logic operators such as *retrieve* and *store*, and dynamic logic operators such as *necessity* over actions. This is a fundamental result of great importance, because it allows us to deal with hybrid-dynamic logic programs (formalized as sets of Horn clauses). To that end, we have studied a variant of Herbrand’s theorem – which is central to the logic-programming paradigm – for hybrid-dynamic first-order logic.

This work is part of a much broader research agenda on the rigorous formal development of reconfigurable systems. There are several clear tasks that we aim to pursue further. One of those is to generalize the results obtained here to an institution-independent setting so that we could apply them with ease to other logical systems, such as those of order-sorted algebra and of preordered algebra. The most convenient way to achieve that is by taking into account the relationship between the basic level and the upper level discussed in Section 3; the former is necessarily logic-dependent, whereas the latter can be developed for arbitrary logics. A second important task is to study Birkhoff completeness and resolution-based inference rules for hybrid-dynamic logics, for which we aim to follow [6, 15]. This will provide a minimal framework for the analysis of reconfigurable systems, and a foundation for developing appropriate tool support and a number of concrete case studies.

References

- 1 Brandon Bohrer and André Platzer. A hybrid, dynamic logic for hybrid-dynamic information flow. In *Proceedings of the 33rd Annual ACM/IEEE Symposium on Logic in Computer Science, LICS 2018, Oxford, UK, July 09–12, 2018*, pages 115–124. ACM, 2018.
- 2 Torben Braüner. *Hybrid Logic and its Proof-Theory*, volume 37 of *Applied Logic Series*. Springer, 2011.
- 3 Răzvan Diaconescu. Herbrand theorems in arbitrary institutions. *Information Processing Letters*, 90(1):29–37, 2004.
- 4 Răzvan Diaconescu. Quasi-varieties and initial semantics for hybridized institutions. *Journal of Logic and Computation*, 26(3):855–891, 2016.
- 5 Răzvan Diaconescu and Alexandre Madeira. Encoding hybridized institutions into first-order logic. *Mathematical Structures in Computer Science*, 26(5):745–788, 2016.
- 6 Daniel Găină. Birkhoff style calculi for hybrid logics. *Formal Aspects of Computing*, 29(5):805–832, 2017.
- 7 Daniel Găină. Foundations of logic programming in hybrid logics with user-defined sharing. *Theoretical Computer Science*, 686:1–24, 2017.
- 8 Daniel Găină, Ionuț Țuțu, and Adrián Riesco. Specification and verification of invariant properties of transition systems. In *25th Asia-Pacific Software Engineering Conference, APSEC 2018, Nara, Japan, 4–7 December 2018*. IEEE, 2018.
- 9 Joseph A. Goguen and Rod M. Burstall. Institutions: Abstract model theory for specification and programming. *Journal of the ACM*, 39(1):95–146, 1992.
- 10 Jan Friso Groote and Mohammad Reza Mousavi. *Modeling and Analysis of Communicating Systems*. MIT Press, 2014.
- 11 David Harel, Dexter Kozen, and Jerzy Tiuryn. Dynamic logic. *SIGACT News*, 32(1):66–69, 2001.
- 12 Rolf Hennicker, Alexandre Madeira, and Alexander Knapp. A hybrid dynamic logic for event/data-based systems. In *Fundamental Approaches to Software Engineering – 22nd International Conference, FASE 2019, Prague, Czech Republic, April 6–11, 2019, Proceedings*, volume 11424 of *LNCS*, pages 79–97. Springer, 2019.
- 13 Manuel A. Martins, Alexandre Madeira, Răzvan Diaconescu, and Luís Soares Barbosa. Hybridization of institutions. In Andrea Corradini, Bartek Klin, and Corina Cîrstea, editors, *Algebra and Coalgebra in Computer Science – 4th International Conference, CALCO 2011, Winchester, UK, August 30 – September 2, 2011. Proceedings*, volume 6859 of *Lecture Notes in Computer Science*, pages 283–297. Springer, 2011.
- 14 Ionuț Țuțu and José Luiz Fiadeiro. Revisiting the institutional approach to Herbrand’s theorem. In Lawrence S. Moss and Paweł Sobociński, editors, *6th Conference on Algebra and Coalgebra in Computer Science, CALCO 2015, June 24–26, 2015, Nijmegen, The Netherlands*, volume 35 of *LIPICs*, pages 304–319. Schloss Dagstuhl – Leibniz-Zentrum fuer Informatik, 2015.
- 15 Ionuț Țuțu and José Luiz Fiadeiro. From conventional to institution-independent logic programming. *Journal of Logic and Computation*, 27(6):1679–1716, 2017.

A Additional proofs

Proof of Lemma 1. The fact that (W, T^W) is a Δ -model is straightforward. For the ‘freeness’ part, we define $\{h_w : T_w^W \rightarrow M'_{f(w)}\}_{w \in |W|}$ by structural induction on hybrid terms:

1. $h_w(\sigma(\tau)) = M'_{f(w),\sigma}(h_w(\tau))$ for all $w \in |W|$, $\tau \in T_{w,ar}^W$, and $\sigma \in F_{ar \rightarrow s}^r$;
2. $h_w(\sigma(w, \tau)) = M'_{f(w),\sigma}(h_w(\tau))$ for all $w \in |W|$, $\tau \in T_{w,ar}^W$, and $\sigma \in F_{ar \rightarrow s}^f$;
3. $h_{w,s}(\tau) = h_{w_0,s}(\tau)$ for all $w_0, w \in |W|$, $\tau \in T_{w_0,s}^W$, and $s \in S^r$.

It is again straightforward to check that $h = (f, \{h_w\}_{w \in |W|})$ is a Kripke homomorphism $(W, T^W) \rightarrow (W', M')$, and (also by induction) that it is unique with this property. ◀

Proof of Proposition 2. By hypothesis, $h: T_{\Sigma^n} \rightarrow W$ is surjective. Therefore, what we need to prove is that, for every nominal term k , the Σ -homomorphism $h_k: T_k^\Delta \rightarrow T_{h(k)}^W$ is surjective. We proceed by induction on the structure of the hybrid terms over W :

1. Assume that $\tau \in T_{h(k), ar}^W$ and $\sigma \in F_{ar \rightarrow s}^r$. By the induction hypothesis, there exists a hybrid term $t \in T_{k, ar}^\Delta$ such that $h_k(t) = \tau$. Therefore, $h_k(\sigma(t)) = \sigma(h_k(t)) = \sigma(\tau)$.
2. Assume that $\tau \in T_{h(k), ar}^W$ and $\sigma \in F_{ar \rightarrow s}^f$. By the induction hypothesis, there exists a hybrid term $t \in T_{k, ar}^\Delta$ such that $h_k(t) = \tau$. Therefore, $h_k(\sigma(k, t)) = \sigma(h(k), h_k(t)) = \sigma(h(k), \tau)$.
3. Assume that $\tau \in T_{w_0, s}^W$ and $s \in S'$. Since W is reachable, $w_0 = h(k_0)$ for some nominal term k_0 . By the induction hypothesis, there exists $t \in T_{k_0, s}^\Delta$ such that $h_{k_0}(t) = \tau$. Finally, given that the sort s is rigid, we have $t \in T_{k, s}^\Delta$ and $h_k(t) = h_{k_0}(t) = \tau$. \blacktriangleleft

Proof of Proposition 7. Consider the following relation on the first-order term model T_{Σ^n} : $k_1 \equiv k_2$ if and only if $\Gamma_n \models k_1 = k_2$, for any two nominal terms k_1 and k_2 over Δ .

It is straightforward to check that \equiv is a Σ^n -congruence on T_{Σ^n} . Therefore, we can define W^{Γ_n} as the Σ^n -model obtained from T_{Σ^n}/\equiv by interpreting each relation symbol $\lambda \in P^n$ as the set $\{[k] \in T_{\Sigma^n}/\equiv \mid \Gamma_n \models \lambda(k)\}$, where $[k]$ denotes the congruence class of k modulo \equiv .

Now let $(W^{\Gamma_n}, M^{\Gamma_n})$ be the (reachable) hybrid-term model defined over W^{Γ_n} . That is, the possible worlds of $(W^{\Gamma_n}, M^{\Gamma_n})$ are congruence classes of nominal terms modulo \equiv , and for each nominal term k , the elements of $M_{[k]}^{\Gamma_n}$ are obtained from standard hybrid terms (according to the definition in Section 2) by replacing each nominal with its congruence class. For convenience, we denote the interpretation of a term t in $(W^{\Gamma_n}, M^{\Gamma_n})$ by $[t]$.

All is left to prove is the universal property of $(W^{\Gamma_n}, M^{\Gamma_n})$. First, notice that, by construction, we have $(W^{\Gamma_n}, M^{\Gamma_n}) \models \Gamma_n$. Then, for every Kripke model (W, M) that satisfies Γ_n , there exists a unique homomorphism $h: (W^{\Gamma_n}, M^{\Gamma_n}) \rightarrow (W, M)$, where:

- on possible worlds, $h: W^{\Gamma_n} \rightarrow W$ is the unique Σ^n -homomorphism between W^{Γ_n} and W , defined by $h([k]) = W_k$ for all $k \in T_{\Sigma^n}$, and
 - for every $k \in T_{\Sigma^n}$, $h_{[k]}: M_{[k]}^{\Gamma_n} \rightarrow M_{W_k}$ is defined by $h_{[k]}([t]) = (W, M)_t$ for all $t \in T_k^\Delta$.
- It is straightforward to check that the definition of h is consistent, since $(W, M) \models \Gamma_n$. \blacktriangleleft