

Discrete Mathematics

Daniel Gonzalez Cedre

University of Notre Dame
Spring of 2023

Chapter 2

First-Order Logic

2.1 Predicates & Quantification

Definition 2.1 (Universe of Discourse).

Our *universe of discourse*, usually denoted by the letter Ω , denotes the collection of objects under consideration. This specifies the objects that are admissible as inputs to predicates, meaning these are the objects we can actually make concrete, descriptive claims about using our formal language.

Definition 2.2 (Predicate).

Given a universe of discourse Ω and a non-negative integer n , we say that $\varphi(x_1, \dots, x_n)$ is an n -ary *predicate* $:\Leftrightarrow$ the substitution of an object ω_i from Ω for each x_i in φ results in a proposition $\varphi(\omega_1, \dots, \omega_n)$. These are sometimes referred to as *propositional functions*. Notice that if $n = 0$ then the 0-ary predicate is simply a proposition.

The placeholders x_1, \dots, x_n in φ are called *variables*, and the process of assigning objects to these variables is called *instantiation* or *variable assignment*.

Example 2.1.

If we choose a universe of discourse Ω consisting of some collection of people, then

$\mu(x) :=$ “ x is a mathematician.”

$\gamma(x, y) :=$ “ x drinks a Guinness with y .”

$\varepsilon(x) :=$ “ x loves espresso.”

$\alpha(x, y, z) :=$ “ x , y , and z are colleagues.”

are unary, binary, and ternary predicates respectively.

Definition 2.3 (Universal Quantifier).

Given a unary predicate $\varphi(x)$, the *universal quantification* of the variable x in φ is denoted by $\forall x(\varphi(x))$ and expresses that $\varphi(\omega)$ is true for any arbitrary ω from our universe of discourse.

Example 2.1.

Using the definitions from [Example 2.1](#), we can rewrite “Every mathematician loves espresso” as $\forall x(\varepsilon(x))$ or as $\forall x(\mu(x) \rightarrow \varepsilon(x))$ depending on whether our universe Ω is the collection of all mathematicians or of all people.

Definition 2.4 (Existential Quantifier).

Given a unary predicate $\varphi(x)$, the *existential quantification* of the variable x in φ is denoted by $\exists x(\varphi(x))$ and expresses that $\varphi(\omega)$ is true for at least one ω from our universe of discourse.

Example 2.1.

Using the definitions from [Example 2.1](#), we can rewrite “Some mathematician loves espresso” as $\exists x(\varepsilon(x))$ or as $\exists x(\mu(x) \wedge \varepsilon(x))$ depending on whether our universe Ω is the collection of all mathematicians or of all people.

2.2 Rules of Inference

Definition 2.5 (Rule of Inference).

A *rule of inference* is a construction in the meta-language that tells us how we're allowed to take previous statements in our formal language and derive new statements from them. They essentially describe the allowable algebraic manipulations we can make to the symbols in our language, and in this way they function analogously to the instruction set of a computer (if you're familiar with that). Another way to think of this is that they are truth-preserving axioms for our formal system; they are fundamental assumptions about the semantics of our formal language that allow us to reinterpret the statements we formulate in it. They usually take one of two forms: equivalences and inferences. For each of these rules, there is an *underlying tautology*, which is a tautological sentence in our formal language expressing the same idea as the rule.

An *equivalence rule* would take the form $\varphi \Leftrightarrow \psi$, where φ and ψ are sentences, indicating that any instance of φ can be replaced by an instance of ψ . The underlying tautology of such a rule is $p \leftrightarrow q$. A popular alternative notation is $\varphi \equiv \psi$.

An *inference rule* would take the form $\Gamma \Rightarrow \psi$ where Γ is a collection of sentences $\varphi_1, \dots, \varphi_n$ in our formal language and ψ is some other sentence, indicating that whenever we write down all of the sentences in Γ , we can then write down ψ afterward. The underlying tautology of such a rule is $\varphi_1 \wedge \dots \wedge \varphi_n \Rightarrow \psi$. A popular alternative notation is $\Gamma \vdash \psi$, or $\varphi_1, \dots, \varphi_n \vdash \psi$, indicating that ψ is deducible from the sentences in Γ .

We will stick to the \Leftrightarrow and \Rightarrow notation in this course since it is the most popular modern notation in use among practicing mathematicians. However, keep in mind that certain people (in particular, modern computer scientists, logicians, and set/model theorists) may instead use the \equiv and \vdash notation.

Equivalence Rules	
Identity	$p \wedge \top \Leftrightarrow p$ $p \vee \perp \Leftrightarrow p$
Complement (a.k.a. Negation)	$p \wedge \neg p \Leftrightarrow \perp$ $p \vee \neg p \Leftrightarrow \top$
Commutativity	$p \wedge q \Leftrightarrow q \wedge p$ $p \vee q \Leftrightarrow q \vee p$
Associativity	$p \wedge (q \wedge r) \Leftrightarrow (p \wedge q) \wedge r$ $p \vee (q \vee r) \Leftrightarrow (p \vee q) \vee r$
Distributivity	$p \wedge (q \vee r) \Leftrightarrow (p \wedge q) \vee (p \wedge r)$ $p \vee (q \wedge r) \Leftrightarrow (p \vee q) \wedge (p \vee r)$
Conditional Disintegration	$p \rightarrow q \Leftrightarrow \neg p \vee q$
Biconditional Disintegration	$p \leftrightarrow q \Leftrightarrow (p \rightarrow q) \wedge (q \rightarrow p)$

Unquantified Inference Rules		
Truth Axiom	We can always write \top without any prior assumptions	\top
Deduction ----- a.k.a. Conditional Introduction	If by assuming p we are able to derive q , then we can derive $p \rightarrow q$	$\frac{p \Rightarrow q}{p \rightarrow q}$
Modus Ponens	If we know $p \rightarrow q$ and we have p , then we can derive q	$\frac{p \quad p \rightarrow q}{q}$
Modus Tollens	If we know $p \rightarrow q$ and we have $\neg q$, then we can derive $\neg p$	$\frac{\neg q \quad p \rightarrow q}{\neg p}$
Reductio ad Absurdum ----- a.k.a. Proof by Contradiction	If by assuming p we can derive both q and $\neg q$, then we can derive $\neg p$	$\frac{p \Rightarrow q \quad p \Rightarrow \neg q}{\neg p}$

Quantified Inference Rules		
Universal Instantiation (\forall Elimination)	If we know $\forall x(\varphi(x))$, then we can derive $\varphi(t)$ for any t .	$\frac{\forall x(\varphi(x))}{\varphi(t) \text{ for any } t}$
Universal Generalization (\forall Introduction)	If we know $\varphi(t)$ for an arbitrary t , then we can derive $\forall x(\varphi(x))$.	$\frac{\varphi(t) \text{ for an arbitrary } t}{\forall x(\varphi(x))}$
Existential Instantiation (\exists Elimination)	If we know $\exists x(\varphi(x))$, then we can derive $\varphi(t)$ for some t that hasn't yet appeared.	$\frac{\exists x(\varphi(x))}{\varphi(t) \text{ for some } t \text{ that hasn't yet appeared}}$
Existential Generalization (\exists Introduction)	If we know $\varphi(t)$ for some particular t , then we can derive $\exists x(\varphi(x))$.	$\frac{\varphi(t) \text{ for some particular } t}{\exists x(\varphi(x))}$

Lemma 2.1.

Let φ be a predicate. If we know $\forall x(\neg\varphi(x))$, then we have $\neg\exists x(\varphi(x))$.

Proof. Suppose that $\forall x(\neg\varphi(x))$. Assume, towards a contradiction, that $\exists x(\varphi(x))$. Then, we know that $\varphi(t)$ for some t . From our first assumption, we know that $\neg\varphi(s)$ for any s ; in particular, we know that $\neg\varphi(t)$. But then, we obtain $\varphi(t) \wedge \neg\varphi(t)$, which is a contradiction. ⚡ Therefore, we can conclude $\neg\exists x(\varphi(x))$.

Q.E.D.

Lemma 2.2.

Let φ be a predicate. If we know $\neg\exists x(\varphi(x))$, then we have $\forall x(\neg\varphi(x))$.

Proof. Suppose that $\neg\exists x(\varphi(x))$ and let t be an arbitrary object. Towards a contradiction, assume that $\varphi(t)$. That would then mean $\exists x(\varphi(x))$. However, we already assumed $\neg\exists x(\varphi(x))$, which is absurd. ⚡ Therefore, we must have $\neg\varphi(t)$. Since t was arbitrary, we can then conclude $\forall x(\neg\varphi(x))$.

Q.E.D.

Lemma 2.3.

Let φ be a predicate. If we know $\exists x(\neg\varphi(x))$, then we have $\neg\forall x(\varphi(x))$.

Proof. Suppose that $\exists x(\neg\varphi(x))$. Then, we know that $\neg\varphi(t)$ for some t . Assume, towards a contradiction, that $\forall x(\varphi(x))$. Then, we have $\varphi(s)$ for any s ; in particular, we know $\varphi(t)$. But then, we have $\neg\varphi(t) \wedge \varphi(t)$, giving us a contradiction. ⚡ Therefore, we can conclude $\neg\forall x(\varphi(x))$.

Q.E.D.

Lemma 2.4.

Let φ be a predicate. If we know $\neg\forall x(\varphi(x))$, then we have $\exists x(\neg\varphi(x))$.

Proof. Suppose $\neg\forall x(\varphi(x))$. Towards a contradiction, assume that $\neg\exists x(\neg\varphi(x))$. Then by [Lemma 2.2](#), we get $\forall x(\neg\neg\varphi(x))$, which is equivalent to $\forall x(\varphi(x))$. However, this contradicts our assumption that $\neg\forall x(\varphi(x))$. ⚡ Therefore, we can conclude that $\neg\neg\exists x(\neg\varphi(x))$, which is equivalent to $\exists x(\neg\varphi(x))$.

Q.E.D.

Theorem 2.1 (Negation of Quantifiers).

For any predicate φ , we have

$$\begin{aligned}\neg\forall x(\varphi(x)) &\Leftrightarrow \exists x(\neg\varphi(x)) \\ \neg\exists x(\varphi(x)) &\Leftrightarrow \forall x(\neg\varphi(x))\end{aligned}$$

Proof. This follows from [Lemma 2.1](#), [2.2](#), [2.3](#), and [2.4](#). The details are left as an exercise for the reader.

Q.E.D.

Remark (Order of Quantification). Notice that the *order* of quantification often matters when writing or interpreting an expression, meaning $\forall x\exists y(\varphi(x, y)) \not\Leftrightarrow \exists y\forall x(\varphi(x, y))$. However, if two instances of the *same* quantifier occur immediately adjacent to each other, then they will commute: $\forall x\forall y(\varphi(x, y)) \Leftrightarrow \forall y\forall x(\varphi(x, y))$.

2.3 Validity of Arguments

Definition 2.6 (Validity).

Let $\varphi_1, \dots, \varphi_n$ and ψ be sentences in our language. Consider an argument of the form

$$\frac{\begin{array}{c} \varphi_1 \\ \vdots \\ \varphi_n \end{array}}{\therefore \psi}$$

We say that such an argument is *valid* \Leftrightarrow either of the following (equivalent) criteria is satisfied:

- I. The sentence $(\varphi_1 \wedge \dots \varphi_n) \rightarrow \psi$ is a tautology.
- II. We have $\varphi_1, \dots, \varphi_n \Rightarrow \psi$.

Condition **I**. states that *any* truth assignment that makes $\varphi_1 \wedge \dots \varphi_n$ true must also make ψ true. Condition **II**. states that, from the assumption that $\varphi_1, \dots, \varphi_n$ all hold, we *have a derivation* that ψ is the case by using the rules of equivalence and inference.

Definition 2.7 (Invalidity).

Conversely, we say the argument is *invalid* \Leftrightarrow either of the following (equivalent) criteria is satisfied:

- I. The sentence $(\varphi_1 \wedge \dots \varphi_n) \rightarrow \psi$ is *not* a tautology.
- II. We have $\varphi_1, \dots, \varphi_n \not\Rightarrow \psi$.

Condition **I**. states that *there is a* truth assignment that makes $\varphi_1 \wedge \dots \varphi_n$ true and also ψ false. Condition **II**. states that, from the assumption that $\varphi_1, \dots, \varphi_n$ all hold, we *can not have a derivation* that ψ is the case by using the rules of equivalence and inference.

Theorem 2.2 (Church's Theorem).

Validity of arguments in the first-order logic is undecidable.

This means that there is no Turing machine that, given an arbitrary sentence in the first-order logic, is able to correctly decide whether or not it has a proof in a finite amount of time.

Definition 2.8 (Term).

Given a universe of discourse Ω , we say that x is a *term* if it is either a specific member of Ω or a variable symbol denoting a not-yet-determined member of Ω . In class, we have been referring to these as *objects*.

Definition 2.9.

In our language, given a universe of discourse Ω , we have the following symbols:

- *Constant symbols*: terms denoting specific members of Ω (e.g., 7, Socrates, \mathbb{R} , \emptyset).
- *Variable symbols*: terms denoting members of Ω that are not-yet determined, usually written with a single letter from the Latin or Greek alphabets with or without natural number subscripts (e.g., x_1 , φ_{23} , y , β , A , Γ).
- *Predicate symbols*: n -ary predicates $\psi(x_1, \dots, x_n)$ taking the terms x_1, \dots, x_n as arguments and returning a proposition once all variables are eliminated (e.g., $=$, \leq , \in , \subseteq , \cong).
- *Functional symbols*: we haven't encountered any of these yet; they work similarly to predicates in that they take n terms as arguments, but they return a specific object of Ω as their output once all free variables are eliminated (e.g., $+$, \times , $\gcd(x, y)$, $|x|$, “ x is the father of y ”).
- *Grouping symbols*: the simple curved parentheses (and), used for grouping expressions.
- *Logical connectives*: the unary connective \neg and the binary connectives $\wedge, \vee, \rightarrow, \leftrightarrow$, which form a Boolean algebra along with \top and \perp .
- *Quantifier symbols*: the universal quantifier \forall and the existential quantifier \exists .

Once we specify some basic predicate and functional symbols, we can then define new predicates and functionals on the basis of the already-existing ones as the need arises.

Definition 2.10 (Atomic Formulæ).

We say φ is an *atomic formula* $:\Leftrightarrow$ we have $\varphi \Leftrightarrow \psi(x_1, \dots, x_n)$ for some n -ary predicate ψ with x_1, \dots, x_n being terms. Note that atomic formulæ may or may not contain free variables.

Definition 2.11 (Well-Formed Formulæ).

We define *well-formed formulæ* recursively (*i.e.*, inductively) using the atomic formulæ as our basis. We say that φ is a *well-formed formula* (abbreviated *wff*) $:\Leftrightarrow$ any of the following holds:

- I. φ is an atomic formula
- II. $\varphi \Leftrightarrow \neg(\psi)$, where ψ is a *wff*
- III. $\varphi \Leftrightarrow (\psi) \wedge (\chi)$, where ψ is a *wff*
- IV. $\varphi \Leftrightarrow (\psi) \vee (\chi)$, where ψ and χ are *wff*
- V. $\varphi \Leftrightarrow (\psi) \rightarrow (\chi)$, where ψ and χ are *wff*
- VI. $\varphi \Leftrightarrow (\psi) \leftrightarrow (\chi)$, where ψ and χ are *wff*
- VII. $\varphi \Leftrightarrow \forall x(\psi)$, where ψ is *wff*
- VIII. $\varphi \Leftrightarrow \exists x(\psi)$, where ψ is *wff*

Definition 2.12 (Sentence).

We say that a *wff* φ is a *sentence* $:\Leftrightarrow$ it contains no free variables.