

Monotone Catenary Degree of Numerical Monoids

Daniel Gonzalez Florida International University Cameron Wright Carleton College

Jenna Zomback SUNY Geneseo



Motivation

- In general, not much is known about the monotone catenary degree of numerical monoids. In the past, the monotone catenary degree in Krull Monoids has been studied, but for numerical monoids, only the regular catenary degree is well understood.
- We seek to gain a deeper understanding of the equivalent and adjacent catenary degrees in order to characterize the relationship between monotone and regular catenary degrees of numerical monoids.
- It is known that, for a numerical monoid M, $c_{mon}(M) \ge c(M)$. We aim to determine when this inequality is strict, and when the two quantities are equal.

Numerical Monoids

A **Numerical Monoid** is a cofinite subset of the nonnegative integers closed under the operation of addition. It is known that for every numerical monoid M, there exists a minimal set of generators n_1, \ldots, n_k , so for a monoid of this form we will write

$$M = \langle n_1, ..., n_k \rangle = \{a_1 n_1 + ... + a_k n_k \mid (a_1, ... a_k) \in \mathbb{N}^k\}$$

Example

 $M = \langle 4, 9, 11 \rangle = \{0, 4, 8, 9, 11, 12, 13, 15, ...\}$

▶ The **Set of Factorizations** of an element $m \in M$ is defined as

$$\mathcal{Z}(m) = \{(a_1, ..., a_k) \in \mathbb{N}^k \mid a_1 n_1 + ... + a_k n_k = m\}$$

Example

For $M = \langle 4, 9, 11 \rangle$, $\mathcal{Z}(26) = \{(2, 2, 0), (0, 1, 2)\}$

The **Length** of a factorization $z=(z_1,\ldots,z_k)\in\mathcal{Z}(m)$ is defined as $|z|=z_1+\cdots+z_k$

Example


Consider $(2,2,0) \in \mathcal{Z}(26) |2,2,0| = 4$

► We can define a **Distance** between factorizations based on the differences of the coordinates of the factorizations.

Example

d((2,2,0),(0,1,2))=3


Factorization Graph



Catenary Degrees


- ► Two factorizations z and z' are connected by an **N-chain** if there exists a sequence $z_0, ..., z_k \in \mathcal{Z}(m)$ such that $z_0 = z, ..., z_k = z'$ and $d(z_i, z_{i+1}) \leq N$ for all $i \in \{1, ...k 1\}$.
- The **catenary degree** of an element $m \in M$, c(m), is the minimum natural number N such that there is an N-chain between any two factorizations of m.
- Besides the regular catenary degree, we are also concerned with the equivalent, adjacent, and monotone catenary degrees. These behave similarly to the regular catenary degree but with added restrictions.

Catenary Graph



- We can remove the edge of length7 and our graph stays connected.
- If we remove the edge of length 6, our graph will be disconnected
- c(40) = 6

Different Types of Catenary Degrees




- Equivalent Catenary Degree only deals with factorizations of the same length
- Adjacent Catenary Degree deals with moving from one length factorization to the next
- Monotone Catenary Degree is the maximum of these two values

When is $c_{mon}(M) = c(M)$?

For any monoid generated by an arithmetic sequence, $M = \langle a, a+d, ..., a+kd \rangle$, $c_{mon}(M) = c(M)$. Furthermore, for any element $m \in M$, $c_{mon}(m) = c(m)$.

Monoids Generated by Arithmetic Sequences



Do Generalized Arithmetic Monoids Behave the Same Way?

In monoids generated by generalized arithmetic sequences $M = \langle a, ah + d, ah + 2d \rangle$ the monotone catenary is more nuanced. Our research has led us to make the following observations:

- If gcd(h-1,d) > 1, then we have that $c(M) = c_{mon}(M)$.
- if gcd(h-1,d)=1, we have several cases:
- If h < d, then $c(M) < c_{mon}(M)$.
- If $h \ge d$ and $c(M) < c_{eq}(M)$, then $c(M) < c_{mon}(M)$
- If $h \ge d$ and $c(M) = c_{eq}(M)$, then $c(M) = c_{mon}(M)$

When is $c_{mon}(M) > c(M)$?

Given that the following two conditions hold, $c_{mon}(s) > c(s)$ for any element s in any monoid M.

- $ightharpoonup C_{eq}(s) > C_{adj}(s)$
- If s has two factorizations z_1 and z_2 of length I, there exists a factorization z_3 of length $q \neq I$ such that $d(z_1, z_3) < c_{eq}(s)$ and $d(z_2, z_3) < c_{eq}(s)$.

A Case Where $c_{mon}(M) > c(M)$

Let $M = \langle na, na + n, 2na + nx + 1 \rangle$ with $x \geq 2$. Then


- $c_{eq}(M) = na + nx + 1$
- $c_{adj}(M) < na + nx + 1$
- $c_{mon}(M) = na + nx + 1$
- $ightharpoonup c_{mon}(M) > c(M)$

Example

Let $M = \langle 8, 10, 21 \rangle$, so n = 2, a = 4, and x = 2. Then $c_{eq}(M) = c_{mon}(M) = 13$, and c(M) = 5. Then $c_{mon}(M) > c(M)$.

$c_{mon}(M) - c(M)$ can be Arbitrarily Large

Monotone and Regular Catenary Degrees of



- $c_{mon}(M) = a^2 2a 1$ c(M) = 2a - 3
- c(W) 2a = 3 $c_{mon}(M) - c(M) = a^2 - 4a - 4$
- For large *a*, this difference can grow arbitrarily large.

Conclusions

- In some monoids, namely those generated by arithmetic sequences, $c_{mon}(M) = c(M)$.
- In generalized arithmetic monoids, we can have either $c_{mon}(M) > c(M)$ or $c_{mon}(M) = c(M)$.
- In general, we expect that $c_{mon}(M) > c(M)$. In fact, the difference between the two can grow arbitrarily large.

Acknowledgments

- We would like to thank...
- Dr. Brian Wissman and Dr. Bob Pelayo for overseeing our research
- Dr. Chris O'Neill and Dr. Scott Chapman for providing our research question and continuing support throughout this process
- Dr. Brian Loft, Dr. Rebecca Garcia, Dr. Luis Garcia, and PURE Math for hosting us
- ► This work was conducted during the 2014 PURE Math REU which is supported by the National Science Foundation (NSF Proposals DMS-1045147 and DMS-1045082).

References

- Bowles, Craig, Scott T. Chapman, Nathan Kaplan, and Daniel Reiser. "On Delta Sets Of Numerical Monoids."
 Journal of Algebra and Its Applications J. Algebra Appl. 05 (2006), 695-718.
- S. T. Chapman, M. Corrales, A. Miller, C. Miller, and D. Patel. *The Catenary Degrees of Elements in Numerical Monoids Generated by Arithmetic Sequences*. Journal of the Australian Mathematical Society, 2014.
- S. T. Chapman, P. A. García-Sánchez, and D. Llena. "The Catenary and Tame Degree of Numerical Monoids."
 Forum Mathematicum 21 (2009).
- A. Geroldinger and P. Yuan, The Monotone Catenary Degree of Krull Monoids, Results in Mathematics 63(2013), 999-1031.
- M. Omidali. "The Catenary and Tame Degree of Numerical Monoids Generated by Generalized Arithmetic Sequences." Forum Mathematicum. 24 (2012), 627-640.
- A. Philip, A characterization of arithmetical invariants by the monoid of relations II: The monotone catenary degree and applications to semigroup rings, *Semigroup Forum* 81 (2010), 424-434.
 J. C. Rosales and P. A. García-Sánchez, *Finitely generated commutative monoids*, Nova Science Publishers,
- New York, 1999.