

Monotone Catenary Degree in Numerical Monoids

D. Gonzalez, C. Wright, J. Zomback

August 4, 2015

Outline

Outline

- Definitions

Outline

- Definitions
- Arithmetic Monoids

Outline

- Definitions
- Arithmetic Monoids
- Generalized Arithmetic Monoids


Outline

- Definitions
- Arithmetic Monoids
- Generalized Arithmetic Monoids
- Difference between Monotone and Regular Catenary Degree


Outline

- Definitions
- Arithmetic Monoids
- Generalized Arithmetic Monoids
- Difference between Monotone and Regular Catenary Degree

Construction of a Catenary Graph




Construction of a Catenary Graph




- Remove edges with the largest distances

Construction of a Catenary Graph




- Remove edges with the largest distances
- Continue until disconnected

Construction of a Catenary Graph




- Remove edges with the largest distances
- Continue until disconnected
- Final distance removed is catenary degree

Construction of a Catenary Graph




- Remove edges with the largest distances
- Continue until disconnected
- Final distance removed is catenary degree

Construction of a Catenary Graph




- Remove edges with the largest distances
- Continue until disconnected
- Final distance removed is catenary degree

Construction of a Catenary Graph




- Remove edges with the largest distances
- Continue until disconnected
- Final distance removed is catenary degree

Construction of a Catenary Graph




- Remove edges with the largest distances
- Continue until disconnected
- Final distance removed is catenary degree

Construction of a Catenary Graph




- Remove edges with the largest distances
- Continue until disconnected
- Final distance removed is catenary degree

Construction of a Catenary Graph




- Remove edges with the largest distances
- Continue until disconnected
- Final distance removed is catenary degree

Construction of a Catenary Graph




- Remove edges with the largest distances
- Continue until disconnected
- Final distance removed is catenary degree

Construction of a Catenary Graph




- Remove edges with the largest distances
- Continue until disconnected
- Final distance removed is catenary degree

Construction of a Catenary Graph



- Remove edges with the largest distances
- Continue until disconnected
- Final distance removed is catenary degree

Construction of a Catenary Graph



- Remove edges with the largest distances
- Continue until disconnected
- Final distance removed is catenary degree

$$c(m) = 2$$

Equivalent Catenary Degree

How does it differ?

Equivalent Catenary Degree

How does it differ?

- Only factorizations of the same length

Equivalent Catenary Degree

How does it differ?

- Only factorizations of the same length
- Minimum N -chain within a given length

Equivalent Catenary Degree


How does it differ?

- Only factorizations of the same length
- Minimum N -chain within a given length
- Maximum over all such minima

Equivalent Catenary Degree

How does it differ?


- Only factorizations of the same length
- Minimum N -chain within a given length
- Maximum over all such minima



Equivalent Catenary Degree

How does it differ?


- Only factorizations of the same length
- Minimum N -chain within a given length
- Maximum over all such minima



Equivalent Catenary Degree

How does it differ?


- Only factorizations of the same length
- Minimum N -chain within a given length
- Maximum over all such minima



Equivalent Catenary Degree

How does it differ?


- Only factorizations of the same length
- Minimum N -chain within a given length
- Maximum over all such minima



Equivalent Catenary Degree

How does it differ?

- Only factorizations of the same length
- Minimum N -chain within a given length
- Maximum over all such minima



$$c_{eq}(m) = \max\{1, 2\} = 2$$

Adjacent Catenary Degree

How does it differ?

Adjacent Catenary Degree

How does it differ?

- Only factorizations of adjacent lengths

Adjacent Catenary Degree

How does it differ?

- Only factorizations of adjacent lengths
- Minimum distance between adjacent lengths

Adjacent Catenary Degree


How does it differ?

- Only factorizations of adjacent lengths
- Minimum distance between adjacent lengths
- Maximum over all such minima

Adjacent Catenary Degree

How does it differ?


- Only factorizations of adjacent lengths
- Minimum distance between adjacent lengths
- Maximum over all such minima



Adjacent Catenary Degree

How does it differ?


- Only factorizations of adjacent lengths
- Minimum distance between adjacent lengths
- Maximum over all such minima



Adjacent Catenary Degree

How does it differ?


- Only factorizations of adjacent lengths
- Minimum distance between adjacent lengths
- Maximum over all such minima



Adjacent Catenary Degree

How does it differ?


- Only factorizations of adjacent lengths
- Minimum distance between adjacent lengths
- Maximum over all such minima



Adjacent Catenary Degree

How does it differ?


- Only factorizations of adjacent lengths
- Minimum distance between adjacent lengths
- Maximum over all such minima



Adjacent Catenary Degree

How does it differ?

- Only factorizations of adjacent lengths
- Minimum distance between adjacent lengths
- Maximum over all such minima



$$c_{\text{adj}}(m) = \max\{3, 2, 2\} = 3$$

Monotone Catenary Degree

How does it differ?

Monotone Catenary Degree

How does it differ?

- Most similar to Catenary Degree

Monotone Catenary Degree


How does it differ?

- Most similar to Catenary Degree
- Only monotone chains between elements

Monotone Catenary Degree

How does it differ?


- Most similar to Catenary Degree
- Only monotone chains between elements



Monotone Catenary Degree

How does it differ?


- Most similar to Catenary Degree
- Only monotone chains between elements



Monotone Catenary Degree

How does it differ?


- Most similar to Catenary Degree
- Only monotone chains between elements



Monotone Catenary Degree

How does it differ?


- Most similar to Catenary Degree
- Only monotone chains between elements



Monotone Catenary Degree

How does it differ?


- Most similar to Catenary Degree
- Only monotone chains between elements



Monotone Catenary Degree

How does it differ?


- Most similar to Catenary Degree
- Only monotone chains between elements



Monotone Catenary Degree

How does it differ?


- Most similar to Catenary Degree
- Only monotone chains between elements



Monotone Catenary Degree

How does it differ?


- Most similar to Catenary Degree
- Only monotone chains between elements



Monotone Catenary Degree

How does it differ?

- Most similar to Catenary Degree
- Only monotone chains between elements



$$c_{mon}(m) = 3$$

Motivation

What is known about the monotone catenary degree?

Motivation

What is known about the monotone catenary degree?

- For elements of a numerical monoid M , $c(m) \leq c_{\text{mon}}(m)$.

Motivation

What is known about the monotone catenary degree?

- For elements of a numerical monoid M , $c(m) \leq c_{\text{mon}}(m)$.
- $c_{\text{mon}}(M) = \max\{c_{\text{eq}}(M), c_{\text{adj}}(M)\}$

Motivation

What is known about the monotone catenary degree?

- For elements of a numerical monoid M , $c(m) \leq c_{\text{mon}}(m)$.
- $c_{\text{mon}}(M) = \max\{c_{\text{eq}}(M), c_{\text{adj}}(M)\}$
- Not much else

Motivation

What is known about the monotone catenary degree?

- For elements of a numerical monoid M , $c(m) \leq c_{\text{mon}}(m)$.
- $c_{\text{mon}}(M) = \max\{c_{\text{eq}}(M), c_{\text{adj}}(M)\}$
- Not much else

We seek to investigate the inequality between monotone and regular catenary degree.

Arithmetic Monoids

Consider a monoid M generated by a finite arithmetic sequence $a, a + d, \dots, a + kd$, so $M = \langle a, a + d, \dots, a + kd \rangle$. We will refer to a monoid of this form as an *arithmetic monoid*.

Arithmetic Monoids

Consider a monoid M generated by a finite arithmetic sequence $a, a + d, \dots, a + kd$, so $M = \langle a, a + d, \dots, a + kd \rangle$. We will refer to a monoid of this form as an *arithmetic monoid*.

Examples:

Arithmetic Monoids

Consider a monoid M generated by a finite arithmetic sequence $a, a + d, \dots, a + kd$, so $M = \langle a, a + d, \dots, a + kd \rangle$. We will refer to a monoid of this form as an *arithmetic monoid*.

Examples:

- $\langle 5, 7, 9 \rangle$

Arithmetic Monoids

Consider a monoid M generated by a finite arithmetic sequence $a, a + d, \dots, a + kd$, so $M = \langle a, a + d, \dots, a + kd \rangle$. We will refer to a monoid of this form as an *arithmetic monoid*.

Examples:


- $\langle 5, 7, 9 \rangle$
- $\langle 11, 15, 19, 23 \rangle$

Monotone vs. Regular Catenary Graph

Factorizations of 96 in $\langle 11, 15, 19, 23, 27 \rangle$


Monotone vs. Regular Catenary Graph

Factorizations of 96 in $\langle 11, 15, 19, 23, 27 \rangle$




Monotone vs. Regular Catenary Graph

Factorizations of 96 in $\langle 11, 15, 19, 23, 27 \rangle$




Monotone vs. Regular Catenary Graph

Factorizations of 96 in $\langle 11, 15, 19, 23, 27 \rangle$




Monotone vs. Regular Catenary Graph

Factorizations of 96 in $\langle 11, 15, 19, 23, 27 \rangle$




Monotone vs. Regular Catenary Graph

Factorizations of 96 in $\langle 11, 15, 19, 23, 27 \rangle$




Monotone vs. Regular Catenary Graph

Factorizations of 96 in $\langle 11, 15, 19, 23, 27 \rangle$



Monotone vs. Regular Catenary Graph

Factorizations of 96 in $\langle 11, 15, 19, 23, 27 \rangle$



$$c(96) = 7$$

Monotone vs. Regular Catenary Graph

Factorizations of 96 in $\langle 11, 15, 19, 23, 27 \rangle$

$$\begin{array}{lcl}
 l = 8 : & (6, 2, 0, 0, 0) & \xrightarrow{2} (7, 0, 1, 0, 0) \\
 & \quad \quad \quad \downarrow 7 & \quad \quad \quad \downarrow 7 \\
 l = 4 : & (0, 0, 0, 3, 1) & \xrightarrow{3} (0, 1, 0, 0, 3) \xrightarrow{2} (0, 0, 0, 1, 2)
 \end{array}$$

Monotone vs. Regular Catenary Graph

Factorizations of 96 in $\langle 11, 15, 19, 23, 27 \rangle$

$$\begin{array}{lcl}
l = 8 : & (6, 2, 0, 0, 0) & \xrightarrow{2} (7, 0, 1, 0, 0) \\
& & \begin{array}{c} | \quad \quad | \\ 7 \quad \quad 7 \end{array} \\
l = 4 : & (0, 0, 0, 3, 1) & \xrightarrow{3} (0, 1, 0, 0, 3) \xrightarrow{2} (0, 0, 0, 1, 2) \\
& & c_{mon}(96) = c(96) = 7.
\end{array}$$

Main Result for Arithmetic Sequences

Theorem

Given an arithmetic monoid $M = \langle a, a + d, \dots, a + kd \rangle$, for all elements $m \in M$,

$$c_{\text{mon}}(m) = c(m).$$

As a result of this, $c_{\text{mon}}(M) = c(M)$.

Generalized Arithmetic Sequences

Consider $M = \langle a, ah + d, \dots, ah + kd \rangle$. We will refer to a monoid of this form as a *generalized arithmetic monoid*. We will discuss generalized arithmetic monoids in embedding dimension three:

$$M = \langle a, ah + d, ah + 2d \rangle.$$

Generalized Arithmetic Sequences

Consider $M = \langle a, ah + d, \dots, ah + kd \rangle$. We will refer to a monoid of this form as a *generalized arithmetic monoid*. We will discuss generalized arithmetic monoids in embedding dimension three:

$$M = \langle a, ah + d, ah + 2d \rangle.$$

Examples:

Generalized Arithmetic Sequences

Consider $M = \langle a, ah + d, \dots, ah + kd \rangle$. We will refer to a monoid of this form as a *generalized arithmetic monoid*. We will discuss generalized arithmetic monoids in embedding dimension three:

$$M = \langle a, ah + d, ah + 2d \rangle.$$

Examples:

- $\langle 3, 11, 16 \rangle$

Generalized Arithmetic Sequences

Consider $M = \langle a, ah + d, \dots, ah + kd \rangle$. We will refer to a monoid of this form as a *generalized arithmetic monoid*. We will discuss generalized arithmetic monoids in embedding dimension three:

$$M = \langle a, ah + d, ah + 2d \rangle.$$

Examples:

- $\langle 3, 11, 16 \rangle$

$$a = 3, h = 2, d = 5$$

Generalized Arithmetic Sequences

Consider $M = \langle a, ah + d, \dots, ah + kd \rangle$. We will refer to a monoid of this form as a *generalized arithmetic monoid*. We will discuss generalized arithmetic monoids in embedding dimension three:

$$M = \langle a, ah + d, ah + 2d \rangle.$$

Examples:

- $\langle 3, 11, 16 \rangle$
 $a = 3, h = 2, d = 5$
- $\langle 5, 38, 41 \rangle$

Generalized Arithmetic Sequences


Consider $M = \langle a, ah + d, \dots, ah + kd \rangle$. We will refer to a monoid of this form as a *generalized arithmetic monoid*. We will discuss generalized arithmetic monoids in embedding dimension three:

$$M = \langle a, ah + d, ah + 2d \rangle.$$



Examples:

- $\langle 3, 11, 16 \rangle$
 $a = 3, h = 2, d = 5$
- $\langle 5, 38, 41 \rangle$
 $a = 5, h = 7, d = 3$

Generalized Arithmetic Sequences




Generalized Arithmetic Sequences



$$\bullet \quad c(m) = c_{mon}(m)$$



Generalized Arithmetic Sequences





- $c(m) = c_{mon}(m)$

- $c(M) = c_{mon}(M)$

Generalized Arithmetic Sequences





Generalized Arithmetic Sequences





• $c(m) \neq c_{mon}(m)$

Generalized Arithmetic Sequences





- $c(m) \neq c_{mon}(m)$
- $c(M) = c_{mon}(M)$

Generalized Arithmetic Sequences





Generalized Arithmetic Sequences



• $c(m) \neq c_{mon}(m)$

Generalized Arithmetic Sequences



- $c(m) \neq c_{mon}(m)$
- $c(M) < c_{mon}(M)$

Generalized Arithmetic Sequences

Conjecture

If $\gcd(h - 1, d) > 1$, then $c_{\text{mon}}(M) = c(M)$.

Generalized Arithmetic Sequences

Conjecture

If $\gcd(h - 1, d) > 1$, then $c_{\text{mon}}(M) = c(M)$.

Conjecture

If $\gcd(h - 1, d) = 1$, then there are several cases.

Generalized Arithmetic Sequences

Conjecture

If $\gcd(h - 1, d) > 1$, then $c_{\text{mon}}(M) = c(M)$.

Conjecture

If $\gcd(h - 1, d) = 1$, then there are several cases.

Case 1: $h < d$

$$c(M) < c_{\text{mon}}(M)$$

Generalized Arithmetic Sequences

Conjecture

If $\gcd(h - 1, d) > 1$, then $c_{mon}(M) = c(M)$.

Conjecture

If $\gcd(h - 1, d) = 1$, then there are several cases.

Case 1: $h < d$

$$c(M) < c_{mon}(M)$$

Case 2: $h \geq d$ and $c(M) < c_{eq}(M)$

$$c(M) < c_{mon}(M)$$

Generalized Arithmetic Sequences

Conjecture

If $\gcd(h - 1, d) > 1$, then $c_{mon}(M) = c(M)$.

Conjecture

If $\gcd(h - 1, d) = 1$, then there are several cases.

Case 1: $h < d$ $c(M) < c_{mon}(M)$

Case 2: $h \geq d$ and $c(M) < c_{eq}(M)$ $c(M) < c_{mon}(M)$

Case 3: $h \geq d$ and $c(M) = c_{eq}(M)$ $c(M) = c_{mon}(M)$

Generalized Arithmetic Sequences


•

•

- Recall that $c_{mon}(m) = \max\{c_{eq}(m), c_{adj}(m)\}$.


•

Generalized Arithmetic Sequences




- Recall that $c_{mon}(m) = \max\{c_{eq}(m), c_{adj}(m)\}$.

Generalized Arithmetic Sequences




- Recall that $c_{\text{mon}}(m) = \max\{c_{\text{eq}}(m), c_{\text{adj}}(m)\}$.
- $c_{\text{eq}}(M) = \frac{ah+2d-a}{\gcd(h-1,d)}$

Generalized Arithmetic Sequences



- Recall that $c_{mon}(m) = \max\{c_{eq}(m), c_{adj}(m)\}$.
- $c_{eq}(M) = \frac{ah+2d-a}{\gcd(h-1,d)}$
- if $\gcd(h-1, d) = 1$, then $c_{eq}(m) \geq c_{adj}(m)$

Generalized Arithmetic Sequences



- Recall that $c_{mon}(m) = \max\{c_{eq}(m), c_{adj}(m)\}$.
- $c_{eq}(M) = \frac{ah+2d-a}{\gcd(h-1,d)}$
- if $\gcd(h-1, d) = 1$, then $c_{eq}(m) \geq c_{adj}(m)$
- if $\gcd(h-1, d) > 1$, then $c_{eq}(m) < c_{adj}(m)$

Recall that $c_{\text{mon}}(M) \geq c(M)$.

Recall that $c_{\text{mon}}(M) \geq c(M)$.

When is $c_{\text{mon}}(M) > c(M)$?

Recall that $c_{mon}(M) \geq c(M)$.

When is $c_{mon}(M) > c(M)$?

If we have

- $c_{eq}(M) > c_{adj}(M)$

Recall that $c_{mon}(M) \geq c(M)$.

When is $c_{mon}(M) > c(M)$?

If we have

- $c_{eq}(M) > c_{adj}(M)$
- Whenever $|z_1| = |z_2| = I$ we can find $|z_3| < I$ such that $d(z_1, z_3) < c_{eq}(M)$ and $d(z_2, z_3) < c_{eq}(M)$

Recall that $c_{mon}(M) \geq c(M)$.


When is $c_{mon}(M) > c(M)$?

If we have


- $c_{eq}(M) > c_{adj}(M)$
- Whenever $|z_1| = |z_2| = I$ we can find $|z_3| < I$ such that $d(z_1, z_3) < c_{eq}(M)$ and $d(z_2, z_3) < c_{eq}(M)$

Then $c_{mon}(M) > c(M)$.

Is $c_{\text{mon}}(M) > c(M)$?




Is $c_{mon}(M) > c(M)$?




- $c_{mon}(M) = c_{eq}(M)$.
- Under our restrictions,
 $c_{eq}(M) > c_{adj}(M)$

Is $c_{mon}(M) > c(M)$?




- $c_{mon}(M) = c_{eq}(M)$
- Under our restrictions,
 $c_{eq}(M) > c_{adj}(M)$
- We can move from two factorizations of length l to one of a lower length with $d < c_{eq}(M)$

Is $c_{mon}(M) > c(M)$?




- $c_{mon}(M) = c_{eq}(M)$
- Under our restrictions,
 $c_{eq}(M) > c_{adj}(M)$
- We can move from two factorizations of length l to one of a lower length with $d < c_{eq}(M)$
- $c_{mon}(M) > c(M)$

Equivalent Catenary Degree in $\langle 8, 9, 19 \rangle$



Equivalent Catenary Degree in $\langle 8, 9, 19 \rangle$



Equivalent Catenary Degree in $\langle n_1, n_2, n_3 \rangle$

Theorem

Let M be a minimally generated numerical monoid $\langle n_1, n_2, n_3 \rangle$.

Then $c_{\text{eq}}(M) = \frac{n_3 - n_1}{\gcd(n_2 - n_1, n_3 - n_1)}$

Equivalent Catenary Degree in $\langle n_1, n_2, n_3 \rangle$

Theorem

Let M be a minimally generated numerical monoid $\langle n_1, n_2, n_3 \rangle$.

Then $c_{\text{eq}}(M) = \frac{n_3 - n_1}{\gcd(n_2 - n_1, n_3 - n_1)}$

- If $m = n_2 \left(\frac{n_3 - n_1}{\gcd(n_2 - n_1, n_3 - n_1)} \right) + x$ for some $x \in M$, then $c_{\text{eq}}(m) = c_{\text{eq}}(M)$.
- If $m = n_2 \left(\frac{n_3 - n_1}{\gcd(n_2 - n_1, n_3 - n_1)} \right) + x$ for some $x \notin M$, then $c_{\text{eq}}(m) = 0$.

Equivalent Catenary Degree in $\langle n_1, n_2, n_3 \rangle$

Theorem

For all $m \in M$ such that $m > n_2 \left(\frac{n_3 - n_1}{\gcd(n_2 - n_1, n_3 - n_1)} \right) + \mathcal{F}(M)$, $c_{\text{eq}}(m) = c_{\text{eq}}(M)$. Furthermore, $c_{\text{eq}}(m)$ can take on only two values: 0 and $c_{\text{eq}}(M)$.

Moving Between Factorizations

$$M = \langle n_1, n_2, n_3 \rangle$$

To calculate c_{eq} we needed a way to move between factorizations of the same length.

Moving Between Factorizations

$$M = \langle n_1, n_2, n_3 \rangle$$

To calculate c_{eq} we needed a way to move between factorizations of the same length.

Let $z_1 = (b, c, d)$ and let $|z_1| = |z_2|$. Then

$$z_2 = \left(b + \frac{k(n_3 - n_1)}{n_3 - n_1} - k, c - \frac{k(n_3 - n_1)}{n_3 - n_1}, d + k \right)$$

Moving Between Factorizations

$$M = \langle n_1, n_2, n_3 \rangle$$

To calculate c_{eq} we needed a way to move between factorizations of the same length.

Let $z_1 = (b, c, d)$ and let $|z_1| = |z_2|$. Then

$$z_2 = \left(b + \frac{k(n_3 - n_1)}{n_3 - n_1} - k, c - \frac{k(n_3 - n_1)}{n_3 - n_1}, d + k \right)$$

What if we want to move between factorization lengths?

Moving Between Factorizations

$$M = \langle n_1, n_2, n_3 \rangle$$

To calculate c_{eq} we needed a way to move between factorizations of the same length.

Let $z_1 = (b, c, d)$ and let $|z_1| = |z_2|$. Then

$$z_2 = \left(b + \frac{k(n_3 - n_1)}{n_3 - n_1} - k, c - \frac{k(n_3 - n_1)}{n_3 - n_1}, d + k \right)$$

What if we want to move between factorization lengths?

Theorem

Consider a factorization $z = (b, c, d)$ of an element $s \in M$ with $|z| = x$. We can write any factorization z_i of s where $|z_i| = x - l$ as $z_i = \left(b + \frac{-l n_1 + k(n_3 - n_1)}{n_3 - n_1} - k - l, c + \frac{l n_1 - k(n_3 - n_1)}{n_3 - n_1}, d + k \right)$.

When is $c_{\text{mon}}(M) > c(M)$?

For certain families of monoids, we can guarantee that our two conditions for $c_{\text{mon}}(M) > c(M)$ hold.

When is $c_{\text{mon}}(M) > c(M)$?

For certain families of monoids, we can guarantee that our two conditions for $c_{\text{mon}}(M) > c(M)$ hold.

$$M = \langle na, na + n, 2na + nx + 1 \rangle \text{ with } x \geq 2$$

When is $c_{mon}(M) > c(M)$?

For certain families of monoids, we can guarantee that our two conditions for $c_{mon}(M) > c(M)$ hold.

$$M = \langle na, na + n, 2na + nx + 1 \rangle \text{ with } x \geq 2$$

Recall that $c_{mon}(M) = \max\{c_{eq}(M), c_{adj}(M)\}$.

When is $c_{mon}(M) > c(M)$?

For certain families of monoids, we can guarantee that our two conditions for $c_{mon}(M) > c(M)$ hold.

$$M = \langle na, na + n, 2na + nx + 1 \rangle \text{ with } x \geq 2$$

Recall that $c_{mon}(M) = \max\{c_{eq}(M), c_{adj}(M)\}$.

- $c_{eq}(M) = na + nx + 1$

When is $c_{\text{mon}}(M) > c(M)$?

For certain families of monoids, we can guarantee that our two conditions for $c_{\text{mon}}(M) > c(M)$ hold.

$$M = \langle na, na + n, 2na + nx + 1 \rangle \text{ with } x \geq 2$$

Recall that $c_{\text{mon}}(M) = \max\{c_{\text{eq}}(M), c_{\text{adj}}(M)\}$.


- $c_{\text{eq}}(M) = na + nx + 1$
- $c_{\text{adj}}(M) < na + nx + 1$

$$\langle na, na + n, 2na + nx + 1 \rangle$$

$$M = \langle na, na + n, 2na + nx + 1 \rangle \text{ with } x \geq 2$$


$$\langle na, na + n, 2na + nx + 1 \rangle$$

$$M = \langle na, na + n, 2na + nx + 1 \rangle \text{ with } x \geq 2$$



$$\langle na, na + n, 2na + nx + 1 \rangle$$

$$M = \langle na, na + n, 2na + nx + 1 \rangle \text{ with } x \geq 2$$



Theorem

Let $M = \langle na, na + n, 2na + nx + 1 \rangle$ for $n \in \mathbb{N}$ and $x \geq 2$. Then $c_{\text{mon}}(M) > c(M)$.

$$\langle a, a + 1, \mathcal{F}\langle a, a + 1 \rangle \rangle$$

We have shown that in many cases, $c_{\text{mon}}(M) > c(M)$.

$$\langle a, a + 1, \mathcal{F}\langle a, a + 1 \rangle \rangle$$

We have shown that in many cases, $c_{\text{mon}}(M) > c(M)$.

How big can the difference be between $c_{\text{mon}}(M)$ and $c(M)$?

$$\langle a, a + 1, \mathcal{F}\langle a, a + 1 \rangle \rangle$$

We have shown that in many cases, $c_{\text{mon}}(M) > c(M)$.

How big can the difference be between $c_{\text{mon}}(M)$ and $c(M)$?

$$\langle a, a + 1, \mathcal{F}\langle a, a + 1 \rangle \rangle$$

- $c_{\text{mon}}(M) = a^2 - 2a - 1$
- $c(M) = 2a - 3$

$$\langle a, a + 1, \mathcal{F}\langle a, a + 1 \rangle \rangle$$


We have shown that in many cases, $c_{\text{mon}}(M) > c(M)$.

How big can the difference be between $c_{\text{mon}}(M)$ and $c(M)$?

$$\langle a, a + 1, \mathcal{F}\langle a, a + 1 \rangle \rangle$$

- $c_{\text{mon}}(M) = a^2 - 2a - 1$
- $c(M) = 2a - 3$
- $c_{\text{mon}}(M) - c(M) = a^2 - 4a - 4$

$$\langle a, a + 1, \mathcal{F}\langle a, a + 1 \rangle \rangle$$



$$c_{\text{mon}}(M) - c(M)$$

Theorem

The difference between the monotone and regular catenary degrees of a monoid can be arbitrarily large.

Takeaways

- In some monoids, namely those generated by arithmetic sequences, $c_{\text{mon}}(M) = c(M)$.

Takeaways

- In some monoids, namely those generated by arithmetic sequences, $c_{\text{mon}}(M) = c(M)$.
- In generalized arithmetic monoids, we can have either $c_{\text{mon}}(M) > c(M)$ or $c_{\text{mon}}(M) = c(M)$.

Takeaways

- In some monoids, namely those generated by arithmetic sequences, $c_{\text{mon}}(M) = c(M)$.
- In generalized arithmetic monoids, we can have either $c_{\text{mon}}(M) > c(M)$ or $c_{\text{mon}}(M) = c(M)$.
- In general, we expect that $c_{\text{mon}}(M) > c(M)$. In fact, the difference between the two can grow arbitrarily large.

Thank You

We would like to thank:

- The NSF and NSA
- Sam Houston State University, University of Hawai'i at Hilo, and PURE Math
- Dr. Brian Loft, Dr. Rebecca García, and Dr. Luis García
- Dr. Brian Wissman, Dr. Bob Pelayo
- Dr. Scott Chapman and Dr. Chris O'Neill
- Felix Gotti and Marly Cormar

References

- Bowles, Craig, Scott T. Chapman, Nathan Kaplan, and Daniel Reiser. "On Delta Sets Of Numerical Monoids." *Journal of Algebra and Its Applications J. Algebra Appl.* **05** (2006), 695-718.
- S. T. Chapman, M. Corrales, A. Miller, C. Miller, and D. Patel. *The Catenary Degrees of Elements in Numerical Monoids Generated by Arithmetic Sequences*. Journal of the Australian Mathematical Society, 2014.
- S. T. Chapman, P. A. Garca-Snchez, and D. Llena. "The Catenary and Tame Degree of Numerical Monoids." *Forum Mathematicum* **21** (2009).
- A. Geroldinger and P. Yuan, The Monotone Catenary Degree of Krull Monoids, *Results in Mathematics* **63**(2013), 999-1031.

References

- M. Omidali. "The Catenary and Tame Degree of Numerical Monoids Generated by Generalized Arithmetic Sequences." *Forum Mathematicum*. **24** (2012), 627-640.
- A. Philip, A characterization of arithmetical invariants by the monoid of relations II: The monotone catenary degree and applications to semigroup rings, *Semigroup Forum* **81** (2010), 424-434.
- J. C. Rosales and P. A. García-Sánchez, *Finitely generated commutative monoids*, Nova Science Publishers, NewYork, 1999.