Roccella phycopsis the Correct Name for R. fucoides (Lichenized Fungi)
Author(s): Anders Tehler
Source: *Taxon*, Vol. 52, No. 3 (Aug., 2003), p. 621
Published by: International Association for Plant Taxonomy (IAPT)
Stable URL: http://www.jstor.org/stable/3647467
Accessed: 13-01-2016 15:00 UTC

REFERENCES
Linked references are available on JSTOR for this article:
http://www.jstor.org/stable/3647467?seq=1&cid=pdf-reference#references_tab_contents

You may need to log in to JSTOR to access the linked references.

Your use of the JSTOR archive indicates your acceptance of the Terms & Conditions of Use, available at http://www.jstor.org/page/info/about/policies/terms.jsp

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact support@jstor.org.

International Association for Plant Taxonomy (IAPT) is collaborating with JSTOR to digitize, preserve and extend access to *Taxon*.
http://www.jstor.org

This content downloaded from 128.123.44.23 on Wed, 13 Jan 2016 15:00:10 UTC
All use subject to JSTOR Terms and Conditions

Roccella phycopsis the correct name for *R. fucoides* (lichenized fungi)

Anders Tehler

Sektionen för kryptogambotanik, Naturhistoriska riksmuseet, Box 50007, S-104 05 Stockholm, Sweden. anders.tehler@nrm.se

In a recent publication on the typification of two *Roccella* names (Tehler, 2002), the names *Lichen fucoides* Neck. and *L. fucoides* With. were overlooked.

Unfortunately, that oversight had a profound impact on the application of the correct name for the species concerned. The name *Roccella fucoides* cannot be used since *Lichen fucoides* Dicks. (Dickson, 1790), regarded as its basionym, was illegitimate when published, being a later homonym of *Lichen fucoides* Neck. (Necker, 1771), an *Umbilicaria* species. Vainio did not publish *R. fucoides* as a legitimate name until 1901, whereas in 1804 Acharius had published the legitimate *Parmelia phycopsis*, based on *L. fucoides* Dicks., which makes *phycopsis* the next available epithet. *Roccella phycopsis* (Ach.) Ach. is thus the correct name for the species. *Lichen fucoides* With., illegitimate for the same reason as *L. fucoides* Dicks., applies to the species *Cornicularia normoerica* (Gunn.) Du Rietz (Laundon, 1984).

Roccella phycopsis (Ach.) Ach., Lich. univ.: 440. 1810. (Cf Art. 33.2) ≡ *Lichen fucoides* Dicks., Pl. Crypt. Brit. 2: 22. 1790, nom illeg., non *Lichen fucoides* Neck., Meth. Musc.: 76. 1771, nec *Lichen fucoides* With., Bot. Arrang. 1: 726. 1776 ≡ *Parmelia phycopsis* Ach., Arch. Syst. Naturgesch. 1(1): 110. 1804 ≡ *Roccella fucoides* Vain., in Hiern & al., Cat. Afr. Pl. 2: 433. 1901. – Type: "Habitat in rupibus et lignis, locis maritimis, prope Gosport". (not found). – Neotype: "*Lichen fucoides*", *J. Dickson s.n.* (E!); isoneotype (E!).

= *Roccella pygmaea* Durieu & Mont., Flore d'Alger 226. 1847. ≡ *Roccella pusilla* de Notaris, Giorn. Bot. Ital. 2: 222. 1846. ≡ *Roccella phycopsis* f. *pygmaea* (Durieu et Mont.) Nyl. ex Zahlbr., Catal. Lich. Univ. 2: 519. 1923. ≡ *Roccella fucoides* f. *pygmaea* (Durieu & Mont.) Zahlbr., Catal. Lich. Univ. 2: 519. 1923. – Lectotype: "Algeria. ad ficos et *Micromeriam inodoram*", *M. Durieu s.n.* (PC!); isolectotypes: (PC!).

= *Roccella phycopsis* f. *tenuior* Leight., Lich. Flor. Brit. ed 3:74. 1879. ≡ *Roccella fucoides* f. *tenuior* (Leight.) Zahlbr., Catal. Lich. Univ. 2: 519. 1923. – Lectotype: [Great Britain], Channels Islands, Jersey, La Moye, s.d., *C. Larbalestier s.n.* (BM!).

= *Roccella phycopsis* var. *ceciliametella* Rabenh., Flecht. Europ. 36: no 958. 1879, nom. nud. ≡ *Roccella fucoides* var. *ceciliametella* (Rabenh.) Zahlbr., Catal. Lich. Univ. 2: 519. 1923, nom. nud. – Coll. orig.: Italy, Sur le tombeau de Cecilia Metella, 1874, Vito Beltrani Pisani in Rabenhorst, Lich. Eur. exs. 958 (PC, S).

= *Roccella fucoides* var. *corticola* Sambo, Nouvo Giorn. Bot. Ital. 34: 832. 1927. – Type: [Greece, Rhodes] "Sugli alberi a Cattavia", 1926, *Senni s.n.* (not found).

= *Roccella fucoides* var. *farinosa* Delise ex Sambo, Nouvo Giorn. Bot. Ital. 34: 832. 1927. – Lectotype: St. Malo, *collector unknown* (PC-Delise).

ACKNOWLEDGEMENTS

I want to thank Rolf Santesson for drawing my attention to the problem.

LITERATURE CITED

Dickson, J. 1790. *Fasciculus Secundus Plantarum Cryptogamicarum Britanniae*. G. Nicol, London.

Laundon, J. R. 1984. The typification of Withering's neglected lichen. *Lichenologist* 16: 211–239.

Necker de, N. J. d. 1771. *Methodus Muscorum*. Academ. Scient. Elect., Mannheim.

Tehler, A. 2002. On the typification of two misunderstood *Roccella* names, *R. tinctoria* and *R. fucoides*. *Taxon* 51: 787–790.

This content downloaded from 128.123.44.23 on Wed, 13 Jan 2016 15:00:10 UTC
All use subject to JSTOR Terms and Conditions