# Estimating missing trait data in comparative biology using Bayesian Augmentation by Chained Equations: the BACE R package

Daniel W. A. Noble[1], Szymon M. Drobniak[2,3], Ayumi Mizuno[4], Patrice Pottier[5], Shinichi Nakagawa[3,4]

[1] Division of Ecology and Evolution, Research School of Biology, The Australian National University, Canberra, ACT, Australia

[2] Institute of Environmental Sciences, Faculty of Biology, Jagiellonian University, Kraków, Poland

[3] School of Biological, Earth and Environmental Sciences, University of New South Wales, Sydney, NSW, Australia

[4] Department of Biological Sciences, Faculty of Science, University of Alberta, Edmonton, Alberta, Canada

[5] Department of Biological and Environmental Sciences, University of Gothenburg, Gothenburg, Sweden

## Abstract

1. Missing trait data are pervasive in comparative datasets limiting the types of questions that can be tested and compromising the power of comparative analyses. Standard approaches, such as complete-case analyses and single imputation, can bias inference and underestimate uncertainty, particularly when missing data is non-random. While robust imputation methods exits, many existing software are restricted to imputing only continuous or count variables and lack the capacity to impute discrete / categorical traits (e.g., multinomial, ordered categorical), which are common trait types in comparative datasets. Such limitations have prevented the widespread use of imputation approaches in comparative biology.
2. We introduce *BACE*, an R package implementing Bayesian Augmentation by Chained Equations for phylogenetic comparative data. *BACE* enables imputation with phylogeny-informed covariance that can account for both phylogenetic and non-phylogenetic species effects, along with mixed data types, in a fully Bayesian pipeline. *BACE* provides substantial flexibility in the imputation models fit, can decouple imputation models from analysis models, and propagates both parameter and imputation uncertainty in subsequent downstream analyses.

3. We show using simulations that *BACE* recovers missing trait values with a high degree of accuracy. Across 18 different scenarios that vary in sample size (n = 100, 300, 1000 species), phylogenetic signal (Pagel's $\lambda = 0.3, 0.7, 1$) and trait correlation ($\rho = 0, 0.5$), we show *BACE* recovers model parameters with little bias and nominal coverage. *BACE* outperformed other comparable frequentist approaches. We illustrate *BACE* using two case studies. The first imputes amphibian heat tolerance (continuous) under standardised conditions across 5,203 species using 2,661 experimental estimates from 524 species; 89.9% of species lack experimental measurements. The second uses AVONET data for 10,563 bird species to impute migratory behaviour (ordered categorical) and hand-wing index (continuous), addressing 17 naturally missing migration records (0.16%) and testing imputation after withholding approximately 20% of observed values for each trait.
4.

# Introduction

Phylogenetic comparative methods (PCMs) provide a framework for understanding how traits evolve and covary across species (Felsenstein, 1985; Grafen, 1989; Martins & Hansen, 1997; Pagel, 1994). Expanding trait databases, increasingly comprehensive phylogenies, and greater access to environmental data are providing opportunities to investigate how evolutionary history and environmental conditions jointly shape variation and covariation among traits allowing a rich set of new evolutionary hypotheses to be tested (Blomberg & Todorov, 2025; Halliwell et al., 2025). However, missing trait data limit the questions that can be addressed, and excluding species with incomplete records can reduce sample sizes, statistical power and precision (Blomberg & Todorov, 2025; Little et al., 2024; Nakagawa & Freckleton, 2008). When the availability of trait data depends on species' characteristics, this exclusion can also bias estimates of trait relationships with significant consequences for inferences (Garamszegi & Møller, 2011; González-Suárez et al., 2012; Nakagawa & Freckleton, 2008). For example, incomplete trait coverage has been shown to alter inferred relationships between mammalian traits and extinction risk (González-Suárez et al., 2012). As such, there is a need for versatile and rigorous statistical approaches to deal with missing data in comparative biology.

Methods for handling missing data have a long history in statistics, and two broad approaches are commonly used (Blomberg & Todorov, 2025; Little & Rubin, 2002; Noble & Nakagawa, 2021; Rubin, 1987; Schafer & Graham, 2002). First, missing data can be accommodated during maximum-likelihood or Bayesian model fitting. For example, Bayesian data augmentation samples missing values alongside model parameters (Tanner & Wong, 1987). Alternatively, multiple-imputation can be used to create, $m$ completed datasets and each dataset analysed separately. Results are then pooled to account for within- and between-imputation uncertainty, using Rubin's rules for estimates and variances or combining posterior draws for Bayesian analyses (Noble & Nakagawa, 2021; Rubin, 1987; Zhou & Reiter, 2010). These approaches are not mutually exclusive. Data augmentation can also generate the completed datasets used in multiple imputation. Imputations can be generated using a joint model for all variables or separate conditional models linked through chained equations. Chained equations cycle through models, drawing missing values for each variable using the latest values of the others. This approach, also called fully conditional specification, allows different models for continuous and categorical traits (Buuren & Groothuis-Oudshoorn, 2011). Standard multiple-imputation analyses assume data are missing at random (MAR), conditional on the variables included in the imputation models. As such, including traits associated with both the missing

values and the probability of missingness helps make this assumption more plausible (Gelman & Hill, 2007, 25; Noble & Nakagawa, 2021; Schafer & Graham, 2002). In comparative biology, both phylogenetic prediction of missing trait values (Bruggeman et al., 2009; Goolsby et al., 2017) and multiple imputation (Blomberg & Todorov, 2025; Pottier et al., 2025) have been used to deal with missing trait data. However, because comparative traits differ in their distributions and relationships, imputation requires flexible models for individual traits for it to be widely used.

While a broad array of statistical methods and packages exist to deal with missing data, many are not suited to the datasets typical of comparative biology (Table 1). A few packages can use the phylogeny in their imputation models (e.g., *Rphylopars*, *phylomice*, *TDIP*), but most cannot (e.g., *mice*, *miceadds*, *jomo*, *pan*; Table 1). Moreover, the phylogeny-aware packages generally return a single completed dataset or single predicted values rather than multiple imputations, and most impute only continuous traits. A notable exception is *TDIP* (Gendre et al., 2024), which combines phylogenetic and machine-learning methods in an ensemble and can impute discrete traits. However, it too returns single completed datasets, and it cannot handle multi-level data structures, count data, or ordered categorical traits (Table 1). As such, there is a need for a statistically robust and flexible imputation approach that can handle multi-level data structures and a diversity of trait types, while ensuring imputation uncertainty is correctly accounted for in any downstream analyses and interpretations.

Here, we introduce a new Bayesian imputation software package for comparative biology, named *BACE* (**B**ayesian **A**ugmentation by **C**hained **E**quations), which uses the versatile Bayesian software *MCMCglmm* (Hadfield, 2010) to conduct multiple imputation through chained equations. *BACE* provides substantial flexibility in the imputation / analysis models fit, as well as a simple user interface, which makes it easy to apply in practice. *BACE* also provides options to decouple imputation and analysis models. Given its Bayesian approach, imputation and analysis models can be fit jointly and a final analysis model allows for pooling of posterior distributions to account for within and across imputation uncertainty. Additionally, samples from the posterior predictive distribution can be used to generate a set of $m$ complete datasets and these datasets can be used with any modelling approach and pooled using Rubin's rules. We describe how *BACE* works, test its robustness using simulations, and demonstrate how to apply it using two case studies.

Table 1: R tools for imputing missing data, against the needs of phylogenetic comparative analysis. *Phylogeny* (Phylo.): a tree-derived covariance informs the imputation. *Proper multiple imputation* (MI): multiple imputed datasets (or an integrated Bayesian posterior) are produced, so downstream analyses reflect imputation uncertainty. *Multilevel* (Multi.): non-independence beyond the phylogeny (random effects / clustered rows). Remaining columns: an explicit imputation model for continuous (Cont.), count, binary (Bin.), ordinal (Ord., treated as ordered) and nominal (Nom., unordered multinomial) variables. ✓ = documented in the package's reference documentation (October 2026); (✓) = partial support; ✗ = not documented.

| Package | Phylo. | MI | Multi. | Cont. | Count | Bin. | Ord. | Nom. | Reference |
|---|---|---|---|---|---|---|---|---|---|
| **BACE** | ✓ | ✓ | (✓)[a] | ✓ | ✓ | ✓ | ✓ | ✓ | this paper |
| Rphylopars | ✓ | ✗ | (✓) | ✓ | ✗ | ✗ | ✗ | ✗ | Goolsby et al. (2017) |
| phylomice[c] | ✓ | ✓ | ✗ | ✓ | ✗ | ✗ | ✗ | ✗ | Drhlik (2016) |

| Package | Phylo. | MI | Multi. | Cont. | Count | Bin. | Ord. | Nom. | Reference |
|---|---|---|---|---|---|---|---|---|---|
| TDIP[c] | ✓ | ✗ | ✗ | ✓ | ✗ | ✓ | ✗ | ✓ | Gendre et al. (2024) |
| BHPMF[c] | (✓)[b] | ✗ | (✓) | ✓ | ✗ | ✗ | ✗ | ✗ | Schrodt et al. (2015) |
| mice | ✗ | ✓ | (✓) | ✓ | ✗ | ✓ | ✓ | ✓ | Buuren & Groothuis-Oudshoorn (2011) |
| miceadds | ✗ | ✓ | ✓ | ✓ | ✗ | ✓ | (✓)[d] | ✗[d] | Robitzsch & Grund (2026) |
| micemd | ✗ | ✓ | ✓ | ✓ | ✓ | ✓ | (✓)[d] | (✓)[d] | Audigier et al. (2018) |
| pan | ✗ | ✓ | ✓ | ✓ | ✗ | ✗ | ✗ | ✗ | Schafer & Yucel (2002) |
| jomo | ✗ | ✓ | ✓ | ✓ | ✗ | ✓ | (✓)[d] | ✓ | Carpenter & Kenward (2013) |
| mitml | ✗ | ✓ | ✓ | ✓ | ✗ | ✓ | ✗ | ✓ | Grund et al. (2023) |
| hmi[c] | ✗ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | Speidel et al. (2020) |
| mdmb | ✗ | ✓ | (✓) | ✓ | ✗ | ✓ | ✓ | ✗ | Robitzsch & Lüdtke (2024) |
| JointAI | ✗ | (✓)[e] | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | Erler et al. (2021) |
| smcfcs | ✗ | ✓ | ✗ | ✓ | ✓ | ✓ | ✓ | ✓ | Bartlett et al. (2015) |
| missForest | ✗ | ✗ | ✗ | ✓ | ✗ | ✓ | ✗ | ✓ | Stekhoven & Bühlmann (2012) |

[a] Phylogenetic and species (replicated taxa) random effects; other grouping factors are not supported. [b] Taxonomic hierarchy (species/genus/family), not a tree with branch lengths. [c] Not on CRAN: *phylomice* and *TDIP* (GitHub only), *BHPMF* (archived 2017), *hmi* (removed 2022). [d] Ordinal/nominal support is limited: single-level methods only (*micemd*), predictive mean matching rather than an ordinal model (*miceadds*, whose multilevel methods also cannot impute nominal variables), or substantive-model-compatible outcome models only (*jomo*). [e] Imputation is integrated with the analysis model (fully Bayesian); imputed datasets can be exported.

## The *BACE* R package: How it works?

*BACE* generates multiple completed datasets or pooled model(s) from comparative data with missing trait values (Figure 1). Users provide a data frame with species identifiers, a phylogeny with branch lengths, and model formulas specifying the predictors of each incomplete trait (Figure 1,





Figure 1: **Overview of the BACE imputation workflow**. (1) Inputs are an incomplete trait dataset and a phylogeny covering the species. (2) Chained imputation: a separate phylogenetic mixed model is fitted for each incomplete trait, and missing values are updated with point estimates until they stabilise. (3) Final imputation: models are refitted and missing values are drawn from their posterior predictive distributions. Repeated runs produce multiple completed datasets while retaining the observed values. (4) Downstream analysis: models are fitted to each completed dataset and posterior draws are pooled across imputations. Alternatively, parameter estimates and their variances can be combined using Rubin's rules to include imputation uncertainty.

step 1). Continuous, count, binary, ordered categorical and unordered categorical traits can be included in the same dataset. Each trait's type is declared by how its column is stored in the data frame (numeric for continuous, integer for counts, factors — ordered or unordered — for categorical traits). *BACE* then selects the correct error distribution for the model automatically. For example, if the response variable is a count a Poisson error distribution (log link) is used whereas if the response is a two-level categorical variable it will use a Bernoulli error distribution (logit link). Because the phylogenetic tree is user-supplied, evolutionary models other than Brownian motion can also be accommodated. For example, any model expressible as a branch-length transformation — Pagel's  and , early burst, or single-optimum Ornstein–Uhlenbeck on an ultrametric tree — is fitted by transforming the tree before imputation, with the transform parameter estimated externally (Blomberg et al., 2003; Hansen, 1997; Pagel, 1999).

*BACE* fits a separate phylogenetic mixed model for each incomplete trait with *MCMCglmm* (Hadfield, 2010). Each model uses relationships among traits and shared evolutionary history to inform probable missing values. When species have repeated observations, a separate species effect can account for similarities not explained by the phylogeny (Cinar et al., 2022; Hadfield & Nakagawa, 2010; Nakagawa & Santos, 2012). Imputation has two phases. First, the models are fitted repeatedly, updating missing values with point estimates until these estimates stabilise (Figure 1, step 2). Once the chained equations stabilise, *BACE* cycles through the models again, drawing plausible missing values from their posterior predictive distributions (Figure 1, step 3). Repeating this process produces $m$ completed datasets, each retaining the original observed values.

The main function `bace()` runs this workflow and returns a single object containing the $m$ completed datasets, pooled model posteriors, individual model fits and convergence diagnostics. Users can extract the completed datasets with `get_imputed_data()` and a pooled model for a particular trait with `get_pooled_model()`. Imputed values are returned on the scale of the supplied data. Model coefficients retain the fitted model's scale: Gaussian variables are standardised internally, count responses use a log link, and binary and ordinal responses use a probit link. The pooled posterior can be used directly when that model's parameters are of interest. Alternatively, users can fit a different analysis model to each completed dataset and combine its results. Both routes carry imputation uncertainty into downstream analyses (Figure 1, step 4).

## Typical *BACE* workflow

We demonstrate the typical user workflow in the code below, and Figure 2 maps the package's relevant functions onto the workflow steps, including the alternative Rubin's-rules pathway. A detailed tutorial with simulated data, and demonstrating our case studies, is also provided in the package documentation (https://daniel1noble.github.io/BACE/).

For users wishing to test out *BACE* or conduct simulations based on characteristics of their data, we also provide a simulation function `sim_bace()`. `sim_bace()` allows users to control the number of species, replicate observations per species, level of missing trait data, the distribution type of response and predictor variables, and the strength of the phylogenetic signal in each trait. Such simulations can help in planning analyses and identifying their power.

```
# 1) Prepare data and tree (rows = taxa; factors declared; counts as integers)
# 2) Run the full workflow: chained imputation -> convergence check ->
#    n_final posterior-predictive imputations -> pooled posteriors
```

```
res <- bace(fixformula     = "y ~ x1 + x2",
            ran_phylo_form = "~ 1 | Species",
            phylo          = tree,
            data           = dat,
            runs           = 10, n_final = 50)

# 3) Diagnostics
plot(assess_convergence(res$initial_results))

# 4a) Bayesian route: pooled MCMCglmm posteriors
summary(get_pooled_model(res, "y"))

# 4b) Rubin's-rules route: any downstream model per imputed dataset
fits <- with_imputations(res, function(d) lm(y ~ x1 + x2, data = d))
pool_mi(fits)
```

*BACE* users with existing species trait datasets can apply the *BACE* workflow by specifying the imputation and/or analysis model(s) in `bace()`. Model formulas can be provided as a single formula or a list of formulas, one for each missing response variable, defined by the argument `fixformula`. Users then supply the name of species-level random effect as a formula (using argument `ran_phylo_form`). If users have repeated species, then the random effect structure can be adjusted by setting the `species = TRUE` argument, which will add an additional non-phylogenetic random effect for species (Cinar et al., 2022; Hadfield & Nakagawa, 2010). If provided an analysis model with missing predictors, *BACE* will create equations for all variables with missing trait data automatically.

*BACE* provides a series of convergence diagnostics which help to identify possible issues during model fitting that need to be checked prior to inference. For example, within-chain MCMC convergence of each per-variable model as well as convergence of the chained-equations loop (the imputed-data sequence) are provided as output when `bace()` is run or which can be investigated after the imputation workflow is comeplete. `assess_convergence()` and `plot_*_convergence()` are two functions that provide users with various diagnositics on the change in imputed values as chained equations proceed. MCMC chains of any final model and parameter can be investigated using standard R functions (e.g., coda:). Such convergence information is impportant to identify when different priors or a different number of MCMC iterations are needed. In `bace()`, `runs` controls the initial chained-imputation passes, and `n_final` sets the number of completed datasets (default 50). Within each model, `nitt`, `burnin` and `thin` control the total MCMC iterations, initial iterations discarded and interval between retained samples, respectively. For `nitt`, `burnin` and `thin`, users can supply one value for all models or a list with one value per equation, in formula order, to tune sampling separately for each response.

*BACE* provides two routes for combining results across imputations. The default, `pool_posteriors()`, stacks posterior draws from the final *MCMCglmm* fits for each trait to approximate its marginal posterior (Zhou & Reiter, 2010). The returned models contain pooled draws for regression coefficients and variance components, from which users can obtain posterior means and credible intervals. Alternatively, `with_imputations()` fits a user-specified analysis model to each completed dataset and `pool_mi()` combines its coefficient estimates and variances using Rubin's rules (Rubin, 1987). `pool_mi()` returns a table with one row per coefficient, including the pooled estimate,





Figure 2: **Map of the core user-facing *BACE* functions**. Solid arrows show the main workflow from raw comparative data through chained imputation (`bace_imp()`), convergence assessment of the chained loop (`assess_convergence()`), posterior-predictive final imputations (`bace_final_imp()`) and posterior pooling (`pool_posteriors()`), with accessors (`get_pooled_model()`, `get_imputed_data()`) reading the finished object. The shaded band marks the steps run end-to-end by `bace()`, which automatically retries the chained phase on non-convergence. The lower-right branch is the alternative Rubin's-rules pathway: `with_imputations()` fits any downstream analysis model to each completed dataset and `pool_mi()` combines the estimates. Dashed boxes mark optional routes: simulating benchmark data (`sim_bace()`), pre-flight phylogenetic-signal screening (`phylo_signal_summary()`), per-model MCMC and prior control, and convergence visualisations.

standard error, confidence interval, test statistic, degrees of freedom and $p$-value. It also reports the **fraction of missing information** (`fmi`), which estimates how much information about each coefficient is lost because of missing data, and the **relative increase in variance** (`riv`), which measures the extra variance from imputation uncertainty relative to the average within-imputation variance of that coefficient (Rubin, 1987). Larger values indicate that missing data have a greater effect on the precision of that coefficient.

# Evaluating the performance of BACE with simulations

## Simulation design

*(Structure only. Sources: `dev/12_recovery_simulation.R`, `dev/13_response_type_simulation.R`, `dev/simulation_results/SIMULATION_REPORT.qmd`; design details in `.agents/simulation-report.html`.)*
- Study A: parameter recovery (gaussian slope) under MCAR/MAR; oracle and complete-case comparators. - Study B: five response types x MCAR/MAR; per-type accuracy/coverage metrics and their finite-draw/metric ceilings (`cover95_t`, prob-based sets, Bayes ceiling for categorical).

## Parameter recovery and inferential calibration (Study A)

## Imputation accuracy across response types (Study B)

# Worked examples

We now turn to two case studies to demonstrate application of *BACE*. The first case study involves data from Pottier et al. (2025). Pottier et al. (2025) imputed heat tolerance (continuous trait) under standardised conditions for 5,203 amphibian species of which only 524 species (10.1%) have trait data. The second case study, uses BIRDBASE and AVONET to impute social (categocial and continuous data), for 10,563 bird species.

# Discussion

*(Structure only.)* - What BACE adds relative to existing MI and phylogenetic-prediction tools. - Honest limitations - Practical guidance (pre-flight signal screening; when BACE is the wrong tool).

# Acknowledgements

*(To write.)*

## Author contributions

Author contributions are reported using the CRediT taxonomy. **DWAN:** conceptualization, methodology, software, formal analysis, validation, visualization, writing — original draft, writing — review and editing, project administration. **SMD:** conceptualization, methodology, software, writing — review and editing. **AM:** software, validation, investigation (AVONET case study and tutorial), visualization, writing — review and editing. **PP:** data curation, investigation (amphibian heat-tolerance case study), writing — review and editing. **SN:** conceptualization, methodology, formal analysis, validation (independent simulation study), writing — review and editing. All authors approved the final version.

## Data and code availability

*BACE* (version 0.1.0) is free and open-source software released under the GPL ( 3). The source code and development version are available on GitHub (https://github.com/daniel1noble/BACE), where issues and feature requests can also be reported. A full tutorial — covering installation, when *BACE* is appropriate relative to other imputation tools, the complete workflow, and interpretation of pooled output and diagnostics — is available at https://daniel1noble.github.io/BACE/. A CRAN submission is in preparation; once accepted, the package will be installable directly via `install.packages("BACE")` (https://CRAN.R-project.org/package=BACE). Code reproducing all simulations reported here is included in the repository (`dev/`).

## References


Audigier, V., White, I. R., Jolani, S., Debray, T. P. A., Quartagno, M., Carpenter, J. R., Buuren, S. van, & Resche-Rigon, M. (2018). Multiple imputation for multilevel data with continuous and binary variables. *Statistical Science*, *33*(2), 160–183. https://doi.org/10.1214/18-STS646

Bartlett, J. W., Seaman, S. R., White, I. R., & Carpenter, J. R. (2015). Multiple imputation of covariates by fully conditional specification: Accommodating the substantive model. *Statistical Methods in Medical Research*, *24*(4), 462–487. https://doi.org/10.1177/0962280214521348

Blomberg, S. P., Garland, T., & Ives, A. R. (2003). Testing for phylogenetic signal in comparative data: Behavioral traits are more labile. *Evolution*, *57*(4), 717–745. https://doi.org/10.1111/j.0014-3820.2003.tb00285.x

Blomberg, S. P., & Todorov, O. S. (2025). The fallacy of single imputation for trait databases: Use multiple imputation instead. *Methods in Ecology and Evolution*, *16*(4), 658–667. https://doi.org/10.1111/2041-210X.14494

Bruggeman, J., Heringa, J., & Brandt, B. W. (2009). PhyloPars: Estimation of missing parameter values using phylogeny. *Nucleic Acids Research*, *37*(suppl_2), W179–W184. https://doi.org/10.1093/nar/gkp370

Buuren, S. van, & Groothuis-Oudshoorn, K. (2011). Mice: Multivariate imputation by chained equations in r. *Journal of Statistical Software*, *45*(3), 1–67. https://doi.org/10.18637/jss.v045.i03

Carpenter, J. R., & Kenward, M. G. (2013). *Multiple imputation and its application*. Wiley. https://doi.org/10.1002/9781119942283

Cinar, O., Nakagawa, S., & Viechtbauer, W. (2022). Phylogenetic multilevel meta-analysis: A simulation study on the importance of modelling the phylogeny. *Methods in Ecology and Evolution*, *13*(2), 383–395. https://doi.org/10.1111/2041-210X.13760

Drhlik, P. (2016). *Phylomice: Phylogenetic imputation methods for mice*. https://github.com/pdrhlik/phylomice

Erler, N. S., Rizopoulos, D., & Lesaffre, E. M. E. H. (2021). JointAI: Joint analysis and imputation of incomplete data in r. *Journal of Statistical Software*, *100*(20), 1–56. https://doi.org/10.18637/jss.v100.i20

Felsenstein, J. (1985). Phylogenies and the comparative method. *The American Naturalist*, *125*(1), 1–15. https://doi.org/10.1086/284325

Garamszegi, L. Z., & Møller, A. P. (2011). Nonrandom variation in within-species sample size and missing data in phylogenetic comparative studies. *Systematic Biology*, *60*(6), 876–880. https://doi.org/10.1093/sysbio/syr060

Gelman, A., & Hill, J. (2007). *Data analysis using regression and multilevel/hierarchical models*. Cambridge University Press. https://doi.org/10.1017/CBO9780511790942

Gendre, M., Hauffe, T., Pimiento, C., & Silvestro, D. (2024). Benchmarking imputation methods for categorical biological data. *Methods in Ecology and Evolution*, *15*(9), 1624–1638. https://doi.org/10.1111/2041-210X.14339

González-Suárez, M., Lucas, P. M., & Revilla, E. (2012). Biases in comparative analyses of extinction risk: Mind the gap. *Journal of Animal Ecology*, *81*(6), 1211–1222. https://doi.org/10.1111/j.1365-2656.2012.01999.x

Goolsby, E. W., Bruggeman, J., & Ané, C. (2017). Rphylopars: Fast multivariate phylogenetic comparative methods for missing data and within-species variation. *Methods in Ecology and Evolution*, *8*(1), 22–27. https://doi.org/10.1111/2041-210X.12612

Grafen, A. (1989). The phylogenetic regression. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, *326*(1233), 119–157. https://doi.org/10.1098/rstb.1989.0106

Grund, S., Robitzsch, A., & Lüdtke, O. (2023). *Mitml: Tools for multiple imputation in multilevel modeling*. https://CRAN.R-project.org/package=mitml

Hadfield, J. D. (2010). MCMC methods for multi-response generalized linear mixed models: The MCMCglmm r package. *Journal of Statistical Software*, *33*(2), 1–22. https://doi.org/10.18637/jss.v033.i02

Hadfield, J. D., & Nakagawa, S. (2010). General quantitative genetic methods for comparative biology: Phylogenies, taxonomies and multi-trait models for continuous and categorical characters. *Journal of Evolutionary Biology*, *23*(3), 494–508. https://doi.org/10.1111/j.1420-9101.2009.01915.x

Halliwell, B., Holland, B. R., & Yates, L. A. (2025). Multi-response phylogenetic mixed models: Concepts and application. *Biological Reviews of the Cambridge Philosophical Society*, *100*(3), 1294–1316. https://doi.org/10.1111/brv.70001

Hansen, T. F. (1997). Stabilizing selection and the comparative analysis of adaptation. *Evolution*, *51*(5), 1341–1351. https://doi.org/10.1111/j.1558-5646.1997.tb01457.x

Little, R. J. A., Carpenter, J. R., & Lee, K. J. (2024). A comparison of three popular methods for handling missing data: Complete-case analysis, inverse probability weighting, and multiple imputation. *Sociological Methods & Research*, *53*(3), 1105–1135. https://doi.org/10.1177/00491241221113873

Little, R. J. A., & Rubin, D. B. (2002). *Statistical analysis with missing data* (2nd ed.). Wiley. https://doi.org/10.1002/9781119013563

Martins, E. P., & Hansen, T. F. (1997). Phylogenies and the comparative method: A general

approach to incorporating phylogenetic information into the analysis of interspecific data. *The American Naturalist*, *149*(4), 646–667. https://doi.org/10.1086/286013

Nakagawa, S., & Freckleton, R. P. (2008). Missing inaction: The dangers of ignoring missing data. *Trends in Ecology & Evolution*, *23*(11), 592–596. https://doi.org/10.1016/j.tree.2008.06.014

Nakagawa, S., & Santos, E. S. A. (2012). Methodological issues and advances in biological meta-analysis. *Evolutionary Ecology*, *26*(5), 1253–1274. https://doi.org/10.1007/s10682-012-9555-5

Noble, D. W. A., & Nakagawa, S. (2021). Planned missing data designs and methods: Options for strengthening inference, increasing research efficiency and improving animal welfare in ecological and evolutionary research. *Evolutionary Applications*, *14*(8), 1958–1968. https://doi.org/10.1111/eva.13273

Pagel, M. (1994). Detecting correlated evolution on phylogenies: A general method for the comparative analysis of discrete characters. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, *255*(1342), 37–45. https://doi.org/10.1098/rspb.1994.0006

Pagel, M. (1999). Inferring the historical patterns of biological evolution. *Nature*, *401*(6756), 877–884. https://doi.org/10.1038/44766

Pottier, P., Kearney, M. R., Wu, N. C., Gunderson, A. R., Rej, J. E., Rivera-Villanueva, A. N., Pollo, P., Burke, S., Drobniak, S. M., & Nakagawa, S. (2025). Vulnerability of amphibians to global warming. *Nature*, *639*(8056), 954–961. https://doi.org/10.1038/s41586-025-08665-0

Robitzsch, A., & Grund, S. (2026). *Miceadds: Some additional multiple imputation functions, especially for mice*. https://CRAN.R-project.org/package=miceadds

Robitzsch, A., & Lüdtke, O. (2024). *Mdmb: Model based treatment of missing data*. https://CRAN.R-project.org/package=mdmb

Rubin, D. B. (1987). *Multiple imputation for nonresponse in surveys*. John Wiley & Sons. https://doi.org/10.1002/9780470316696

Schafer, J. L., & Graham, J. W. (2002). Missing data: Our view of the state of the art. *Psychological Methods*, *7*(2), 147–177. https://doi.org/10.1037/1082-989X.7.2.147

Schafer, J. L., & Yucel, R. M. (2002). Computational strategies for multivariate linear mixed-effects models with missing values. *Journal of Computational and Graphical Statistics*, *11*(2), 437–457. https://doi.org/10.1198/106186002760180608

Schrodt, F., Kattge, J., Shan, H., Fazayeli, F., Joswig, J., Banerjee, A., Reichstein, M., Bönisch, G., Díaz, S., Dickie, J., Gillison, A., Karpatne, A., Lavorel, S., Leadley, P., Wirth, C. B., Wright, I. J., Wright, S. J., & Reich, P. B. (2015). BHPMF – a hierarchical bayesian approach to gap-filling and trait prediction for macroecology and functional biogeography. *Global Ecology and Biogeography*, *24*(12), 1510–1521. https://doi.org/10.1111/geb.12335

Speidel, M., Drechsler, J., & Jolani, S. (2020). The r package hmi: A convenient tool for hierarchical multiple imputation and beyond. *Journal of Statistical Software*, *95*(9), 1–48. https://doi.org/10.18637/jss.v095.i09

Stekhoven, D. J., & Bühlmann, P. (2012). MissForest—non-parametric missing value imputation for mixed-type data. *Bioinformatics*, *28*(1), 112–118. https://doi.org/10.1093/bioinformatics/btr597

Tanner, M. A., & Wong, W. H. (1987). The calculation of posterior distributions by data augmentation. *Journal of the American Statistical Association*, *82*(398), 528–540. https://doi.org/10.1080/01621459.1987.10478458

Zhou, X., & Reiter, J. P. (2010). A note on Bayesian inference after multiple imputation. *The American Statistician*, *64*(2), 159–163. https://doi.org/10.1198/tast.2010.09109