Supplementary Analysis, Figures, and Tables

Default priors for all Bayesian models were used. For all population-level (i.e. Fixed effects), the default prior for the intercept is a normal distribution with a mean 0 and standard deviation 10. The default prior for the shape parameter of the intercept was a Student-t distribution with mean 0, scale 2.5, and 3 degrees of freedom. The default prior for residuals (sigma) was a Student-t distribution with mean 0, scale 2.5, and 3 degrees of freedom. The Cholesky factor was used as the default prior for correlations between random effects.

To determine if standard metabolic rate (SMR) resulted in different conclusions compared to if we used all metabolic measurements taken over night, we refit our models using only SMR. We defined SMR as the lowest 10% of values of oxygen consumption rate during our overnight trials. For both species this resulted in the removal of nearly 90% of our data and resulted in higher sampling error (unsurprisingly). Nonetheless, this did not change the overall results. Below we provide the detailed results and corresponding figures and tables using SMR for each species.

Bassiana duperreyi - Once SMR data (lowest 10% of metabolic rate) were removed, we had a total of 83 measurements for 40 individuals (male_{SR} XX: n = 13, female XX: n = 15, male XY: n = 12). There was a strong scaling relationship between log metabolic rate and log mass (Table S1). Sex-reversed male XX B. duperreyi had a scaling relationship that was most like their phenotypic counterparts (male XY - male_{SR} XX; pMCMC = 0.26; Table S1; Fig. S1) compared to their genotypic counterparts (female XX - male_{SR} XX; pMCMC = 0.07). The homogeneous variance model was the most parsimonious ([heteroscedastic model – homoscedastic model] loo: -1.31, SE = 2), accounting for 73% (95% CI:0.63 - 0.8) of the variation in metabolic rate.

Pogona vitticeps - Once SMR data (lowest 10% of metabolic rate) were filtered we had a total of total of 146 measurements for 96 individuals (female_{SR} ZZ: n=28, female ZW: n=30, male ZZ: n=38) were recorded. There was a strong scaling relationship between log metabolic rate and log mass (Table S2). Sex-reversed female *P. vitticeps* (female_{SR} ZZ) had a scaling relationship that was overall higher than their genotypic counterparts (male ZZ - female_{SR} ZZ; pMCMC = 0.61; Fig S1), but lower than their phenotypic counterparts (female ZW - female_{SR} ZZ; pMCMC = 0.64; Table 2). The heteroscedasticity variance model was the most parsimonious ([heteroscedastic model – homoscedastic model] loo: -16.4, SE = 5.03), accounting for 86% (95% CI:0.79 - 0.92) of the variation in metabolic rate.

Formatted: Font: Not Italic




Figure S1. Comparison of mean nest temperatures between Piccadilly Circus and Mt Ginini.




Figure \$2. Comparison of log standard metabolic rate ($\dot{V}O2 \text{ mL min}^{-1}$) across log body mass (g) by sex class for *Bassiana duperreyi* (A-B) and *Pogona vitticeps* (C-D). Sex-reversed individuals (male_{SR} XX or female_{SR} ZZ) are denoted by red colour, phenotypic females (female XX or female ZW) are denoted in black, phenotypic males (male XY or male ZZ) are denoted in blue. Fitted lines were obtained from predicted values from the brms model for each species and confidence bands were constructed from the SE of prediction values for each sex (A,C). Density plots above each regression plot denote the distribution in body mass (log mass) by sex for each species. To visualize how log metabolic rate changes across log body mass, panels (B and D) show the distribution of predicted metabolic rate at three areas of log body mass (mean, +1.5SD, and -1.5SD) denoted by the dash-dot line in panels A and C for each sex and species

Deleted: S1

Table S1. Model coefficients for testing whether sex affects the slope of standard metabolic rate for *Bassiana duperreyi*, where the intercept is concordant females. Metabolic rate and mass were log transformed and time was z-transformed. Columns l-95% CI and u-95% CI, are the lower and upper bound of the 95% credible interval for each parameter, estimated from the posterior distribution.

Parameter	Estimate	1-95% CI	u-95% CI			
Fixed Effects						
Intercept (FemaleXX)	-4.78	-5.02	-4.52			
Male _{SR} XX	-0.06	-0.22	0.10			
MaleXY	-0.12	-0.28	0.05			
logMass	1.70	0.93	2.52			
ztime	0.01	-0.06	0.08			
Male _{SR} XX:logMass	-0.70	-1.52	0.08			
MaleXY:logMass	-0.14	-1.09	0.85			
Random Effects	Random Effects					
Lizard Identity (id)						
Intercept	0.08	0.00	0.19			
Slope	0.07	0.00	0.18			
Sampling Session (day)						
Intercept	0.38	0.17	0.83			
Residuals	0.26	0.21	0.32			

Table S2. Model coefficients form hetero testing whether sex affects the slope of metabolic rate for *Pogona vitticeps*, which heteroscedasticity was accounted for within the data. The intercept is concordant females. Metabolic rate and mass were log transformed and time was z-transformed. Columns 1-95% CI and u-95% CI, are the lower and upper bound of the 95% credible interval for each parameter, estimated from the posterior distribution.

Parameter	Estimate	1-95% CI	u-95% CI
Fixed Effects			
Intercept (FemaleZW)	-2.05	-2.22	-1.89
$Female_{SR}ZZ$	-0.17	-0.34	0.02
MaleZZ	-0.08	-0.22	0.06
logMass	1.41	1.08	1.76
ztime	0.00	-0.07	0.07
Female _{SR} ZZ:logMass	-0.08	-0.42	0.31
MaleZZ:logMass	-0.17	-0.53	0.19
Random Effects			
Lizard Identity (id)			
Intercept	0.18	0.11	0.26
Slope	0.11	0.01	0.23
Sampling Session (day)			
Intercept	0.22	0.13	0.35
Residuals			
Sigma_Intercept	-1.67	-2.01	-1.39
Sigma_logMass	-1.51	-2.35	-0.53
Sigma ztime	-0.11	-0.39	0.15

Table 3: BRMS model coefficients for each respective species when testing mass differences across sex class for animals used in respirometry experiments. Mass was log-transformed and lower and upper bounds were derived from the 95% credible interval for each parameter, estimated from the posterior samples.

Species	Sex	n	Estimate	1-95% CI	u-95% CI	Contrast	pMCMC Value
	Female XX	15	0.04	-0.02	0.11	Female XX-MalesR XX	0.45
B. duperreyi	MalesR XX	13	-0.02	-0.10	0.05	Male _{SR} XX-Male XY	0.91
	Male XY	12	0.02	-0.06	0.09	Male XY-Female XX	0.70
	Female ZW	30	0.01	-0.10	0.11	Female ZW-FemalesR ZZ	0.48
P. vitticeps	FemalesR ZZ	28	0.06	-0.04	0.16	Female _{SR} ZZ-Male ZZ	0.44
	Male ZZ	38	0.01	-0.08	0.10	Male ZZ-Female ZW	0.96

Deleted: Table S3. Like phenotype/Like genotype contrast comparisons of the distribution of predicted metabolic rate at three areas of log body mass (mean, +1.5SD, and -1.5SD) for *Bassiana duperreyi* and *Pogona vitticeps*. Estimate error denotes the lower and upper 95% CI and bold values indicate pMCMC values <0.05.

Species	([1]
Deleted: S3	
Deleted: 4	

Table \$4: BRMS Model coefficients for SVL and mass growth rate estimates across sex class and metabolism for Bassiana duperreyi and Pogona vitticeps. Growth rate was calculated by dividing the change in growth (SVL or mass) between the initial measurement and subsequent remeasurement by the total number of days elapsed. Due to the small size and rate of change in grams, mass was converted to centigrams for B. duperreyi (cg). Animals were remeasured between 3 and 6 months post hatch. Metabolism was estimated by the mean log O2 measurement for each individual.

Deleted: S4
Deleted: 5
Deleted: class
Deleted: duperreyi

Deleted: ¶
Deleted: ¶
Growth rate

(... [2]

<u>Species</u>	Growth rate	Covariate	Estimate	<u>1-95% CI</u>	<u>u-95% CI</u>	1
		Intercept (O2_SexFemaleXX)	-4.62	-5.02,	-4.21,	
		Growth Rate (SVLmm)	6.20,	-2.26	14.35	1
		O2_SexMaleSRXX	-0.14	-0.73	0.42	1
	<u>SVL</u> (mm/d)	O2_SexMaleXY _v	0.10	-0.44	0.64	*
		Growth Rate (SVLmm):O2_SexMaleSRXX	<u>-4.07</u> ,	-16.08,	7.37,	*
B. duperreyi		Growth Rate (SVLmm):O2 SexMaleXY,	<u>-7.84</u>	-19.83 _e	4.18,	1
		Intercept (O2_SexFemaleXX)	-4.61,	<u>-4.97</u> ,	-4.24	4
		Growth Rate (mass cg/d)	1.06	-0.98	3.04	4
	Mara	O2_SexMaleSRXX	0.20	-0.47 _y	0.87,	4
	Mass (cg/d)	O2 SexMaleXY,	0.18	-0.45,	0.82	4
		Growth Rate (mass cg/d):O2 SexMaleSRXX	-2.59	-6.49 _e	1.25,	1
		Intercept (O2_SexFemaleXX)	<u>-4.61</u>	<u>-4.97</u>	<u>-4.24</u>	4
		Intercept (O2_SexFemaleZW)	<u>-2.17</u>	<u>-2.77</u>	<u>-1.57</u>	4
		Growth Rate (SVLmm)	1.47	<u>-0.84</u>	3.79	4
		O2_SexFemaleSRZZ	0.17	<u>-0.63</u>	0.95	4
	$\frac{SVL}{(mm/d)}$	O2_SexMaleZZ	0.01	<u>-0.73</u>	0.74	¥
		Growth Rate (SVLmm):O2_SexFemaleSRZZ	<u>-1.51</u>	<u>-4.74</u>	<u>1.66</u>	*
D ::::		Growth Rate (SVLmm):O2 SexMaleZZ	<u>-0.35</u>	<u>-3.11</u>	2.45	**
P. vitticeps,		Intercept (O2_SexFemaleZW)	-1.79	<u>-2.30</u>	-1.27	4
		Growth Rate (mass g/d)	<u>-0.07</u>	<u>-2.55</u>	<u>2.41</u>	4
		O2_SexFemaleSRZZ	0.00	<u>-0.77</u>	0.75	4
	<u>Mass</u> (g/d)	O2_SexMaleZZ	<u>-0.27</u>	<u>-0.91</u>	0.37	4
	15/4/	Growth Rate (mass g/d):O2 SexFemaleSRZZ	<u>-0.90</u>	<u>-4.48</u>	<u>2.78</u>	**
		Growth Rate (mass g/d):O2 SexMaleZZ	0.92	<u>-2.08</u>	3.83	4

Formatted	[4]
Formatted	[6]
Formatted	[7]
Formatted	[8]
Formatted Table	([3])
Formatted	[5]
Deleted: Intercept (O2_SexFemaleXX)	
Deleted: -4.62	
Deleted: -5.02	
Formatted	([10])
Formatted	[11]
Formatted	[12]
Deleted: -4.21	
Formatted	[13]
Deleted: Growth Rate (SVLmm)	
Deleted: 6.20	
Deleted: -2.26	
Deleted: 14.35	
Formatted	[14]
Formatted	[15]
Deleted: O2_SexMaleSRXX	
Deleted: -0.14	
Deleted: -0.73	
Deleted: 0.42	
Formatted	[16]
Formatted	[17]
Formatted	[18]
Formatted	[9]
Deleted: O2_SexMaleXY	
Deleted: 0.10	
Formatted	[19]
Formatted	[20]
Formatted	[21]
Deleted: -0.44	
Deleted: 0.64	
Deleted: SVL(mm/d)	
Formatted	[22]
Deleted: -4.07	
Deleted: -16.08	
Deleted: 7.37	
Formatted	[23]
Formatted	[24]
Deleted: Growth Rate (SVLmm):O2_SexMaleSRX	X
Formatted	[25]
Deleted: -7.84	
Deleted: -19.83	
Deleted: 4.18	
Formatted	[26]
Formatted	[27]
Deleted: Growth Rate (SVLmm):O2_SexMaleXY	
Deleted: Intercept (O2_SexFemaleXX)	
Deleted: -4.61	
Deleted: -4.97	
Deleted: -4.24	
Formatted	[29]
Formatted	[30]
Formatted	[31]
Deleted: Growth Rate (mass cg/d)	
Dalata da 1 06	1

Deleted: 1.06

Deleted:

Table S5S6: BRMS Model coefficients for SVL and mass growth rate estimates across sex class for and metabolism *Pogona vitticeps*. Growth rate was calculated by dividing the change in growth (SVL or mass) between the initial measurement and subsequent remeasurement by the total number of days elapsed. Animals were remeasured between 3 and 6 months post hatch. Metabolism was estimated by the mean log O2 measurement for each individual.

Growth rate

(... [86])

Page 6: [1] Deleted	Author	
Page 7: [2] Deleted	Author	
Page 8: [3] Formatted Table	Author	
Formatted Table		-
Page 8: [4] Formatted	Author	
Right: 0.58 cm		-
Page 8: [5] Formatted	Author	
Font: 11 pt, Font color	ır: Black	
Page 8: [6] Formatted	Author	
Left		
Page 8: [7] Formatted	Author	
Left, Right: -0.44 cm		
Page 8: [8] Formatted	Author	
Left, Indent: Left: 0.1	cm, Right: -0.44 cm	
Page 8: [9] Formatted	Author	
Right: 0.58 cm		
Page 8: [10] Formatted	Author	
Left, Right: -0.55 cm		
Page 8: [11] Formatted	Author	
Left		
Page 8: [12] Formatted	Author	
Left, Right: -0.44 cm		
Page 8: [13] Formatted	Author	
Left, Right: -0.55 cm		
Page 8: [14] Formatted	Author	
Left		
Page 8: [15] Formatted	Author	
Left, Right: -0.44 cm		-
Page 8: [16] Formatted	Author	
Left, Right: -0.55 cm		-
Page 8: [17] Formatted	Author	
Left		
Page 8: [18] Formatted	Author	
Left, Right: -0.44 cm		
Page 8: [19] Formatted	Author	
Left, Right: -0.55 cm		-
Page 8: [20] Formatted	Author	
Left		•
Page 8: [21] Formatted	Author	
Left, Right: -0.44 cm		

ı

I

Left, Right: -0.55 cm		
Page 8: [23] Formatted	Author	
Left		
Page 8: [24] Formatted	Author	
Left, Right: -0.44 cm		
Page 8: [25] Formatted	Author	
Left, Right: -0.55 cm		
Page 8: [26] Formatted	Author	
Left		
Page 8: [27] Formatted	Author	
Left, Right: -0.44 cm		
Page 8: [28] Formatted	Author	
Right: 0.58 cm		
Page 8: [29] Formatted	Author	
Left, Right: -0.55 cm		
Page 8: [30] Formatted	Author	
Left		
Page 8: [31] Formatted	Author	
Left, Right: -0.44 cm		
Page 8: [32] Formatted	Author	
Left, Right: -0.55 cm		
Page 8: [33] Formatted	Author	
Left		
Page 8: [34] Formatted	Author	
Left, Right: -0.44 cm		
Page 8: [35] Formatted	Author	
Left, Right: -0.55 cm		
Page 8: [36] Formatted	Author	
Left		
Page 8: [37] Formatted	Author	
Left, Right: -0.44 cm		
Page 8: [38] Formatted	Author	
Left, Right: -0.55 cm		
Page 8: [39] Formatted	Author	
Left		
Page 8: [40] Formatted	Author	
Left, Right: -0.44 cm		
Page 8: [41] Formatted	Author	
Left, Right: -0.55 cm		
Page 8: [42] Formatted	Author	
Left		

(

ı

I

ı

ı

ı

ı

I

I

I

I

I

ı

I

ı

I

ı

ı

ı

ı

		_
Page 8: [43] Formatted	Author	
Left, Right: -0.44 cn	n	
Page 8: [44] Formatted	Author	
Left, Right: -0.55 cn	n	
Page 8: [45] Formatted	Author	
Left		
Page 8: [46] Formatted	Author	
Left, Right: -0.44 cn	n	_
Page 8: [47] Formatted	Author	
Font: Italic		
Page 8: [48] Formatted	Author	
Right: 0.58 cm		
Page 8: [49] Formatted	Author	1
Right: -0.55 cm		_
Page 8: [50] Formatted	Author	1
Left		•
Page 8: [51] Formatted	Author	1
Left, Right: -0.44 cn		_
Page 8: [52] Formatted	Author	<u> </u>
Right: -0.55 cm		-
Page 8: [53] Formatted	Author	1
Left		-
Page 8: [54] Formatted	Author	1
Left, Right: -0.44 cn		-
Page 8: [55] Formatted	Author	1
Right: -0.55 cm		-
Page 8: [56] Formatted	Author	1
Left		-
Page 8: [57] Formatted	Author	
Left, Right: -0.44 cn		_
Page 8: [58] Formatted	Author	1
Right: -0.55 cm	THE PARTY OF THE P	_
Page 8: [59] Formatted	Author	1
Left	Author	_
Page 8: [60] Formatted	Author	1
Left, Right: -0.44 cn		_
Page 8: [61] Formatted	Author	1
Right: -0.55 cm	Author	
	Author	1
Page 8: [62] Formatted Left	Author	
<u> </u>		1
Page 8: [63] Formatted	Author	
Left, Right: -0.44 cn	П	

(

(

(

(.

(.

(

(

(.

(.

(.

(

(

(

(.

(.

(.

I

I

I

I

I

I

I

I

I

I

I

I

I

I

I

I

I

I

Page 8: [64] Formatted	Author	
Right: -0.55 cm	Author	
Page 8: [65] Formatted	Ath.o.	
Left	Author	
<u> </u>	A 4	
Page 8: [66] Formatted Left, Right: -0.44 cn	Author	
Page 8: [67] Formatted	Author	
Right: 0.58 cm		_
Page 8: [68] Formatted	Author	
Right: -0.55 cm		
Page 8: [69] Formatted	Author	
Left		
Page 8: [70] Formatted	Author	
Left, Right: -0.44 cn	n	
Page 8: [71] Formatted	Author	
Right: -0.55 cm		
Page 8: [72] Formatted	Author	
Left		
Page 8: [73] Formatted	Author	
Left, Right: -0.44 cn	n	
Page 8: [74] Formatted	Author	
Right: -0.55 cm		
Page 8: [75] Formatted	Author	
Left		
Page 8: [76] Formatted	Author	
Left, Right: -0.44 cn		
Page 8: [77] Formatted	Author	
Right: -0.55 cm		_
Page 8: [78] Formatted	Author	
Left		
Page 8: [79] Formatted	Author	
Left, Right: -0.44 cn		
Page 8: [80] Formatted	Author	
Right: -0.55 cm	Author	
	Ath.o.	
Page 8: [81] Formatted Left	Author	
<u> </u>		
Page 8: [82] Formatted	Author	
Left, Right: -0.44 cn		-
Page 8: [83] Formatted	Author	
Left		
Page 8: [84] Formatted	Author	

(

(.

(.

(.

(.

(.

(

(.

(.

(.

(.

(

(.

(

I

I

I

I

I

I

I

I

I

I

I

Left, Right: -0.44 cm

Page 8: [85] Deleted	Author	
Page 9: [86] Deleted	Author	