

CORRIGENDUM

Nakagawa, S., Lagisz, M., Jennions, M. D., Koricheva, J., Noble, D. W., Parker, T. H., Sánchez-Tójar, A., Yang, Y., & O'Dea, R. E. (2022). Methods for testing publication bias in ecological and evolutionary meta-analyses. *Methods in Ecology and Evolution*, 13, 4–21. <https://doi.org/10.1111/2041-210X.13724>

In this review by Nakagawa et al. (2022) there were two errors in Section 4.2 'Methods for dependent effect sizes', where β_1 was given instead of β_0 :

'Stanley and Doucouliagos (2012, 2014) have shown that, with statistically significant β_1 , β_0 provides an adjusted estimate that is downwardly biased, when a true positive or a null effect exists (Figure 6e). They also state that with non-statistically significant β_1 , β_0 provides the best estimate of an adjusted mean. If the slope of SE (β_1) is statistically significant then fitting sampling variance instead of SE is recommended according to the following equation:'

has been corrected to:

'Stanley and Doucouliagos (2012, 2014) have shown that, with statistically significant β_0 , β_0 provides an adjusted estimate that is downwardly biased, when a true positive or a non-zero effect exists (Figure 6e). They also state that with non-statistically significant β_0 , β_0 provides the best estimate of an adjusted mean. If the intercept of SE (β_0) is statistically significant then fitting sampling variance instead of SE is recommended according to the following equation:'

and

'We recommend using Equation 27 to check the statistical significance of funnel asymmetry (small-study effects) because it has greater statistical power than Equation 28. Equation 27 can also be used to obtain an adjusted mean when β_1 is not statistically significant. This is because β_0 represents an adjusted overall mean when $\sqrt{\frac{1}{n_i}} = 0$. In other words, the predicted overall mean when a study has an infinitely large sample size, $\tilde{n}_i$, and therefore little to no sampling variance. In contrast, when β_1 is statistically significant in Equation 27, we recommend using Equation 28 to obtain an overall estimate adjusted for publication bias because it is less biased.'

has been corrected to:

'We recommend using Equation 27 to check the statistical significance of funnel asymmetry (small-study effects) because it has greater statistical power than Equation 28. Equation 27 can also be used to obtain an adjusted mean when β_0 is not statistically significant. This is because β_0 represents an adjusted overall mean when $\sqrt{\frac{1}{n_i}} = 0$. In other words, the predicted overall mean when a study has an infinitely large sample size, $\tilde{n}_i$, and therefore little to no sampling variance. In contrast, when β_0 is statistically significant in Equation 27, we recommend using Equation 28 to obtain an overall estimate adjusted for publication bias because it is less biased.'

The authors thank Tom Stanley for pointing out the error and apologise for any confusion or inconvenience caused.