

CPS I — Exercise Sheet 7

University of Freiburg · Winter Term 2025/26

Author: Zhanyu Dong, Xin Pu

1	2	Σ
6/6	5/9	11/15

Exercise 1: Linear-Time Properties

- Formalize P_i as a set of traces using set comprehension

- $P_1 = \{A_0 A_1 \dots \in (2^{AP})^\omega \mid \forall i \in \mathbb{N}. a \in A_i \vee b \in A_i\}$ ✓
- $P_2 = \{A_0 A_1 \dots \in (2^{AP})^\omega \mid (\forall i \in \mathbb{N}. (b \in A_i \implies \exists j \leq i. a \in A_j))\}$ ✓
- $P_3 = \{A_0 A_1 \dots \in (2^{AP})^\omega \mid \forall i \in \mathbb{N}. (a \in A_i \implies \exists j > i. b \in A_j)\}$ ✓
- $P_4 = \{A_0 A_1 \dots \in (2^{AP})^\omega \mid |\{i \in \mathbb{N} \mid a \in A_i\}| = 3\}$ ✓
- $P_5 = \{A_0 A_1 \dots \in (2^{AP})^\omega \mid (|\{i \in \mathbb{N} \mid a \in A_i\}| \neq \infty) \rightarrow (|\{i \in \mathbb{N} \mid b \in A_i\}| \neq \infty)\}$ ✓
 ∞ is not a number...
- $P_6 = \{A_0 A_1 \dots \in (2^{AP})^\omega \mid |\{i \in \mathbb{N} \mid a \in A_i\}| < \infty\}$ ✓
This is ok, but from the lecture we know better ways to express that: use $\exists$

- satisfies P_i

- $(\{a\}\{b\})^\omega$ ✓
- $\{a\}\{a, b\}^\omega$ ✓
- $(\{a\}\{b\})^\omega$ ✓
- $\{a\}\{a\}\{a\}^\omega$ ✓
- $(\{a\}\{b\})^\omega$ ✓
- $\{a\}\emptyset^\omega$ ✓

- does not satisfy P_i

- $\emptyset(\{a\}\{b\})^\omega$ ✓
- $(\{b\}\{a\})^\omega$ ✓
- $\{a\}\emptyset^\omega$ ✓
- $\{b\}^\omega$ ✓
- $\emptyset\{a\}^\omega$ ✓
- $\{a\}^\omega$ ✓

- Explain whether or not the transition system below satisfies P

- o false
- o true
- o true
- o false
- o true
- o false

no explanation, no points...


sorry

0/1.5

Exercise 2: Trace Inclusion

5/9

1. 0/3



2. First we find the Traces.

3/4

- $\text{Traces}(\mathcal{T}_{P_1}): \{\{x > 0\}^n \{x = 0\}^\omega \mid n \in \{0, 1, \dots, 5\}\}$
- $\text{Traces}(\mathcal{T}_{P_2}): \{\{x > 0\}^n \{x = 0\}^\omega \mid n \in \mathbb{N}_0\}$
- $\text{Traces}(\mathcal{T}_{P_{3a} \parallel P_{3b}}): \{\{x > 0\}^\omega\} \cup \{\{x > 0\}^k \{x = 0\}^\omega \mid k \in \mathbb{N}, k \neq 0\}$
- $\text{Traces}(\mathcal{T}_4): \{\{x > 0\}^\omega\} \cup \{\{x > 0\}^n \{x = 0\}^\omega \mid n \in \mathbb{N}_0\}$

$n \neq 0$

1. $\mathcal{T}_{P_1} \subseteq \mathcal{T}_{P_2}$ ✓
2. $\mathcal{T}_{P_2} \not\subseteq \mathcal{T}_{P_1}$ ✓
3. $\mathcal{T}_{P_1} \not\subseteq \mathcal{T}_{P_{3a} \parallel P_{3b}}$ ✓
4. $\mathcal{T}_{P_{3a} \parallel P_{3b}} \not\subseteq \mathcal{T}_{P_1}$ ✓
5. $\mathcal{T}_{P_1} \not\subseteq \mathcal{T}_4$
6. $\mathcal{T}_4 \not\subseteq \mathcal{T}_{P_1}$ ✓
7. $\mathcal{T}_{P_2} \not\subseteq \mathcal{T}_{P_{3a} \parallel P_{3b}}$ ✓

8. $\mathcal{T}_{P_{3a} \parallel P_{3b}} \not\subseteq \mathcal{T}_{P_2}$ ✓
9. $\mathcal{T}_{P_2} \not\subseteq \mathcal{T}_4$
10. $\mathcal{T}_4 \not\subseteq \mathcal{T}_{P_2}$ ✓
11. $\mathcal{T}_{P_{3a} \parallel P_{3b}} \subseteq \mathcal{T}_4$ ✓
12. $\mathcal{T}_4 \not\subseteq \mathcal{T}_{P_{3a} \parallel P_{3b}}$
 $\subseteq$

3. Eventually x=0

$$E = \{A_0 A_1 \dots \in (2^{\text{AP}})^\omega \mid \exists^\infty i \in \mathbb{N}. x = 0 \in A_i\}$$

As i mentioned the Traces in $(2)\mathcal{T}_{P_1} \models E\mathcal{T}_{P_2} \models E$

- $\mathcal{T}_{P_{3a} \parallel P_{3b}} \not\models E \text{ Exist Trace}\{x > 0\}^\omega$
- $\mathcal{T}_4 \not\models E \text{ Exist Trace}\{x > 0\}^\omega$ ✓