

CPS I — Exercise Sheet 1

University of Freiburg · Winter Term 2025/26

Author: Zhanyu Dong, XinPu

1	2	Σ
4.25/6	8/9	12.25/15

1. Propositional Logic

4.25/6

Atomic Propositions

- a Alice joins group 1
- b Bob joins group 1
- c Claire joins group 1

Assign the value true if the corresponding person joins group 1, and false else

Given Conditions

1. If Alice joins group 1, the tutor refuses to accept Bob.

Formula: $a \rightarrow \neg b$ ✓

2. At least one of Bob and Claire cannot go to group 1.

Formula: $\neg(b \wedge c)$ ✓

3. Claire hates Alice and doesn't want to be in the same group.

Formula: $a \leftrightarrow \neg c$ ✓

4. Alice wants to submit solutions with either Bob or Claire.

Formula: $a \rightarrow ((a \leftrightarrow b) \vee (a \leftrightarrow c))$ - 0.5

Truth Table

a	b	c	1	2	3	4	Overall Logic
T	T	T	F	F	F	T	F
T	T	F	F	T	T	T	F
T	F	T	T	T	F	T	F
T	F	F	T	T	T	F	F

Since a is F, your implication would be T - 0.25

a	b	c	1	2	3	4	Overall Logic
F	T	T	T	F	T	F	F
F	T	F	T	T	T	T	F
F	F	T	T	T	T	T	T
F	F	F	T	T	F	T	F


Conclusion: $a = \text{False}$, $b = \text{False}$, $c = \text{True}$, ✓ meaning


Alice & Bob join group 2, Claire joins group 1 - 1

8/9 2. Finite Automata

- (a) $L_1 = \{x_0x_1 \dots x_n \mid n \in \mathbb{N}, n \geq 1, (\forall i \leq n, x_i \in \{a, b\}), x_{n-1} = a\}$ ✓
- (b) $L_2 = \{x_0x_1 \dots x_n \mid n \in \mathbb{N}, (\forall i \leq n, x_i \in \{a, b\}), x_0 = x_n\}$ ✓
-

L1:





This automaton accepts $abbb$ for example, which is not in L_2 .

-1

• L1

- $Q: \{Q_0, Q_1, Q_2, Q_3\}$
- $\Sigma: \{a, b\}$
- $\delta: \left\{ \begin{array}{l} (Q_0, a, Q_1), (Q_0, b, Q_0) \\ (Q_1, a, Q_3), (Q_1, b, Q_2) \\ (Q_3, a, Q_3), (Q_3, b, Q_2) \\ (Q_2, a, Q_1), (Q_2, b, Q_0) \end{array} \right\}$
- $Q_{\text{init}}: \{Q_0\}$
- $F: \{Q_2, Q_3\}$

✓

• L2

- $Q: \{Q_0, Q_1, Q_2, Q_3\}$
- $\Sigma: \{a, b\}$
- $\delta: \left\{ \begin{array}{l} (Q_0, a, Q_2), (Q_0, b, Q_3) \\ (Q_2, a, Q_2), (Q_2, b, Q_1) \\ (Q_1, a, Q_2), (Q_1, b, Q_3) \\ (Q_3, a, Q_1), (Q_3, b, Q_3) \end{array} \right\}$
- $Q_{\text{init}}: \{Q_0\}$
- $F: \{Q_2, Q_3\}$

✓