

# Optimization for Machine Learning

Antonio Fuduli

Dipartimento di Matematica e Informatica - Università della Calabria

Email: [antonio.fuduli@unical.it](mailto:antonio.fuduli@unical.it)

Home page: <http://www.mat.unical.it/~fuduli>

January 2, 2022



UNIVERSITÀ DELLA CALABRIA

DIPARTIMENTO DI **MATEMATICA**  
**E INFORMATICA**

# OUTLINE

## ► PART I: PRELIMINARIES

1

► Sign of square matrices

2

► Norm of a vector

# OUTLINE

## ▸ PART II: ELEMENTS OF NONLINEAR PROGRAMMING

- 1 ▸ The optimization problems
- 2 ▸ Global and local minima
- 3 ▸ Convexity
- 4 ▸ Optimality conditions
- 5 ▸ The Wolfe dual
- 6 ▸ Some notions on the algorithms

# OUTLINE

## ▸ PART III: THE LAGRANGIAN RELAXATION

- 1 ▸ Relaxed problems
- 2 ▸ The Lagrangian relaxation problem
- 3 ▸ The Lagrangian dual

# OUTLINE

## ▶ PART IV: NUMERICAL OPTIMIZATION AND MACHINE LEARNING

1

▶ Introduction to machine learning

2

▶ Optimization in machine learning

# OUTLINE

## ▸ PART V: BINARY SUPERVISED CLASSIFICATION

- 1 ▸ Linear separation
- 2 ▸ Polyhedral separation
- 3 ▸ Spherical separation
- 4 ▸ Support Vector Machine
- 5 ▸ Fixed-center spherical separation with kernel
- 6 ▸ Proximal Support Vector Machine
- 7 ▸ Spherical separation with margin

# OUTLINE

## ▶ PART VI: UNSUPERVISED CLASSIFICATION

1

▶ A mixed integer model

2

▶ A nonsmooth model



# OUTLINE

## ▸ PART VII: BINARY SEMISUPERVISED CLASSIFICATION

1 ▸ Transductive Support Vector Machine

2 ▸ Semisupervised polyhedral separation

3 ▸ Semisupervised spherical separation



# OUTLINE

## ▶ PART VIII: MULTIPLE INSTANCE LEARNING

1

▶ Introduction to Multiple Instance Learning

2

▶ Support Vector Machine for Multiple Instance Learning

# OUTLINE

## ► PART IX: EVALUATION OF A CLASSIFIER

- 1 ► 10-fold cross-validation
- 2 ► Performance indicators
- 3 ► Leave-One-Out
- 4 ► Model selection

# OUTLINE

► PART IX: REFERENCES

# PART I

## PRELIMINARIES

# Sign of square matrices

# Sign of square matrices

## Definition (Positive semidefinite matrix)

A matrix  $A \in \mathbb{R}^{n \times n}$  is **positive semidefinite** if

$$x^T A x \geq 0 \quad \text{for any } x \in \mathbb{R}^n.$$

## Definition (Positive definite matrix)

A matrix  $A \in \mathbb{R}^{n \times n}$  is **positive definite** if

$$x^T A x > 0 \quad \text{for any } x \in \mathbb{R}^n, \text{ such that } x \neq 0.$$

# Sign of square matrices

## Definition (Negative semidefinite matrix)

A matrix  $A \in \mathbb{R}^{n \times n}$  is **negative semidefinite** if

$$x^T A x \leq 0 \quad \text{for any } x \in \mathbb{R}^n.$$

## Definition (Negative definite matrix)

A matrix  $A \in \mathbb{R}^{n \times n}$  is **negative definite** if

$$x^T A x < 0 \quad \text{for any } x \in \mathbb{R}^n, \text{ such that } x \neq 0.$$

**NOTE:** In all the other cases the matrix  $A$  is said to be **indefinite**.

# Sign of square matrices: characterizations

**NOTE 1:** A matrix  $A \in \mathbb{R}^{n \times n}$  is positive semidefinite if and only if all the eigenvalues are  $\geq 0$ .

**NOTE 2:** A matrix  $A \in \mathbb{R}^{n \times n}$  is positive definite if and only if all the eigenvalues are  $> 0$ .

**NOTE 3:** A matrix  $A \in \mathbb{R}^{n \times n}$  is negative semidefinite if and only if all the eigenvalues are  $\leq 0$ .

**NOTE 4:** A matrix  $A \in \mathbb{R}^{n \times n}$  is negative definite if and only if all the eigenvalues are  $< 0$ .



# Eigenvectors and eigenvalues

## Definition (Eigenvector and eigenvalue)

Letting  $A \in \mathbb{R}^{n \times n}$ ,  $\lambda \in \mathbb{R}$  and  $x \in \mathbb{R}^n$  such that  $x \neq 0$ , the vector  $x$  is an **eigenvector** of  $A$  and  $\lambda$  is the corresponding **eigenvalue** if

$$Ax = \lambda x.$$

# Computing the eigenvalues

$$Ax = \lambda x, \quad x \neq 0$$

$$\Downarrow$$

$$(Ax - \lambda x) = 0, \quad x \neq 0$$

$$\Downarrow$$

$$(A - \lambda I)x = 0, \quad x \neq 0$$

$$\Downarrow$$

The columns of the matrix  $A - \lambda I$  are linearly dependent, i.e. the matrix  $A - \lambda I$  is singular, i.e.

$$\underbrace{\det(A - \lambda I) = 0}.$$

characteristic polynomial

# Norm of a vector



# Definition

The norm, denoted by  $\| \cdot \|$ , is a map

$$\| \cdot \| : \mathbb{R}^n \mapsto \mathbb{R}_+$$

such that

1

$$\|x\| = 0 \Rightarrow x = 0;$$

2

$$\|x + y\| \leq \|x\| + \|y\| \quad \text{for any } x, y \in \mathbb{R}^n;$$

3

$$\|\alpha x\| = |\alpha| \|x\| \quad \text{for any } \alpha \in \mathbb{R} \text{ and } x \in \mathbb{R}^n.$$

# Some norms

1  $L_1$ -norm:  $\|x\|_1 = \sum_{j=1}^n |x_j|;$

2  $L_2$ -norm (Euclidean):  $\|x\|_2 = \sqrt{\sum_{j=1}^n x_j^2};$

3  $L_\infty$ -norm:  $\|x\|_\infty = \max_{1 \leq j \leq n} |x_j|.$

**NOTE 1:** If not differently specified, by  $\|x\|$  we mean the Euclidean norm of vector  $x$ .

**NOTE 2:** The norm is a convex function.

**NOTE 3:** The norm is a nonsmooth function. In fact, in case  $n = 1$ , we have  $\|x\|_1 = \|x\|_2 = \|x\|_\infty = |x|.$

# A note on the Euclidean norm

1

$$x \in \mathbb{R}^n \Rightarrow \|x\|_2 = \sqrt{\sum_{j=1}^n x_j^2}$$

- 2 Since the Euclidean norm is a nonsmooth function, we adopt the following trick:

$$\min_x \|x\|_2 \Leftrightarrow \min_x \frac{1}{2} \|x\|_2^2.$$

# PART II

## ELEMENTS OF NONLINEAR PROGRAMMING

# The optimization problems



# Optimization problems: some definitions


$$P \left\{ \begin{array}{ll} \min_x & f(x) \\ & x \in X, \end{array} \right.$$

where  $f : \mathbb{R}^n \mapsto \mathbb{R}$  and  $X \subseteq \mathbb{R}^n$ .


**NOTE:**

$$\left\{ \begin{array}{ll} \max_x & f(x) \\ & x \in X, \end{array} \right. \Leftrightarrow \left\{ \begin{array}{ll} -\min_x & -f(x) \\ & x \in X, \end{array} \right.$$


# Optimization problems: some definitions




# Optimization problems: some definitions



# Optimization problems: some definitions



# Optimization problems: some definitions



# Global and local minima

# Global and local minima

$$P \left\{ \begin{array}{ll} \min_x & f(x) \\ & x \in X \end{array} \right.$$

## Definition (Global minimum)

A point  $x^*$  is a global minimum for  $P$  if

- $x^* \in X$ ;
- $f(x^*) \leq f(x)$  for any  $x \in X$ .

# Global and local minima

$$P \left\{ \begin{array}{l} \min_x f(x) \\ x \in X, \end{array} \right.$$

## Definition (Local minimum)

A point  $x^*$  is a local minimum for  $P$  if

- $x^* \in X$ ;
- there exists a neighbourhood  $N$  of  $x^*$ , such that  $f(x^*) \leq f(x)$  for any  $x \in N \cap X$ .



# Global and local minima

$$P \left\{ \begin{array}{ll} \min_x & f(x) \\ & x \in X \end{array} \right.$$

## Definition (Strict local minimum)


A point  $x^*$  is a strict local minimum for  $P$  if

- $x^* \in X$ ;
- there exists a neighbourhood  $N$  of  $x^*$ , such that  $f(x^*) < f(x)$  for any  $x \in N \cap X$ , with  $x \neq x^*$ .

**NOTE:**  $x^*$  is a global minimum  $\Rightarrow x^*$  is a local minimum.


# Global and local minima

$$P \left\{ \min_x f(x) \right.$$




# Global and local minima

$$P \left\{ \min_x f(x) \right.$$




# Global and local minima

$$P \left\{ \min_x f(x) \right.$$




# Global and local minima

$$P \left\{ \min_x f(x) \right.$$



# Global and local minima

$$P \left\{ \min_x f(x) \right.$$



# Convexity

# Convex combination of two vectors

## Definition (Convex combination of two vectors)

Let  $x_1, x_2 \in R^n$ . The convex combination of  $x_1$  and  $x_2$  is the vector

$$w = \lambda x_1 + (1 - \lambda)x_2,$$

with  $\lambda \in [0, 1]$ .


$$\begin{aligned}\lambda &= 0 \\ w &= x_2\end{aligned}$$



# Convex combination of two vectors

## Definition (Convex combination of two vectors)

Let  $x_1, x_2 \in \mathbb{R}^n$ . The convex combination of  $x_1$  and  $x_2$  is the vector

$$w = \lambda x_1 + (1 - \lambda)x_2,$$

with  $\lambda \in [0, 1]$ .


$$\begin{aligned}\lambda &= 0 \\ w &= x_2\end{aligned}$$


$$\begin{aligned}\lambda &= 1 \\ w &= x_1\end{aligned}$$


# Convex combination of two vectors

## Definition (Convex combination of two vectors)

Let  $x_1, x_2 \in \mathbb{R}^n$ . The convex combination of  $x_1$  and  $x_2$  is the vector

$$w = \lambda x_1 + (1 - \lambda)x_2,$$

with  $\lambda \in [0, 1]$ .



# Convex functions

## Definition (Convex function)

A function  $f : \mathbb{R}^n \mapsto \mathbb{R}$  is **convex** if for any  $x_1, x_2 \in \mathbb{R}^n$ ,

$$f(\lambda x_1 + (1 - \lambda)x_2) \leq \lambda f(x_1) + (1 - \lambda)f(x_2),$$

for any  $\lambda \in [0, 1]$ .

## Definition (Strictly convex function)

A function  $f : \mathbb{R}^n \mapsto \mathbb{R}$  is **strictly convex** if for any  $x_1, x_2 \in \mathbb{R}^n$ , with  $x_1 \neq x_2$ ,

$$f(\lambda x_1 + (1 - \lambda)x_2) < \lambda f(x_1) + (1 - \lambda)f(x_2),$$

for any  $\lambda \in (0, 1)$ .

**NOTE:** The **sum** of convex functions is a convex function.

# Convex functions

## Definition (Concave function)


A function  $f : \mathbb{R}^n \mapsto \mathbb{R}$  is **concave** if  $-f(x)$  is convex.

## Definition (Strictly concave function)


A function  $f : \mathbb{R}^n \mapsto \mathbb{R}$  is **strictly concave** if  $-f(x)$  is strictly convex.

**NOTE:** A **linear** function is at the same time convex and concave.


# Convex functions




# Convex functions



# Convex functions




# Convex functions




$$\underbrace{f(\lambda x_1 + (1 - \lambda)x_2)}_{\text{arc}} \leq \underbrace{\lambda f(x_1) + (1 - \lambda)f(x_2)}_{\text{chord}}, \text{ for any } \lambda \in [0, 1]$$




# Convex functions




# Convex functions



# Convex functions




# Convex functions




$$\underbrace{f(\lambda x_1 + (1 - \lambda)x_2)}_{\text{arc}} \leq \underbrace{\lambda f(x_1) + (1 - \lambda)f(x_2)}_{\text{chord}}, \text{ for any } \lambda \in [0, 1]$$


# Convex functions




# Convex functions



# Convex functions



# Convex functions



$$f(\lambda x_1 + (1 - \lambda)x_2) \not\leq \lambda f(x_1) + (1 - \lambda)f(x_2), \text{ for any } \lambda \in [0, 1]$$



# Convex sets


## Definition (Convex set)

A set  $X \subseteq \mathbb{R}^n$  is **convex** if for any  $x_1, x_2 \in X$ , the vector


$$w = \lambda x_1 + (1 - \lambda)x_2 \in X$$

for any  $\lambda \in [0, 1]$ .


# Convex sets



# Convex sets




# Convex sets




$$w = \lambda x_1 + (1 - \lambda)x_2 \in X \text{ for any } \lambda \in [0, 1]$$

# Convex sets




# Convex sets




$$w = \lambda x_1 + (1 - \lambda)x_2 \in X \text{ for any } \lambda \in [0, 1]$$

# Convex sets




# Convex sets




$$w = \lambda x_1 + (1 - \lambda)x_2 \in X \text{ for any } \lambda \in [0, 1]$$




# Convex sets

 $X$

# Convex sets



# Convex sets



$$w = \lambda x_1 + (1 - \lambda)x_2 \notin X \text{ for any } \lambda \in [0, 1]$$


# Convex sets

## Definition (Convex hull)


Given a set  $X \subset \mathbb{R}^n$ , the convex hull of  $X$  is the smallest convex set containing  $X$ . It is indicated by  $\text{conv}(X)$ .

**NOTE:** If  $X$  is convex, then  $\text{conv}(X) = X$ .

# Convex hull


 $X$

# Convex hull




$\text{conv}(X)$

# Convex hull



# Convex hull





$$P \left\{ \begin{array}{ll} \min_x & f(x) \\ & x \in X, \end{array} \right.$$

where  $f : \mathbb{R}^n \mapsto \mathbb{R}$  and  $X \subseteq \mathbb{R}^n$ .

**NOTE:** If  $f$  is a convex function and  $X$  is a convex set, then  $P$  is a **convex program**.

# Optimality conditions

# The unconstrained case: optimality conditions

If  $X = \mathbb{R}^n$ , then we have the following **unconstrained optimization problem**:

$$P \left\{ \min_{x \in \mathbb{R}^n} f(x) \right.$$

where  $f : \mathbb{R}^n \mapsto \mathbb{R}$ .

# The unconstrained case: optimality conditions

**Assumption:**  $f \in C^2$ , i.e. the first and second order derivatives exist and are continuous.

**Gradient:**  $\nabla f(x) = \begin{bmatrix} \frac{\partial f}{\partial x_1} \\ \frac{\partial f}{\partial x_2} \\ \vdots \\ \frac{\partial f}{\partial x_n} \end{bmatrix}$

# The unconstrained case: optimality conditions

Assumption:  $f \in C^2$ , i.e. the first and second order derivatives exist and are continuous.

Hessian matrix:  $\nabla^2 f(x) =$

$$\begin{bmatrix} \frac{\partial^2 f}{\partial x_1^2} & \frac{\partial^2 f}{\partial x_1 \partial x_2} & \cdots & \frac{\partial^2 f}{\partial x_1 \partial x_n} \\ \frac{\partial^2 f}{\partial x_2 \partial x_1} & \frac{\partial^2 f}{\partial x_2^2} & \cdots & \frac{\partial^2 f}{\partial x_2 \partial x_n} \\ \vdots & \vdots & \vdots & \vdots \\ \frac{\partial^2 f}{\partial x_n \partial x_1} & \frac{\partial^2 f}{\partial x_n \partial x_2} & \cdots & \frac{\partial^2 f}{\partial x_n^2} \end{bmatrix}$$

# The unconstrained case: optimality conditions

$$P \left\{ \min_x f(x) \right.$$

## Theorem (First order necessary condition)

$x^*$  is a local minimum  $\Rightarrow \nabla f(x^*) = 0$ .

**NOTE 1:** We call  $x^*$  a **stationary point** if  $\nabla f(x^*) = 0$ .

**NOTE 2:** If  $f$  is **convex** then  $x^*$  is a global minimum  $\Leftrightarrow \nabla f(x^*) = 0$ .

## Theorem (Second order necessary condition)

$x^*$  is a local minimum  $\Rightarrow \nabla f(x^*) = 0$  and  $\nabla^2 f(x^*)$  is positive semidefinite.

# The unconstrained case: optimality conditions

$$P \left\{ \min_x f(x) \right.$$

## Theorem (Second order sufficient condition)

*If  $\nabla f(x^*) = 0$  and  $\nabla^2 f(x^*)$  is positive definite  $\Rightarrow x^*$  is a strict local minimum.*

# The constrained case: optimality conditions

$$P \left\{ \begin{array}{ll} \min_x f(x) \\ g_i(x) = 0 & i \in E \\ g_i(x) \geq 0 & i \in I \end{array} \right\} \triangleq X \text{ (feasible region)}$$

where  $f, g_i : \mathbb{R}^n \mapsto \mathbb{R}, i \in E \cup I$ .

**NOTE 1:** If  $f(x) = \frac{1}{2}x^T Hx + c^T x$ , with  $H \in \mathbb{R}^{n \times n}$  is symmetric, and all the functions  $g_i$  are linear, then  $P$  is a **quadratic program**.

**NOTE 2:** If function  $f$  is linear (i.e.  $f = c^T x$ ) and all the functions  $g_i$  are linear, then  $P$  is a **linear program**.



# The constrained case: optimality conditions

$$P \left\{ \begin{array}{ll} \min_x & f(x) \\ & g_i(x) = 0 \quad i \in E \\ & g_i(x) \geq 0 \quad i \in I \end{array} \right\} \triangleq X \text{ (feasible region)}$$

where  $f, g_i : \mathbb{R}^n \mapsto \mathbb{R}, i \in E \cup I$ .

## SOME PRELIMINARIES

### Definition (Active constraint)

Given a point  $\bar{x} \in X$ , a constraint  $g_i, i \in E \cup I$ , is **active** at  $\bar{x}$  if  $g_i(\bar{x}) = 0$ .

$A(\bar{x}) \triangleq$  index set of the active constraints at  $\bar{x}$ .

# The constrained case: optimality conditions

$$P \left\{ \begin{array}{ll} \min_x & f(x) \\ & g_i(x) = 0 \quad i \in E \\ & g_i(x) \geq 0 \quad i \in I \end{array} \right\} \triangleq X \text{ (feasible region)}$$

where  $f, g_i : \mathbb{R}^n \mapsto \mathbb{R}, i \in E \cup I$ .

$A(\bar{x}) \triangleq$  index set of the active constraints at  $\bar{x}$ .

**Definition (Linear Independence Constraint Qualification - LICQ)**

Given a point  $\bar{x} \in X$ , we say that the **Linear Independence Constraint Qualification** holds at  $\bar{x}$ , if the set

$$\{\nabla g_i(\bar{x}) \mid i \in A(\bar{x})\}$$

is linearly independent.

# The constrained case: optimality conditions

$$P \left\{ \begin{array}{ll} \min_x & f(x) \\ & g_i(x) = 0 \quad i \in E \\ & g_i(x) \geq 0 \quad i \in I \end{array} \right\} \triangleq X \text{ (feasible region)}$$

where  $f, g_i : \mathbb{R}^n \mapsto \mathbb{R}, i \in E \cup I$ .

## Definition (Lagrangian function)

The **Lagrangian function** of problem  $P$  is the following:

$$\mathcal{L}(x, \lambda) = f(x) - \sum_{i \in E} \lambda_i g_i(x) - \sum_{i \in I} \lambda_i g_i(x),$$

with  $\lambda_i \geq 0, \quad i \in I$ .

**NOTE:** The variables  $\lambda_i, i \in E \cup I$  are the **Lagrangian multipliers**.

# The constrained case: optimality conditions

Assumptions:  $f \in C^1$ ;  $g_i \in C^1$ ,  $i \in E \cup I$

## Theorem (Karush Kuhn Tucker conditions - KKT)

*Let  $x^*$  be a local minimum of  $P$  and let LICQ hold at  $x^*$ . Then there exist  $\lambda^*$  such that*

$$KKT \left\{ \begin{array}{lll} \nabla_x \mathcal{L}(x^*, \lambda^*) = 0 & & \\ g_i(x^*) = 0 & i \in E & \leftarrow \text{feasibility} \\ g_i(x^*) \geq 0 & i \in I & \leftarrow \text{feasibility} \\ \lambda_i^* \geq 0 & i \in I & \\ \lambda_i^* g_i(x^*) = 0 & i \in E \cup I & \leftarrow \text{complementarity conditions.} \end{array} \right.$$

# The constrained case: optimality conditions

## NOTE

$$\mathcal{L}(x, \lambda) = f(x) - \sum_{i \in E} \lambda_i g_i(x) - \sum_{i \in I} \lambda_i g_i(x)$$

$$\nabla_x \mathcal{L}(x^*, \lambda^*) = 0$$

$$\Updownarrow$$

$$\nabla f(x^*) - \sum_{i \in E} \lambda_i^* \nabla g_i(x^*) - \sum_{i \in I} \lambda_i^* \nabla g_i(x^*) = 0$$

$$\Updownarrow$$

$$\nabla f(x^*) = \sum_{i \in E} \lambda_i^* \nabla g_i(x^*) + \sum_{i \in I} \lambda_i^* \nabla g_i(x^*).$$

$$\Updownarrow \text{ since } \lambda_i^* g_i(x^*) = 0 \quad i \in E \cup I$$

$$\nabla f(x^*) = \sum_{i \in A(x^*)} \lambda_i^* \nabla g_i(x^*).$$

# The Wolfe dual

# The Wolfe dual (Wolfe, 1961 [Wol61])

## Definition (Dual)

Given an optimization problem (called **primal**), its **dual** is another optimization problem associated to the primal (by means of suitable rules).

# The Wolfe dual (Wolfe, 1961 [Wol61])

## PRIMAL

$$P \left\{ \begin{array}{ll} \min_x & f(x) \\ & g_i(x) = 0 \quad i \in E \\ & g_i(x) \geq 0 \quad i \in I \end{array} \right\} \text{feasible region } X$$

where  $f, g_i : \mathbb{R}^n \mapsto \mathbb{R}, i \in E \cup I$ .

Assumptions:  $f \in C^1; g_i \in C^1, i \in E \cup I$

The **Wolfe dual** of problem  $P$  is defined as follows:

$$D \left\{ \begin{array}{ll} \max_{x, \lambda} & \mathcal{L}(x, \lambda) \\ & \nabla_x \mathcal{L}(x, \lambda) = 0 \\ & \lambda_i \geq 0, \quad i \in I \end{array} \right.$$



# The Wolfe dual of a linear program

## Theorem

Given the problem

$$P \left\{ \begin{array}{ll} \min_x & c^T x \\ & Ax \geq b \\ & x \geq 0, \end{array} \right.$$

the Wolfe dual of  $P$  is the ordinary dual.

## Proof.

$$\mathcal{L}(x, \lambda, \mu) = \overbrace{c^T x + \lambda^T (b - Ax) - \mu^T x}^{(c - A^T \lambda - \mu)^T x + \lambda^T b} \quad \text{and} \quad \nabla_x \mathcal{L} = c - A^T \lambda - \mu$$

$\Downarrow$

$$D \left\{ \begin{array}{ll} \max_{x, \lambda, \mu} & \overbrace{(c - A^T \lambda - \mu)^T x + \lambda^T b}^0 \\ & c - A^T \lambda - \mu = 0 \\ & \lambda, \mu \geq 0. \end{array} \right. \Leftrightarrow D \left\{ \begin{array}{ll} \max_{x, \lambda} & \lambda^T b \\ & \underbrace{c - A^T \lambda}_{A^T \lambda \leq c} \geq 0 \\ & \lambda \geq 0. \end{array} \right. \Leftrightarrow D \left\{ \begin{array}{ll} \max_{x, \lambda} & \lambda^T b \\ & A^T \lambda \leq c \\ & \lambda \geq 0, \end{array} \right.$$

□

# Some notions on the algorithms

# Sketch of the algorithms

- $x_0$ : starting point;
- $x_1, x_2, \dots$ : next iterates;
- unconstrained case: **stop** in case a stationary point is generated;
- constrained case: **stop** in case a KKT point is generated.

# Line search methods



$$x_{k+1} = x_k + \alpha_k d_k,$$

where  $\alpha_k > 0$  is the **stepsize** and  $d_k$  is the **search direction**.

- Once a search direction  $d_k$  is **computed**, the stepsize  $\alpha_k$  is determined by solving the following univariate problem:

$$LS \left\{ \min_{\alpha} f(x_k + \alpha d_k) \right.$$

- Exact line search** if problem  $LS$  is exactly solved.
- Inexact line search** if problem  $LS$  is approximately solved.

# Trust region methods



$$x_{k+1} = x_k + d_k,$$

where  $d_k$  is the **search direction**, obtained by solving the following problem:

$$TR \left\{ \min_d m_k(x_k + d), \right.$$

where  $m_k$  is a “model function”, well approximating  $f$  in a neighbourhood of  $x_k$ .

- Generally:

$$TR \left\{ \begin{array}{l} \min_d m_k(x_k + d) \\ \|d\| \leq \Delta_k, \end{array} \right.$$

with  $d_k$  being the **radius** of the trust region.

# The unconstrained case: descent directions

$$P \left\{ \min_x f(x), \right.$$

where  $f : \mathbb{R}^n \mapsto \mathbb{R}$  and  $f \in C^2$ .

## Definition (Descent direction)

Let  $\bar{x} \in \mathbb{R}^n$ . The vector  $\bar{d}$  is a descent direction for problem  $P$  at  $\bar{x}$  if there exists  $\bar{\alpha} > 0$  such that

$$f(\bar{x} + \alpha \bar{d}) < f(\bar{x}), \quad \text{for any } \alpha \in ]0, \bar{\alpha}].$$

**NOTE 1:** If  $\nabla f(\bar{x})^T \bar{d} < 0$ , then  $\bar{d}$  is a descent direction at  $\bar{x}$ .

**NOTE 2:** In case  $f$  is convex, if  $\bar{d}$  is a descent direction at  $\bar{x}$ , then  $\nabla f(\bar{x})^T \bar{d} < 0$ .

# The unconstrained case: the steepest descent method

By the Taylor theorem:

$$f(x_k + d) \approx f(x_k) + \nabla f(x_k)^T d$$

$$\Downarrow$$

$$P_k \left\{ \min_d f(x_k) + \nabla f(x_k)^T d \right.$$

$$\Downarrow$$

$$P_k \left\{ \min_d \begin{array}{l} \nabla f(x_k)^T d \\ \|d\| = 1 \end{array} \right.$$

$$\Downarrow$$

$$P_k \left\{ \min_d \begin{array}{l} \nabla f(x_k)^T d \\ \frac{1}{2} \|d\|^2 = \frac{1}{2} \end{array} \right.$$

# The unconstrained case: the steepest descent method

$$\mathcal{L}(d, \lambda) = \nabla f(x_k)^T d - \lambda \left( \frac{1}{2} \|d\|^2 - \frac{1}{2} \right)$$

$\Downarrow$

$$\nabla_d \mathcal{L}(d, \lambda) = \nabla f(x_k) - \lambda d = 0 \stackrel{\text{if } \lambda \neq 0}{\Rightarrow} d = \frac{\nabla f(x_k)}{\lambda}$$

$$\frac{1}{2} \|d\|^2 = \frac{1}{2} \Rightarrow \|d\|^2 = 1 \Rightarrow \frac{\|\nabla f(x_k)\|^2}{\lambda^2} = 1 \Rightarrow \lambda = -\|\nabla f(x_k)\|$$

$\Downarrow$

$$d = -\frac{\nabla f(x_k)}{\|\nabla f(x_k)\|},$$

which is a descent direction.



# The unconstrained case: **Newton method**

By the Taylor theorem:

$$f(x_k + d) \approx f(x_k) + \nabla f(x_k)^T d + \frac{1}{2} d^T \nabla^2 f(x_k) d$$

$\Downarrow$  Assumption:  $\nabla^2 f(x_k)$  positive definite

$$P_k \left\{ \min_d \underbrace{f(x_k) + \nabla f(x_k)^T d + \frac{1}{2} d^T \nabla^2 f(x_k) d}_{m_k(d)} \right.$$

$\Downarrow$

# The unconstrained case: Newton method

$$\nabla m_k(d) = \nabla f(x_k) + \nabla^2 f(x_k)d$$

$$\Downarrow$$

$$d = -\nabla^2 f(x_k)^{-1} \nabla f(x_k),$$

which is a descent direction.

## NOTE:

- $m_k(0) = f(x_k)$ ;
- $\nabla m_k(0) = \nabla f(x_k)$ ;
- $\nabla^2 m_k(0) = \nabla^2 f(x_k)$ .

# PART III

## THE LAGRANGIAN RELAXATION

# Relaxed problems

# Relaxed problems

## Definition (Relaxed problem)

Given

$$P \left\{ \begin{array}{l} \min_x f(x) \\ x \in X_P \end{array} \right. \quad \text{and} \quad R \left\{ \begin{array}{l} \min_x g(x) \\ x \in X_R, \end{array} \right.$$

$R$  is a relaxed problem with respect to  $P$  if

- 1  $X_R \supseteq X_P$ ;
- 2  $g(x) \leq f(x)$ , for any  $x \in X_P$ .

# Relaxed problems

## Theorem (Properties of the relaxed problems)

- 1  $R$  infeasible  $\Rightarrow P$  infeasible.
- 2 Let  $x_P^*$  be an optimal solution to  $P$  and  $x_R^*$  be an optimal solution to  $R$ . Then  $g(x_R^*) \leq f(x_P^*)$ .
- 3 Let  $x_R^*$  be an optimal solution to  $R$ . If  $x_R^* \in X_P$  and  $f(x_R^*) = g(x_R^*)$ , then  $x_R^*$  is optimal to  $P$ .

# Relaxed problems

## Proof.

$$1 \quad X_R = \emptyset \Rightarrow X_P = \emptyset.$$

2

$$g(x_R^*) \leq g(x) \quad \text{for any } x \in X_R$$

$\Downarrow$

$$g(x_R^*) \leq g(x) \quad \text{for any } x \in X_P.$$

Moreover, by the definition of relaxed problem:

$$g(x) \leq f(x) \quad \text{for any } x \in X_P.$$

Combining the last two inequalities, we have:

$$g(x_R^*) \leq f(x) \quad \text{for any } x \in X_P$$

and then:

$$g(x_R^*) \leq f(x_P^*).$$

3 As above,

$$\underbrace{g(x_R^*)}_{=f(x_R^*)} \leq f(x) \quad \text{for any } x \in X_P.$$

□

# The Lagrangian relaxation problem



# The Lagrangian relaxation problem

Let  $ILP$  be the following integer program:

$$ILP \left\{ \begin{array}{l} \min_x c^T x \\ \quad \quad \quad m \text{ difficult constraints} \\ \quad \quad \quad \overbrace{Ax \geq b} \\ Bx \geq d \\ x \geq 0 \\ x \text{ int} \end{array} \right\} \triangleq X \text{ (feasible region)}$$

# The Lagrangian relaxation problem

## Definition (Lagrangian relaxation)

Let  $\lambda \in \mathbb{R}^m$  such that  $\lambda \geq 0$ . The **Lagrangian relaxation** of *ILP*, with respect to the constraints  $Ax \geq b$ , is the following problem:

$$LR(\lambda) \left\{ \begin{array}{l} z_{LR}^*(\lambda) = \min_x \overbrace{c^T x - \lambda^T (Ax - b)}^{\mathcal{L}(x, \lambda)} \\ Bx \geq d \\ x \geq 0 \\ x \text{ int} \end{array} \right\} \triangleq X_{LR}$$

# The Lagrangian relaxation problem

## Theorem

For any  $\lambda \geq 0$ ,  $LR(\lambda)$  is a relaxed problem with respect to ILP.

## Proof.

- 1  $X_{LR} \supseteq X$ .
- 2 Let  $\bar{x} \in X$ . Then:

$$\begin{aligned}
 A\bar{x} \geq b &\Rightarrow A\bar{x} - b \geq 0 \\
 &\Downarrow \\
 c^T \bar{x} - \underbrace{\lambda^T}_{\geq 0} \underbrace{(A\bar{x} - b)}_{\geq 0} &\leq c^T \bar{x} \\
 \underbrace{\hspace{10em}}_{g(\bar{x})} &\hspace{1em} f(\bar{x})
 \end{aligned}$$



E INFORMATICA

# The Lagrangian relaxation problem

## Theorem

*Let  $x_{LR}^*(\lambda)$  be an optimal solution to  $LR(\lambda)$ . If  $x_{LR}^*(\lambda) \in X$  and  $\lambda^T(Ax_{LR}^*(\lambda) - b) = 0$ , then  $x_{LR}^*(\lambda)$  is optimal to ILP.*

## Proof.

See property 3 of the theorem relative to the properties of the relaxed problems. □

# The Lagrangian dual

# The Lagrangian dual

The **Lagrangian dual** of *ILP* is the following problem:

$$LD \left\{ z_{LD}^* = \max_{\lambda \geq 0} z_{LR}(\lambda), \right.$$

i.e.

$$LD \left\{ \begin{array}{l} z_{LD}^* = \max_{\lambda \geq 0} \min_x \overbrace{c^T x - \lambda^T (Ax - b)}^{\mathcal{L}(x, \lambda)} \\ \left. \begin{array}{l} Bx \geq d \\ x \geq 0 \\ x \text{ int} \end{array} \right\} \triangleq X_{LR} \end{array} \right.$$

# The Lagrangian dual

i.e.

$$LD \left\{ \begin{array}{l} z_{LD}^*(\lambda) = \max_{\lambda \geq 0} \min_x \overbrace{c^T x - \lambda^T (Ax - b)}^{\mathcal{L}(x, \lambda)} \\ x \in \text{conv}(X_{LR}) \end{array} \right.$$

## Theorem

$z_{LD}^*$  is the optimal objective function value of the following problem:

$$\overline{LD} \left\{ \begin{array}{l} z_{LD}^* = \min_x c^T x \\ x \in \text{conv}(X_{LR}) \cap X_b, \end{array} \right.$$

where

$$X_b \triangleq \{x \in \mathbb{R}^n \mid Ax \geq b\}.$$

# The Lagrangian dual and the continuous relaxation

On one hand:

$$\overline{LD} \left\{ \begin{array}{l} z_{LD}^* = \min_x c^T x \\ x \in \text{conv}(X_{LR}) \cap X_b. \end{array} \right.$$

On the other hand, the continuous relaxation of  $ILP$  is:

$$LP \left\{ \begin{array}{l} z_{LP}^* = \min_x c^T x \\ \underbrace{X_b}_{Ax \geq b} \\ Bx \geq d \\ x \geq 0 \end{array} \right\} \triangleq X_{d0} \supseteq \text{conv}(X_{LR}),$$

As a consequence,  $LP$  is a relaxed problem with respect to  $\overline{LD}$ . Then:

$$z_{LP}^* \leq z_{LD}^*.$$



# The integrality property

We say that the **integrality property** holds if the extreme points of  $X_{d0}$  are integer. In such case:

$$\text{conv}(X_{RL}) = X_{d0}.$$

As a consequence,  $LP$  and  $\overline{LD}$  coincide and  $z_{LP}^* = z_{LD}^*$ .

# The Lagrangian dual of a linear program

## Theorem

Given the problem

$$P \left\{ \begin{array}{ll} \min_x & c^T x \\ & Ax \geq b \\ & x \geq 0, \end{array} \right.$$

the Lagrangian dual of  $P$  is the ordinary dual.

## Proof.

Relaxing the constraints  $Ax \geq b$ , the Lagrangian dual of  $P$  is

$$LD \left\{ \begin{array}{l} \max_{\lambda \geq 0} \lambda^T b + \min_{x \geq 0} (c - A^T \lambda)^T x. \end{array} \right.$$

If there exists  $j$  such that  $c_j - A_j^T \lambda < 0$ , with  $A_j$  being the  $j$ th column of  $A$ , then the min-problem is unbounded. Then we need to impose  $c - A^T \lambda \geq 0$  and in such case  $x_{LR}^*(\lambda) = 0$ . As a consequence:

$$LD \left\{ \begin{array}{ll} \max_{\lambda \geq 0} & \lambda^T b \\ & A^T \lambda \leq c. \end{array} \right.$$



## PART IV

# NUMERICAL OPTIMIZATION AND MACHINE LEARNING

# Introduction to Machine Learning

# Machine Learning

Definition (Arthur Samuel (1901-1990), 1959)



**Machine Learning** is the field of study that gives computers the ability **to learn** without being explicitly programmed.

Definition (Tom Mitchell, Machine Learning, McGraw Hill, 1997)

The field of **Machine Learning** is concerned with the question of how to construct computer programs that automatically **improve with experience**.

# Machine Learning and pattern classification

- A relevant part of Machine Learning is constituted by the **pattern classification**, whose objective is to categorize different objects into two or more classes, on the basis of their similarities.
- From the mathematical point of view, the objects can be represented as vectors of  $n$  real numbers (**points in  $\mathbb{R}^n$** ), where each number describes a **feature** of the object (**feature vector**).
- Constructing a **classifier** means to generate one or more **surfaces**, which separate the objects into two or more different classes.
- The generation of the surfaces is performed by **learning** from some objects (**training set**) whose class is known (for example on the basis of the experience).
- Why? The aim is **to predict** the class of any new object, after **training** the classifier on the training set.




# Pattern classification: an example

The aim is **to predict** the class of any new object (after **training** the classifier on the **training set**).

## EXAMPLE


- A bank needs a criterion to decide whether to loan money or not.
- Starting from the past experience, the analyst tries to analyze the data relative to the past clients on the basis of their salary and of the size of the loan (**two features**, i.e.  $n = 2$ ).

# Pattern classification: an example






# Pattern classification: an example




18 past solvent clients

# Pattern classification: an example



18 past solvent clients and  
14 past insolvent ones.


# Pattern classification: an example



TRAINING SET:

18 past solvent clients and  
14 past insolvent ones.


# Pattern classification: an example



TRAINING SET:

18 past solvent clients and  
14 past insolvent ones.

# Pattern classification: an example




NEW CLIENT:

30.000 euros (loan size)

50.000 euros (annual salary).

# Pattern classification: an example



NEW CLIENT:

70.000 euros (loan size)

40.000 euros (annual salary).

# Pattern classification: an example

In the example, given the separating hyperplane,

$$H(v, \gamma) \triangleq \{x \in \mathbb{R}^n \mid v^T x = \gamma\}, \text{ with } v \in \mathbb{R}^n \text{ and } \gamma \in \mathbb{R},$$

for classifying the new client  $\bar{x}$ , we have used the following decision function:

$$\text{sign}(v^T \bar{x} - \gamma),$$

i.e.

$$\text{if } v^T \bar{x} - \gamma \begin{cases} \geq 0, & \text{the client is classified as solvent} \\ < 0, & \text{the client is classified as insolvent} \end{cases}$$

# Pattern classification: some applications

- Text and web classification.
- Object recognition of machine vision.
- Gene expression profile analysis.
- DNA and protein analysis.
- Medical diagnosis.




# Optimization in machine learning

Question:


Where does optimization intervene?

# Machine Learning and Optimization



Separable case by a hyperplane.

# Machine Learning and Optimization



This set of points is **not separable** by a hyperplane!!

**Answer:**  
To **minimize** a measure of the  
number of misclassified points.

# Wordplay...

## Classification of classification approaches...

# Classification of classification approaches...

At each client, we have “attached” a **label**:

client  $\rightarrow$   $\begin{cases} \text{solvent} \\ \text{insolvent} \end{cases}$



SUPERVISED CLASSIFICATION



On the basis of the **labelled** objects, we would like to predict the class of any new future object.

# Supervised, unsupervised and semisupervised classification

- **Supervised classification:** on the basis of the **labelled objects**, we would like to predict the class of any new future object.
- **Unsupervised classification:** we have only **unlabelled objects** that we would like to cluster on the basis of their similarities.
- **Semisupervised classification:** on the basis of the **labelled and unlabelled objects**, we would like to predict the class of the unlabelled objects.



## PART V

# BINARY SUPERVISED CLASSIFICATION

# Binary supervised classification

In the **binary classification**, we would like to discriminate only between **two classes** of objects (points in  $\mathbb{R}^n$ ).

We have two nonempty, disjoint, finite point sets in  $\mathbb{R}^n$ :



$$\mathcal{A} = \{a_1, \dots, a_m\}, \quad \text{with } a_i \in \mathbb{R}^n, \quad i = 1, \dots, m$$



$$\mathcal{B} = \{b_1, \dots, b_k\}, \quad \text{with } b_l \in \mathbb{R}^n, \quad l = 1, \dots, k.$$

- The objective is to construct a criterion for discriminating between the elements of the two sets. Then the classifier can be utilized for classifying any new object point  $\bar{x} \in \mathbb{R}^n$  as a point belonging to the set  $\mathcal{A}$  or, alternatively, to the set  $\mathcal{B}$ .

# Linear separation

# Linear separation (Mangasarian, 1965 [Man65])

- The sets  $\mathcal{A}$  and  $\mathcal{B}$  are **linearly separable** if and only if there exists a hyperplane

$$H(v, \gamma) \triangleq \{x \in \mathbb{R}^n | v^T x = \gamma\}, \text{ with } v \in \mathbb{R}^n \text{ and } \gamma \in \mathbb{R},$$

such that

- 

$$v^T a_i \geq \gamma + 1, \quad i = 1, \dots, m$$


and

- 


$$v^T b_l \leq \gamma - 1, \quad l = 1, \dots, k.$$

- NOTE:**  $\mathcal{A}$  and  $\mathcal{B}$  are linearly separable if and only if  $\text{conv}(\mathcal{A}) \cap \text{conv}(\mathcal{B}) = \emptyset$ .


# Linear separation: first example



# Linear separation: first example




# Linear separation: first example




# Linear separation: first example

$$\text{conv}(\mathcal{A}) \cap \text{conv}(\mathcal{B}) = \emptyset$$


 $\mathcal{A}$ 



# Linear separation: second example




# Linear separation: second example



# Linear separation: second example

$$\text{conv}(\mathcal{A}) \cap \text{conv}(\mathcal{B}) \neq \emptyset$$

 $\mathcal{A}$ 

# Linear separation: error function

What can we do when  $\mathcal{A}$  and  $\mathcal{B}$  are not linearly separable?

- A point  $a_i \in \mathcal{A}$  is correctly classified if

$$v^T a_i \geq \gamma + 1, \text{ i.e. if } v^T a_i - \gamma - 1 \geq 0$$

- As a consequence, a point  $a_i \in \mathcal{A}$  is misclassified if

$$v^T a_i - \gamma - 1 < 0, \text{ i.e. if } -v^T a_i + \gamma + 1 > 0.$$

- Then, for a point  $a_i \in \mathcal{A}$ , the classification error is

$$\max\{0, -v^T a_i + \gamma + 1\}.$$

# Linear separation: error function

- A point  $b_l \in \mathcal{B}$  is correctly classified if

$$v^T b_l \leq \gamma - 1, \text{ i.e. if } v^T b_l - \gamma + 1 \leq 0.$$

- As a consequence, a point  $b_l \in \mathcal{B}$  is misclassified if

$$v^T b_l - \gamma + 1 > 0.$$

- Then, for a point  $b_l \in \mathcal{B}$ , the classification error is

$$\max\{0, v^T b_l - \gamma + 1\}.$$

- Then we minimize the following classification error function:

$$f(v, \gamma) \triangleq \frac{1}{m} \sum_{i=1}^m \max\{0, -v^T a_i + \gamma + 1\} + \frac{1}{k} \sum_{l=1}^k \max\{0, v^T b_l - \gamma + 1\}.$$

# Linear separation

$$f(v, \gamma) \triangleq \frac{1}{m} \sum_{i=1}^m \overbrace{\max\{0, -v^T a_i + \gamma + 1\}}^{\xi_i} + \frac{1}{k} \sum_{l=1}^k \overbrace{\max\{0, v^T b_l - \gamma + 1\}}^{\psi_l}.$$

- Function  $f$  is a **convex nonsmooth function**;
- Minimizing  $f$  corresponds to solving the following **linear program**:

$$\left\{ \begin{array}{ll} \min_{v, \gamma, \xi, \psi} & \frac{1}{m} \sum_{i=1}^m \xi_i + \frac{1}{k} \sum_{l=1}^k \psi_l \\ & \xi_i \geq -v^T a_i + \gamma + 1 \quad i = 1, \dots, m \\ & \psi_l \geq v^T b_l - \gamma + 1 \quad l = 1, \dots, k \\ & \xi_i \geq 0 \quad i = 1, \dots, m \\ & \psi_l \geq 0 \quad l = 1, \dots, k. \end{array} \right.$$

# Polyhedral separation

# Polyhedral separation - (Megiddo, 1988 [Meg88])

- The set  $\mathcal{A}$  is  **$h$ -polyhedrally separable** from  $\mathcal{B}$  if there exists a set of  $h$  hyperplanes

$$H(v_j, \gamma_j) \triangleq \{x \in \mathbb{R}^n | v_j^T x = \gamma_j\}, \text{ with } v_j \in \mathbb{R}^n \text{ and } \gamma_j \in \mathbb{R}, j = 1, \dots, h,$$

such that



$$v_j^T a_i \leq \gamma_j - 1, \quad i = 1, \dots, m, \quad j = 1, \dots, h$$

and

- for any  $l = 1, \dots, k$ , there exists an index  $j \in \{1, \dots, h\}$  such that


$$v_j^T b_l \geq \gamma_j + 1.$$

- NOTE:**  $\mathcal{A}$  is  $h$ -polyhedrally separable from  $\mathcal{B}$  if and only if


$$\text{conv}(\mathcal{A}) \cap \mathcal{B} = \emptyset.$$




# Polyhedral separation




# Polyhedral separation: first example




# Polyhedral separation: first example




# Polyhedral separation: first example




# Polyhedral separation: first example




# Polyhedral separation: first example



# Polyhedral separation: first example




# Polyhedral separation: first example







# Polyhedral separation: second example




# Polyhedral separation: second example



# Polyhedral separation: second example



# Polyhedral separation: second example



# Polyhedral separation: error function

What can we do if  $\mathcal{A}$  is not polyhedrally separable from  $\mathcal{B}$ ?

- A point  $a_i \in \mathcal{A}$  is correctly classified if

$$v_j^T a_i - \gamma_j + 1 \leq 0, \quad \text{for all } j = 1, \dots, h,$$

i.e. if

$$\max_{j=1, \dots, h} v_j^T a_i - \gamma_j + 1 \leq 0.$$

- As a consequence, a point  $a_i \in \mathcal{A}$  is misclassified if

$$\max_{j=1, \dots, h} v_j^T a_i - \gamma_j + 1 > 0.$$

- Then the classification error, in correspondence to a point  $a_i \in \mathcal{A}$ , is
- $$\max\{0, \max_{j=1, \dots, h} v_j^T a_i - \gamma_j + 1\} = \max_{j=1, \dots, h} \{0, v_j^T a_i - \gamma_j + 1\}.$$

# Polyhedral separation: error function

- A point  $b_l \in \mathcal{B}$  is correctly classified if there exists an index  $j \in \{1, \dots, h\}$  such that

$$v_j^T b_l \geq \gamma_j + 1, \text{ i.e. } -v_j^T b_l + \gamma_j + 1 \leq 0.$$

- As a consequence, a point  $b_l \in \mathcal{B}$  is misclassified if

$$\text{for all } j = 1, \dots, h, \quad -v_j^T b_l + \gamma_j + 1 > 0,$$

- i.e. if


$$\min_{j=1, \dots, h} -v_j^T b_l + \gamma_j + 1 > 0.$$

- Then the classification error, in correspondence to a point  $b_l \in \mathcal{B}$ , is

$$\max\{0, \min_{j=1, \dots, h} -v_j^T b_l + \gamma_j + 1\}.$$


# Polyhedral separation: error function

$$f_l^+ = \max\{0, \underbrace{\min_{j=1,\dots,h} -v_j^T b_l + \gamma_j + 1}_{f_l}\}.$$



# Polyhedral separation: error function


$$f_l^+ = \max\{0, \underbrace{\min_{j=1,\dots,h} -v_j^T b_l + \gamma_j + 1}_{f_l}\}.$$






# Polyhedral separation: error function

$$f_l^+ = \max\{0, \underbrace{\min_{j=1,\dots,h} -v_j^T b_l + \gamma_j + 1}_{f_l}\}.$$




# Polyhedral separation: error function

$$f_l^+ = \max\{0, \underbrace{\min_{j=1,\dots,h} -v_j^T b_l + \gamma_j + 1}_{f_l}\}.$$




# Polyhedral separation: error function

$$f_l^+ = \max\{0, \underbrace{\min_{j=1,\dots,h} -v_j^T b_l + \gamma_j + 1}_{f_l}\}.$$



# Polyhedral separation: error function

$$f_l^+ = \max\{0, \underbrace{\min_{j=1,\dots,h} -v_j^T b_l + \gamma_j + 1}_{f_l}\}.$$



# Polyhedral separation: error function

We obtain the following classification error function:

$$f(v_1, \dots, v_h; \gamma_1, \dots, \gamma_h) \triangleq \frac{1}{m} \sum_{i=1}^m \max_{1 \leq j \leq h} \{0, v_j^T a_i - \gamma_j + 1\} + \frac{1}{k} \sum_{l=1}^k \max\{0, \min_{1 \leq j \leq h} -v_j^T b_l + \gamma_j + 1\}.$$

- Function  $f$  is nonsmooth and nonconvex.

# Spherical separation

# Spherical separation - (Tax and Duin, 1999 [TD99])

- The set  $\mathcal{A}$  is **spherically separable** from the set  $\mathcal{B}$  if there exists a sphere

$$S(x_0, R) \triangleq \{x \in \mathbb{R}^n \mid \|x - x_0\|^2 = R^2\},$$

with  $x_0 \in \mathbb{R}^n$  and  $R \in \mathbb{R}$ ,

such that

•

$$\|a_i - x_0\|^2 \leq R^2, \quad i = 1, \dots, m$$

and


•

$$\|b_l - x_0\|^2 \geq R^2, \quad l = 1, \dots, k$$

- NOTE 1:** The role played by  $\mathcal{A}$  and  $\mathcal{B}$  is not symmetric.
- NOTE 2:**  $\mathcal{A}$  is spherically separable from  $\mathcal{B} \Rightarrow \text{conv}(\mathcal{A}) \cap \mathcal{B} = \emptyset$ .




# Spherical separation: first example







# Spherical separation: first example




# Spherical separation: first example




# Spherical separation: first example




# Spherical separation: second example




# Spherical separation: second example



# Spherical separation: second example



# Spherical separation: second example



# Spherical separation: error function

What can we do if  $\mathcal{A}$  is not spherically separable from  $\mathcal{B}$ ?

- A point  $a_i \in \mathcal{A}$  is correctly classified if

$$\|a_i - x_0\|^2 - R^2 \leq 0.$$

- As a consequence, a point  $a_i \in \mathcal{A}$  is misclassified if

$$\|a_i - x_0\|^2 - R^2 > 0.$$

- Then the classification error, in correspondence to a point  $a_i \in \mathcal{A}$ , is

$$\max\{0, \|a_i - x_0\|^2 - R^2\}.$$



# Spherical separation: error function

- A point  $b_l \in \mathcal{B}$  is correctly classified if

$$R^2 - \|b_l - x_0\|^2 \leq 0.$$

- As a consequence, a point  $b_l \in \mathcal{B}$  is misclassified if

$$R^2 - \|b_l - x_0\|^2 > 0.$$

- Then the classification error, in correspondence to a point  $b_l \in \mathcal{B}$ , is

$$\max\{0, R^2 - \|b_l - x_0\|^2\}.$$

# Spherical separation: error function

- We obtain the following classification error function:

$$f(x_0, R) \triangleq R^2 + C \sum_{i=1}^m \max\{0, \|a_i - x_0\|^2 - R^2\} + \\ + C \sum_{l=1}^k \max\{0, R^2 - \|b_l - x_0\|^2\},$$

with  $C > 0$ , tuning the trade-off between the minimization of the volume of the sphere and the minimization of the misclassification error.

- Function  $f$  is nonsmooth and nonconvex.

# Spherical separation: fixing the center (Astorino and Gaudioso, 2009 [AG09])

$$f(x_0, R) \triangleq R^2 + C \sum_{i=1}^m \max\{0, \|a_i - x_0\|^2 - R^2\} + C \sum_{l=1}^k \max\{0, R^2 - \|b_l - x_0\|^2\},$$

**NOTE:** If  $x_0$  is fixed, setting  $z \triangleq R^2 \geq 0$ , then function  $f$  is convex in  $z$ .

# Spherical separation: fixing the center

$$f(z) \triangleq z + C \sum_{i=1}^m \overbrace{\max\{0, \|a_i - x_0\|^2 - z\}}^{\xi_i} + C \sum_{l=1}^k \overbrace{\max\{0, z - \|b_l - x_0\|^2\}}^{\psi_l}.$$

In this case, minimization of  $f$  corresponds to solve the following **linear program**:

$$\left\{ \begin{array}{ll} \min_{z, \xi, \psi} & z + C \sum_{i=1}^m \xi_i + C \sum_{l=1}^k \psi_l \\ & \xi_i \geq \|a_i - x_0\|^2 - z & i = 1, \dots, m \\ & \psi_l \geq z - \|b_l - x_0\|^2 & l = 1, \dots, k \\ & \xi_i \geq 0 & i = 1, \dots, m \\ & \psi_l \geq 0 & l = 1, \dots, k. \\ & z \geq 0 \end{array} \right.$$

# Support Vector Machine

# Support Vector Machine (SVM) (Vapnik, 1995 [Vap95])

- **Motivation:** To maximize the **generalization capability** of the classifier, i.e. to maximize the probability that a new point is correctly classified.
- This minimizes also possible **overfitting** phenomena.

## OVERFITTING




There is overfitting, when the classifier **fits too much** the training set.




Bad performance on the classification of new points.

# SVM: an example




# SVM: an example







# SVM: an example




# SVM: an example



# SVM: an example



# SVM: an example




# Support Vector Machine (SVM)

## Definition (Supporting hyperplane)

Let  $X \subset \mathbb{R}^n$ . A **supporting hyperplane** of  $X$  is a hyperplane such that:

- $X$  is entirely contained in one of the two half-spaces generated by the hyperplane;
- at least one boundary point of  $X$  is on the hyperplane.




$X$

# Support Vector Machine (SVM)

## Definition (Supporting hyperplane)

Let  $X \subset \mathbb{R}^n$ . A **supporting hyperplane** of  $X$  is a hyperplane such that:

- $X$  is entirely contained in one of the two half-spaces generated by the hyperplane;
- at least one boundary point of  $X$  is on the hyperplane.




# Support Vector Machine (SVM)

## Definition (Supporting hyperplane)

Let  $X \subset \mathbb{R}^n$ . A **supporting hyperplane** of  $X$  is a hyperplane such that:

- $X$  is entirely contained in one of the two half-spaces generated by the hyperplane;
- at least one boundary point of  $X$  is on the hyperplane.




# Support Vector Machine (SVM)

## Definition (Supporting hyperplane)

Let  $X \subset \mathbb{R}^n$ . A **supporting hyperplane** of  $X$  is a hyperplane such that:

- $X$  is entirely contained in one of the two half-spaces generated by the hyperplane;
- at least one boundary point of  $X$  is on the hyperplane.






# Support Vector Machine (SVM)

## Definition (Supporting hyperplane)

Let  $X \subset \mathbb{R}^n$ . A **supporting hyperplane** of  $X$  is a hyperplane such that:

- $X$  is entirely contained in one of the two half-spaces generated by the hyperplane;
- at least one boundary point of  $X$  is on the hyperplane.




# Support Vector Machine (SVM)

## Definition (Supporting hyperplane)

Let  $X \subset \mathbb{R}^n$ . A **supporting hyperplane** of  $X$  is a hyperplane such that:

- $X$  is entirely contained in one of the two half-spaces generated by the hyperplane;
- at least one boundary point of  $X$  is on the hyperplane.




# Support Vector Machine (SVM)


## Definition (Supporting hyperplane)

Let  $X \subset \mathbb{R}^n$ . A **supporting hyperplane** of  $X$  is a hyperplane such that:


- $X$  is entirely contained in one of the two half-spaces generated by the hyperplane;
- at least one boundary point of  $X$  is on the hyperplane.




# SVM: an example



# SVM: an example



# SVM: an example



# SVM

- The **margin** is the area between the two parallel hyperplanes


$$H_+(v, \gamma) \triangleq \{x \in \mathbb{R}^n | v^T x = \gamma + 1\}, \text{ with } v \in \mathbb{R}^n \text{ and } \gamma \in \mathbb{R},$$

and

$$H_-(v, \gamma) \triangleq \{x \in \mathbb{R}^n | v^T x = \gamma - 1\}, \text{ with } v \in \mathbb{R}^n \text{ and } \gamma \in \mathbb{R},$$


which are the **supporting hyperplanes** of  $\mathcal{A}$  and  $\mathcal{B}$ , respectively.

# SVM: an example






# SVM: an example



# SVM: an example




# SVM: the margin

How to compute the margin? We solve the following problem:

$$P \left\{ \begin{array}{l} \min_x \frac{1}{2} \|x - \bar{x}\|^2 \\ v^T x = \gamma + 1 \end{array} \right. ,$$

with  $\bar{x}$ , such that  $v^T \bar{x} = \gamma$ .

# SVM: an example



$$P \left\{ \begin{array}{l} \min_x \frac{1}{2} \|x - \bar{x}\|^2 \\ v^T x = \gamma + 1 \end{array} \right. \quad \text{with } \bar{x} \text{ such that } v^T \bar{x} = \gamma.$$

# SVM: the margin

## KKT conditions

$$L(x, \lambda) = \frac{1}{2} \|x - \bar{x}\|^2 - \lambda(v^T x - \gamma - 1), \text{ with } \lambda \in \mathbb{R}.$$



$$\nabla_x L(x, \lambda) = \frac{1}{2} 2(x - \bar{x}) - \lambda v = 0, \text{ i.e.}$$

$$x = \bar{x} + \lambda v.$$



$$\underbrace{v^T x}_{\gamma+1} = \underbrace{v^T \bar{x}}_{\gamma} + \lambda \|v\|^2 \Rightarrow \lambda = \frac{1}{\|v\|^2}$$

# SVM: the margin

We have obtained:

$$\begin{cases} x - \bar{x} = \lambda v \\ \lambda = \frac{1}{\|v\|^2} \end{cases}$$

$\Downarrow$

$$\|x - \bar{x}\| = |\lambda| \|v\| = \lambda \|v\| = \frac{1}{\|v\|^2} \|v\| = \frac{1}{\|v\|}.$$

$\Downarrow$

$$\text{MARGIN} = \frac{2}{\|v\|}$$

$\Downarrow$

$$\max \text{MARGIN} \Leftrightarrow \min_v \|v\| \Leftrightarrow \min_v \frac{1}{2} \|v\|^2.$$



# Support Vector Machine (SVM) (Vapnik, 1995 [Vap95])

To summarize:

- The sets

$$\mathcal{A} = \{a_1, \dots, a_m\}, \quad \text{with } a_i \in \mathbb{R}^n, \quad i = 1, \dots, m$$

and

$$\mathcal{B} = \{b_1, \dots, b_k\}, \quad \text{with } b_l \in \mathbb{R}^n, \quad l = 1, \dots, k$$

are given.

- We compute a separating hyperplane

$$H(v, \gamma) \triangleq \{x \in \mathbb{R}^n \mid v^T x = \gamma\}, \quad \text{with } v \in \mathbb{R}^n \text{ and } \gamma \in \mathbb{R},$$

called the **support vector machine**, which is furthest from the closest points in the two sets.

# Support Vector Machine (SVM)

The separation hyperplane (the **support vector machine**) is constructed by minimizing the following **nonsmooth** error function:

$$f(v, \gamma) \triangleq \frac{1}{2} \|v\|^2 + C \sum_{i=1}^m \max\{0, -v^T a_i + \gamma + 1\} + \\ + C \sum_{l=1}^k \max\{0, v^T b_l - \gamma + 1\}.$$

- The first term maximizes the margin.
- By minimizing the last two terms we minimize the misclassification measure of the points of the two sets  $\mathcal{A}$  and  $\mathcal{B}$ , respectively.
- Parameter  $C > 0$  tunes the weight of the two objectives.



# Smoothing...

$$f(v, \gamma) \triangleq \frac{1}{2} \|v\|^2 + C \sum_{i=1}^m \overbrace{\max\{0, -v^T a_i + \gamma + 1\}}^{\xi_i} + C \sum_{l=1}^k \overbrace{\max\{0, v^T b_l - \gamma + 1\}}^{\psi_l}$$

Minimization of  $f$  corresponds to solve the following **quadratic program**:

$$\left\{ \begin{array}{ll} \min_{v, \gamma, \xi, \psi} & \frac{1}{2} \|v\|^2 + C \sum_{i=1}^m \xi_i + C \sum_{l=1}^k \psi_l \\ & \xi_i \geq -v^T a_i + \gamma + 1 & i = 1, \dots, m \\ & \psi_l \geq v^T b_l - \gamma + 1 & l = 1, \dots, k \\ & \xi_i \geq 0 & i = 1, \dots, m \\ & \psi_l \geq 0 & l = 1, \dots, k. \end{array} \right.$$

# The SVM Wolfe dual

## PRIMAL

$$P \left\{ \begin{array}{ll} \min_{v, \gamma, \xi, \psi} & \frac{1}{2} \|v\|^2 + C \sum_{i=1}^m \xi_i + C \sum_{l=1}^k \psi_l \\ & \xi_i \geq -v^T a_i + \gamma + 1 & i = 1, \dots, m \\ & \psi_l \geq v^T b_l - \gamma + 1 & l = 1, \dots, k \\ & \xi_i \geq 0 & i = 1, \dots, m \\ & \psi_l \geq 0 & l = 1, \dots, k \end{array} \right.$$

# The Wolfe dual

The **Wolfe** dual is defined as follows:

$$D \left\{ \begin{array}{l} \max_{x, \lambda} \quad L(x, \lambda) \\ \nabla_x L(x, \lambda) = 0 \\ \lambda_i \geq 0, \quad i \in I \end{array} \right.$$

# The SVM Wolfe dual

## PRIMAL

$$P \left\{ \begin{array}{lll} \min_{v, \gamma, \xi, \psi} & \frac{1}{2} \|v\|^2 + C \sum_{i=1}^m \xi_i + C \sum_{l=1}^k \psi_l & \\ \xi_i \geq -v^T a_i + \gamma + 1 & i = 1, \dots, m & \lambda_i \\ \psi_l \geq v^T b_l - \gamma + 1 & l = 1, \dots, k & \mu_l \\ \xi_i \geq 0 & i = 1, \dots, m & \alpha_i \\ \psi_l \geq 0 & l = 1, \dots, k & \beta_l \end{array} \right.$$

# The SVM Wolfe dual

Objective function (max)

$$\begin{aligned}
 L(v, \xi, \psi, \lambda, \mu, \alpha, \beta) = & \frac{1}{2} \|v\|^2 + C \sum_{i=1}^m \xi_i + C \sum_{l=1}^k \psi_l \\
 & - \sum_{i=1}^m \lambda_i (\xi_i + v^T a_i - \gamma - 1) \\
 & - \sum_{l=1}^k \mu_l (\psi_l - v^T b_l + \gamma - 1) \\
 & - \sum_{i=1}^m \alpha_i \xi_i - \sum_{l=1}^k \beta_l \psi_l
 \end{aligned}$$

# The SVM Wolfe dual

## Objective function (max)

$$\begin{aligned}
 L(v, \xi, \psi, \lambda, \mu, \alpha, \beta) &= \frac{1}{2} \|v\|^2 - v^T \left( \sum_{i=1}^m \lambda_i a_i - \sum_{l=1}^k \mu_l b_l \right) \\
 &+ \gamma \left( \sum_{i=1}^m \lambda_i - \sum_{l=1}^k \mu_l \right) \\
 &+ \sum_{i=1}^m \xi_i (C - \lambda_i - \alpha_i) + \sum_{l=1}^k \psi_l (C - \mu_l - \beta_l) \\
 &+ \sum_{i=1}^m \lambda_i + \sum_{l=1}^k \mu_l
 \end{aligned}$$

# The SVM Wolfe dual

## Constraints

$$\nabla L_v = v + \sum_{l=1}^k \mu_l b_l - \sum_{i=1}^m \lambda_i a_i = 0 \Leftrightarrow v = \sum_{i=1}^m \lambda_i a_i - \sum_{l=1}^k \mu_l b_l$$

$$\nabla L_\gamma = \sum_{i=1}^m \lambda_i - \sum_{l=1}^k \mu_l = 0$$

$$\nabla L_{\xi_i} = C - \lambda_i - \alpha_i = 0 \Leftrightarrow \lambda_i = C - \alpha_i \Leftrightarrow \lambda_i \leq C \quad i = 1, \dots, m$$

$$\nabla L_{\psi_l} = C - \mu_l - \beta_l = 0 \Leftrightarrow \mu_l = C - \beta_l \Leftrightarrow \mu_l \leq C \quad l = 1, \dots, k$$

$$\lambda, \mu, \alpha, \beta \geq 0$$

# The SVM Wolfe dual

Objective function (max)

$$\begin{aligned}
 L(v, \xi, \psi, \lambda, \mu, \alpha, \beta) = & \frac{1}{2} \|v\|^2 - v^T \overbrace{\left( \sum_{i=1}^m \lambda_i a_i - \sum_{l=1}^k \mu_l b_l \right)}^v \\
 & + \gamma \overbrace{\left( \sum_{i=1}^m \lambda_i - \sum_{l=1}^k \mu_l \right)}^0 \\
 & + \sum_{i=1}^m \xi_i \overbrace{(C - \lambda_i - \alpha_i)}^0 + \sum_{l=1}^k \psi_l \overbrace{(C - \mu_l - \beta_l)}^0 \\
 & + \sum_{i=1}^m \lambda_i + \sum_{l=1}^k \mu_l
 \end{aligned}$$



# The SVM Wolfe dual

Objective function (max)

$$L(v, \lambda, \mu) = \frac{1}{2} \|v\|^2 - \overbrace{v^T v}^{\|v\|^2} + \sum_{i=1}^m \lambda_i + \sum_{l=1}^k \mu_l = -\frac{1}{2} \|v\|^2 + \sum_{i=1}^m \lambda_i + \sum_{l=1}^k \mu_l$$

Constraints

$$v = \sum_{i=1}^m \lambda_i a_i - \sum_{l=1}^k \mu_l b_l$$

$$\sum_{i=1}^m \lambda_i - \sum_{l=1}^k \mu_l = 0$$

$$0 \leq \lambda_i \leq C \quad i = 1, \dots, m$$

$$0 \leq \mu_l \leq C \quad l = 1, \dots, k$$

# The SVM Wolfe dual

$$D \left\{ \begin{array}{l} \max_{\lambda, \mu} \quad -\frac{1}{2} \left\| \sum_{i=1}^m \lambda_i a_i - \sum_{l=1}^k \mu_l b_l \right\|^2 + \sum_{i=1}^m \lambda_i + \sum_{l=1}^k \mu_l \\ \sum_{i=1}^m \lambda_i - \sum_{l=1}^k \mu_l = 0 \\ 0 \leq \lambda_i \leq C \quad i = 1, \dots, m \\ 0 \leq \mu_l \leq C \quad l = 1, \dots, k \end{array} \right.$$

# The SVM Wolfe dual

$$\max \overbrace{\quad}^{\quad} \Rightarrow \min$$

$$D \left\{ \begin{array}{l} \min_{\lambda, \mu} \quad \frac{1}{2} \left\| \sum_{i=1}^m \lambda_i a_i - \sum_{l=1}^k \mu_l b_l \right\|^2 - \sum_{i=1}^m \lambda_i - \sum_{l=1}^k \mu_l \\ \sum_{i=1}^m \lambda_i - \sum_{l=1}^k \mu_l = 0 \\ 0 \leq \lambda_i \leq C \quad i = 1, \dots, m \\ 0 \leq \mu_l \leq C \quad l = 1, \dots, k \end{array} \right.$$

**NOTE:** Quadratic program with one constraint and  $m + k$  box constraints.

# The SVM Wolfe dual

$$(\lambda^*, \mu^*)$$



$$v^* = \sum_{i=1}^m \lambda_i^* a_i - \sum_{l=1}^k \mu_l^* b_l$$

$$\gamma^* = v^{*T} a_i - 1, \text{ with } i \text{ such that } 0 < \lambda_i^* < C$$

or

$$\gamma^* = v^{*T} b_l + 1, \text{ with } l \text{ such that } 0 < \mu_l^* < C$$

## The kernel trick

**Motivation:** To separate  $\mathcal{A}$  and  $\mathcal{B}$  by means of a **nonlinear** surface, using the SVM approach.

- We indicate by  $X_I \subseteq \mathbb{R}^n$  the so-called **input space**, such that  $\mathcal{A}, \mathcal{B} \subset X_I$ .
- We define the so-called **feature space**  $X_F \subseteq \mathbb{R}^N$ , with generally  $N > n$ .
- Given a map

$$\phi : X_I \mapsto X_F,$$

the **kernel function** is defined as:

$$K : X_I \times X_I \mapsto \mathbb{R}$$

such that

$$K(x_1, x_2) = \phi(x_1)^T \phi(x_2).$$

# The kernel trick

## Some kernel functions

- Linear:

$$K(x_1, x_2) = x_1^T x_2$$

- RBF (Radial Basis Function) or Gaussian:

$$K(x_1, x_2) = \exp(-\|x_1 - x_2\|^2 / 2\sigma), \text{ for some value of } \sigma$$

- Hyperbolic tangent:

$$K(x_1, x_2) = \tanh(\beta x_1^T x_2 + \gamma), \text{ for some values of } \beta \text{ and } \gamma.$$

# The kernel trick

## The linear kernel

$$D \left\{ \begin{array}{l} \min_{\lambda, \mu} \quad \frac{1}{2} \left( \sum_{i=1}^m \sum_{j=1}^m \lambda_i \lambda_j \underbrace{a_i^T a_j}_{K(a_i, a_j)} + \sum_{l=1}^k \sum_{j=1}^k \mu_l \mu_j \underbrace{b_l^T b_j}_{K(b_l, b_j)} - 2 \sum_{i=1}^m \sum_{l=1}^k \lambda_i \mu_l \underbrace{a_i^T b_l}_{K(a_i, b_l)} \right) \\ - \sum_{i=1}^m \lambda_i - \sum_{l=1}^k \mu_l \\ \sum_{i=1}^m \lambda_i - \sum_{l=1}^k \mu_l = 0 \\ 0 \leq \lambda_i \leq C \quad i = 1, \dots, m \\ 0 \leq \mu_l \leq C \quad l = 1, \dots, k \end{array} \right.$$

# The kernel trick

## The general case

$$D \left\{ \begin{array}{l} \min_{\lambda, \mu} \frac{1}{2} \left( \sum_{i=1}^m \sum_{j=1}^m \lambda_i \lambda_j \overbrace{\phi(a_i)^T \phi(a_j)}^{K(a_i, a_j)} + \sum_{l=1}^k \sum_{j=1}^k \mu_l \mu_j \overbrace{\phi(b_l)^T \phi(b_j)}^{K(b_l, b_j)} \right. \\ \left. - 2 \sum_{i=1}^m \sum_{l=1}^k \lambda_i \mu_l \overbrace{\phi(a_i)^T \phi(b_l)}^{K(a_i, b_l)} \right) - \sum_{i=1}^m \lambda_i - \sum_{l=1}^k \mu_l \\ \\ \sum_{i=1}^m \lambda_i - \sum_{l=1}^k \mu_l = 0 \\ 0 \leq \lambda_i \leq C \quad i = 1, \dots, m \\ 0 \leq \mu_l \leq C \quad l = 1, \dots, k \end{array} \right.$$

**NOTE:** There is **no need to know explicitly** the map  $\phi$ .



# The kernel trick

$$(\lambda^*, \mu^*)$$



$$v^* = \sum_{i=1}^m \lambda_i^* \phi(a_i) - \sum_{l=1}^k \mu_l^* \phi(b_l) \quad \text{not needed explicitly}$$

$$\gamma^* = v^{*T} \phi(a_i) - 1, \text{ with } i \text{ such that } 0 < \lambda_i^* < C$$

or

$$\gamma^* = v^{*T} \phi(b_l) + 1, \text{ with } l \text{ such that } 0 < \mu_l^* < C$$

**NOTE:** Substituting  $v^*$  in the expression of  $\gamma^*$ ,  $\gamma^*$  is expressed in terms of the kernel function  $K$ .

# The kernel trick

## The decision function

In correspondence to any new point  $\bar{x}$ , we compute:

$$\begin{aligned}
 \underbrace{v^{*T} \phi(\bar{x})}_{\text{linear in } X_F} - \gamma^* &= \left( \sum_{i=1}^m \lambda_i^* \phi(a_i) - \sum_{l=1}^k \mu_l^* \phi(b_l) \right)^T \phi(\bar{x}) - \gamma^* \\
 &= \sum_{i=1}^m \lambda_i^* \phi(a_i)^T \phi(\bar{x}) - \sum_{l=1}^k \mu_l^* \phi(b_l)^T \phi(\bar{x}) - \gamma^* \\
 &= \underbrace{\sum_{i=1}^m \lambda_i^* K(a_i, \bar{x}) - \sum_{l=1}^k \mu_l^* K(b_l, \bar{x})}_{\text{nonlinear in } X_I, \text{ if } K \text{ is nonlinear}} - \gamma^*.
 \end{aligned}$$

# Fixed-center spherical separation with kernel

# Fixed-center spherical separation with kernel (Astorino and Gaudioso, 2009 [AG09])

$$f(x_0, R) \triangleq R^2 + C \sum_{i=1}^m \max\{0, \|a_i - x_0\|^2 - R^2\} + C \sum_{l=1}^k \max\{0, R^2 - \|b_l - x_0\|^2\},$$

**NOTE:** If  $x_0$  is fixed, setting  $z \triangleq R^2 \geq 0$ , then function  $f$  is convex in  $z$ .

# Fixed-center spherical separation with kernel

$$f(z) \triangleq z + C \sum_{i=1}^m \overbrace{\max\{0, \|a_i - x_0\|^2 - z\}}^{\xi_i} + C \sum_{l=1}^k \overbrace{\max\{0, z - \|b_l - x_0\|^2\}}^{\psi_l}.$$

In this case, minimization of  $f$  corresponds to solve the following **linear program**:

$$\left\{ \begin{array}{ll} \min_{z, \xi, \psi} & z + C \sum_{i=1}^m \xi_i + C \sum_{l=1}^k \psi_l \\ & \xi_i \geq \|a_i - x_0\|^2 - z & i = 1, \dots, m \\ & \psi_l \geq z - \|b_l - x_0\|^2 & l = 1, \dots, k \\ & \xi_i \geq 0 & i = 1, \dots, m \\ & \psi_l \geq 0 & l = 1, \dots, k. \\ & z \geq 0 \end{array} \right.$$

# Fixed-center spherical separation with kernel

## NOTE

$$\|a_i - x_0\|^2 = \|a_i\|^2 + \|x_0\|^2 - 2a_i^T x_0,$$

i.e.

$$\|a_i - x_0\|^2 = \underbrace{a_i^T a_i}_{K(a_i, a_i)} + \underbrace{x_0^T x_0}_{K(x_0, x_0)} - 2 \underbrace{a_i^T x_0}_{K(a_i, x_0)}.$$

Moreover:

$$\|b_l - x_0\|^2 = \|b_l\|^2 + \|x_0\|^2 - 2b_l^T x_0,$$

i.e.

$$\|b_l - x_0\|^2 = \underbrace{b_l^T b_l}_{K(b_l, b_l)} + \underbrace{x_0^T x_0}_{K(x_0, x_0)} - 2 \underbrace{b_l^T x_0}_{K(b_l, x_0)}.$$

# Fixed-center spherical separation with kernel

$$f(z) \triangleq z + C \sum_{i=1}^m \overbrace{\max\{0, K(a_i, a_i) + K(x_0, x_0) - 2K(a_i, x_0) - z\}}^{\xi_i} \\ + C \sum_{l=1}^k \underbrace{\max\{0, z - K(b_l, b_l) - K(x_0, x_0) + 2K(b_l, x_0)\}}_{\psi_l}.$$

Minimization of  $f$  corresponds to solve the following **linear program**:

$$\left\{ \begin{array}{ll} \min_{z, \xi, \psi} & z + C \sum_{i=1}^m \xi_i + C \sum_{l=1}^k \psi_l \\ & \xi_i \geq K(a_i, a_i) + K(x_0, x_0) - 2K(a_i, x_0) - z \quad i = 1, \dots, m \\ & \psi_l \geq z - K(b_l, b_l) - K(x_0, x_0) + 2K(b_l, x_0) \quad l = 1, \dots, k \\ & \xi_i \geq 0 \quad i = 1, \dots, m \\ & \psi_l \geq 0 \quad l = 1, \dots, k. \\ & z \geq 0 \end{array} \right.$$

# The kernel trick

## The decision function

In correspondence to any new point  $\bar{x}$ , given  $z^*$ , we compute:

$$\|\phi(\bar{x}) - \phi(x_0)\|^2 = \|\phi(\bar{x})\|^2 + \|\phi(x_0)\|^2 - 2\phi(\bar{x})^T \phi(x_0),$$

i.e.

$$\|\phi(\bar{x}) - \phi(x_0)\|^2 = \underbrace{\phi(\bar{x})^T \phi(\bar{x})}_{K(\bar{x}, \bar{x})} + \underbrace{\phi(x_0)^T \phi(x_0)}_{K(x_0, x_0)} - 2 \underbrace{\phi(\bar{x})^T \phi(x_0)}_{K(\bar{x}, x_0)}.$$

- if  $K(\bar{x}, \bar{x}) + K(x_0, x_0) - 2K(\bar{x}, x_0) \leq z^*$  then  $\bar{x}$  is classified as a point of  $\mathcal{A}$ ;
- if  $K(\bar{x}, \bar{x}) + K(x_0, x_0) - 2K(\bar{x}, x_0) > z^*$  then  $\bar{x}$  is classified as a point of  $\mathcal{B}$ .



# Proximal Support Vector Machine

# Proximal Support Vector Machine (PSVM) (Fung and Mangasarian, 2001 [FM01])

Given  $\mathcal{A}$  and  $\mathcal{B}$ , the standard SVM model is:

$$\left\{ \begin{array}{ll} \min_{v, \gamma, \xi, \psi} & \frac{1}{2} \|v\|^2 + C \sum_{i=1}^m \xi_i + C \sum_{l=1}^k \psi_l \\ & \xi_i \geq -v^T a_i + \gamma + 1 & i = 1, \dots, m \\ & \psi_l \geq v^T b_l - \gamma + 1 & l = 1, \dots, k \\ & \xi_i \geq 0 & i = 1, \dots, m \\ & \psi_l \geq 0 & l = 1, \dots, k. \end{array} \right.$$

$\Downarrow$

The two hyperplanes


$$H_+(v, \gamma) \triangleq \{x \in \mathbb{R}^n | v^T x = \gamma + 1\}, \text{ with } v \in \mathbb{R}^n \text{ and } \gamma \in \mathbb{R},$$

and


$$H_-(v, \gamma) \triangleq \{x \in \mathbb{R}^n | v^T x = \gamma - 1\}, \text{ with } v \in \mathbb{R}^n \text{ and } \gamma \in \mathbb{R}.$$

are the **supporting** hyperplanes.

# SVM example



# SVM example



$H_+$  and  $H_-$  are **supporting** hyperplanes

# Proximal Support Vector Machine (PSVM)

Given the sets  $\mathcal{A}$  and  $\mathcal{B}$ , the standard SVM model is:

$$\left\{ \begin{array}{ll} \min_{v, \gamma, \xi, \psi} & \frac{1}{2} \|v\|^2 + C \sum_{i=1}^m \xi_i + C \sum_{l=1}^k \psi_l \\ & \xi_i \geq -v^T a_i + \gamma + 1 & i = 1, \dots, m \\ & \psi_l \geq v^T b_l - \gamma + 1 & l = 1, \dots, k \\ & \xi_i \geq 0 & i = 1, \dots, m \\ & \psi_l \geq 0 & l = 1, \dots, k. \end{array} \right.$$

i.e.

$$\left\{ \begin{array}{ll} \min_{v, \gamma, \xi, \psi} & \frac{1}{2} \|v\|^2 + C \sum_{i=1}^m |\xi_i| + C \sum_{l=1}^k |\psi_l| \\ & \xi_i \geq -v^T a_i + \gamma + 1 & i = 1, \dots, m \\ & \psi_l \geq v^T b_l - \gamma + 1 & l = 1, \dots, k \\ & \xi_i \geq 0 & i = 1, \dots, m \\ & \psi_l \geq 0 & l = 1, \dots, k. \end{array} \right.$$



# Proximal Support Vector Machine (PSVM)

Instead of

$$\left\{ \begin{array}{ll} \min_{v, \gamma, \xi, \psi} & \frac{1}{2} \|v\|^2 + C \sum_{i=1}^m |\xi_i| + C \sum_{l=1}^k |\psi_l| \\ & \xi_i \geq -v^T a_i + \gamma + 1 & i = 1, \dots, m \\ & \psi_l \geq v^T b_l - \gamma + 1 & l = 1, \dots, k \\ & \xi_i \geq 0 & i = 1, \dots, m \\ & \psi_l \geq 0 & l = 1, \dots, k. \end{array} \right.$$

consider

$$\left\{ \begin{array}{ll} \min_{v, \gamma, \xi, \psi} & \frac{1}{2} \left\| \begin{pmatrix} v \\ \gamma \end{pmatrix} \right\|^2 + \frac{C}{2} \|\xi\|^2 + \frac{C}{2} \|\psi\|^2 \\ & \xi_i \geq -v^T a_i + \gamma + 1 & i = 1, \dots, m \\ & \psi_l \geq v^T b_l - \gamma + 1 & l = 1, \dots, k \end{array} \right.$$

# Proximal Support Vector Machine (PSVM)

Instead of


$$\left\{ \begin{array}{ll} \min_{v, \gamma, \xi, \psi} & \frac{1}{2} \left\| \begin{array}{c} v \\ \gamma \end{array} \right\|^2 + \frac{C}{2} \|\xi\|^2 + \frac{C}{2} \|\psi\|^2 \\ & \xi_i \geq -v^T a_i + \gamma + 1 & i = 1, \dots, m \\ & \psi_l \geq v^T b_l - \gamma + 1 & l = 1, \dots, k \end{array} \right.$$

consider

$$\left\{ \begin{array}{ll} \min_{v, \gamma, \xi, \psi} & \frac{1}{2} \left\| \begin{array}{c} v \\ \gamma \end{array} \right\|^2 + \frac{C}{2} \|\xi\|^2 + \frac{C}{2} \|\psi\|^2 \\ & \xi_i = -v^T a_i + \gamma + 1 & i = 1, \dots, m \\ & \psi_l = v^T b_l - \gamma + 1 & l = 1, \dots, k \end{array} \right.$$


which can be solved **very quickly in a closed form.**

# PSVM example





# PSVM example



$H_+$  and  $H_-$  are proximal hyperplanes

# Spherical separation with margin

# Spherical separation with margin (Astorino et al., 2012 [AFG12])

**Motivation:** To extend the concept of margin to spherical separation.  
We recall that:

The set  $\mathcal{A}$  is **spherically separable** from the set  $\mathcal{B}$  if there exists a sphere

$$S(x_0, R) \triangleq \{x \in \mathbb{R}^n \mid \|x - x_0\|^2 = R^2\},$$

with  $x_0 \in \mathbb{R}^n$  and  $R \in \mathbb{R}$ ,

such that

$$\|a_i - x_0\|^2 \leq R^2, \quad i = 1, \dots, m$$

and

$$\|b_l - x_0\|^2 \geq R^2, \quad l = 1, \dots, k.$$

# Spherical separation with margin

The set  $\mathcal{A}$  is **strictly spherically separable** from the set  $\mathcal{B}$  if there exists a sphere

$$S(x_0, R) \triangleq \{x \in \mathbb{R}^n \mid \|x - x_0\|^2 = R^2\},$$

with  $x_0 \in \mathbb{R}^n$  and  $R \in \mathbb{R}$ ,

such that


$$\|a_i - x_0\|^2 \leq (R - M)^2, \quad i = 1, \dots, m$$

and


$$\|b_l - x_0\|^2 \geq (R + M)^2, \quad l = 1, \dots, k$$

for some  $M$  with  $0 < M \leq R$ .


# Spherical separation with margin




# Spherical separation with margin



# Spherical separation with margin



# Spherical separation with margin



Margin =  $2M$ .



# Spherical separation with margin

## Error function

- A point  $a_i \in \mathcal{A}$  is correctly classified if

$$\|a_i - x_0\|^2 \leq (R - M)^2.$$

- As a consequence, a point  $a_i \in \mathcal{A}$  is misclassified if

$$\|a_i - x_0\|^2 - (R - M)^2 > 0.$$

- Then the classification error, in correspondence to a point  $a_i \in \mathcal{A}$ , is

$$\max\{0, \|a_i - x_0\|^2 - (R - M)^2\}.$$

# Spherical separation with margin

## Error function

- A point  $b_l \in \mathcal{B}$  is correctly classified if

$$\|b_l - x_0\|^2 \geq (R + M)^2.$$

- As a consequence, a point  $b_l \in \mathcal{B}$  is misclassified if

$$(R + M)^2 - \|b_l - x_0\|^2 > 0.$$

- Then the classification error, in correspondence to a point  $b_l \in \mathcal{B}$ , is

$$\max\{0, (R + M)^2 - \|b_l - x_0\|^2\}.$$

# Spherical separation with margin

## Error function

$$e(x_0, R, M) \triangleq \sum_{i=1}^m \max \{0, \|a_i - x_0\|^2 - (R - M)^2\} \\ + \sum_{l=1}^k \max \{0, (R + M)^2 - \|b_l - x_0\|^2\}.$$

Setting  $z \triangleq R^2 + M^2 \geq 0$  and  $q \triangleq 2RM \geq 0$ , we have:

$$e(x_0, z, q) = \sum_{i=1}^m \max \{0, q - z + \|a_i - x_0\|^2\} \\ + \sum_{l=1}^k \max \{0, q + z - \|b_l - x_0\|^2\}$$

} nonsmooth and nonconvex

# Spherical separation with margin

Then we solve the following nonsmooth nonconvex optimization problem:

$$P \left\{ \begin{array}{ll} \min_{x_0, z, q} & f(x_0, z, q) \\ & 0 \leq q \leq z, \end{array} \right.$$

where

$$f(x_0, z, q) \triangleq Ce(x_0, z, q) - q$$

with  $C > 0$ .

**NOTE 1:** Minimizing  $-q$  corresponds to maximize the margin.

**NOTE 2:** The parameter  $C > 0$  tunes the weight of the two objectives.

# Spherical separation with margin: fixing the center

**NOTE:** If  $x_0$  is fixed, function  $f$  is convex in  $z$  and  $q$ .

$$f(z, q) \triangleq -q + C \sum_{i=1}^m \overbrace{\max\{0, q - z + \|a_i - x_0\|^2\}}^{\xi_i} + C \sum_{l=1}^k \overbrace{\max\{0, q + z - \|b_l - x_0\|^2\}}^{\psi_l}.$$

In this case, minimization of  $f$  corresponds to solve the following linear program:

$$\left\{ \begin{array}{ll} \min_{z, q, \xi, \psi} & -q + C \sum_{i=1}^m \xi_i + C \sum_{l=1}^k \psi_l \\ & \xi_i \geq q - z + \|a_i - x_0\|^2 & i = 1, \dots, m \\ & \psi_l \geq q + z - \|b_l - x_0\|^2 & l = 1, \dots, k \\ & \xi_i \geq 0 & i = 1, \dots, m \\ & \psi_l \geq 0 & l = 1, \dots, k. \\ & 0 \leq q \leq z. \end{array} \right.$$

# PART VI

## UNSUPERVISED CLASSIFICATION

# Unsupervised classification

**Unsupervised classification:** we have only **unlabelled objects**, that we would like to cluster on the basis of their similarities.

# The clustering problem

- A set

$$\mathcal{X} = \{x_1, \dots, x_q\}$$

of  $q$  unlabelled points is given.

- The objective is to group the points into  $h$  clusters, with  $h \leq q$ , on the basis of their similarities.
- Criterion: for each cluster  $j$ ,  $j = 1 \dots, h$ , compute the center  $x_{0j}$  of the cluster such that each point  $x_p$ ,  $p = 1, \dots, q$ , is assigned to the cluster with the closest center.



# A mixed integer model

# A mixed integer model (Bagirov and Yearwood, 2006 [BY06])

A constrained optimization model is:

$$\left\{ \begin{array}{l} \min_{x_0, w} \quad \frac{1}{q} \sum_{p=1}^q \sum_{j=1}^h w_{pj} \|x_p - x_{0_j}\|^2 \\ \sum_{j=1}^h w_{pj} = 1 \quad p = 1, \dots, q \\ w_{pj} \in \{0, 1\} \quad p = 1, \dots, q \quad j = 1, \dots, h \end{array} \right.$$

where  $x_{0_j}$  is the center of the cluster  $j$ , for  $j = 1, \dots, h$  and

$$w_{pj} = \begin{cases} 1 & \text{if the point } x_p \text{ is assigned to cluster } j \\ 0 & \text{otherwise} \end{cases}$$

**NOTE:** It is a mixed integer nonlinear nonconvex program.

# A mixed integer model

## KKT conditions

$$L(x_0, w, \lambda) = \frac{1}{q} \sum_{p=1}^q \sum_{j=1}^h w_{pj} \|x_p - x_{0j}\|^2 - \sum_{p=1}^q \lambda_p \left( \sum_{j=1}^h w_{pj} - 1 \right)$$

$$\nabla L_{x_{0j}} = \frac{1}{q} \sum_{p=1}^q 2w_{pj}(x_{0j} - x_p) = 0, \quad j = 1 \dots, h$$

$$\sum_{p=1}^q w_{pj} x_{0j} = \sum_{p=1}^q w_{pj} x_p, \quad j = 1 \dots, h$$

$$\text{barycenter} \rightarrow x_{0j} = \frac{\sum_{p=1}^q w_{pj} x_p}{\sum_{p=1}^q w_{pj}}, \quad j = 1 \dots, h.$$

# An integer model

The model becomes an integer program:

$$\left\{ \begin{array}{l} \min_w \quad \frac{1}{q} \sum_{p=1}^q \sum_{j=1}^h w_{pj} \|x_p - \sum_{r=1}^q w_{rj} x_r / \sum_{r=1}^q w_{rj}\|^2 \\ \sum_{j=1}^h w_{pj} = 1 \quad p = 1, \dots, q \\ w_{pj} \in \{0, 1\} \quad p = 1, \dots, q \quad j = 1 \dots, h \end{array} \right.$$

where

$$w_{pj} = \begin{cases} 1 & \text{if the point } x_p \text{ is assigned to cluster } j \\ 0 & \text{otherwise} \end{cases}$$

# A nonsmooth model

# A nonsmooth model (Bagirov and Yearwood, 2006 [BY06])

An **unconstrained** optimization **model** is:

$$\min_{x_0} \frac{1}{q} \sum_{p=1}^q \min_{1 \leq j \leq h} \|x_p - x_{0_j}\|^2,$$


where  $x_{0_j}$  is the center of the cluster  $j$ , for  $j = 1, \dots, h$ .

**NOTE:** If  $h > 1$ , the objective function is nonconvex and nonsmooth.

# Unsupervised classification

**Example:**  $f(x) = \min_{1 \leq j \leq 4} f_j(x)$ , with  $f_j(x)$  convex, for  $j = 1, \dots, 4$ .


# Unsupervised classification



**Example:**  $f(x) = \min_{1 \leq j \leq 4} f_j(x)$ , with  $f_j(x)$  convex, for  $j = 1, \dots, 4$ .




# Unsupervised classification




**Example:**  $f(x) = \min_{1 \leq j \leq 4} f_j(x)$ , with  $f_j(x)$  convex, for  $j = 1, \dots, 4$ .

# Unsupervised classification




**Example:**  $f(x) = \min_{1 \leq j \leq 4} f_j(x)$ , with  $f_j(x)$  convex, for  $j = 1, \dots, 4$ .

# Unsupervised classification



**Example:**  $f(x) = \min_{1 \leq j \leq 4} f_j(x)$ , with  $f_j(x)$  convex, for  $j = 1, \dots, 4$ .

# Unsupervised classification



**Example:**  $f(x) = \min_{1 \leq j \leq 4} f_j(x)$ , with  $f_j(x)$  convex, for  $j = 1, \dots, 4$ .

## PART VII

# BINARY SEMISUPERVISED CLASSIFICATION

# Semisupervised classification

**Semisupervised classification:** on the basis of the **labelled and unlabelled objects**, we would like to predict the class of the unlabelled objects.

# Transductive Support Vector Machine

# Transductive Support Vector Machine (TSVM)

## (Chapelle and Zien, 2005 [CZ05])

- The TSVM (Transductive Support Vector Machine) technique is the **semisupervised version of the SVM** approach.
- We compute the best support vector machine, on the basis of the **labelled points** (i.e. the sets  $\mathcal{A}$  and  $\mathcal{B}$ ) and some **unlabelled points**.
- The objective is to classify the unlabelled points.



# Transductive Support Vector Machine (TSVM)

- The sets

$$\mathcal{A} = \{a_1, \dots, a_m\}, \quad \text{with } a_i \in \mathbb{R}^n, \quad i = 1, \dots, m$$

and

$$\mathcal{B} = \{b_1, \dots, b_k\}, \quad \text{with } b_l \in \mathbb{R}^n, \quad l = 1, \dots, k$$

are given.

- Another set


$$\mathcal{X} = \{x_1, \dots, x_q\}$$

of  $q$  **unlabelled** points is given.


- The objective is to obtain the best SVM having **as few unlabelled points as possible in the margin**.
- **NOTE:** Number  $q$  in the practical cases is very large.




# SVM: an example




# TSVM: an example




# TSVM: an example



# TSVM: an example



# TSVM: an example



# TSVM: the error function

**Question:** How to minimize the number of unlabelled points in the margin?

# TSVM: the error function

The **margin** is the area between the two supporting hyperplanes

$$H_+(v, \gamma) \triangleq \{x \in \mathbb{R}^n | v^T x = \gamma + 1\}, \text{ with } v \in \mathbb{R}^n \text{ and } \gamma \in \mathbb{R},$$

and

$$H_-(v, \gamma) \triangleq \{x \in \mathbb{R}^n | v^T x = \gamma - 1\}, \text{ with } v \in \mathbb{R}^n \text{ and } \gamma \in \mathbb{R}.$$

Then, a point  $x \in \mathcal{X}$  belongs to the margin if

$$v^T x < \gamma + 1 \text{ and } v^T x > \gamma - 1,$$

i.e. if

$$-1 < v^T x - \gamma < 1,$$

i.e. if

$$|v^T x - \gamma| < 1,$$

i.e. if

$$1 - |v^T x - \gamma| > 0.$$



# TSVM: the error function

We want to find a separating hyperplane  $H(v, \gamma)$ , with  $v \in \mathbb{R}^n$  and  $\gamma \in \mathbb{R}$ , by minimizing the following function:

$$f(v, \gamma) \triangleq \frac{1}{2} \|v\|^2 + C_1 \left[ \sum_{i=1}^m \max\{0, -v^T a_i + \gamma + 1\} + \sum_{l=1}^k \max\{0, v^T b_l - \gamma + 1\} \right] + C_2 \sum_{p=1}^q \max\{0, 1 - |v^T x_p - \gamma|\}.$$

- $f$  is nonsmooth;
- $f$  is nonconvex, due to the last term involving the unlabelled points;
- $C_1, C_2 > 0$  tune the weights of the three objectives (generally  $C_2 \leq C_1$ ).

# Semisupervised polyhedral separation

# Semisupervised polyhedral separation (Astorino and Fuduli, 2015 [AF15b])

- The sets

$$\mathcal{A} = \{a_1, \dots, a_m\}, \quad \text{with } a_i \in \mathbb{R}^n, \quad i = 1, \dots, m$$

and

$$\mathcal{B} = \{b_1, \dots, b_k\}, \quad \text{with } b_l \in \mathbb{R}^n, \quad l = 1, \dots, k$$

are given.

- Another set

$$\mathcal{X} = \{x_1, \dots, x_q\}$$

of  $q$  **unlabelled** points is given.


- The objective is to obtain the best polyhedral separation having **as few unlabelled points as possible in the margin**.

# Semisupervised polyhedral separation


In the standard polyhedral separation we minimize the following function:

$$f(v_1, \dots, v_h; \gamma_1, \dots, \gamma_h) \triangleq \frac{1}{m} \sum_{i=1}^m \max_{1 \leq j \leq h} \{0, v_j^T a_i - \gamma_j + 1\} + \frac{1}{k} \sum_{l=1}^k \max\{0, \min_{1 \leq j \leq h} -v_j^T b_l + \gamma_j + 1\}.$$

# Semisupervised polyhedral separation



# Semisupervised polyhedral separation



# Semisupervised polyhedral separation

Then, combining the TSVM approach and the polyhedral separation, we obtain the following error function:

$$\begin{aligned}
 f(v_1, \dots, v_h; \gamma_1, \dots, \gamma_h) &\triangleq \frac{1}{2} \sum_{j=1}^h \|v_j\|^2 + C_1 \sum_{i=1}^m \max_{1 \leq j \leq h} \{0, v_j^T a_i - \gamma_j + 1\} + \\
 &+ C_1 \sum_{l=1}^k \max\{0, \min_{1 \leq j \leq h} -v_j^T b_l + \gamma_j + 1\} \\
 &+ C_2 \sum_{j=1}^h \sum_{p=1}^q \max\{0, 1 - |v_j^T x_p - \gamma_j|\}.
 \end{aligned}$$

- $f$  is nonsmooth and nonconvex;
- $C_1, C_2 > 0$  tune the weights of the three objectives (generally  $C_2 \leq C_1$ ).

# Semisupervised spherical separation



# Semisupervised spherical separation (Astorino and Fuduli, 2015 [AF15a])

- In the semisupervised spherical separation approach, we compute a separating sphere, on the basis of the **labelled points** (i.e. the sets  $\mathcal{A}$  and  $\mathcal{B}$ ) and some **unlabelled points**.
- The objective is to classify the unlabelled points.

# Semisupervised spherical separation

- The sets

$$\mathcal{A} = \{a_1, \dots, a_m\}, \quad \text{with } a_i \in \mathbb{R}^n, \quad i = 1, \dots, m$$

and

$$\mathcal{B} = \{b_1, \dots, b_k\}, \quad \text{with } b_l \in \mathbb{R}^n, \quad l = 1, \dots, k$$

are given.


- Another set

$$\mathcal{X} = \{x_1, \dots, x_q\}$$


of  $q$  **unlabelled** points is given.

- The objective is to obtain a separating sphere having **as few unlabelled points as possible in the margin**.

# Semisupervised spherical separation: an example



# Semisupervised spherical separation: an example



# The error function

**Question:** How to minimize the number of unlabelled points in the margin?

# The error function

A point  $x \in \mathcal{X}$  belongs to the margin if

$$\|x - x_0\|^2 < (R + M)^2 \text{ and } \|x - x_0\|^2 > (R - M)^2,$$

i.e. if

$$(R + M)^2 - \|x - x_0\|^2 > 0 \text{ and } \|x - x_0\|^2 - (R - M)^2 > 0,$$

i.e. if

$$\min\{(R + M)^2 - \|x - x_0\|^2, \|x - x_0\|^2 - (R - M)^2\} > 0.$$

Setting  $z \triangleq R^2 + M^2 \geq 0$  and  $q \triangleq 2RM \geq 0$ , we have that  $x$  belongs to the margin if:

$$\min\{q + z - \|x - x_0\|^2, \|x - x_0\|^2 - z + q\} > 0. \quad \text{UNIVERSITÀ DELLA CALABRIA}$$

DIPARTIMENTO DI MATEMATICA  
E INFORMATICA

# The error function

Then we minimize the following function:

$$\begin{aligned}
 f(x_0, z, q) = & -q \\
 & + C_1 \sum_{i=1}^m \max \{0, q - z + \|a_i - x_0\|^2\} \\
 & + C_1 \sum_{l=1}^k \max \{0, q + z - \|b_l - x_0\|^2\} \\
 & + C_2 \sum_{p=1}^q \max \{0, \min[q + z - \|x_p - x_0\|^2, \|x_p - x_0\|^2 - z + q]\}
 \end{aligned}$$

such that  $0 \leq q \leq z$ .

- $f$  is nonsmooth and nonconvex;
- $C_1, C_2 > 0$  tune the weights of the three objectives (generally  $C_2 \leq C_1$ ).

# PART VIII

## MULTIPLE INSTANCE LEARNING



# Introduction to Multiple Instance Learning

# Multiple instance learning (MIL)

- **Supervised learning**: the objective is to categorize points into different classes, on the basis of labelled points.
- **Multiple instance learning (MIL)**: the objective is to classify **bags** of points, each point being an **instance**.
- **NOTE**: In the learning phase of a MIL approach, **we know** the label of each bag, but the label of each instance inside the bags is **unknown**.

# The first MIL problem (Dietterich et al., 1997 [DLLP97])

- Drug design problem: we want to discriminate between **active** and **non-active** drug molecules;
- a drug molecule is **active** if it is able to bind to a particular target site (typically a larger protein molecule);
- each molecule can assume different conformations;
- ...but indeed **it is not known** which conformation makes a molecule active;
- in the MIL perspective, each molecule is a **bag** and the conformations of the molecules are the **instances**.

# MIL: the binary case

- **Binary case:** we would like to discriminate between two classes of bags (positive and negative) and to predict the class label of new bags.
- **NOTE:** Even in the binary case, we can have more than two classes of instances.

# Example n. 1

- We have some images and we would like to discriminate between **beach** and **non-beach**;
- each image is a **bag** containing some “subregions” (**instances**):  
sea, countryside, cities, cars, offices, sky, sand, trees, mountains, etc.;
- an image is **positive** (i.e. a beach) if it contains both sea and sand;
- an image is **negative** if it does not contain both sea and sand.

## Example n. 2

- Objective: to discriminate between **non-healthy** and **healthy** patients on the basis of their medical scan (**bag**);
- a patient is **positive** if he/she presents at least an abnormal subregion (**instance**) in his/her medical scan;
- a patient is **negative** if all the subregions (**instances**) in his/her medical scan are healthy.

# Multiple instance learning (MIL)

**NOTE:** In both previous examples, **only some portions** of the image (or medical scan) make the image positive.



The MIL approach can be interpreted as a **weakly supervised** approach.

**NOTE:** In the binary case, a crucial issue is to specify **what a positive bag is**.

# Possible applications of MIL

- Classification of images;
- drug discovery;
- classification of text documents;
- bankruptcy prediction;
- speaker identification.



## Classification of the MIL approaches

# The binary case: bag-space learning

We have two classes of bags: positive and negative.

- In the bag-space learning we separate directly the positive bags from the negative ones, considering each bag as a whole entity.
- This approach is necessary when there is no class of instances appearing only in positive bags.

# The binary case: instance-space learning

We have two classes of bags: positive and negative.

- In the instance-space learning **we separate the instances** belonging to the positive bags from the instances belonging to the negative ones.
- Then **the class label information of a bag** is obtained as aggregation of the instance-space responses.
- This approach is possible when some classes of instances appear only in positive bags.

# The binary case: embedding-space learning

We have two classes of bags: positive and negative.

- In the embedding-space learning **we map each bag to a single feature vector** (typically the most representative instance belonging to the bag), resulting in a classical supervised classification problem to be solved in the instance space.

# MIL surveys



J. FOULDS AND E. FRANK, [A review of multi-instance learning assumptions](#), Knowledge Engineering Review, 25 (2010), pp. 1–25.



J. AMORES, [Multiple instance classification: Review, taxonomy and comparative study](#), Artificial Intelligence, 201 (2013), pp. 81–105.



M. CARBONNEAU, V. CHEPLYGINA, E. GRANGER AND G. GAGNON, [Multiple instance learning: a survey of problem characteristics and applications](#), Pattern Recognition, 77 (2018), pp. 329 – 353.



G. QUELLEC, G. CAZUGUEL, B. COCHENER AND M. LAMARD, [Multiple-Instance Learning for Medical Image and Video Analysis](#), IEEE Reviews in Biomedical Engineering, 10 (2017), pp. 213–234.



# Binary MIL problem: assumptions


## MIL STANDARD ASSUMPTION

- Two classes of bags: positive and negative;
- two classes of instances: positive and negative.




- A bag is **positive** if it contains at least a positive instance;
- a bag is **negative** if all its instances are negative.

# Standard MIL assumption: an example



- A bag is classified **positive** if at least one of its instances is classified positive.
- A bag is classified **negative** if all its instances are classified negative.


# Standard MIL assumption: an example



- A bag is classified **positive** if at least one of its instances is classified positive.
- A bag is classified **negative** if all its instances are classified negative.



# Standard MIL assumption: an example



- A bag is classified **positive** if at least one of its instances is classified positive.
- A bag is classified **negative** if all its instances are classified negative.

# Support Vector Machine for Multiple Instance Learning

# A MIL SVM model (Andrews et al., 2003 [ATH03])

## NOTATION

- $\mathcal{A}_1, \dots, \mathcal{A}_m$ :  $m$  positive bags;
- $\mathcal{B}_1, \dots, \mathcal{B}_k$ :  $k$  negative bags;
- $J_i^+$ : index set corresponding to  $\mathcal{A}_i$ ,  $i = 1, \dots, m$ ;
- $J_l^-$ : index set corresponding to  $\mathcal{B}_l$ ,  $l = 1, \dots, k$ ;
- $x_j$ : the  $j$ th instance;
- $y_j \in \{1, -1\}$ : the class label of the instance  $x_j$ , when  $x_j$  belongs to a positive bag.



$$H(v, \gamma) \triangleq \{x \in \mathbb{R}^n \mid v^T x = \gamma\}.$$

# A MIL SVM model

Minimize  $f(v, \gamma, \mathbf{y})$ , where

$$f(v, \gamma, \mathbf{y}) \triangleq \frac{1}{2} \|v\|^2 + C \sum_{i=1}^m \sum_{j \in J_i^+} \max\{0, 1 + \mathbf{y}_j (-v^T x_j + \gamma)\} \\ + C \sum_{l=1}^k \sum_{j \in J_l^-} \max\{0, 1 + (v^T x_j - \gamma)\},$$

such that:

$$\sum_{j \in J_i^+} \frac{\mathbf{y}_j + 1}{2} \geq 1, \quad i = 1, \dots, m$$

and

$$\mathbf{y}_j \in \{-1, 1\}, \quad j \in J_i^+, \quad i = \dots, m.$$

# A MIL SVM model

$$\left\{ \begin{array}{l} f^* = \min_{v, \gamma, \mathbf{y}} f(v, \gamma, \mathbf{y}) \\ \sum_{j \in J_i^+} \frac{y_j + 1}{2} \geq 1 \quad i = 1, \dots, m \\ y_j \in \{-1, 1\}, \quad j \in J_i^+, \quad i = 1, \dots, m. \end{array} \right.$$

**NOTE:** Constrained, nonlinear, nonconvex, mixed integer problem.

# A MIL SVM model

$$\text{MIL - SVM} \left\{ \begin{array}{l}
 \min_{v, \gamma, \mathbf{y}, \xi, \psi} \quad \frac{1}{2} \|v\|^2 + C \sum_{i=1}^m \sum_{j \in J_i^+} \xi_j + C \sum_{l=1}^k \sum_{j \in J_l^-} \psi_j \\
 \xi_j \geq 1 + \mathbf{y}_j (-v^T x_j + \gamma) \quad j \in J_i^+, \quad i = 1, \dots, m \\
 \psi_j \geq 1 + (v^T x_j - \gamma) \quad j \in J_l^-, \quad l = 1, \dots, k \\
 \sum_{j \in J_i^+} \frac{\mathbf{y}_j + 1}{2} \geq 1 \quad i = 1, \dots, m \\
 \mathbf{y}_j \in \{-1, +1\} \quad j \in J_i^+, \quad i = 1, \dots, m \\
 \xi_j \geq 0 \quad j \in J_i^+, \quad i = 1, \dots, m \\
 \psi_j \geq 0 \quad j \in J_l^-, \quad l = 1, \dots, k.
 \end{array} \right.$$

# A MIL SVM model: the BCD approach

- BCD = Block Coordinate Descent method.
- Once  $y_j$  is fixed, solve the **SVM** problem to compute  $v$  and  $\gamma$ .
- Once  $v$  and  $\gamma$  are fixed, compute  $y_j$  **by inspection**.

# A MIL SVM model: the BCD approach

## COMPUTING $y_j$ BY INSPECTION

$$z_j \triangleq \max\{0, y_j L_j + 1\}, \text{ where } L_j \triangleq -v^T x_j + \gamma$$

$$\begin{cases} \text{if } L_j > 0 \Rightarrow y_j^* = -1 \\ \text{if } L_j \leq 0 \Rightarrow y_j^* = +1 \end{cases}$$

**NOTE 1:**  $L_j > 0 \Rightarrow -v^T x_j + \gamma > 0 \Rightarrow v^T x_j - \gamma > 0 \Rightarrow v^T x_j < \gamma$

**NOTE 2:**  $L_j \leq 0 \Rightarrow -v^T x_j + \gamma \leq 0 \Rightarrow v^T x_j - \gamma \geq 0 \Rightarrow v^T x_j \geq \gamma$



# A MIL SVM model: the BCD approach

- 1 Set  $\bar{y}_j := +1$ , for any  $j \in J_i^+$ ,  $i = 1, \dots, m$ .
- 2 Solve  $MIL - SVM$  with  $y = \bar{y}$ , to compute  $\bar{v}$  and  $\bar{\gamma}$ .
- 3 If  $\bar{v}^T x_j \geq \bar{\gamma}$  set  $\bar{y}_j := +1$ , else set  $\bar{y}_j := -1$ .
- 4 For any  $i \in \{1, 2, \dots, m\}$  such that

$$\sum_{j \in J_i^+} \frac{\bar{y}_j + 1}{2} = 0,$$

compute  $k_i$  such that

$$\bar{v}^T x_{k_i} - \bar{\gamma} = \max_{j \in J_i^+} \{\bar{v}^T x_j - \bar{\gamma}\}$$

and set  $\bar{y}_{k_i} := +1$ .

- 5 If  $\bar{y}$  has changed go to Step 2, else STOP.

# A MIL SVM model: a Lagrangian relaxation approach

$$LR(\lambda) \begin{cases} z_{LR}^*(\lambda) = \min_{v, \gamma, y} \mathcal{L}(v, \gamma, y, \lambda) \\ y_j \in \{-1, 1\}, \quad j \in J_i^+, \quad i = \dots, m, \end{cases}$$

where

$$\begin{aligned} \mathcal{L}(v, \gamma, y, \lambda) \triangleq & \frac{1}{2} \|v\|^2 + C \sum_{i=1}^m \sum_{j \in J_i^+} \max\{0, 1 + y_j(-v^T x_j + \gamma)\} \\ & + C \sum_{l=1}^k \sum_{j \in J_l^-} \max\{0, 1 + (v^T x_j - \gamma)\} \\ & - \sum_{i=1}^m \lambda_i \left( \sum_{j \in J_i^+} \frac{y_j + 1}{2} - 1 \right). \end{aligned}$$

# A MIL SVM model: a Lagrangian relaxation approach

**BCD** APPROACH FOR SOLVING  $LR(\lambda)$ , when  $\lambda \geq 0$  is fixed

$$\begin{aligned} \mathcal{L}(v, \gamma, y, \lambda) \triangleq & \frac{1}{2} \|v\|^2 + C \sum_{i=1}^m \sum_{j \in J_i^+} \max\{0, 1 + y_j(-v^T x_j + \gamma)\} \\ & + C \sum_{l=1}^k \sum_{j \in J_l^-} \max\{0, 1 + (v^T x_j - \gamma)\} \\ & + \sum_{i=1}^m \lambda_i - \sum_{i=1}^m \sum_{j \in J_i^+} \lambda_i \frac{y_j + 1}{2} \end{aligned}$$

- Once  $y_j$  is fixed, solve the **SVM** problem to compute  $v$  and  $\gamma$ .
- Once  $v$  and  $\gamma$  are fixed, compute  $y_j$  **by inspection**.

# A MIL SVM model: a Lagrangian relaxation approach

## COMPUTING $y_j$ BY INSPECTION

$$z_j \triangleq \max\{0, y_j L_j + 1\} - \lambda_i \frac{y_j + 1}{2},$$

where  $L_j \triangleq -v^T x_j + \gamma$  and  $\lambda_i$  is the Lagrangian multiplier such that  $j \in J_i^+$ .



3 cases, on the basis of the value of  $L_j$ .

# A MIL SVM model: a Lagrangian relaxation approach

## CASE 1 ( $L_j \leq -1$ )

- $y_j = +1 \Rightarrow y_j L_j + 1 = L_j + 1 \leq 0 \Rightarrow z_j = -\lambda_i \leq 0$ .
- $y_j = -1 \Rightarrow y_j L_j + 1 = \underbrace{-L_j + 1}_{\geq 1} \geq 2 \Rightarrow z_j = C \underbrace{(-L_j + 1)}_{\geq 2} \geq 2C > 0$ .



If  $L_j \leq 1$ , then  $y_j^* = +1$ .

# A MIL SVM model: a Lagrangian relaxation approach

## CASE 2 ( $L_j \geq 1$ )

- $y_j = +1 \Rightarrow y_j L_j + 1 = L_j + 1 \geq 2 \Rightarrow z_j = C(L_j + 1) - \lambda_i.$
- $y_j = -1 \Rightarrow y_j L_j + 1 = \underbrace{-L_j + 1}_{\leq -1} \leq 0 \Rightarrow z_j = 0.$

$\Downarrow$

If  $L_j \geq 1$ , then  $\begin{cases} \text{if } C(L_j + 1) - \lambda_i \leq 0, & \text{then } y_j^* = +1 \\ \text{if } C(L_j + 1) - \lambda_i > 0, & \text{then } y_j^* = -1. \end{cases}$

$\Downarrow$

If  $L_j \geq 1$ , then  $\begin{cases} \text{if } \lambda_i \geq C(L_j + 1), & \text{then } y_j^* = +1 \\ \text{if } \lambda_i < C(L_j + 1), & \text{then } y_j^* = -1. \end{cases}$

# A MIL SVM model: a Lagrangian relaxation approach

## CASE 3 ( $-1 < L_j < 1$ )

- $y_j = +1 \Rightarrow y_j L_j + 1 = L_j + 1 > 0 \Rightarrow z_j = C(L_j + 1) - \lambda_i.$
- $y_j = -1 \Rightarrow y_j L_j + 1 = -L_j + 1 > 0 \Rightarrow z_j = C(-L_j + 1).$

$\Downarrow$

If  $-1 < L_j < 1$ , then  $\begin{cases} \text{if } C(L_j + 1) - \lambda_i \leq C(-L_j + 1), & \text{then } y_j^* = +1 \\ \text{if } C(L_j + 1) - \lambda_i > C(-L_j + 1), & \text{then } y_j^* = -1. \end{cases}$

$\Downarrow$

If  $-1 < L_j < 1$ , then  $\begin{cases} \text{if } \lambda_i \geq 2CL_j, & \text{then } y_j^* = +1 \\ \text{if } \lambda_i < 2CL_j, & \text{then } y_j^* = -1. \end{cases}$

# PART IX

## EVALUATION OF A CLASSIFIER



# 10-fold cross-validation

# Evaluation of a classifier

**Question:** How to evaluate the **quality** of a binary classifier?

**Answer:** A possibility is to use a **10-fold cross-validation**, which consists in randomly generating 10 folds, each of them constituted by a **training set** and a **testing set**.


- 1 The **training set**: (90% of the data) is used to construct (**to learn**) the classifier, i.e. the separation surface (such as a hyperplane, a sphere, and so on). It corresponds to the  $m$  positive points of  $\mathcal{A}$  and to the  $k$  negative points of  $\mathcal{B}$ .
- 2 The **testing set** (10% of the data) simulates the unknown data to be classified.

# Evaluation of a classifier

## 10 fold cross-validation (first level)



INITIAL DATASET



RANDOM SPLIT

# Evaluation of a classifier

## 10 fold cross-validation



INITIAL DATASET

1	2	3	4	5	6	7	8	9	10
---	---	---	---	---	---	---	---	---	----

FOLD 1  $\left\{ \begin{array}{ll} \text{testing set:} & 1 \\ \text{training set:} & 2, 3, \dots, 9, 10 \end{array} \right.$

# Evaluation of a classifier

10 fold cross-validation (first level)



INITIAL DATASET

1	2	3	4	5	6	7	8	9	10
---	---	---	---	---	---	---	---	---	----

FOLD 2  $\left\{ \begin{array}{ll} \text{testing set:} & 2 \\ \text{training set:} & 1, 3, \dots, 9, 10 \end{array} \right.$

# Evaluation of a classifier

## 10 fold cross-validation



INITIAL DATASET

1	2	3	4	5	6	7	8	9	10
---	---	---	---	---	---	---	---	---	----

FOLD 3 { testing set: 3  
training set: 1, 2, 4, ..., 9, 10

# Evaluation of a classifier

...and so on...

# Evaluation of a classifier

## 10 fold cross-validation



INITIAL DATASET

1	2	3	4	5	6	7	8	9	10
---	---	---	---	---	---	---	---	---	----

FOLD 10  $\left\{ \begin{array}{ll} \text{testing set:} & 10 \\ \text{training set:} & 1, 2, 3, \dots, 9 \end{array} \right.$



# Performance indicators

# Evaluation of a classifier



For each fold, we compute the **testing correctness**:

$$\frac{\text{\# points correctly classified in the testing set}}{\text{\# total points in the testing set}}$$



**Average testing correctness = accuracy of the classifier.**

**NOTE:** The **average testing correctness** measures the generalization capability of a classifier, i.e. the capability to correctly classify the new data.

# Evaluation of a classifier

For each fold, we can compute also the **training correctness**:

$$\frac{\text{\# points correctly classified in the training set}}{\text{\# number of total points in the training set}}$$



**Average training correctness**: measures the quality of the **optimization process** in the learning phase.

# Evaluation of a classifier

## OTHER INDICATORS (Testing/Training)



$\mathcal{A}$ : set of positive points

$\mathcal{B}$ : set of negative points

$$\text{Sensitivity} = \frac{\# \text{ points of } \mathcal{A} \text{ correctly classified}}{\# \text{ points of } \mathcal{A}}$$

**NOTE:** The **sensitivity** is called also the **true positive rate** or **recall**. It measures the proportion of positive points correctly identified.

# Evaluation of a classifier

## OTHER INDICATORS (Testing/Training)



$\mathcal{A}$ : set of positive points

$\mathcal{B}$ : set of negative points

$$\text{Specificity} = \frac{\# \text{ points of } \mathcal{B} \text{ correctly classified}}{\# \text{ points of } \mathcal{B}}$$

**NOTE:** The **specificity** is called also the **true negative rate**. It measures the proportion of negative points correctly identified.

# Evaluation of a classifier

OTHER INDICATORS (Testing/Training)



$\mathcal{A}$ : set of positive points

$\mathcal{B}$ : set of negative points

$$\begin{aligned}\text{Precision} &= \frac{\# \text{ points of } \mathcal{A} \text{ correctly classified}}{\# \text{ points of } \mathcal{A} \text{ correctly classified} + \# \text{ points of } \mathcal{B} \text{ misclassified}} \\ &= \frac{\# \text{ points of } \mathcal{A} \text{ correctly classified}}{\# \text{ total points classified as positive}}\end{aligned}$$

# Evaluation of a classifier

## OTHER INDICATORS (Testing/Training)



$\mathcal{A}$ : set of positive points

$\mathcal{B}$ : set of negative points

$$\begin{aligned}\text{F-score or F1-Score} &= \frac{2}{\frac{1}{\text{sensitivity}} + \frac{1}{\text{precision}}} \\ &= 2 \frac{\text{sensitivity} \cdot \text{precision}}{\text{sensitivity} + \text{precision}}\end{aligned}$$

# Leave-One-Out



# Leave-One-Out

## Leave-One-Out

Each time, the **testing set** is constituted by a single point. The remaining points of the dataset constitute the **training set**.

# Model selection

# Model selection

**Question:** How to compute the suitable values of the parameters  $C$ ,  $C_1$ ,  $C_2$ ,  $\sigma$ , and so on...?

# Model selection

Simple case: computing  $C$ .

# Model selection - 10 fold Cross Validation

- 1 The data set is randomly split into ten different pieces (**tenfold cross-validation - first level**).
- 2 For ten times, each time, nine pieces form the **first level training set**.
- 3 The tenth piece forms **the first level testing set**, which simulates the new unknown data to be classified.
- 4 Then we have ten training sets and ten corresponding testing sets.
- 5 We fix a grid of possible values for  $C$  (for example 1, 10, 100, 1000).
- 6 For each first level training set, we perform a **fivefold cross-validation - second level**, testing each value of  $C$ .



# Model selection


To compute the best value of  $C$  for the  $i$ -th first level fold, we perform a 5 fold cross-validation (second level) on the first level training set

# Model selection

5 fold cross-validation (second level) on fold  $i$



TRAINING SET (FIRST LEVEL)




SPLIT

# Model selection

## 5 fold cross-validation (second level)



TRAINING SET (FIRST LEVEL)




SPLIT

FOLD 1 (second level)  $\left\{ \begin{array}{ll} \text{testing set (second level):} & 1 \\ \text{training set (second level):} & 2, 3, 4, 5 \end{array} \right.$



# Model selection

## 5 fold cross-validation (second level)



FOLD 2 (second level)  $\left\{ \begin{array}{ll} \text{testing set (second level):} & 2 \\ \text{training set (second level):} & 1, 3, 4, 5 \end{array} \right.$

# Model selection


...and so on...

# Model selection

## 5 fold cross-validation (second level)



TRAINING SET (FIRST LEVEL)



SPLIT

FOLD 5 (second level)  $\left\{ \begin{array}{ll} \text{testing set (second level):} & 5 \\ \text{training set (second level):} & 1, 2, 3, 4 \end{array} \right.$

# Model selection

For any single first level fold:

- 1 For each prefixed value of  $C$  in the grid we come out with a **second level average testing correctness** (average of 5 values).
- 2 Among the values of  $C$  in the grid, we take the best value  $C^*$  such that the second level average testing correctness is minimum.

# PART X

## REFERENCES

# References I



A. Astorino and A. Fuduli, Semisupervised spherical separation, Applied Mathematical Modelling **39** (2015), no. 20, 6351–6358.



———, Support vector machine polyhedral separability in semisupervised learning, Journal of Optimization Theory and Applications **164** (2015), no. 3, 1039–1050.



A. Astorino, A. Fuduli, and M. Gaudioso, Margin maximization in spherical separation, Computational Optimization and Applications **53** (2012), no. 2, 301–322.



A. Astorino, A. Fuduli, P. Veltri, and E. Vocaturo, Melanoma detection by means of multiple instance learning, Interdisciplinary Sciences: Computational Life Sciences **12** (2020), no. 1, 24–31.



# References II



A. Astorino and M. Gaudioso, A fixed-center spherical separation algorithm with kernel transformations for classification problems, Computational Management Science **6** (2009), no. 3, 357–372.



S. Andrews, I. Tsochantaridis, and T. Hofmann, Support vector machines for multiple-instance learning, Advances in Neural Information Processing Systems (S. Becker, S. Thrun, and K. Obermayer, eds.), MIT Press, Cambridge, 2003, pp. 561–568.



P. S. Bradley and O. L. Mangasarian, Feature selection via concave minimization and support vector machines, Machine Learning Proceedings of the Fifteenth International Conference (ICML 98), Morgan Kaufmann, 1998, pp. 82–90.



## References III



A. M. Bagirov and J. Yearwood, A new nonsmooth optimization algorithm for minimum sum-of-squares clustering problems, *European Journal of Operational Research* **170** (2006), no. 2, 578–596.



O. Chapelle and A. Zien, Semi-supervised classification by low density separation, *AISTATS 2005 - Proceedings of the 10th International Workshop on Artificial Intelligence and Statistics*, 2005, pp. 57–64.



T. G. Dietterich, R. H. Lathrop, and T. Lozano-Pérez, Solving the multiple instance problem with axis-parallel rectangles, *Artificial Intelligence* **89** (1997), no. 1-2, 31–71.





# References IV



G. Fung and O. L. Mangasarian, Proximal support vector machine classifiers, Proceedings KDD-2001: Knowledge discovery and data mining (F. Provost and R. Srikant, eds.), ACM, 2001, pp. 77–86.



O. L. Mangasarian, Linear and nonlinear separation of patterns by linear programming, Operations Research **13** (1965), 444–452.



N. Megiddo, On the complexity of polyhedral separability, Discrete and Computational Geometry **3** (1988), 325–337.



S. Maldonado, J. Pérez, R. Weber, and M. Labbé, Feature selection for support vector machines via mixed integer linear programming, Information Sciences **279** (2014), 163–175.



# References V



D. M. J. Tax and R. P. W. Duin, Data domain description using support vectors, ESANN'1999 proceedings Bruges (Belgium), April 1999, pp. 251–256.



V. Vapnik, The nature of the statistical learning theory, Springer Verlag, New York, 1995.



P. Wolfe, A duality theorem for non-linear programming, Quarterly of Applied Mathematics **19** (1961), no. 3, 239–244.

