

UNIVERSITY NAME

DOCTORAL THESIS

Hierarchical deterministic wallet

Author:

Daniele FORNARO

Supervisor:

Daniele MARAZZINA

*A thesis submitted in fulfillment of the requirements
for the degree of Mathematical Engineering*

in the

Research Group Name
Department or School Name

January 10, 2018

Declaration of Authorship

I, Daniele FORNARO, declare that this thesis titled, “Hierarchical deterministic wallet” and the work presented in it are my own. I confirm that:

- This work was done wholly or mainly while in candidature for a research degree at this University.
- Where any part of this thesis has previously been submitted for a degree or any other qualification at this University or any other institution, this has been clearly stated.
- Where I have consulted the published work of others, this is always clearly attributed.
- Where I have quoted from the work of others, the source is always given. With the exception of such quotations, this thesis is entirely my own work.
- I have acknowledged all main sources of help.
- Where the thesis is based on work done by myself jointly with others, I have made clear exactly what was done by others and what I have contributed myself.

Signed:

Date:

“Thanks to my solid academic training, today I can write hundreds of words on virtually any topic without possessing a shred of information, which is how I got a good job in journalism.”

Dave Barry

UNIVERSITY NAME

Abstract

Faculty Name
Department or School Name

Mathematical Engineering

Hierarchical deterministic wallet

by Daniele FORNARO

The Thesis Abstract is written here (and usually kept to just this page). The page is kept centered vertically so can expand into the blank space above the title too...

Acknowledgements

The acknowledgments and the people to thank go here, don't forget to include your project advisor...

Contents

Declaration of Authorship	iii
Abstract	vii
Acknowledgements	ix
1 Elliptic Curve Geometry	1
1.1 Introduction	1
1.2 Elliptic Curve over \mathbb{F}_p	1
1.2.1 Operations	1
Symmetry	1
Point addition	2
Scalar multiplication	3
1.2.2 Group order	3
Cyclic subgroups	3
1.2.3 Bitcoin Elliptic Curve	4
1.3 Bitcoin private-public key cryptography	4
A Frequently Asked Questions	5
A.1 How do I change the colors of links?	5
Bibliography	7

List of Figures

1.1	Points on the Elliptic Curve $y^2 = x^3 - 7x + 10 \pmod{p}$, with $p =$ 19, 97, 127, 487	2
1.2	Elliptic Curve $y^2 = x^3 - 7x + 10 \pmod{97}$	2

List of Tables

List of Abbreviations

LAH List Abbreviations **Here**
WSF What (it) Stands **For**

Physical Constants

Speed of Light $c_0 = 2.997\,924\,58 \times 10^8 \text{ m s}^{-1}$ (exact)

List of Symbols

a	distance	m
P	power	W (J s ⁻¹)
ω	angular frequency	rad

For/Dedicated to/To my...

Chapter 1

Elliptic Curve Geometry

1.1 Introduction

Bitcoin security is based on public and private key cryptography. The main concept is that it is simple to compute the public key, knowing the private, but it is infeasible to calculate the private key, knowing the public.

In order to obtain this result a particular Elliptic Curve is used.

1.2 Elliptic Curve over \mathbb{F}_p

A point Q , which coordinates are x and y , belong to an Elliptic Curve if and only if Q satisfies the following equation:

$$y^2 = x^3 + ax + b \quad \text{over } \mathbb{F}_p \quad (1.1)$$

Where \mathbb{F}_p is the finite field defined over the set of integers modulo p and a and b are the coefficients of the curve.

We can rewrite the equation 1.1 in the following way:

$$y^2 = x^3 + ax + b \quad \text{mod } p \quad (1.2)$$

Figure 1.1 shows some examples of Elliptic Curve over \mathbb{F}_p with $a = -7$ and $b = 10$

1.2.1 Operations

A point on the Elliptic Curve has some particular properties:

- Symmetry
- Point addition
- Scalar multiplication

Symmetry

For every point in the x axis exists two points in the y axis. Suppose that a point $P(x, y)$ belongs to the Elliptic Curve, then it must satisfy the equation 1.1. So it is easy to prove that the point $Q(x, p - y)$ belongs to the curve too.

Furthermore we have $P = -Q$, from the moment that $P + Q = 0$ (see addition below).



FIGURE 1.1: Points on the Elliptic Curve $y^2 = x^3 - 7x + 10 \pmod{p}$, with $p = 19, 97, 127, 487$

Point addition

We need to change our definition of addition in order to make it works in \mathbb{F}_p . In this framework we claim that if three points are aligned over the finite field \mathbb{F}_p , then they have zero sum.

So $P + Q = R$ if and only if P, Q and $-R$ are aligned, in the sense shown in figure 1.2



FIGURE 1.2: Elliptic Curve $y^2 = x^3 - 7x + 10 \pmod{97}$

The equations for calculating point additions are the follow:
Suppose that A and B belong to the Elliptic Curve.

$$A = (x_1, y_1) \quad B = (x_2, y_2)$$

Let's defined $A + B := (x_3, y_3)$

So we have:

$$s = \begin{cases} \frac{y_2 - y_1}{x_2 - x_1}, & \text{if } x_1 \neq x_2 \\ \frac{3x_1^2 + a}{2y_1}, & \text{if } x_1 = x_2 \end{cases}$$

$$\begin{aligned} x_3 &= s^2 - x_1 - x_2 \pmod{p} \\ y_3 &= s(x_1 - x_3) - y_1 \pmod{p} \end{aligned}$$

Scalar multiplication

Once defined the addition, any multiplication can be defined as:

$$nP = \underbrace{P + P + \dots + P}_{n \text{ times}}$$

When n is a very large number can be difficult or even infeasible to compute nP in this way, but we can use the *double and add algorithm* in order to perform multiplication in $\mathcal{O}(\log n)$ steps.

1.2.2 Group order

An elliptic curve defined over a finite field is a group and so it has a finite number of points. This number is called order of the group.

If the prime order is a very large number, it is impossible to count all the point in that field, but there is an algorithm that allows to calculate the order of a group in a fast and efficient way: Schoof's algorithm.

Cyclic subgroups

Let's consider a generic point P , we have:

$$nP + mP = \underbrace{P + \dots + P}_{n \text{ times}} + \underbrace{P + \dots + P}_{m \text{ times}} = \underbrace{P + \dots + P}_{n+m \text{ times}} = (n+m)P$$

So multiple of P are closed under addition and this is enough to prove that the set of the multiples of P is a cyclic subgroup of the group formed by the elliptic curve.

The point P is called generator of the cyclic subgroup. The order of a subgroup is a divisor of the order of the parent group.

Remark The order of P is linked to the order of the elliptic curve by Lagrange's theorem, which states that the order of a subgroup is a divisor of the order of the parent group.

1.2.3 Bitcoin Elliptic Curve

Bitcoin uses a specific Elliptic Curve defined over the finite field of the natural numbers, where $a = 0$ and $b = 7$.

The equation 1.1 becomes:

$$y^2 = x^3 + 7 \pmod{p} \quad (1.3)$$

The \pmod{p} (modulo prime number) indicates that this curve is over a finite field of prime order $p = 2^{256} - 2^{32} - 2^9 - 2^8 - 2^7 - 2^6 - 2^4 - 1$.

1.3 Bitcoin private-public key cryptography

Appendix A

Frequently Asked Questions

A.1 How do I change the colors of links?

The color of links can be changed to your liking using:

```
\hypersetup{urlcolor=red}, or  
\hypersetup{citecolor=green}, or  
\hypersetup{allcolor=blue}.
```

If you want to completely hide the links, you can use:

```
\hypersetup{allcolors=.}, or even better:  
\hypersetup{hidelinks}.
```

If you want to have obvious links in the PDF but not the printed text, use:

```
\hypersetup{colorlinks=false}.
```


Bibliography

- Arnold, A. S. et al. (Mar. 1998). "A Simple Extended-Cavity Diode Laser". In: *Review of Scientific Instruments* 69.3, pp. 1236–1239. URL: <http://link.aip.org/link/?RSI/69/1236/1>.
- Hawthorn, C. J., K. P. Weber, and R. E. Scholten (Dec. 2001). "Littrow Configuration Tunable External Cavity Diode Laser with Fixed Direction Output Beam". In: *Review of Scientific Instruments* 72.12, pp. 4477–4479. URL: <http://link.aip.org/link/?RSI/72/4477/1>.
- Wieman, Carl E. and Leo Hollberg (Jan. 1991). "Using Diode Lasers for Atomic Physics". In: *Review of Scientific Instruments* 62.1, pp. 1–20. URL: <http://link.aip.org/link/?RSI/62/1/1>.