

Complex Analysis - X400386

Yoav Eshel

February 1, 2021

Contents

1	Regions in the Complex Plane	2
2	Analytic Functions	2

This notes are based on Complex Variables and Applications by James Ward Brown (9th edition).

1 Regions in the Complex Plane

For $z_0 \in \mathbb{C}$, its *neighborhood* is the set

$$|z - z_0| \leq \varepsilon, \quad \varepsilon > 0.$$

It consists of all the point that lie inside (but not on!) the circle of radius ε centered at z_0 . A *deleted neighborhood* is the same set as the neighborhood minus the origin, i.e

$$0 < |z - z_0| < \varepsilon.$$

A point z_0 is an *interior point* of some set S , if there exists a z_0 -neighborhood that contains only points in S . It is an *exterior point* if there exists a neighborhood containing no points in S . If z_0 is neither, then it is a *boundary point*.

A set is *open* if it does not contain any of its boundary points (\iff every point is an interior point) and *closed* if it contains all of them. A set can be neither or both. The disk $0 < |z| \leq 1$ is neither while the entire complex plane is both.

An open set S is *connected* if there for any two points in the set there is a finite number of straight lines (polygonal line), all in S , that connect them. A nonempty open set that is connected is called a *domain* (e.g. every neighborhood is a domain). A connected set that is not open is called a *region*.

A set is *bounded* if it can be contained in some circle, otherwise it is *unbounded*.

A point z_0 is an *accumulation point* of a set S if every deleted neighborhood of z_0 contains at least one point of S . For example, the point 0 is an accumulation point of the set $S = \{\frac{1}{n} \mid n \in \mathbb{N}\}$.

2 Analytic Functions

We say that $f(z) \rightarrow w_0$ as $z \rightarrow z_0 = (x_0, y_0)$ if for every $\varepsilon > 0$ there exists $\delta > 0$ s.t.

$$0 < |z - z_0| < \delta \implies |f(z) - w_0| < \varepsilon$$

where $|z - z_0| = |x - x_0 + (y - y_0)i| = \sqrt{(x - x_0)^2 + (y - y_0)^2}$. If such w_0 exists then it is unique.

If $f : \mathbb{C} \rightarrow \mathbb{C}$ is given by

$$f(z) = f(x + yi) = u(x, y) + iv(x, y)$$

for u, v real valued and

$$\lim_{(x,y) \rightarrow (x_0,y_0)} u(x, y) = u_0, \quad \lim_{(x,y) \rightarrow (x_0,y_0)} v(x, y) = v_0$$

exist, then

$$\lim_{(x,y) \rightarrow (x_0,y_0)} f(x+yi) = u_0 + v_0i.$$

Therefore the limit rules for real valued functions are easily applied to complex functions.

If we center a unit sphere on the origin of the complex plane, then every point z on the plane is mapped to a unique point P on the surface of the sphere, where P is the intersection of the line from z to the north pole of the sphere. Note that points outside the unit circle are mapped to the northern hemisphere while points inside the unit circle are mapped to the southern hemisphere. Then the unit circle is mapped to the equator and the origin is mapped to the south pole. The *point of infinity* is defined as the point that is mapped to the north pole. Then for small ε the set $|z| > \frac{1}{\varepsilon}$ is called a *neighborhood of ∞* . Thus we can simply replace the neighborhoods in the definition of a limit by neighborhoods of ∞ .