

Obsvd. $t_g$	$v_1$ (i.e., $1/t_g$ )	Obsvd. $t_F$	$v_2$ (i.e., $1/t_F$ )	$(v'_2 - v_2)$	$\sqrt{v_1}(v'_2 - v_2)$
7.2 sec	0.1389	2.1 sec	0.4762	0.1984	0.0739
		3.6	0.2778	0.1978	0.0737
		12.5	0.0800		
5.8 sec	0.1724	1.9	0.5263	0.3792	0.1575
		6.8	0.1471	0.8529	0.3541
		1.0	1.000	0.7059	0.2931
		3.4	0.2941		

TABLE VII

$n$	$4.917 \times n$	Observed Charge	$n$	$4.917 \times n$	Observed Charge
1	4.917	...	10	49.17	49.41
2	9.834	...	11	54.09	53.91
3	14.75	...	12	59.00	59.12
4	19.66	19.66	13	63.92	63.68
5	24.59	24.60	14	68.84	68.65
6	29.50	29.62	15	73.75	...
7	34.42	34.47	16	78.67	78.34
8	39.34	39.38	17	83.59	83.22
9	44.25	44.42	18	88.51	...

	$d = 0.5\text{cm}$		$d = 0.5\text{cm}$	Charge on ion			Frictional charge					
$t_g$	$v_1(= d/t_g)$ (cm/sec)	$t_F$	$v_2(= d/t_F)$ (cm/sec)	$(v'_2-v_2)$	$n'$	$\frac{v'_2-v_2}{n'}$	$v_1 + v_2$	$n$	$\frac{v_1+v_2}{n}$			
18.2	.00286 <i>avr</i>	3.8	.01316	.00470	1	.00470	.01602	3	.00534			
18.6		2.8	.01786									
19.2			.01561	3	.00520							
18.0		22.2								.00225		
17.2			.00544	1	.00544							
15.4		6.5								.00769		
16.7			.00541	1	.00541							
18.0		21.9								.00228		
15.4			.01123	2	.00562							
17.3		3.7								.01351		
18.4												
17.5 <i>avr</i>						.00527 <i>avr</i>			.00534			

TABLE VI<sup>a</sup>

$t_g$ Sec.	$t_F$ Sec.	$\frac{1}{t_F}$	$\frac{1}{t'_F} - \frac{1}{t_F}$	$n$	$\frac{1}{n'}(\frac{1}{t'_F} - \frac{1}{t_F})$	$\frac{1}{t_g} + \frac{1}{t_F}$	n	$\frac{1}{n}(\frac{1}{t_g} + \frac{1}{t_F})$
11.848	80.708	.01236	.03234	6	.005390	.09655	18	.005366
11.890	22.366	.04470					7	
11.908	22.390		.007192	1	.005348	.09138		17
11.904	22.368	.005348					3	
11.882	140.565		.01254	5	.005375	.11833		22
11.906	79.600	.02870					5	
11.838	34.748		.02870	5	.005375	.11833		22
11.816	34.762	.03414					5	
11.776	34.846		.03414	5	.005375	.11833		22
11.840	29.286	.03414					5	
11.904	29.236		.026872	5	.005375	.11833		22
11.870	137.308	.007268					5	
11.952	34.638		.02884	5	.005375	.11833		22
11.860		.02884					5	
11.846	22.104		.04507	5	.005375	.11833		22
11.912	22.268	.04507					5	
11.910	500.1		.002000	5	.005375	.11833		22
11.918	19.704	.05079					5	
11.870	19.668		.05079	5	.005375	.11833		22
11.888	77.630	.03874					5	
11.894	77.806		.01285	5	.005375	.11833		22
11.878	42.302	.01079					5	
11.880			.02364	5	.005375	.11833		22
11.880		.02364					5	

Duration of exp.	= 45 min.	Pressure	= 75.62 cm.
Plate distance	= 16 mm.	Oil density	= .9199
Fall distance	= 10.21 mm.	Air viscosity	= $1,824 \times 10^{-7}$ [poise]
Initial volts	= 5,088.8	Radius ( $a$ )	= .000276 cm.
Final volts	= 5,081.2	$\frac{l}{a}$ [mean free path $\div a$ ]	= .034
Temperature	= 22.82°C.	Speed of fall	= .08584 cm./sec.

$$e_i = 4.991 \times 10^{-10} \text{ [statcoulomb]}^b$$

<sup>a</sup>[The bracketed numbers are our corrections of errors in the original paper.]

<sup>b</sup>[The value presently accepted is  $4.802 \times 10^{-10}$  statcoulombs.]

1	2	3	4
5	6	7	8
9	10	11	12

1	2	3	4
5	6	7	8
9	10	11	12

- First line
- Second line
- Third line, which is quite long and seemingly tedious in the extreme
- Fourth line, which isn't as long as the third
- Fifth line

$$P(\mathbf{a}, \mathbf{b}) = \int d\lambda \rho(\lambda) A(\mathbf{a}, \lambda) B(\mathbf{b}, \lambda).^1 \tag{1}$$


---

<sup>1</sup>[According to the hypothesis Bell is investigating, the "more complete specification" of the particles' state, the extra information that would tell the observer who knew it which way the particle would respond to a given measurement, is given by  $\lambda$ . Equation (2) considers all the possible  $\lambda$ 's and weights them according to how likely each is. Mathematically, that means integrating over all the possible values of  $\lambda$  and weighting each by its likelihood, represented by the probability distribution  $\rho(\lambda)$ . It may be helpful to note that  $P$  was likely chosen to stand for "product," since the term we are integrating  $\rho$  against is the product of  $A$  and  $B$ . Because we've defined  $A$  and  $B$  such that the results of measurements yield the values  $+1$  and  $-1$ , the product for any given value of  $\lambda$  will always be either  $+1$  (meaning that the two particles would both be measured to be pointing either in the selected directions  $\mathbf{a}$  and  $\mathbf{b}$  or in directions opposed to them) or  $-1$  (meaning that the directions of  $\sigma_1$  and  $\sigma_2$  would be measured to be opposed, whether because  $\sigma_1$  would *not* be measured to point in direction  $\mathbf{a}$  and  $\sigma_2$  *would* be measured to point in direction  $\mathbf{b}$  or because  $\sigma_1$  *would* be measured to point in direction  $\mathbf{a}$  and  $\sigma_2$  would *not* be measured to point in direction  $\mathbf{b}$ . Since it is weighted sum of all the positive and negative terms,  $P(\mathbf{a}, \mathbf{b})$  is thus a measure of the *correlation* between the spin-direction of the two particles relative to the angles  $\mathbf{a}$  and  $\mathbf{b}$ .]

## Note

### Single-slit Diffraction and the Limit of Optical Resolution

1. *Single-slit diffraction.* Let there be a narrow slit of width  $d$ , illuminated uniformly along the axis perpendicular to the plane of the slit. As in Huygens' treatment, consider the plane of the slit to be populated by infinitely many centers of wavelets, expanding in all forward directions. Thus the light beam will spread out as it leaves the slit.




There will be some angle, say  $\theta$ , at which the wavelets that were emitted from *one edge* of the slit and from the *center* of the slit, respectively, will cover distances that differ by an integral number of half-wavelengths and arrive together at some distant point—where they will mutually cancel, being exactly out of phase with one another. The innumerable remaining wavelets may be similarly paired for mutual cancellation; so that the overall result will be a dark spot at angle  $\theta$  from the perpendicular axis.

In the right triangle thus formed with hypotenuse  $d/2$  and one leg an integral multiple of  $\lambda/2$ , the marked angle will be equal to  $\theta$  and will have sine given by

$$\sin \theta = \frac{n\lambda/2}{d/2} = \frac{n\lambda}{d}.$$

Since  $n$  is any integer, there will be alternating bands of illumination and darkness for all angles up to  $90^\circ$  on either side of the perpendicular axis. But in practice the greatest illumination



is found within the “central maximum,” that region bounded by the two minima for which  $n = 1$  (called “first-order” minima). Suppose these minima appear at angles  $\theta_1$  on either side of the axis. Then  $\theta_1$  will be the *angular half-width* of the central maximum, and will be given by


$$\sin \theta_1 = \frac{\lambda}{d}.$$

2. *Optical resolution.*<sup>2</sup> Consider points  $P_1$  and  $P_2$  on the surface of some object. Let their angular separation be  $\beta$ , and let it be supposed that both points emit or reflect light of wavelength  $\lambda$ . For the sake of simplicity, we will let the aperture through which light is admitted to the instrument be a slit of width  $a$  (a circular aperture can be analogously treated, but numerous complications arise for that case). Light from each point forms its own independent diffraction pattern with a central maximum flanked by pairs of minima. As shown in the previous section 1, each central maximum will have an angular half-width equal to  $\theta_1$  such that

$$\sin \theta_1 = \lambda/a.$$

Now the central maxima of the two patterns lie in the focal plane of the instrument lens and (since rays passing through the center of a lens are not bent) must be separated from one another by the angle  $\beta$ . But as aperture  $a$  is made smaller, the angular half-width of each central maximum increases, until eventually  $\theta_1 = \beta$ —that is, the central maximum of each pattern coincides with a first-order minimum of the other pattern. The central bright maxima will then be immediately adjacent to one another. Inspection of the sum of the two intensity graphs under these conditions shows that instead of forming two distinct bands, the central maxima will merge into a *single band*.

Someone viewing such an image through the instrument would be quite unable to distinguish either of these maxima from the other; and thus the points  $P_1$  and  $P_2$  would become *indistinguishable*. No increase of magnifying power, so



<sup>2</sup>After Curtis Wilson, c. 1980.

long as the same aperture width is retained, can remedy this limitation. However, if the aperture width  $a$  is increased even the slightest amount, so that the separation between the two patterns increases by any degree at all, there will be found a dip between the two maxima when the sum of the intensity graphs is plotted as before. Thus, that the central maximum of each pattern shall coincide with the first-order minimum of the other is a *limiting condition* for the resolution of the images of two points; it is called *Rayleigh's criterion*.

Heisenberg's equation (16) (put page number) derives directly from Rayleigh's criterion. For suppose  $\theta_1 = \beta$  as described above; that is, let

$$\sin \beta = \lambda/a. \quad (2)$$

Let the object distance be  $R$ , and suppose also that angle  $\beta$  is very small. Then  $\Delta x$  is the chord of a circle with center at the vertex of  $\beta$ , and so it very nearly equals the arc which it subtends. This small arc equals  $R\beta$  (radians); while in its turn a very small angle  $\beta$  (in radians) nearly equals  $\sin \beta$ . This gives  $\Delta x \approx R \times \sin \beta$ , so that

$$\sin \beta \approx \Delta x/R. \quad (3)$$

From equations (1) and (2) it follows that

$$\frac{\Delta x}{R} = \frac{\lambda}{a}. \quad (4)$$

Similarly, the aperture of width  $a$  located at distance  $R$  from a point will subtend an angle  $\varepsilon$  such that, if  $\varepsilon$  is very small,

$$\sin \varepsilon \approx a/R,$$


from which

$$a \approx R \sin \varepsilon. \quad (5)$$

Substitution of equation (4) into equation (3) above yields


$$\frac{\Delta x}{R} \approx \frac{\lambda}{R \sin \varepsilon}$$

which becomes Heisenberg's equation (16), when  $R$  is canceled from both sides. Q.E.D.



## SINGLE-SLIT DIFFRACTION AND LIMIT OF OPTICAL RESOLUTION

1. *Single-slit diffraction.* Let there be a narrow slit of width  $d$ , illuminated uniformly along the axis perpendicular to the plane of the slit. As in Huygens' treatment, consider the plane of the slit to be populated by infinitely many centers of wavelets, expanding in all forward directions. Thus the light beam will spread out as it leaves the slit.




There will be some angle, say  $\theta$ , at which the wavelets that were emitted from *one edge* of the slit and from the *center* of the slit, respectively, will cover distances that differ by an integral number of half-wavelengths and arrive together at some distant point—where they will mutually cancel, being exactly out of phase with one another. The innumerable remaining wavelets may be similarly paired for mutual cancellation; so that the overall result will be a dark spot at angle  $\theta$  from the perpendicular axis.

In the right triangle thus formed with hypotenuse  $d/2$  and one leg an integral multiple of  $\lambda/2$ , the marked angle will be equal to  $\theta$  and will have sine given by


$$\sin \theta = \frac{n\lambda/2}{d/2} = \frac{n\lambda}{d}.$$

Since  $n$  is any integer, there will be alternating bands of illumination and darkness for all angles up to  $90^\circ$  either side of the perpendicular axis. But in practice the greatest illumination is found within the “central maximum,” that region bounded by the two minima for which  $n = 1$  (called “first-order” minima).



Suppose these minima appear at angles  $\theta_1$  on either side of the axis. Then  $\theta_1$  will be the *angular half-width* of the central maximum, and will be given by

$$\sin \theta_1 = \frac{\lambda}{d}.$$



2. *Optical resolution.*<sup>1</sup> Consider points  $P_1$  and  $P_2$  on the surface of some object. Let their angular separation be  $\beta$ , and let it be supposed that both points emit or reflect light of wavelength  $\lambda$ . For the sake of simplicity, we will let the aperture through which light is admitted to the instrument be a slit of width  $a$  (a circular aperture can be analogously treated, but numerous complications arise for that case). Light from each point forms its own independent diffraction pattern with a central maximum flanked by pairs of minima. As shown in the previous section 1, each central maximum will have an angular half-width equal to  $\theta_1$  such that

$$\sin \theta_1 = \lambda / a .$$

Now the central maxima of the two patterns lie in the focal plane of the instrument lens and (since rays passing through the center of a lens are not bent) must be separated from one another by the angle  $\beta$ . But as aperture  $a$  is made smaller, the angular half-width of each central maximum increases, until eventually  $\theta_1 = \beta$ —that is, the central maximum of each pattern coincides with a first-order minimum of the other pattern. The central bright maxima will then be immediately adjacent to one another. Inspection of the sum of the two intensity graphs

<sup>1</sup> After Curtis Wilson, c. 1980.



under these conditions shows that instead of forming two distinct bands, the central maxima will merge into a *single band*.

Someone viewing such an image through the instrument would be quite unable to distinguish either of these maxima from the other; and thus the points  $P_1$  and  $P_2$  would become *indistinguishable*. No increase of magnifying power, so long as the same aperture width is retained, can remedy this limitation. However, if the aperture width  $a$  is increased even the slightest amount, so that the separation between the two patterns increases by any degree at all, there will be found a dip between the two maxima when the sum of the intensity graphs is plotted as before. Thus, that the central maximum of each pattern shall coincide with the first-order minimum of the other is a *limiting condition* for the resolution of the images of two points; it is called *Rayleigh's criterion*.

Heisenberg's equation (16), p. 202 above, derives directly from Rayleigh's criterion. For suppose  $\theta_1 = \beta$  as described above; that is, let

$$\sin \beta = \lambda/a . \quad (1)$$

Let the object distance be  $R$ , and suppose also that angle  $\beta$  is very small. Then  $\Delta x$  is the chord of a circle with center at the vertex of  $\beta$ , and so it very nearly equals the arc which it subtends. This small arc equals  $R \times \beta$  (radians); while in its turn a very small angle  $\beta$  (in radians) nearly equals  $\sin \beta$ . This gives  $\Delta x \approx R \times \sin \beta$ , so that

$$\sin \beta \approx \Delta x/R . \quad (2)$$

From equations (1) and (2) it follows that

$$\frac{\Delta x}{R} \approx \frac{\lambda}{a} . \quad (3)$$

Similarly, the aperture of width  $a$  located at distance  $R$  from a point will subtend an angle  $\varepsilon$  such that, if  $\varepsilon$  is very small,

$$\sin \varepsilon \approx a/R ,$$

from which

$$a \approx R \sin \varepsilon . \quad (4)$$

Substitution of equation (4) into equation (3) above yields

$$\frac{\Delta x}{R} \approx \frac{\lambda}{R \sin \varepsilon}$$

which becomes Heisenberg's equation (16), when  $R$  is canceled from both sides. Q.E.D.

