CORE PURE / DIFFERENTIAL EQUATIONS

Systems of Differential Equations

Questions by topic 12 questions · 171 marks

Questions

Question 1	*10 marks*	2
Question 2	*11 marks*	3
Question 3	*11 marks*	4
Question 4	*13 marks*	5
Question 5	*10 marks*	6
Question 6	*11 marks*	7
Question 7	*18 marks*	8
Question 8	*17 marks*	9
Question 9	*16 marks*	10
Question 10	*16 marks*	11
Question 11	*15 marks*	12
Question 12	*23 marks*	13

1. The functions x and y satisfy

$$\frac{\mathrm{d}x}{\mathrm{d}t} + \ y = 5x$$
$$\frac{\mathrm{d}y}{\mathrm{d}t} + 2y = 6x$$

(a) Differentiate the first equation and eliminate y to show that x satisfies

$$\frac{\mathrm{d}^2x}{\mathrm{d}t^2} - 3\frac{\mathrm{d}x}{\mathrm{d}t} - 4x = 0 \qquad \textbf{[3]}$$

(b) Given that $x = 1$ and $y = 3$ when $t = 0$, find explicit expressions for x and y in terms of t. **[7]**

2. The functions x and y satisfy the simultaneous differential equations

$$\frac{\mathrm{d}x}{\mathrm{d}t} + y = 0$$
$$\frac{\mathrm{d}y}{\mathrm{d}t} - x = 2e^t$$

and when $t = 0$, $x = 0$ and $y = 1$.

Solve the system, giving x and y explicitly in terms of t **[11]**

3. The functions x and y satisfy

$$\frac{\mathrm{d}x}{\mathrm{d}t} = 3x + 4y + 7$$
$$\frac{\mathrm{d}y}{\mathrm{d}t} = -x - y - 1$$

Given that $x = 5$ and $y = -3$ when $t = 0$, find x and y as functions of t **[11]**

4. The functions x and y satisfy the system of differential equations

$$\frac{\mathrm{d}x}{\mathrm{d}t} = x + 2y - 7$$
$$\frac{\mathrm{d}y}{\mathrm{d}t} = 2x + y - 8$$

(a) Find the stationary point of the system.
That is, find the values of x and y such that $\frac{\mathrm{d}x}{\mathrm{d}t} = 0$ and $\frac{\mathrm{d}y}{\mathrm{d}t} = 0$ **[2]**

(b) Given that $x = 4$ and $y = 0$ when $t = 0$, find the particular solutions for x and y in terms of t **[11]**

5. For $t \geq 0$, the functions $x = f(t)$ and $y = g(t)$ satisfy the coupled first order differential equations

$$\frac{\mathrm{d}x}{\mathrm{d}t} = 2x - y$$
$$\frac{\mathrm{d}y}{\mathrm{d}t} = x + 2y$$

Given that $x = 1$ and $y = 2$ when $t = 0$, determine exact simplified expressions for $f(t)$ and $g(t)$ **[10]**

6. The functions x and y satisfy the simultaneous differential equations

$$\frac{\mathrm{d}x}{\mathrm{d}t} + y = 0$$
$$\frac{\mathrm{d}y}{\mathrm{d}t} - x = 2$$

and when $t = \frac{\pi}{2}$, $x = 1$ and $y = 2$.

Solve the system, giving x and y explicitly in terms of t **[11]**

7. An ecologist is studying the effect of introducing a population of weed-eating snails into a lake containing an invasive waterweed. During the first few months, a temporary pollutant also affects the snail population.

At time t months, the amount of waterweed, w, and the number of snails, s, are modelled by the differential equations

$$\frac{\mathrm{d}w}{\mathrm{d}t} = 4w - s$$

$$\frac{\mathrm{d}s}{\mathrm{d}t} = 6w - s - 48e^{-t}$$

(a) Show that

$$\frac{\mathrm{d}^2w}{\mathrm{d}t^2} - 3\frac{\mathrm{d}w}{\mathrm{d}t} + 2w = 48e^{-t}$$ **[3]**

(b) Find a general solution for the amount of waterweed at time t months **[6]**

(c) Find a general solution for the number of snails at time t months **[2]**

The model predicts that, at time T months, the waterweed will have been eliminated.
Given that $w = 30$ and $s = 110$ when $t = 0$

(d) Find the value of T, giving your answer to 3 decimal places **[6]**

(e) Suggest one limitation of the model **[1]**

8. At the start of 2021, a survey began of the number of mature trees and saplings in a managed woodland.

At time t years after the survey began, the number of mature trees, m, and the number of saplings, s, are modelled by the differential equations

$$\frac{\mathrm{d}m}{\mathrm{d}t} = \frac{1}{4}s$$
$$\frac{\mathrm{d}s}{\mathrm{d}t} = -\frac{1}{20}m - \frac{1}{10}s$$

(a) Show that

$$\frac{\mathrm{d}^2s}{\mathrm{d}t^2} + \frac{1}{10}\frac{\mathrm{d}s}{\mathrm{d}t} + \frac{1}{80}s = 0$$ **[3]**

(b) Find a general solution for the number of saplings in the woodland at time t years **[4]**


(c) Hence find a general solution for the number of mature trees in the woodland at time t years **[3]**

At the start of 2021 there were 20 mature trees and 20 saplings.

(d) (i) According to this model, find the first value of t for which no saplings remain, and state the year.

(ii) According to this model, how many mature trees are there at that time?

(iii) Use your answers to parts (i) and (ii) to comment on the suitability of the model. **[7]**



9. A treatment system consists of two well-stirred tanks, A and B. Tank A has capacity 300 litres and tank B has capacity 500 litres.

At time $t = 0$, tank A is full and contains 300 grams of salt, while tank B is full and contains 1200 grams of salt.

When $t > 0$, liquid flows in the following ways:

- Brine with a salt concentration of μ grams per litre flows into tank B at a constant rate
- Liquid flows from tank A to tank B at a rate of 3 litres per minute
- Liquid flows from tank B to tank A at a rate of r litres per minute
- Liquid flows out of tank A through a waste pipe
- The amount of liquid in tank A remains at 300 litres and the amount of liquid in tank B remains at 500 litres

At time t minutes ($t \geq 0$) there are x grams of salt in tank A and y grams of salt in tank B.

This system is represented by the coupled differential equations

$$\frac{\mathrm{d}x}{\mathrm{d}t} = -\frac{1}{20}x + \frac{3}{100}y$$
$$\frac{\mathrm{d}y}{\mathrm{d}t} = \frac{1}{100}x - \frac{3}{100}y + 24$$

(a) Find the value of r **[2]**

(b) Show that $\mu = 2$ **[3]**

(c) Solve the coupled differential equations to find both x and y in terms of t **[11]**

10. A water-treatment plant uses two storage tanks, tank A and tank B, joined by a balancing pipe. At any given instant, water can be transferred between the two tanks. It is assumed that these transfers happen instantaneously.

The volume of water in tank A at time t hours is denoted by A and the corresponding volume in tank B is B. Because the volumes are large for most of the time, A and B are modelled as being continuous.

When the volumes in the two tanks are different, water is pumped from the tank with more water to the tank with less water. In addition, water flows directly into tank A at a rate proportional to t^2. An engineer therefore decides to model A and B with the following coupled differential equations.

$$\frac{\mathrm{d}A}{\mathrm{d}t} = k(B - A) + \gamma t^2$$
$$\frac{\mathrm{d}B}{\mathrm{d}t} = k(A - B)$$

(a) Explain why k must be positive. **[1]**

After examining data, the engineer chooses $k = 1$ and $\gamma = 4$.

(b) Show that A satisfies the second order differential equation

$$\frac{\mathrm{d}^2A}{\mathrm{d}t^2} + 2\frac{\mathrm{d}A}{\mathrm{d}t} = 4t^2 + 8t$$

[3]

(c) (i) Find the complementary function for the differential equation from part (b). **[2]**

(ii) Explain why a particular integral of the form $A = at^2 + bt + c$ will not work in this situation. **[1]**

(iii) Using a particular integral of the form $A = at^3 + bt^2 + ct$, find the general solution of the differential equation from part (b). **[3]**

(d) At time $t = 0$, tank A contains $52\,\mathrm{m}^3$ of water and the volume in tank A is increasing at $13\,\mathrm{m}^3\,\mathrm{h}^{-1}$. Use the model to estimate the volume of water in tank A 30 minutes later. **[5]**

(e) Explain why the model becomes unreliable as t gets very large. **[1]**

11. A gardener releases ladybirds into a large greenhouse to control the aphids living there. At time t weeks after the ladybirds are released, there are x thousand aphids and y hundred ladybirds in the greenhouse.

Aphids fly into the greenhouse at a constant rate. Aphids die from natural causes at a rate proportional to the number of aphids, and they are eaten at a rate proportional to the number of ladybirds. The number of ladybirds increases at a rate proportional to the number of aphids and decreases at a rate proportional to the number of ladybirds.

For $t \geq 0$, the two populations are modelled by the differential equations

$$\frac{\mathrm{d}x}{\mathrm{d}t} = 40 - 6x - 4y$$
$$\frac{\mathrm{d}y}{\mathrm{d}t} = x - y$$

At time $t = 0$, there are 1000 aphids and 100 ladybirds in the greenhouse.

(a) Show that

$$\frac{\mathrm{d}^2x}{\mathrm{d}t^2} + 7\frac{\mathrm{d}x}{\mathrm{d}t} + 10x = 40$$ **[3]**

(b) Find x in terms of t. **[7]**

(c) Find the exact value of t at which the number of aphids in the greenhouse is greatest. **[2]**

(d) Hence find the number of ladybirds in the greenhouse at this time. **[2]**

(e) With reference to the first differential equation, explain why the model is unrealistic when there are very few aphids in the greenhouse. **[1]**

12. Two invasive plant species, X and Y, are spreading through a wetland and compete for light and nutrients. A model for this competition assumes, in one particular wetland, the following.

- In the absence of the other species, species X would increase at a rate proportional to the number present, with constant of proportionality p.
- In the absence of the other species, species Y would increase at a rate proportional to the number present, with constant of proportionality q.
- Competition reduces the rate of increase of species X by an amount proportional to the number of species Y present.
- Competition reduces the rate of increase of species Y by an amount proportional to the number of species X present.

If the numbers of species X and Y present at time t years are x and y respectively, the model gives the differential equations

$$\frac{\mathrm{d}x}{\mathrm{d}t} = px - ay \qquad \text{and} \qquad \frac{\mathrm{d}y}{\mathrm{d}t} = qy - bx$$

(a) (i) Show that x satisfies

$$\frac{\mathrm{d}^2x}{\mathrm{d}t^2} - (p+q)\frac{\mathrm{d}x}{\mathrm{d}t} + (pq - ab)x = 0$$

and hence that the general solution for x is

$$x = Ae^{\lambda_1 t} + Be^{\lambda_2 t}$$

where λ_1 and λ_2 are the roots of

$$\lambda^2 - (p+q)\lambda + (pq - ab) = 0$$ **[5]**

(ii) Hence show that the general solution for y can be written in the form

$$y = \frac{p-\lambda_1}{a}Ae^{\lambda_1 t} + \frac{p-\lambda_2}{a}Be^{\lambda_2 t}$$ **[2]**

Observations suggest that suitable values for the model are $p = 0.035$, $q = 0.005$, $a = 0.03$ and $b = 0.0225$. You should use these values in the rest of this question.

(b) When $t = 0$, the numbers present of species X and Y in this wetland are x_0 and y_0 respectively.

(i) Show that

$$x = \frac{1}{4}(3x_0 - 2y_0)e^{0.05t} + \frac{1}{4}(x_0 + 2y_0)e^{-0.01t}$$ **[3]**

(ii) Hence show that

$$y = -\frac{1}{8}(3x_0 - 2y_0)e^{0.05t} + \frac{3}{8}(x_0 + 2y_0)e^{-0.01t}$$ **[1]**

(c) Use initial values $x_0 = 400$ and $y_0 = 500$ with the results in part (b) to determine what the model predicts for each of the following.

(i) What numbers of each species will be present after 25 years? **[2]**
(ii) When will the numbers of the two species be equal? **[4]**
(iii) Does either species ever disappear from the wetland? Justify your answer **[3]**

(d) Different initial values will apply in other wetlands where the two species compete, but previous studies indicate that in most wetlands one population eventually increases while the other decreases.

(i) Find a relation between x_0 and y_0 for which the model does not predict this outcome **[1]**
(ii) Explain what the model predicts in the long term for this exceptional case **[2]**