FURTHER MECHANICS / MOMENTUM & COLLISIONS

Vector Momentum and Impulse

Worked solutions 8 questions · 49 marks

EXAM BOARDS Edexcel FM1 9FM0/3C · AQA 7367/3M · OCR A Y543 · OCR MEI Mech Minor Y431 OCR MEI Mech Major Y421

Instructions

Take $g = 9.8\,\mathrm{m\,s^{-2}}$ unless a question gives a value. Where a question leaves g unspecified and a final answer depends on it, the solutions also give that answer for $g = 10\,\mathrm{m\,s^{-2}}$.

Questions

Question 1	*6 marks*	2
Question 2	*6 marks*	4
Question 3	*5 marks*	6
Question 4	*6 marks*	8
Question 5	*7 marks*	10
Question 6	*6 marks*	12
Question 7	*6 marks*	14
Question 8	*7 marks*	15

1. A particle P of mass $5\,\text{kg}$ is moving with speed $13\,\text{m s}^{-1}$ in the direction of the vector $5\mathbf{i} - 12\mathbf{j}$ when it receives an impulse $(45\mathbf{i} + 90\mathbf{j})\,\text{N s}$.

(a) Find the speed of P immediately after receiving the impulse **[4]**

The direction of motion of P is then changed by $\alpha°$.

(b) Find the value of α **[2]**

Solution

(a) The given direction vector is $5\mathbf{i} - 12\mathbf{j}$, whose magnitude is

$$\sqrt{5^2 + (-12)^2} = \sqrt{25 + 144} = 13$$

Since the speed is $13\,\text{m s}^{-1}$, the initial velocity is

$$\mathbf{u} = 13\left(\frac{5\mathbf{i} - 12\mathbf{j}}{13}\right) = 5\mathbf{i} - 12\mathbf{j}$$

Using impulse = change in momentum:

$$\mathbf{I} = m\mathbf{v} - m\mathbf{u}$$

Substituting $m = 5$ and $\mathbf{I} = 45\mathbf{i} + 90\mathbf{j}$:

$$45\mathbf{i} + 90\mathbf{j} = 5\mathbf{v} - 5(5\mathbf{i} - 12\mathbf{j})$$

Hence

$$5\mathbf{v} = (45\mathbf{i} + 90\mathbf{j}) + (25\mathbf{i} - 60\mathbf{j}) = 70\mathbf{i} + 30\mathbf{j}$$

Therefore the final velocity is

$$\mathbf{v} = \frac{70\mathbf{i} + 30\mathbf{j}}{5} = 14\mathbf{i} + 6\mathbf{j}$$

So the speed immediately after the impulse is

$$|\mathbf{v}| = \sqrt{14^2 + 6^2} = \sqrt{196 + 36} = \sqrt{232} = 2\sqrt{58}$$

Therefore the speed is $2\sqrt{58}\,\text{m s}^{-1} \approx 15.2\,\text{m s}^{-1}$.

(b) The angle α is the angle between the initial and final velocities:

$$\mathbf{u} = 5\mathbf{i} - 12\mathbf{j}, \qquad \mathbf{v} = 14\mathbf{i} + 6\mathbf{j}$$

Using the scalar product:

$$\cos\alpha = \frac{\mathbf{u} \cdot \mathbf{v}}{|\mathbf{u}||\mathbf{v}|}$$

Now

$$\mathbf{u} \cdot \mathbf{v} = (5)(14) + (-12)(6) = 70 - 72 = -2$$

The magnitudes are

$$|\mathbf{u}| = 13, \qquad |\mathbf{v}| = 2\sqrt{58}$$

So

$$\cos\alpha = -\frac{2}{13(2\sqrt{58})} = -\frac{1}{13\sqrt{58}}$$

Hence

$$\alpha = \cos^{-1}\left(-\frac{1}{13\sqrt{58}}\right) \approx 90.6$$

2. A particle P of mass 2 kg is moving with speed $15\,\text{m s}^{-1}$ in the direction of $3\mathbf{i}+4\mathbf{j}$ when it receives an impulse $\mathbf{J}$ N s. Immediately after receiving the impulse, P is moving with speed $17\,\text{m s}^{-1}$ in the direction of $8\mathbf{i}+15\mathbf{j}$.

(a) Find the magnitude of $\mathbf{J}$ **[4]**

The angle between the direction of the impulse and the direction of motion of P immediately before receiving the impulse is $\alpha°$.

(b) Find the value of α **[2]**

Solution

(a) The initial direction is $3\mathbf{i}+4\mathbf{j}$, whose magnitude is

$$\sqrt{3^2+4^2}=5$$

So the initial velocity is

$$\mathbf{u}=15\left(\frac{3\mathbf{i}+4\mathbf{j}}{5}\right)=9\mathbf{i}+12\mathbf{j}$$

The final direction is $8\mathbf{i}+15\mathbf{j}$, whose magnitude is

$$\sqrt{8^2+15^2}=17$$

So the final velocity is

$$\mathbf{v}=17\left(\frac{8\mathbf{i}+15\mathbf{j}}{17}\right)=8\mathbf{i}+15\mathbf{j}$$

Using impulse = change in momentum:

$$\mathbf{J}=m\mathbf{v}-m\mathbf{u}$$

Substituting $m=2$:

$$\begin{aligned}\mathbf{J}&=2\big((8-9)\mathbf{i}+(15-12)\mathbf{j}\big)\\&=2(-\mathbf{i}+3\mathbf{j})\\&=-2\mathbf{i}+6\mathbf{j}\end{aligned}$$

Therefore its magnitude is

$$\begin{aligned}|\mathbf{J}|&=\sqrt{(-2)^2+6^2}\\&=\sqrt{40}\\&=2\sqrt{10}\end{aligned}$$

So the magnitude of the impulse is $2\sqrt{10}\,\text{N s}$.

(b) The direction of motion before the impulse is the direction of the initial velocity $\mathbf{u}=9\mathbf{i}+12\mathbf{j}$, and the impulse vector is

$$\mathbf{J}=-2\mathbf{i}+6\mathbf{j}$$

Using the scalar product:

$$\cos\alpha = \frac{\mathbf{J}\cdot\mathbf{u}}{|\mathbf{J}||\mathbf{u}|}$$

Now

$$\begin{aligned}\mathbf{J}\cdot\mathbf{u} &= (-2)(9) + (6)(12)\\ &= -18 + 72\\ &= 54\end{aligned}$$

The magnitudes are


$$|\mathbf{J}| = 2\sqrt{10}, \qquad |\mathbf{u}| = 15$$

So

$$\begin{aligned}\cos\alpha &= \frac{54}{(2\sqrt{10})(15)}\\ &= \frac{9}{5\sqrt{10}}\end{aligned}$$

Hence

$$\alpha = \cos^{-1}\left(\frac{9}{5\sqrt{10}}\right) \approx 55.3$$



3. A particle P of mass 0.4 kg is moving due east with speed $15\,\text{m s}^{-1}$ on a smooth horizontal plane. The particle receives a horizontal impulse of magnitude J N s.

Immediately afterwards, P is moving with speed $9\,\text{m s}^{-1}$ on a bearing of 330°, as shown in the diagram.

Find the value of J **[5]**

Solution

Take east as the positive x-direction and north as the positive y-direction.

Initially, the particle is moving due east at $15\,\text{m s}^{-1}$, so its initial velocity is

$$\mathbf{u} = \begin{pmatrix} 15 \\ 0 \end{pmatrix}$$

After the impulse, the particle moves with speed $9\,\text{m s}^{-1}$ on a bearing of 330°.

A bearing of 330° means 30° west of north, so the final velocity has components

$$\mathbf{v} = \begin{pmatrix} -9\sin 30° \\ 9\cos 30° \end{pmatrix} = \begin{pmatrix} -\frac{9}{2} \\ \frac{9\sqrt{3}}{2} \end{pmatrix}$$

Using impulse = change in momentum:

$$\mathbf{J} = m\mathbf{v} - m\mathbf{u}$$

Substituting $m = 0.4$, the impulse vector is

$$\mathbf{J} = 0.4\begin{pmatrix} -\frac{9}{2} \\ \frac{9\sqrt{3}}{2} \end{pmatrix} - 0.4\begin{pmatrix} 15 \\ 0 \end{pmatrix} = \begin{pmatrix} -\frac{9}{5} \\ \frac{9\sqrt{3}}{5} \end{pmatrix} - \begin{pmatrix} 6 \\ 0 \end{pmatrix} = \begin{pmatrix} -\frac{39}{5} \\ \frac{9\sqrt{3}}{5} \end{pmatrix}$$

Its magnitude is

$$\begin{aligned} J &= |\mathbf{J}| \\ &= \sqrt{\left(-\frac{39}{5}\right)^2 + \left(\frac{9\sqrt{3}}{5}\right)^2} \\ &= \sqrt{\frac{1521}{25} + \frac{243}{25}} \\ &= \sqrt{\frac{1764}{25}} \\ &= \frac{42}{5} \end{aligned}$$

Hence

$$J = 8.4\,\mathrm{N\,s}$$

4. A particle Q of mass $0.5\,\text{kg}$ is moving on a smooth horizontal surface with speed $10\,\text{m s}^{-1}$ in the direction of the vector $4\mathbf{i} - 3\mathbf{j}$. The particle then receives an impulse of magnitude $5\,\text{N s}$ in the direction of the vector $3\mathbf{i} + 4\mathbf{j}$.

(a) Find the speed of Q immediately after receiving the impulse **[4]**

As a result of receiving the impulse, the direction of motion of Q is turned through an angle $\theta°$.

(b) Find the value of θ **[2]**

Solution

(a) The given direction vector for the initial motion is $4\mathbf{i} - 3\mathbf{j}$, whose magnitude is

$$\sqrt{4^2 + (-3)^2} = 5$$

So the unit vector in this direction is

$$\frac{4\mathbf{i} - 3\mathbf{j}}{5}$$

Hence the initial velocity is

$$\mathbf{u} = 10\left(\frac{4\mathbf{i} - 3\mathbf{j}}{5}\right)$$

$$= 8\mathbf{i} - 6\mathbf{j}$$

The impulse has magnitude $5\,\text{N s}$ in the direction $3\mathbf{i} + 4\mathbf{j}$. Since

$$|3\mathbf{i} + 4\mathbf{j}| = 5$$

the unit vector in this direction is

$$\frac{3\mathbf{i} + 4\mathbf{j}}{5}$$

So the impulse vector is

$$\mathbf{I} = 5\left(\frac{3\mathbf{i} + 4\mathbf{j}}{5}\right)$$

$$= 3\mathbf{i} + 4\mathbf{j}$$

Using impulse = change in momentum:

$$\mathbf{I} = m\mathbf{v} - m\mathbf{u}$$

Substituting $m = 0.5$:

$$3\mathbf{i} + 4\mathbf{j} = 0.5\mathbf{v} - 0.5(8\mathbf{i} - 6\mathbf{j})$$

so

$$0.5\mathbf{v} = (3\mathbf{i} + 4\mathbf{j}) + (4\mathbf{i} - 3\mathbf{j})$$

$$= 7\mathbf{i} + \mathbf{j}$$

Hence the final velocity is

$$\begin{aligned}\mathbf{v} &= \frac{7\mathbf{i}+\mathbf{j}}{0.5} \\ &= 14\mathbf{i}+2\mathbf{j}\end{aligned}$$

Therefore the speed immediately after the impulse is

$$\begin{aligned}|\mathbf{v}| &= \sqrt{14^2+2^2} \\ &= \sqrt{196+4} \\ &= \sqrt{200} \\ &= 10\sqrt{2}\end{aligned}$$

So the speed is $10\sqrt{2}\,\mathrm{m\,s^{-1}}$.

(b) The angle turned is the angle between the initial and final velocities, $\mathbf{u} = 8\mathbf{i} - 6\mathbf{j}$ and $\mathbf{v} = 14\mathbf{i} + 2\mathbf{j}$.

Using the scalar product:

$$\cos\theta = \frac{\mathbf{u}\cdot\mathbf{v}}{|\mathbf{u}||\mathbf{v}|}$$

Now

$$\mathbf{u}\cdot\mathbf{v} = (8)(14) + (-6)(2) = 112 - 12 = 100$$

The magnitudes are

$$|\mathbf{u}| = 10, \qquad |\mathbf{v}| = 10\sqrt{2}$$

So

$$\begin{aligned}\cos\theta &= \frac{100}{10(10\sqrt{2})} \\ &= \frac{1}{\sqrt{2}}\end{aligned}$$

Hence

$$\theta = 45$$

5. A particle of mass 0.8 kg is moving with speed $5\,\text{m}\,\text{s}^{-1}$ in the direction of the vector $3\mathbf{i}+4\mathbf{j}$. It is then acted on by a constant force of $(8\mathbf{i}-6\mathbf{j})$ N for 0.4 s.

(a) Find the speed of the particle at the end of the 0.4 s interval **[5]**

(b) Find the size of the angle between the direction of motion of the particle before the force acts and the direction of motion of the particle at the end of the 0.4 s interval **[2]**

Solution

(a) The given direction vector is $3\mathbf{i}+4\mathbf{j}$, whose magnitude is

$$\sqrt{3^2+4^2}=5$$

So the initial velocity is

$$\mathbf{u}=5\left(\frac{3\mathbf{i}+4\mathbf{j}}{5}\right)=3\mathbf{i}+4\mathbf{j}$$

Using $\mathbf{F}=m\mathbf{a}$, the acceleration is

$$\mathbf{a}=\frac{8\mathbf{i}-6\mathbf{j}}{0.8}=10\mathbf{i}-7.5\mathbf{j}$$

Since the force acts for 0.4 s, the change in velocity is

$$\mathbf{a}t=(10\mathbf{i}-7.5\mathbf{j})(0.4)=4\mathbf{i}-3\mathbf{j}$$

Hence the final velocity is

$$\begin{aligned}\mathbf{v}&=\mathbf{u}+\mathbf{a}t\\&=(3\mathbf{i}+4\mathbf{j})+(4\mathbf{i}-3\mathbf{j})\\&=7\mathbf{i}+\mathbf{j}\end{aligned}$$

Therefore the final speed is

$$|\mathbf{v}|=\sqrt{7^2+1^2}=\sqrt{50}=5\sqrt{2}$$

So the speed at the end of the interval is $5\sqrt{2}\,\text{m}\,\text{s}^{-1}\approx 7.07\,\text{m}\,\text{s}^{-1}$.

(b) The angle between the two directions of motion is the angle between the initial and final velocity vectors.

Initial velocity:

$$\mathbf{u}=3\mathbf{i}+4\mathbf{j}$$

Final velocity:

$$\mathbf{v}=7\mathbf{i}+\mathbf{j}$$

Using the scalar product:

$$\cos\theta=\frac{\mathbf{u}\cdot\mathbf{v}}{|\mathbf{u}||\mathbf{v}|}$$

Now

$$\begin{aligned}\mathbf{u}\cdot\mathbf{v} &= (3)(7)+(4)(1)\\ &= 25\end{aligned}$$

The magnitudes are

$$|\mathbf{u}| = 5, \qquad |\mathbf{v}| = \sqrt{50} = 5\sqrt{2}$$

So

$$\begin{aligned}\cos\theta &= \frac{25}{5(5\sqrt{2})}\\ &= \frac{1}{\sqrt{2}}\end{aligned}$$

Hence

$$\theta = 45^\circ$$

6. A particle P of mass 0.8 kg is moving with velocity $\lambda(\mathbf{i}+2\mathbf{j})\,\mathrm{m\,s^{-1}}$ when P receives an impulse of magnitude $\frac{8\sqrt{2}}{5}\,\mathrm{N\,s}$.

Immediately after receiving the impulse, P is moving at speed $5\,\mathrm{m\,s^{-1}}$ in the direction of $3\mathbf{i}+4\mathbf{j}$.

Given that λ is a constant, find the two possible values of λ **[6]**

Solution

Let the initial velocity be $\mathbf{u}$ and the final velocity be $\mathbf{v}$.

The particle initially has velocity

$$\mathbf{u} = \lambda(\mathbf{i}+2\mathbf{j})$$

After the impulse, its speed is $5\,\mathrm{m\,s^{-1}}$ in the direction of $3\mathbf{i}+4\mathbf{j}$.

Since

$$|3\mathbf{i}+4\mathbf{j}| = \sqrt{3^2+4^2} = 5$$

the unit vector in that direction is

$$\frac{1}{5}(3\mathbf{i}+4\mathbf{j})$$

so the final velocity is

$$\mathbf{v} = 5\times\frac{1}{5}(3\mathbf{i}+4\mathbf{j}) = 3\mathbf{i}+4\mathbf{j}$$

Using impulse = change in momentum:

$$\mathbf{I} = m\mathbf{v} - m\mathbf{u}$$

Since the mass is $0.8\,\mathrm{kg} = \frac{4}{5}\,\mathrm{kg}$, the impulse is

$$\mathbf{I} = \frac{4}{5}\mathbf{v} - \frac{4}{5}\mathbf{u}$$

So

$$\begin{aligned}\mathbf{I} &= \frac{4}{5}[(3\mathbf{i}+4\mathbf{j}) - \lambda(\mathbf{i}+2\mathbf{j})]\\ &= \frac{4}{5}[(3-\lambda)\mathbf{i} + (4-2\lambda)\mathbf{j}]\end{aligned}$$

We are given that the magnitude of the impulse is $\frac{8\sqrt{2}}{5}\,\mathrm{N\,s}$, so

$$\left|\frac{4}{5}[(3-\lambda)\mathbf{i} + (4-2\lambda)\mathbf{j}]\right| = \frac{8\sqrt{2}}{5}$$

Squaring both sides:

$$\left(\frac{4}{5}\right)^2\left[(3-\lambda)^2 + (4-2\lambda)^2\right] = \left(\frac{8\sqrt{2}}{5}\right)^2$$

Hence

$$\frac{16}{25}\left[(3-\lambda)^2 + (4-2\lambda)^2\right] = \frac{128}{25}$$

Multiply by $\frac{25}{16}$:

$$(3-\lambda)^2+(4-2\lambda)^2=8$$

Expand:

$$\begin{aligned}(3-\lambda)^2+(4-2\lambda)^2&=8\\(\lambda^2-6\lambda+9)+(4\lambda^2-16\lambda+16)&=8\\5\lambda^2-22\lambda+25&=8\\5\lambda^2-22\lambda+17&=0\end{aligned}$$

Factorising:

$$(5\lambda-17)(\lambda-1)=0$$

Therefore

$$\lambda=\frac{17}{5}\quad\text{or}\quad\lambda=1$$

7. A particle P of mass 1 kg is moving with velocity $(3\mathbf{i}+\mathbf{j}+\mathbf{k})\,\mathrm{m\,s^{-1}}$.

The particle receives an impulse $\lambda(2\mathbf{i}+\mathbf{j}+2\mathbf{k})\,\mathrm{N\,s}$, where λ is a constant.

Immediately after receiving the impulse, the kinetic energy of P is 19 J.

Find the possible values of λ **[6]**

Solution

Using impulse = change in momentum:

$$\mathbf{I} = m\mathbf{v} - m\mathbf{u}$$

Here $m = 1$, $\mathbf{u} = 3\mathbf{i}+\mathbf{j}+\mathbf{k}$ and $\mathbf{I} = \lambda(2\mathbf{i}+\mathbf{j}+2\mathbf{k})$, so

$$\lambda(2\mathbf{i}+\mathbf{j}+2\mathbf{k}) = \mathbf{v} - (3\mathbf{i}+\mathbf{j}+\mathbf{k})$$

Hence the velocity immediately after the impulse is

$$\mathbf{v} = (3+2\lambda)\mathbf{i} + (1+\lambda)\mathbf{j} + (1+2\lambda)\mathbf{k}$$

The kinetic energy after the impulse is 19 J, so

$$\frac{1}{2}(1)|\mathbf{v}|^2 = 19$$

hence

$$|\mathbf{v}|^2 = 38$$

Now

$$|\mathbf{v}|^2 = (3+2\lambda)^2 + (1+\lambda)^2 + (1+2\lambda)^2$$

So

$$(3+2\lambda)^2 + (1+\lambda)^2 + (1+2\lambda)^2 = 38$$

$$(9+12\lambda+4\lambda^2) + (1+2\lambda+\lambda^2) + (1+4\lambda+4\lambda^2) = 38$$

$$11+18\lambda+9\lambda^2 = 38$$

Therefore

$$9\lambda^2+18\lambda-27=0$$

$$\lambda^2+2\lambda-3=0$$

$$(\lambda+3)(\lambda-1)=0$$

Hence the possible values of λ are

$$\lambda = 1 \quad \text{or} \quad \lambda = -3$$

8. A particle of mass 1 kg is moving with velocity $(\mathbf{i}+3\mathbf{j}-\mathbf{k})\,\mathrm{m\,s^{-1}}$ when it receives an impulse $\mathbf{I}\,\mathrm{N\,s}$.

As a result, its kinetic energy increases by 6.5 J and the particle then moves in the direction $(\mathbf{i}+2\mathbf{j}+\mathbf{k})$.

Find $\mathbf{I}$ **[7]**

Solution

Let the initial velocity be

$$\mathbf{u}=\mathbf{i}+3\mathbf{j}-\mathbf{k}$$

Since the mass is 1 kg, the initial kinetic energy is

$$\frac{1}{2}(1)|\mathbf{u}|^2$$

Now

$$\begin{aligned}|\mathbf{u}|^2&=1^2+3^2+(-1)^2\\&=1+9+1\\&=11\end{aligned}$$

So the initial kinetic energy is

$$\frac{1}{2}(1)(11)=\frac{11}{2}\,\mathrm{J}$$

The kinetic energy increases by $6.5\,\mathrm{J}=\frac{13}{2}\,\mathrm{J}$, so the final kinetic energy is

$$\begin{aligned}\frac{11}{2}+\frac{13}{2}&=\frac{24}{2}\\&=12\,\mathrm{J}\end{aligned}$$

If $\mathbf{v}$ is the final velocity, then

$$\frac{1}{2}(1)|\mathbf{v}|^2=12$$

so

$$|\mathbf{v}|^2=24$$

We are told the particle then moves in the direction $(\mathbf{i}+2\mathbf{j}+\mathbf{k})$, so

$$\mathbf{v}=\lambda(\mathbf{i}+2\mathbf{j}+\mathbf{k})$$

for some positive scalar λ.

Hence

$$\begin{aligned}|\mathbf{v}|^2&=\lambda^2(1^2+2^2+1^2)\\&=\lambda^2(6)\end{aligned}$$

But $|\mathbf{v}|^2=24$, so

$$\begin{aligned}6\lambda^2&=24\\\lambda^2&=4\end{aligned}$$

Since the motion is in the direction $\mathbf{i} + 2\mathbf{j} + \mathbf{k}$, we take $\lambda = 2$.

Therefore

$$\mathbf{v} = 2(\mathbf{i} + 2\mathbf{j} + \mathbf{k}) = 2\mathbf{i} + 4\mathbf{j} + 2\mathbf{k}$$

Using impulse = change in momentum:

$$\mathbf{I} = m\mathbf{v} - m\mathbf{u}$$

Since $m = 1$, this is

$$\mathbf{I} = \mathbf{v} - \mathbf{u}$$

So

$$\begin{aligned}\mathbf{I} &= (2\mathbf{i} + 4\mathbf{j} + 2\mathbf{k}) - (\mathbf{i} + 3\mathbf{j} - \mathbf{k}) \\ &= \mathbf{i} + \mathbf{j} + 3\mathbf{k}\end{aligned}$$

Hence the impulse is

$$\mathbf{I} = (\mathbf{i} + \mathbf{j} + 3\mathbf{k})\,\mathrm{N\,s}$$