TMUA / PRACTICE PAPER

Set A Paper 2

Mathematical Reasoning 20 questions · 75 minutes

Questions

Question 1	*Proof & Error Analysis*	2
Question 2	*Trigonometry*	3
Question 3	*Proof & Error Analysis*	4
Question 4	*Exponentials & Logarithms*	5
Question 5	*Number*	6
Question 6	*Proof & Error Analysis*	7
Question 7	*Geometry*	8
Question 8	*Coordinate Geometry*	9
Question 9	*Logic*	10
Question 10	*Logic*	11
Question 11	*Probability & Counting*	12
Question 12	*Proof & Error Analysis*	13
Question 13	*Integration*	14
Question 14	*Logic*	15
Question 15	*Proof & Error Analysis*	16
Question 16	*Sequences & Series*	17
Question 17	*Logic*	18
Question 18	*Logic*	19
Question 19	*Algebra*	20
Question 20	*Polynomials*	21

Proof & Error Analysis

1. Let x and y be real numbers.

A student attempts to prove the following claim:

$$\textbf{Claim: If } x + y \geq 2 \text{ then } x^2 + y^2 \geq 2$$

The student's attempt is as follows:

Line 1: Since $x + y \geq 2$, at least one of x and y is at least 1.
Assume $x \geq 1$; the other case is the same with x and y swapped.

Line 2: From $x + y \geq 2$ we get $y \geq 2 - x$

Line 3: Therefore $y^2 \geq (2 - x)^2$

Line 4: Hence $x^2 + y^2 \geq x^2 + (2 - x)^2$

Line 5: Now $x^2 + (2 - x)^2 = 2(x - 1)^2 + 2$, and this is at least 2.

Line 6: Therefore $x^2 + y^2 \geq 2$, as claimed.

Which of the following best describes the argument?

A Line 1 contains the first error.

B Line 2 contains the first error.

C Line 3 contains the first error.

D Line 4 contains the first error.

E Line 5 contains the first error.

F The argument is a correct proof of the claim.

G The argument contains no error, but the claim is false.

Trigonometry

2. In quadrilateral $ABCD$, the diagonal AC divides the figure into two right-angled triangles, $\triangle ABC$ and $\triangle ACD$.

- In $\triangle ABC$, $\angle ABC = 90°$, $\angle BAC = 30°$, and $AB = \sqrt{6}$ cm
- In $\triangle ACD$, $\angle ACD = 90°$, and $\tan(\angle CAD) = \dfrac{1}{\sqrt{2}}$

What is the area of quadrilateral $ABCD$, in cm^2?

A $\sqrt{3} + \sqrt{2}$

B $\sqrt{3} + 2\sqrt{2}$

C $2\sqrt{3} + \sqrt{2}$

D $2\sqrt{3} + 2\sqrt{2}$

E $\sqrt{3} + 4\sqrt{2}$

F $2\sqrt{3} + 4\sqrt{2}$

Proof & Error Analysis

3. Consider the following claim and attempted proof.

Claim: Let $f(x)$ be a quadratic polynomial. If $\displaystyle\int_{-1}^{1} f(x)\,\mathrm{d}x = 0$, then the graph of $y = f(x)$ must intersect the x-axis at two distinct points in the interval $-1 < x < 1$.

Attempted proof:

Line 1 Let $f(x) = ax^2 + bx + c$, where a, b, c are real constants with $a \neq 0$

Line 2 Evaluating the integral gives $\displaystyle\int_{-1}^{1} f(x)\,\mathrm{d}x = \frac{2a}{3} + 2c = 0$, so $c = -\frac{a}{3}$

Line 3 The discriminant is $b^2 - 4ac = b^2 + \frac{4a^2}{3} > 0$, so $f(x) = 0$ has two distinct real roots

Line 4 Writing these roots as x_1 and x_2, we have $x_1 x_2 = \frac{c}{a} = -\frac{1}{3}$

Line 5 Since $|x_1 x_2| = \frac{1}{3} < 1$, it follows that $|x_1| < 1$ and $|x_2| < 1$

Line 6 Therefore, both roots lie in the interval $-1 < x < 1$

Which line of the proof contains the first error?

A Line 1

B Line 2

C Line 3

D Line 4

E Line 5

F Line 6

G There is no error, the proof is correct as written.

Exponentials & Logarithms

4. Let a be a positive real number with $a \neq 1$

The equation

$$\log_a(3x - 1) = 2 + \log_a(x + 1)$$

has exactly one real solution x

What is the complete range of all the possible values of a?

A $0 < a < \sqrt{3}$ and $a \neq 1$

B $a > 1$

C $a > 0$ and $a \neq 1$

D No such a exists

E $a = 3$

F $a = \frac{1}{3}$

G $a > 0$, $a \neq 1$ and $a \neq \sqrt{3}$

Number

5. A positive integer n can be written as $n = 2^a 3^b$ where a and b are non-negative integers.
The number n has exactly 12 positive factors, counting 1 and n itself.
Given that $n < 200$, what is the greatest possible value of n?

A 72

B 96

C 108

D 144

E 162

F 486

Proof & Error Analysis

6. Let a and b be real numbers. A student attempts to prove the following claim:

Claim: If $a+b$ and ab are both integers, then a and b are both integers.

The student's attempt is as follows:

I a and b are the two solutions of $t^2-(a+b)t+ab=0$

II Write $s=a+b$ and $p=ab$, so s and p are integers and the two solutions are $t=\dfrac{s\pm\sqrt{s^2-4p}}{2}$

III Since a and b are real, $s^2-4p\geq 0$

IV Also s^2-4p is an integer, so $\sqrt{s^2-4p}$ is either an integer or an irrational number.

V If $\sqrt{s^2-4p}$ were irrational, then a would be irrational, and so $a+b$ would be irrational, contradicting the fact that $a+b$ is an integer.

VI Hence $\sqrt{s^2-4p}$ is an integer, so $s\pm\sqrt{s^2-4p}$ is an integer, and therefore a and b are both integers.

Which of the following best describes this attempt?

A It is completely correct.

B It is incorrect, and the first error is on line I.

C It is incorrect, and the first error is on line II.

D It is incorrect, and the first error is on line III.

E It is incorrect, and the first error is on line IV.

F It is incorrect, and the first error is on line V.


G It is incorrect, and the first error is on line VI.

Geometry

7. Triangle ABC is right-angled at A, with $AB = 6\,\text{cm}$ and $AC = 8\,\text{cm}$.

Semicircles are drawn with AB, AC and BC as diameters, as shown. The semicircle on BC is on the same side of the line BC as A. The portions of the other two semicircles that lie outside the semicircle on BC are shaded.

What is the total shaded area, in cm^2?



A 12

B 24

C 30


D 40

E 48

F 50

Coordinate Geometry

8. The line $y = 3x + 2$ meets the parabola $y = x^2$ at the points P and Q.
The tangents to the parabola at P and Q meet at R, as shown.
What are the coordinates of R?



A $\left(\frac{3}{2}, -4\right)$

B $\left(\frac{3}{2}, -2\right)$

C $\left(\frac{3}{2}, -1\right)$

D $(3, -2)$

E $(1, -3)$

F $\left(2, -\frac{3}{2}\right)$

Logic

9. Consider the statement $(*)$:

$$\textbf{If } 0 \leq f(x) \leq 1 \text{ for all real } x \text{ with } 0 \leq x \leq 1, \textbf{ then}$$
$$\int_0^1 f(x)\,\mathrm{d}x \leq \int_0^1 (f(x))^2\,\mathrm{d}x \qquad (*)$$

Which of the following functions is a **counterexample** to $(*)$?

A $f(x) = 2x$

B $f(x) = x^2 + \frac{1}{2}$

C $f(x) = 1$

D $f(x) = 1 + x$

E $f(x) = 0$

F $f(x) = 5x(1 - x)$

G $f(x) = \sqrt{x} + \frac{1}{2}$

H $f(x) = \frac{1}{2}$

Logic

10. Consider the following three conditions on a positive real number x

(1) $x^2 - 3x + 2 < 0$

(2) $\log_{10} x > 0$

(3) $2^x > 4$

Which of the following statements is/are true?

I: Condition (2) holds only if at least one of conditions (1) and (3) holds

II: If condition (2) does not hold, then condition (3) does not hold

III: For condition (1) to hold, it is necessary but not sufficient that condition (3) does not hold

A None of them

B I only

C II only

D III only

E I and II only

F I and III only

G II and III only

H I, II and III

Probability & Counting

11. A three-digit code is formed by selecting three different digits from $1, 2, 3, 4, 5, 6$ and placing them in random order.

Each possible code is equally likely.

It is revealed that when the three digits in the code are multiplied together, the result is divisible by 6.

What is the probability that the code contains the digit 6?

A $\frac{1}{3}$

B $\frac{1}{2}$

C $\frac{3}{5}$

D $\frac{2}{3}$

E $\frac{3}{4}$

F $\frac{5}{6}$

Proof & Error Analysis

12. Consider the claim about cubic polynomials:

Claim: If a cubic polynomial f has distinct roots a, b and c, where $a < b < c$, then there exists x with $a < x < b$ such that $f''(x) = 0$

The student attempts to prove the claim and produces the following argument:

I Since $f(a) = f(b) = 0,$ there is a number p such that $a < p < b$ and $f'(p) = 0$

II Similarly, there is a number q such that $b < q < c$ and $f'(q) = 0$

III The polynomial f' is quadratic with roots p and $q,$ so $f''\left(\frac{p+q}{2}\right) = 0$

IV Since $a < p < b < q < c,$ we have $p - b < 0 < q - b,$ and hence
$$\frac{p+q}{2} - b = \frac{(p-b)+(q-b)}{2} < 0$$

V Therefore the claim is true, the claimed x can be chosen as $x = \frac{p+q}{2}$

Which line contains the first error?

A Line **I**

B Line **II**

C Line **III**

D Line **IV**

E Line **V**

F There is no error, the proof is correct as written.

Integration

13. For each positive integer n, define

$$F(n) = \int_0^1 x^n \, \mathrm{d}x$$

and

$$H(n) = \sum_{r=1}^{n} \frac{1}{F(r)}$$

What is the smallest positive integer n such that $H(n) > 15$?

A 3

B 4

C 5

D 6

E 7

Logic

14. Let S be the statement:

$$\text{For all real numbers } x, \text{ if } |x| \geq 1 \text{ then } g(x) > 0$$

Which of the following is a correct **negation** of S?

A There exists a real number x with $|x| \geq 1$ such that $g(x) \leq 0$

B There exists a real number x with $|x| > 1$ such that $g(x) < 0$

C For all real numbers x, either $|x| < 1$ or $g(x) > 0$

D For all real numbers x with $|x| \geq 1$, $g(x) \geq 0$

E There exists a real number x such that $|x| < 1$ and $g(x) \leq 0$

F There exists a real number x such that $|x| < 1$ or $g(x) \leq 0$

Proof & Error Analysis

15. A function f is defined for all real x, and $f'(x)$ exists for every real x.

The function f is also **strictly increasing:** whenever $a < b$, also $f(a) < f(b)$.

A student makes the following claim:

Claim: It follows that $f'(x) > 0$ for every real x

The student argues as follows:

Line 1: Suppose, for a contradiction, that $f'(c) \leq 0$ for some real number c

Line 2: If $f'(c) < 0$, then just to the right of c the graph of f falls below $f(c)$, which contradicts f being strictly increasing. Hence $f'(c) = 0$

Line 3: So the tangent to the graph of f at $x = c$ is horizontal

Line 4: A horizontal tangent at $x = c$ means there are numbers $a < c < b$ with $f(a) \geq f(c)$ or with $f(b) \leq f(c)$

Line 5: Either possibility contradicts f being strictly increasing. Hence $f'(x) > 0$ for every real x

Which of the following best describes the argument?

A The argument is correct, and the claim is true

B The first error occurs in Line 2

C The first error occurs in Line 3

D The first error occurs in Line 4

E The first error occurs in Line 5

F The argument contains an error, but the claim is nevertheless true

Sequences & Series

16. Let n be an integer with $n \geq 6$.

In the expansion of $(1 + x)^n$ the coefficients of x^4, x^5 and x^6, in that order, are consecutive terms of an arithmetic sequence.

What is the sum of all possible values of n?

A 13

B 14

C 20

D 21

E 22

F 28

Logic

17. Let a be a real number. Consider the following statement:

$$\text{For all real numbers } x, \text{ there exists a real number } y \text{ such that}$$
$$a \leq y \leq a + \pi \text{ and } \sin y = \sin x$$

Which condition on a is **necessary and sufficient** for the statement to be true?

A $\sin a = 0$

B $\cos a = 0$

C $\sin 2a = 0$

D $\cos 2a = 0$

E $\sin a = \cos a$

F $\sin a = -\cos a$

Logic

18. Let p and q be real constants, and let

$$f(x) = x^3 + px^2 + qx$$

Consider the statement

S: There exists a real number x such that $f'(x) < 0$

Which of the following conditions is **necessary** but **not sufficient** for S?

A $q < 0$

B $q \leq 0$

C $3q < p^2$

D $3q \leq p^2$

E $p^2 > 0$

F $p^2 \geq 3|q|$

Algebra

19. A real number x satisfies

$$x^2 - 3x + 1 = 0$$

What is the value of $x^6 + \dfrac{1}{x^6}$?

A 288

B 306

C 320

D 322

E 324

Polynomials

20. The equation

$$x^3 - 3x + p = 0$$

where p is a real constant, has three distinct real solutions for x.

Find the complete set of values of p

A $-3 < p < 3$

B $-2 < p < 2$

C $-2 \leq p \leq 2$

D $p < -2$ or $p > 2$

E $0 < p < 4$

F $-4 < p < 0$