

Questions

Question 1	2
Question 2	3
Question 3	4
Question 4	5
Question 5	6
Question 6	7
Question 7	8
Question 8	9
Question 9	10
Question 10	11
Question 11	12

1. (i) In each of the following cases, find a 2×2 matrix that represents
- (a) a reflection in the line $y = x$ [1]
 - (b) a rotation of 60° clockwise about $(0, 0)$ [1]
 - (c) a rotation of 60° clockwise about $(0, 0)$ followed by a reflection in the line $y = x$ [2]

- (ii) The triangle T has vertices at the points $(1, 2)$, $(5, 5)$ and $(k, -1)$, where $k > -3$ is a constant.

Triangle T is transformed onto the triangle T' by the matrix

$$\begin{pmatrix} 2 & 5 \\ 1 & -1 \end{pmatrix}$$

Given that the area of triangle T' is 84 square units, find the value of k . [6]

2.

$$\mathbf{P} = \begin{pmatrix} p+1 & 2 \\ 2 & p+1 \end{pmatrix}$$

where p is a positive constant.

The matrix $\mathbf{P}$ represents a linear transformation U .

The parallelogram S has vertices at the points with coordinates $(1, 1)$, $(4, 2)$, $(6, 5)$ and $(3, 4)$.

The area of the image of S under the linear transformation U is 35.

(a) Determine the value of p . [4]

The transformation V consists of a reflection in the line $y = x$ followed by a stretch of scale factor 2 parallel to the x -axis with the y -axis invariant. The transformation V is represented by the matrix $\mathbf{Q}$.

(b) Write down the matrix $\mathbf{Q}$. [2]

Given that U followed by V is the transformation W , which is represented by the matrix $\mathbf{R}$

(c) find the matrix $\mathbf{R}$. [2]

3. (a) Specify fully the transformation $\mathbf{T}$ of the plane associated with the matrix $\mathbf{M}$, where

$$\mathbf{M} = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}$$

and $0 < \theta < 2\pi$. [2]

- (b) (i) Find $\det \mathbf{M}$. [1]

(ii) Deduce **two properties** of the transformation $\mathbf{T}$ from the value of $\det \mathbf{M}$. [2]

- (c) Prove that

$$\mathbf{M}^n = \begin{pmatrix} \cos n\theta & -\sin n\theta \\ \sin n\theta & \cos n\theta \end{pmatrix}$$

where n is a positive integer. [4]

- (d) Hence specify fully a **single** transformation which is equivalent to n applications of the transformation $\mathbf{T}$. [1]

4. Consider the matrix

$$A = \begin{pmatrix} -\frac{1}{2} & \frac{\sqrt{3}}{2} \\ \frac{\sqrt{3}}{2} & \frac{1}{2} \end{pmatrix}$$

- (i) Calculate A^2 . [1]

You are told that the matrix A represents a reflection in a line through the origin.

- (ii) Explain why your answer to part (i) is consistent with this information. [1]

- (iii) By finding the invariant points of the transformation, determine the equation of the mirror line. [3]

- (iv) Describe fully the transformation represented by

$$B = \begin{pmatrix} -\frac{1}{2} & -\frac{\sqrt{3}}{2} \\ \frac{\sqrt{3}}{2} & -\frac{1}{2} \end{pmatrix} \quad [2]$$

- (v) A composite transformation is formed by the transformation represented by A followed by the transformation represented by B . Find the single matrix that represents this composite transformation. [2]

- (vi) The composite transformation described in part (v) is equivalent to a single reflection. Find the equation of the mirror line of this reflection. [1]

5. The transformation U , represented by the 2×2 matrix P , is a reflection in the x -axis.

- (a) Write down the matrix P . [1]

The transformation V , represented by the 2×2 matrix Q , is a reflection in the line $y = x$.

- (b) Write down the matrix Q . [1]

Given that U followed by V is transformation T , which is represented by the matrix R

- (c) express R in terms of P and Q , [1]
(d) find the matrix R , [2]
(e) give a full geometrical description of T as a single transformation. [2]

6. The matrix $\mathbf{M}$ is given by

$$\mathbf{M} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & k \end{pmatrix}$$

where $k > 0$ and $k \neq 1$.

- (a) The matrix $\mathbf{M}$ represents a sequence of two geometrical transformations. State the type of each transformation, and make clear the order in which they are applied. [2]

The unit square in the x - y plane is transformed by $\mathbf{M}$ onto rectangle $OABC$.

- (b) Find, in terms of k , the area of rectangle $OABC$ and the matrix which transforms $OABC$ onto the unit square. [3]
- (c) Show that the line through the origin with gradient $\frac{1}{\sqrt{k}}$ is invariant under the transformation in the x - y plane represented by $\mathbf{M}$. [3]

7. You are given the matrix $\mathbf{A} = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}$.

(a) Find $\mathbf{A}^5$. [1]

(b) Describe the transformation that $\mathbf{A}$ represents. [2]

The matrix $\mathbf{B}$ represents a reflection in the plane $x = y$.

(c) Write down the matrix $\mathbf{B}$. [1]

The point Q has coordinates $(4, -1, 2)$. The point Q' is the image of Q under the transformation represented by $\mathbf{B}$.

(d) Find the coordinates of Q' . [1]

8. A right-angled triangle T has vertices $A(-2, 1)$, $B(1, 1)$ and $C(1, 5)$. When T is transformed by the matrix

$$P = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$$

the image is T' .

- (a) Find the coordinates of the vertices of T' . [2]

- (b) Describe fully the transformation represented by P . [2]

The matrices

$$Q = \begin{pmatrix} 2 & -1 \\ 0 & 3 \end{pmatrix}$$

and

$$R = \begin{pmatrix} 1 & 2 \\ 1 & 0 \end{pmatrix}$$

represent two transformations. When T is transformed by the matrix QR , the image is T'' .

- (c) Find QR . [2]

- (d) Find the determinant of QR . [2]

- (e) Using your answer to part (d), find the area of T'' . [3]

9. Given that the matrix $\mathbf{M}$, where

$$\mathbf{M} = \begin{pmatrix} -\frac{\sqrt{2}}{2} & 0 & -\frac{\sqrt{2}}{2} \\ 0 & 1 & 0 \\ \frac{\sqrt{2}}{2} & 0 & -\frac{\sqrt{2}}{2} \end{pmatrix}$$

represents a rotation through α° about the y -axis in the positive sense given by the right-hand rule.

(a) determine the value of α . [1]

(b) Hence determine the smallest possible positive integer value of k for which $\mathbf{M}^k = \mathbf{I}$. [2]

The 3×3 matrix $\mathbf{N}$ represents a reflection in the plane with equation $x = 0$.

(c) Write down the matrix $\mathbf{N}$. [1]

The point A has coordinates $(1, 2, 3)$.

The point B is the image of the point A under the transformation represented by matrix $\mathbf{M}$ followed by the transformation represented by matrix $\mathbf{N}$.

(d) Show that the coordinates of B are $(2\sqrt{2}, 2, -\sqrt{2})$. [2]

Given that O is the origin

(e) show that, to 3 significant figures, the size of angle AOB is 79.4° . [2]

(f) Hence determine the area of triangle AOB , giving your answer to 3 significant figures. [2]

10. Bilal has been asked to describe the transformation given by the matrix

$$\begin{pmatrix} \frac{\sqrt{3}}{2} & 0 & -\frac{1}{2} \\ 0 & 1 & 0 \\ \frac{1}{2} & 0 & \frac{\sqrt{3}}{2} \end{pmatrix}$$

He writes his answer as follows:

The transformation is a rotation about the y -axis through an angle of θ , where

$$\cos \theta = \frac{\sqrt{3}}{2} \quad \text{and} \quad \sin \theta = \frac{1}{2}$$
$$\theta = 30^\circ$$

Identify and correct the error in Bilal's work.

[2]

11. Three non-singular 3×3 matrices, $\mathbf{A}$, $\mathbf{B}$ and $\mathbf{R}$, are such that

$$\mathbf{R}\mathbf{A} = \mathbf{B}$$

The matrix $\mathbf{R}$ represents a rotation about the x -axis through an angle θ and

$$\mathbf{B} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & -\sin \theta & \cos \theta \\ 0 & \cos \theta & \sin \theta \end{bmatrix}$$

(a) Show that $\mathbf{A}$ is independent of the value of θ . [3]

(b) Give a full description of the single transformation represented by the matrix $\mathbf{A}$. [1]